

AD-A108 235

UTAH STATE UNIV LOGAN ELECTRO-DYNAMICS LAB

F/6 13/2

DOWN-LOOKING INTERFEROMETER STUDY II. VOLUME II. HANDBOOK OF RE--ETC(U)

MAR 80 A S ZACHOR

F19628-77-C-0203

UNCLASSIFIED


SCIENTIFIC-3

AFGL-TR-80-0237

NL

1 OF 1
AD A
106 2 55

END
DATE
FILMED
01-82
DTIC



MICROCOPY RESOLUTION TEST CHART

NBS 1963-A

4

LEVEL III

AFGL-TR-80-0237

Vol 1-A108 299

(12)

AD A108235

DOWN-LOOKING INTERFEROMETER STUDY II
VOLUME II: HANDBOOK OF RESULTS

ALEXANDER S. ZACHOR

ELECTRO-DYNAMICS LABORATORIES
UTAH STATE UNIVERSITY
LOGAN, UTAH 84322

MARCH 1980

SCIENTIFIC REPORT NO. 3

DTIC
ELECTE
DEC 8 1981
H

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.

DTIC FILE COPY

AIR FORCE GEOPHYSICS LABORATORY
AIR FORCE SYSTEMS COMMAND
UNITED STATES AIR FORCE
HANSCOM AFB, MASSACHUSETTS 01731

81 12 08 296
123 870


Qualified requestors may obtain additional copies from the Defense Technical Information Center. All others should apply to the National Technical Information Service.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFGL-TR-80-0237	2. GOVT ACCESSION NO. AD-A108 235	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) DOWN-LOOKING INTERFEROMETER STUDY II VOLUME II: HANDBOOK OF RESULTS		5. TYPE OF REPORT & PERIOD COVERED Scientific Report No. 3
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Alexander S. Zachor		8. CONTRACT OR GRANT NUMBER(s) F19628-77-C-0203
9. PERFORMING ORGANIZATION NAME AND ADDRESS Atmospheric Radiation Consultants, Inc. 59 High Street Acton, Massachusetts 01720		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 62101F 767010AB
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Geophysics Laboratory (OPR) Hanscom AFB, Massachusetts 01731 Monitor/Dean F. Kimball/OPR		12. REPORT DATE March 1980
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Electro-Dynamics Laboratories Utah State University Logan, Utah 84322		13. NUMBER OF PAGES 76
		15. SECURITY CLASS. (of this report) Unclassified
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Atmospheric transmission, interferometer spectrometers, spectral correlation techniques, minor contaminant species detection.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This volume is a handbook of results for the Down-Looking Interferometer Study II (AFGL-TR-80-0236). Three pages of data are presented in a standard format for each candidate detection band.		

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)



SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

PREFACE

This handbook of results for the Down-Looking Interferometer Study II is described and summarized in Section I of Volume I. Three pages of data are presented for each candidate detection band in a standard format. The following index gives the number of the first page for each band.

<u>Molecule (Transition)</u>	<u>Detection Band (cm⁻¹)</u>	<u>Page</u>
C ₂ H ₄ O ₂	880-1000	3
CH ₄ (v ₃)	3000-3025	6
(v ₄)	1295-1310	9
CH ₃ I	790-990	12
CO(1-0)	2130-2185	15
CO ₂ (blue spike)	2375-2400	18
DF	2500-2700	21
	2700-2900	24
	2900-3000	27
HBr	2450-2650	30
HCl	2500-2700	33
	2700-2900	36
	2900-3000	39
HF	3240-3440	42
HI	2100-2200	45
HNO ₃ (2 v ₉)	887.5-902.5	48

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability	
Notes	
A	

<u>Molecule (Transition)</u>	<u>Detection Band (cm⁻¹)</u>	<u>Page</u>
NH ₃ (v ₂)	915-970	51
N ₂ O (2 v ₁)	2525-2590	54
(v ₃)	2160-2210	57
(v ₁)	1250-1320	60
NO ₂ (v ₁ + v ₃)	2850-2935	63
(v ₂)	710-795	66
SO ₂ (v ₁ + v ₃)	2450-2525	69
(v ₁)	1090-1210	72

$C_2H_4O_2$ 880-1000 cm^{-1} region






Table 1 Detection Parameters for $C_2H_4O_2$ 880-1000 cm^{-1} region

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	880-1000	cm^{-1}
Approximate wavelength	10.6	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.80$)	M = 150	
BAND PHOTON RADIANCE (scene)	6.00×10^{16}	ph/sec cm^2 sr
Maximum of contrast $\tau_{\nu} \alpha_{gv}$	C _{MAX} = 8.38×10^{-20}	$cm^2/molec$
Mean contrast	$\mu' = 5.50 \times 10^{-20}$	$cm^2/molec$
STANDARD DEVIATION OF CONTRAST	$\sigma' = 1.69 \times 10^{-20}$	$cm^2/molec$
* Photon flux density on detector		
* from scene	1.4×10^{15}	phot/sec cm^2
* from internal sources	1.5×10^{15}	phot/sec cm^2
* TOTAL	J = 2.9×10^{15}	phot/sec cm^2
* BLIP $D_{\lambda_c}^*$	4.2×10^{11}	cm $\sqrt{Hz/W}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	min $t_d = 6.54 \times 10^{-4}$	sec
* CORRESPONDING BASELINE NESR	(NESR) ₀ = 2.5×10^{-7}	W/cm^2 sr cm^{-1}
* MINIMUM DETECTABLE QUANTITY D (see figure)	min D = $2.0 - 7.0 \times 10^{12}$	(molec/ cm^2) (W/ cm^2 sr cm^{-1})
* UNCERTAINTY IN D	$\sigma_D = 5.1 \times 10^{11}$	(molec/ cm^2) (W/ cm^2 sr cm^{-1})



CH₄ v₃ band



CH₄ v₃ band







Table 2 Detection Parameters for CH₄ ν_3 band

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	3000-3025	cm ⁻¹
Approximate wavelength	3.3	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 250	
BAND PHOTON RADIANCE (scene)	2.57×10^{12}	ph/sec cm ² sr
Maximum of contrast $\tau_{\nu}^{\alpha}_{gv}$	C _{MAX} = 1.14×10^{-20}	cm ² /molec
Mean contrast	$\mu' = 1.84 \times 10^{-21}$	cm ² /molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 2.71 \times 10^{-21}$	cm ² /molec
* Photon flux density on detector		
* from scene	5.9×10^{10}	phot/sec cm ²
* from internal sources	1.5×10^{11}	phot/sec cm ²
* TOTAL	J = 2.1×10^{11}	phot/sec cm ²
* BLIP $D_{\lambda_c}^*$	1.3×10^{13}	cm $\sqrt{\text{Hz}}/\text{W}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	min $t_d = 7.79$	sec
* CORRESPONDING BASELINE NESR	(NESR) ₀ = 6.0×10^{-10}	W/cm ² sr cm ⁻¹
* MINIMUM DETECTABLE QUANTITY D (see figure)	min D = $2.5 - 8.0 \times 10^{10}$	(molec/cm ²) (W/cm ² sr cm ⁻¹)
* UNCERTAINTY IN D	$\sigma_{D'} = 1.6 \times 10^{10}$	(molec/cm ²) (W/cm ² sr cm ⁻¹)







PRECEDING PAGE BLANK-NOT FILLED

CH₃I 790-990 cm⁻¹ region





Table 4 Detection Parameters for CH_3I 790-990 cm^{-1} region

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	790-990	cm^{-1}
Approximate wavelength	11.2	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 2000	
BAND PHOTON RADIANCE (scene)	1.14×10^{17}	$\text{ph/sec cm}^2 \text{ sr}$
Maximum of contrast $\tau_v \alpha_{gv}$	CNAX = 3.62×10^{-20}	cm^2/molec
Mean contrast.	$\mu' = 4.82 \times 10^{-21}$	cm^2/molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 4.40 \times 10^{-21}$	cm^2/molec
* Photon flux density on detector		
* from scene.	2.6×10^{15}	phot/sec cm^2
* from internal sources	2.7×10^{15}	phot/sec cm^2
* TOTAL	$J = 5.3 \times 10^{15}$	phot/sec cm^2
* BLIP $D_{\lambda_c}^*$	2.9×10^{11}	$\text{cm } \sqrt{\text{Hz/W}}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	$\min t_d = 3.63 \times 10^{-3}$	sec
* CORRESPONDING BASELINE NESR.	$(\text{NESR})_0 = 1.23 \times 10^{-6}$	$\text{W/cm}^2 \text{ sr cm}^{-1}$
* MINIMUM DETECTABLE QUANTITY D (see figure)	$\min D = 1.0 - 3.5 \times 10^{13}$	$(\text{molec/cm}^2)(\text{W/cm}^2 \text{ sr cm}^{-1})$
* UNCERTAINTY IN D	$\sigma_D = 5.96 \times 10^{12}$	$(\text{molec/cm}^2)(\text{W/cm}^2 \text{ sr cm}^{-1})$



CO 1-0 band





Table 5 Detection Parameters for CO 1-0 band

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	2130-2185	cm^{-1}
Approximate wavelength	4.7	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 550	
BAND PHOTON RADIANCE (scene)	3.66×10^{14}	$\text{ph/sec cm}^2 \text{sr}$
Maximum of contrast $\tau_v \alpha_{gv}$	C _{MAX} = 2.01×10^{-19}	cm^2/molec
Mean contrast.	$\mu' = 2.09 \times 10^{-20}$	cm^2/molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 3.64 \times 10^{-20}$	cm^2/molec
* Photon flux density on detector		
* from scene.	8.3×10^{12}	phot/sec cm^2
* from internal sources	1.0×10^{13}	phot/sec cm^2
* TOTAL	J = 1.9×10^{13}	phot/sec cm^2
* BLIP $D_{\lambda_c}^*$	1.9×10^{12}	$\text{cm } \sqrt{\text{Hz}}/\text{W}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	$\min t_d = 0.833$	sec
* CORRESPONDING BASELINE NESR.	$(\text{NESR})_0 = 1.2 \times 10^{-8}$	$\text{W/cm}^2 \text{sr cm}^{-1}$
* MINIMUM DETECTABLE QUANTITY D (see figure)	$\min D = 3.0 - 8.5 \times 10^{10}$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{sr cm}^{-1})$
* UNCERTAINTY IN D	$\sigma_D = 1.8 \times 10^{10}$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{sr cm}^{-1})$



CO₂ 2375-2400 cm⁻¹ region





Table 6 Detection Parameters for CO₂ 2375-2400 cm⁻¹ region

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	2375-2400	cm ⁻¹
Approximate wavelength	4.19	μm
* NO. OF SPECTRAL ELEMENTS (for Δν = 0.10)	M = 250	
BAND PHOTON RADIANCE (scene)	2.75 × 10 ¹³	ph/sec cm ² sr
Maximum of contrast τ _v α _{gv}	C _{MAX} = ** 2.71 × 10 ⁻²⁰	cm ² /molec
Mean contrast.	μ' = ** 9.72 × 10 ⁻²²	cm ² /molec
STANDARD DEVIATION OF CONTRAST	σ' = ** 4.80 × 10 ⁻²¹	cm ² /molec
* Photon flux density on detector		
* from scene.	6.3 × 10 ¹¹	phot/sec cm ²
* from internal sources	1.9 × 10 ¹²	phot/sec cm ²
* TOTAL	J = 2.5 × 10 ¹²	phot/sec cm ²
* BLIP D _{λ_c} [*]	4.7 × 10 ¹²	cm √Hz/W
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	min t _d = 2.24	sec
* CORRESPONDING BASELINE NESR.	(NESR) ₀ = 3.1 × 10 ⁻⁹	W/cm ² sr cm ⁻¹
* MINIMUM DETECTABLE QUANTITY D (see figure)	min D = ** 1.0 - 2.5 × 10 ¹¹	(molec/cm ²) (W/cm ² sr cm ⁻¹)
* UNCERTAINTY IN D	σ _D = ** 5.7 × 10 ¹⁰	(molec/cm ²) (W/cm ² sr cm ⁻¹)

** Assumes CO₂ temperature = 850°K.



DF 2500-2700 cm^{-1} region





Table 7 Detection Parameters for DF 2500-2700 cm^{-1} region

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	2500-2700	cm^{-1}
Approximate wavelength	3.85	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 2000	
BAND PHOTON RADIANCE (scene)	2.81×10^{14}	$\text{ph/sec cm}^2 \text{sr}$
Maximum of contrast $\tau_{\nu} a_{\nu}$	C _{MAX} = 2.62×10^{-19}	cm^2/molec
Mean contrast	$\mu' = 2.86 \times 10^{-22}$	cm^2/molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 6.38 \times 10^{-21}$	cm^2/molec
* Photon flux density on detector		
* from scene	6.4×10^{12}	phot/sec cm^2
* from internal sources	6.8×10^{12}	phot/sec cm^2
* TOTAL	$J = 1.3 \times 10^{13}$	phot/sec cm^2
* BLIP $D_{\lambda_c}^*$	1.9×10^{12}	$\text{cm} \sqrt{\text{Hz/W}}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	$\min t_d = 1.02$	sec
* CORRESPONDING BASELINE NESR	$(\text{NESR})_0 = 1.1 \times 10^{-8}$	$\text{W/cm}^2 \text{sr cm}^{-1}$
* MINIMUM DETECTABLE QUANTITY D (see figure)	$\min D = 0.8 - 2.3 \times 10^{11}$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{sr cm}^{-1})$
* UNCERTAINTY IN D	$\sigma_D = 5.6 \times 10^{10}$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{sr cm}^{-1})$



OF 2700-2900 cm^{-1} region





Table 8 Detection Parameters for DF 2700-2900 cm^{-1} region

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	2700-2900	cm^{-1}
Approximate wavelength	3.57	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 2000	
BAND PHOTON RADIANCE (scene)	1.23×10^{14}	$\text{ph/sec cm}^2 \text{ sr}$
Maximum of contrast $\tau_v \alpha_{gv}$	C MAX = 1.66×10^{-18}	cm^2/molec
Mean contrast	$\mu' = 1.45 \times 10^{-20}$	cm^2/molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 9.39 \times 10^{-20}$	cm^2/molec
* Photon flux density on detector		
* from scene	2.8×10^{12}	phot/sec cm^2
* from internal sources	3.0×10^{12}	phot/sec cm^2
* TOTAL	J = 5.8×10^{12}	phot/sec cm^2
* BLIP $D_{\lambda_c}^*$	2.7×10^{12}	$\text{cm } \sqrt{\text{Hz/W}}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	min $t_d = 1.53$	sec
* CORRESPONDING BASELINE NESR.	$(\text{NESR})_0 = 6.6 \times 10^{-9}$	$\text{W/cm}^2 \text{ sr cm}^{-1}$
* MINIMUM DETECTABLE QUANTITY D (see figure)	min D = $3.0 - 9.0 \times 10^9$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{ sr cm}^{-1})$
* UNCERTAINTY IN D	$\sigma_D = 2.2 \times 10^9$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{ sr cm}^{-1})$



OF 2900-3000 cm^{-1} region





Table 9 Detection Parameters for DF 2900-3000 cm^{-1} region

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	2900-3000	cm^{-1}
Approximate wavelength	3.39	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 1000	
BAND PHOTON RADIANCE (scene)	2.56×10^{13}	$\text{ph/sec cm}^2 \text{sr}$
Maximum of contrast $\tau_{\nu} \alpha_{\text{gv}}$	C MAX = 2.39×10^{-19}	cm^2/molec
Mean contrast	$\mu' = 4.34 \times 10^{-21}$	cm^2/molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 1.80 \times 10^{-20}$	cm^2/molec
* Photon flux density on detector		
* from scene	5.8×10^{11}	phot/sec cm^2
* from internal sources	8.0×10^{11}	phot/sec cm^2
* TOTAL	$J = 1.4 \times 10^{12}$	phot/sec cm^2
* BLIP $D_{\lambda}^* \lambda_c$	5.2×10^{12}	$\text{cm } \sqrt{\text{Hz}}/\text{W}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	$\min t_d = 3.09$	sec
* CORRESPONDING BASELINE NESR	$(\text{NESR})_0 = 2.4 \times 10^{-9}$	$\text{W/cm}^2 \text{sr cm}^{-1}$
* MINIMUM DETECTABLE QUANTITY D (see figure)	$\min D = 1.0 - 2.3 \times 10^{10}$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{sr cm}^{-1})$
* UNCERTAINTY IN D	$\sigma_D = 5.8 \times 10^9$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{sr cm}^{-1})$



HBr 2450-2650 cm^{-1} region





Table 10 Detection Parameters for HBr 2450-2650 cm^{-1} region

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	2450-2650	cm^{-1}
Approximate wavelength	3.9	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 2000	
BAND PHOTON RADIANCE (scene)	3.39×10^{14}	$\text{ph/sec cm}^2 \text{sr}$
Maximum of contrast $\tau_{\nu} \sigma_{\text{gv}}$	C _{MAX} = 2.48×10^{-19}	cm^2/molec
Mean contrast.	$\mu' = 4.19 \times 10^{-21}$	cm^2/molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 2.10 \times 10^{-20}$	cm^2/molec
* Photon flux density on detector		
* from scene.	7.7×10^{12}	phot/sec cm^2
* from internal sources	8.2×10^{12}	phot/sec cm^2
* TOTAL	J = 1.6×10^{13}	phot/sec cm^2
* BLIP $D_{\lambda_c}^*$	1.8×10^{12}	$\text{cm } \sqrt{\text{Hz}}/\text{W}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE		
	min $t_d = 0.930$	sec
* CORRESPONDING BASELINE NESR.	$(\text{NESR})_0 = 1.3 \times 10^{-8}$	$\text{W/cm}^2 \text{sr cm}^{-1}$
* MINIMUM DETECTABLE QUANTITY D (see figure)	min D = $3.0 - 8.0 \times 10^{10}$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{sr cm}^{-1})$
* UNCERTAINTY IN D	$\sigma_D = 1.9 \times 10^{10}$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{sr cm}^{-1})$



HCl 2500-2700 cm^{-1} region





Table 11 Detection Parameters for HCl 2500-2700 cm^{-1} region

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	2500-2700	cm^{-1}
Approximate wavelength	3.85	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 2000	
BAND PHOTON RADIANCE (scene)	2.81×10^{14}	$\text{ph/sec cm}^2 \text{sr}$
Maximum of contrast $\tau_{\nu}^{\alpha_{\text{gv}}}$	$\text{CMAX} = 7.11 \times 10^{-20}$	cm^2/molec
Mean contrast.	$\mu' = 1.16 \times 10^{-22}$	cm^2/molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 1.57 \times 10^{-21}$	cm^2/molec
* Photon flux density on detector		
* from scene.	6.4×10^{12}	phot/sec cm^2
* from internal sources	6.8×10^{12}	phot/sec cm^2
* TOTAL	$J = 1.3 \times 10^{13}$	phot/sec cm^2
* BLIP $D_{\lambda_c}^*$	1.9×10^{12}	$\text{cm } \sqrt{\text{Hz/W}}$
* MINIMUM SCAN TIME FOR RLIP PERFORMANCE	$\min t_d = 1.02$	sec
* CORRESPONDING BASELINE NESR.	$(\text{NESR})_0 = 1.1 \times 10^{-8}$	$\text{W/cm}^2 \text{sr cm}^{-1}$
* MINIMUM DETECTABLE QUANTITY D (see figure)	$\min D = 3.0 - 9.0 \times 10^{11}$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{sr cm}^{-1})$
* UNCERTAINTY IN D	$\sigma_D = 2.3 \times 10^{11}$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{sr cm}^{-1})$



HCl 2700-2900 cm^{-1} region





Table 12 Detection Parameters for HCl 2700-2900 cm^{-1} region

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	2700-2900	cm^{-1}
Approximate wavelength	3.57	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 2000	
BAND PHOTON RADIANCE (scene)	1.23×10^{14}	$\text{ph/sec cm}^2 \text{ sr}$
Maximum of contrast $\tau_v \alpha_{gv}$	C MAX = 1.30×10^{-18}	cm^2/molec
Mean contrast.	$\mu' = 9.28 \times 10^{-21}$	cm^2/molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 6.77 \times 10^{-20}$	cm^2/molec
* Photon flux density on detector		
* from scene.	2.8×10^{12}	phot/sec cm^2
* from internal sources	3.0×10^{12}	phot/sec cm^2
* TOTAL	J = 5.8×10^{12}	phot/sec cm^2
* BLIP $D_{\lambda_c}^*$	2.7×10^{12}	$\text{cm } \sqrt{\text{Hz}}/\text{W}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	min $t_d = 1.53$	sec
* CORRESPONDING BASELINE NESR.	$(\text{NESR})_0 = 6.6 \times 10^{-9}$	$\text{W/cm}^2 \text{ sr cm}^{-1}$
* MINIMUM DETECTABLE QUANTITY D (see figure)	min D = $0.5 - 1.25 \times 10^{10}$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{ sr cm}^{-1})$
* UNCERTAINTY IN D	$\sigma_D = 3.1 \times 10^9$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{ sr cm}^{-1})$



HCl 2900-3000 cm^{-1} region





Table 13 Detection Parameters for HCl 2900-3000 cm^{-1} region

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	2900-3000	cm^{-1}
Approximate wavelength	3.39	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 1000	
BAND PHOTON RADIANCE (scene)	2.56×10^{13}	$\text{ph/sec cm}^2 \text{sr}$
Maximum of contrast $\tau_{\nu}^{\alpha_{\text{gv}}}$	C _{MAX} = 1.64×10^{-18}	cm^2/molec
Mean contrast.	$\mu' = 1.64 \times 10^{-20}$	cm^2/molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 9.81 \times 10^{-20}$	cm^2/molec
* Photon flux density on detector		
* from scene.	5.8×10^{11}	phot/sec cm^2
* from internal sources	8.0×10^{11}	phot/sec cm^2
* TOTAL	$J = 1.4 \times 10^{12}$	phot/sec cm^2
* BLIP $D_{\lambda_c}^*$	5.2×10^{12}	$\text{cm} \sqrt{\text{Hz/W}}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	$\min t_D = 3.09$	sec
* CORRESPONDING BASELINE NESR.	$(\text{NESR})_0 = 2.4 \times 10^{-9}$	$\text{W/cm}^2 \text{sr cm}^{-1}$
* MINIMUM DETECTABLE QUANTITY D (see figure)	$\min D = 2.0 - 4.5 \times 10^9$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{sr cm}^{-1})$
* UNCERTAINTY IN D	$\sigma_D = 1.1 \times 10^9$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{sr cm}^{-1})$



HF 3240-3440 cm^{-1} region





Table 14 Detection Parameters for HF 3240-3440 cm^{-1} region

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	3240-3440	cm^{-1}
Approximate wavelength	3.0	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 2000	
BAND PHOTON RADIANCE (scene)	5.46×10^{12}	$\text{ph/sec cm}^2 \text{sr}$
Maximum of contrast $\tau_{\nu} \alpha_{\text{gv}}$	CMAX = 9.62×10^{-23}	cm^2/molec
Mean contrast	$\mu' = 5.70 \times 10^{-26}$	cm^2/molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 1.72 \times 10^{-24}$	cm^2/molec
* Photon flux density on detector		
* from scene	1.2×10^{11}	phot/sec cm^2
* from internal sources	3.2×10^{11}	phot/sec cm^2
* TOTAL	$J = 4.5 \times 10^{11}$	phot/sec cm^2
* BLIP $D_{\lambda_c}^*$	8.0×10^{12}	$\text{cm } \sqrt{\text{Hz/W}}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	$\min t_d = 5.52$	sec
* CORRESPONDING BASELINE NESR	$(\text{NESR})_0 = 1.2 \times 10^{-9}$	$\text{W/cm}^2 \text{sr cm}^{-1}$
* MINIMUM DETECTABLE QUANTITY D (see figure)	$\min D = 3.0 - 9.0 \times 10^{13}$	$(\text{molec/cm}^2)(\text{W/cm}^2 \text{sr cm}^{-1})$
* UNCERTAINTY IN D	$\sigma_D = 2.1 \times 10^{13}$	$(\text{molec/cm}^2)(\text{W/cm}^2 \text{sr cm}^{-1})$



HI 2100-2200 cm^{-1} region





Table 15 Detection Parameters for HI 2100-2200 cm^{-1} region

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	2100-2200	cm^{-1}
Approximate wavelength	4.65	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 1000	
BAND PHOTON RADIANCE (scene)	6.19×10^{14}	$\text{ph/sec cm}^2 \text{ sr}$
Maximum of contrast $\tau_{\nu}^{\alpha}_{\text{gv}}$	C MAX = 2.13×10^{-21}	cm^2/molec
Mean contrast.	$\mu' = 1.17 \times 10^{-23}$	cm^2/molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 9.51 \times 10^{-23}$	cm^2/molec
* Photon flux density on detector		
* from scene.	1.4×10^{13}	phot/sec cm^2
* from internal sources	2.0×10^{13}	phot/sec cm^2
* TOTAL	J = 3.4×10^{13}	phot/sec cm^2
* BLIP $D_{\lambda_c}^*$	1.4×10^{12}	$\text{cm } \sqrt{\text{Hz/W}}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	$\min t_d = 0.629$	sec
* CORRESPONDING BASELINE NESR.	$(\text{NESR})_0 = 1.9 \times 10^{-8}$	$\text{W/cm}^2 \text{ sr cm}^{-1}$
* MINIMUM DETECTABLE QUANTITY D (see figure)	$\min D = 1.5 - 3.5 \times 10^{13}$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{ sr cm}^{-1})$
* UNCERTAINTY IN D	$\sigma_D = 8.9 \times 10^{12}$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{ sr cm}^{-1})$



HNO_3 $2\nu_9$ band





Table 16 Detection Parameters for HNO_3 $2\nu_g$ band

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	887.5-902.5	cm^{-1}
Approximate wavelength	11.2	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 150	
BAND PHOTON RADIANCE (scene)	8.61×10^{15}	$\text{ph/sec cm}^2 \text{ sr}$
Maximum of contrast $\tau_{\nu g\nu}$	C _{MAX} = 1.66×10^{-19}	cm^2/molec
Mean contrast	$\mu' = 5.38 \times 10^{-20}$	cm^2/molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 6.98 \times 10^{-20}$	cm^2/molec
* Photon flux density on detector		
* from scene	2.0×10^{14}	phot/sec cm^2
* from internal sources	2.1×10^{14}	phot/sec cm^2
* TOTAL	J = 4.1×10^{14}	phot/sec cm^2
* BLIP $D_{\lambda_c}^*$	1.2×10^{12}	$\text{cm } \sqrt{R_z}/W$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	$\min t_d = 1.82 \times 10^{-2}$	sec
* CORRESPONDING BASELINE NESR.	$(\text{NESR})_0 = 1.4 \times 10^{-7}$	$W/\text{cm}^2 \text{ sr cm}^{-1}$
* MINIMUM DETECTABLE QUANTITY D (see figure)	$\min D = 3.0 - 8.0 \times 10^{11}$	$(\text{molec/cm}^2) (W/\text{cm}^2 \text{ sr cm}^{-1})$
* UNCERTAINTY IN D	$\sigma_{D'} = 1.8 \times 10^{11}$	$(\text{molec/cm}^2) (W/\text{cm}^2 \text{ sr cm}^{-1})$







Table 17 Detection Parameters for NH_3 ν_2 band

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	915-970	cm^{-1}
Approximate wavelength	10.7	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 550	
BAND PHOTON RADIANCE (scene)	2.75×10^{16}	$\text{ph/sec cm}^2 \text{ sr}$
Maximum of contrast $\tau_{\nu}^{\alpha}_{\text{gv}}$	C _{MAX} = 9.90×10^{-19}	cm^2/molec
Mean contrast.	$\mu' = 6.78 \times 10^{-20}$	cm^2/molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 1.30 \times 10^{-19}$	cm^2/molec
* Photon flux density on detector		
* from scene.	6.3×10^{14}	phot/sec cm^2
* from internal sources	6.9×10^{14}	phot/sec cm^2
* TOTAL	$J = 1.3 \times 10^{15}$	phot/sec cm^2
* BLIP $D_{\lambda_c}^*$	6.2×10^{11}	$\text{cm } \sqrt{\text{Hz}}/\text{W}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	$\min t_d = 7.46 \times 10^{-3}$	sec
* CORRESPONDING BASELINE NESR.	$(\text{NESR})_0 = 4.1 \times 10^{-7}$	$\text{W/cm}^2 \text{ sr cm}^{-1}$
* MINIMUM DETECTABLE QUANTITY D (see figure)	$\min D = 4.5 - 7.3 \times 10^{11}$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{ sr cm}^{-1})$
* UNCERTAINTY IN D	$\sigma_D = 1.69 \times 10^{11}$	$(\text{molec/cm}^2) (\text{W/cm}^2 \text{ sr cm}^{-1})$







Table 18 Detection Parameters for N_2O $2\nu_1$ band

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	2525-2590	cm^{-1}
Approximate wavelength	4.0	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 650	
BAND PHOTON RADIANCE (scene)	1.04×10^{14}	ph/sec cm^2 sr
Maximum of contrast $\tau_{\nu} a_{gv}$	C _{MAX} = 3.51×10^{-20}	$cm^2/molec$
Mean contrast.	$\mu' = 1.32 \times 10^{-20}$	$cm^2/molec$
STANDARD DEVIATION OF CONTRAST	$\sigma' = 8.95 \times 10^{-21}$	$cm^2/molec$
* Photon flux density on detector		
* from scene.	2.4×10^{12}	phot/sec cm^2
* from internal sources	2.5×10^{12}	phot/sec cm^2
* TOTAL	J = 4.9×10^{12}	phot/sec cm^2
* BLIP $D_{\lambda}^* C$	3.1×10^{12}	$cm \sqrt{Hz/W}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	min $t_d = 1.63$	sec
* CORRESPONDING BASELINE NESR.	$(NESR)_0 = 5.4 \times 10^{-9}$	$W/cm^2 sr cm^{-1}$
* MINIMUM DETECTABLE QUANTITY D (see figure)	min D = $0.5 - 1.4 \times 10^{11}$	$(molec/cm^2)(W/cm^2 sr cm^{-1})$
* UNCERTAINTY IN D	$\sigma_D = 1.9 \times 10^{10}$	$(molec/cm^2)(W/cm^2 sr cm^{-1})$



N₂O v₃ band





Table 19 Detection Parameters for N_2O ν_3 band

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	2160-2210	cm^{-1}
Approximate wavelength	4.5	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	$M = 500$	
BAND PHOTON RADIANCE (scene)	2.23×10^{14}	$ph/sec\ cm^2\ sr$
Maximum of contrast $\tau_{\nu} a_{gv}$	$C_{MAX} = 8.93 \times 10^{-20}$	$cm^2/molec$
Mean contrast.	$\mu' = 3.73 \times 10^{-20}$	$cm^2/molec$
STANDARD DEVIATION OF CONTRAST	$\sigma' = 2.59 \times 10^{-20}$	$cm^2/molec$
* Photon flux density on detector		
* from scene.	5.1×10^{12}	$phot/sec\ cm^2$
* from internal sources	8.5×10^{12}	$phot/sec\ cm^2$
* TOTAL	$J = 1.4 \times 10^{13}$	$phot/sec\ cm^2$
* BLIP $D_{\lambda_c}^*$	2.2×10^{12}	$cm\ \sqrt{Hz}/W$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	$\min t_d = 0.978$	sec
* CORRESPONDING BASELINE NESR.	$(NESR)_0 = 9.9 \times 10^{-9}$	$W/cm^2\ sr\ cm^{-1}$
* MINIMUM DETECTABLE QUANTITY D (see figure)	$\min D = 0.3 - 1.0 \times 10^{11}$	$(molec/cm^2)(W/cm^2\ sr\ cm^{-1})$
* UNCERTAINTY IN D	$\sigma_D = 1.4 \times 10^{10}$	$(molec/cm^2)(W/cm^2\ sr\ cm^{-1})$







Table 20 Detection Parameters for N_2O ν_1 band

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	1250-1320	cm^{-1}
Approximate wavelength	7.8	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 700	
BAND PHOTON RADIANCE (scene)	6.62×10^{15}	ph/sec cm^2 sr
Maximum of contrast $\tau_v \alpha_{gv}$	C _{MAX} = 2.84×10^{-20}	$cm^2/molec$
Mean contrast	$\mu' = 3.40 \times 10^{-21}$	$cm^2/molec$
STANDARD DEVIATION OF CONTRAST	$\sigma' = 5.50 \times 10^{-21}$	$cm^2/molec$
* Photon flux density on detector		
* from scene	1.5×10^{14}	phot/sec cm^2
* from internal sources	3.1×10^{14}	phot/sec cm^2
* TOTAL	$J = 4.6 \times 10^{14}$	phot/sec cm^2
* BLIP $D_{\lambda_c}^*$	7.7×10^{11}	$cm \sqrt{Hz}/W$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	$\min t_d = 1.27 \times 10^{-2}$	sec
* CORRESPONDING BASELINE NESR.	$(NESR)_0 = 2.5 \times 10^{-7}$	W/cm^2 sr cm^{-1}
* MINIMUM DETECTABLE QUANTITY D (see figure)	$\min D = 0.4 - 1.0 \times 10^{13}$	(molec/ cm^2) (W/ cm^2 sr cm^{-1})
* UNCERTAINTY IN D	$\sigma_{D'} = 2.1 \times 10^{12}$	(molec/ cm^2) (W/ cm^2 sr cm^{-1})







Table 21 Detection Parameters for NO₂ $\nu_1 + \nu_3$ band

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	2850-2935	cm ⁻¹
Approximate wavelength	3.44	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu \approx 0.10$)	M = 850	
BAND PHOTON RADIANCE (scene)	3.41×10^{13}	ph/sec cm ² sr
Maximum of contrast $\tau_{\nu}^{\alpha}_{gv}$	C _{MAX} = 1.10×10^{-19}	cm ² /molec
Mean contrast.	$\mu' = 2.28 \times 10^{-20}$	cm ² /molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 1.99 \times 10^{-20}$	cm ² /molec
* Photon flux density on detector		
* from scene.	7.8×10^{11}	phot/sec cm ²
* from internal sources	8.5×10^{11}	phot/sec cm ²
* TOTAL	$J = 1.6 \times 10^{12}$	phot/sec cm ²
* BLIP $D_{\lambda_c}^*$	4.8×10^{12}	cm $\sqrt{\text{Hz/W}}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	min $t_d = 2.84$	sec
* CORRESPONDING BASELINE NESR.	$(\text{NESR})_0 = 2.7 \times 10^{-9}$	W/cm ² sr cm ⁻¹
* MINIMUM DETECTABLE QUANTITY D (see figure)	min D = $1.0 - 2.5 \times 10^{10}$	(molec/cm ²) (W/cm ² sr cm ⁻¹)
* UNCERTAINTY IN D	$\sigma_D = 4.3 \times 10^9$	(molec/cm ²) (W/cm ² sr cm ⁻¹)



NO₂ v₂ band





Table 22 Detection Parameters for NO₂ ν_2 band

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	710-795	cm ⁻¹
Approximate wavelength	13.3	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 850	
BAND PHOTON RADIANCE (scene)	5.32×10^{17}	ph/sec cm ² sr
Maximum of contrast $\tau_v \alpha_{gv}$	C _{MAX} = 1.03×10^{-19}	cm ² /molec
Mean contrast.	$\mu' = 2.95 \times 10^{-21}$	cm ² /molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 6.58 \times 10^{-21}$	cm ² /molec
* Photon flux density on detector		
* from scene.	1.2×10^{15}	phot/sec cm ²
* from internal sources	1.7×10^{15}	phot/sec cm ²
* TOTAL	J = 3.0×10^{15}	phot/sec cm ²
* BLIP $D_{\lambda_c}^*$	5.16×10^{11}	cm $\sqrt{\text{Hz}}/\text{W}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	min $t_d = 5.13 \times 10^{-3}$	sec
* CORRESPONDING BASELINE NESR.	(NESR) ₀ = 5.9×10^{-7}	W/cm ² sr cm ⁻¹
* MINIMUM DETECTABLE QUANTITY D (see figure)	min D = $0.6 - 1.75 \times 10^{15}$	(molec/cm ²) (W/cm ² sr cm ⁻¹)
* UNCERTAINTY IN D	$\sigma_{D'} = 4.0 \times 10^{14}$	(molec/cm ²) (W/cm ² sr cm ⁻¹)



SO₂ $\nu_1 + \nu_3$ band





Table 23 Detection Parameters for SO₂ $\nu_1 + \nu_3$ band

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	2450-2525	cm ⁻¹
Approximate wavelength	4.0	μ m
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 750	
BAND PHOTON RADIANCE (scene)	1.59×10^{14}	ph/sec cm ² sr
Maximum of contrast $\tau_{\nu} \alpha_{gv}$	C _{MAX} = 2.34×10^{-20}	cm ² /molec
Mean contrast.	$\mu' = 7.82 \times 10^{-21}$	cm ² /molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 6.08 \times 10^{-21}$	cm ² /molec
* Photon flux density on detector		
* from scene.	3.6×10^{12}	phot/sec cm ²
* from internal sources	3.9×10^{12}	phot/sec cm ²
* TOTAL	$J = 7.5 \times 10^{12}$	phot/sec cm ²
* BLIP $D_{\lambda_c}^*$	2.6×10^{12}	cm $\sqrt{\text{Hz}}/\text{W}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	min $t_d = 1.32$	sec
* CORRESPONDING BASELINE NESR.	$(\text{NESR})_0 = 7.2 \times 10^{-9}$	W/cm ² sr cm ⁻¹
* MINIMUM DETECTABLE QUANTITY D (see figure)	min D = $0.9 - 2.5 \times 10^{11}$	(molec/cm ²) (W/cm ² sr cm ⁻¹)
* UNCERTAINTY IN D	$\sigma_D = 3.8 \times 10^{10}$	(molec/cm ²) (W/cm ² sr cm ⁻¹)






Table 24 Detection Parameters for SO₂ ν_1 band

QUANTITY	SYMBOL, VALUE	UNITS
DETECTION BAND	1090-1210	cm ⁻¹
Approximate wavelength	8.7	μm
* NO. OF SPECTRAL ELEMENTS (for $\Delta\nu = 0.10$)	M = 1200	
BAND PHOTON RADIANCE (scene)	3.18×10^{16}	ph/sec cm ² sr
Maximum of contrast $\tau_v \alpha_{gv}$	CMAX = 5.13×10^{-20}	cm ² /molec
Mean contrast.	$\mu' = 1.30 \times 10^{-20}$	cm ² /molec
STANDARD DEVIATION OF CONTRAST	$\sigma' = 9.61 \times 10^{-21}$	cm ² /molec
* Photon flux density on detector		
* from scene.	7.3×10^{14}	phot/sec cm ²
* from internal sources	8.2×10^{14}	phot/sec cm ²
* TOTAL	J = 1.5×10^{15}	phot/sec cm ²
* BLIP $D_{\lambda_c}^*$	4.7×10^{11}	cm $\sqrt{\text{Hz/W}}$
* MINIMUM SCAN TIME FOR BLIP PERFORMANCE	min $\tau_d = 7.07 \times 10^{-3}$	sec
* CORRESPONDING BASELINE NESR.	(NESR) ₀ = 5.5×10^{-7}	W/cm ² sr cm ⁻¹
* MINIMUM DETECTABLE QUANTITY D (see figure)	min D = $0.3 - 1.0 \times 10^{13}$	(molec/cm ²) (W/cm ² sr cm ⁻¹)
* UNCERTAINTY IN D	$\sigma_{D'} = 1.4 \times 10^{12}$	(molec/cm ²) (W/cm ² sr cm ⁻¹)

