

LEVEL II

13

AD A108884

OFFICE OF NAVAL RESEARCH

Contract N000014-81-K-0167

Task No. NR 207-275

TECHNICAL REPORT NO. 1

A Role for Cytoplasmic Structural Proteins in the
Transport of Water and Salts in the Intestine

by

Paula T. Beall, Ph.D.

Department of Physiology
Baylor College of Medicine
1206 Moursund
Houston, Texas 77030

December 8, 1981

Reproduction in whole or in part is permitted for
any purpose of the United States Government

*Distribution of this report is unlimited.

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

DTIC
ELECTE
S DEC 28 1981 D

412102
81 12 28 024

Annual Report Office of Naval Research N00014-81-K-0167

A Role for Cytoplasmic Structural Proteins in the
Transport of Water and Salts in the Intestine

February 1, 1981 - January 31, 1984

This contract supports research to test the hypothesis - that the movement of water and salts across the epithelial membrane of the intestine depends not only on the properties of the cellular membranes but also on the structural properties of the cytoplasm of the cells.

Current theories ascribe all controls of cellular transport of water and salts to passive diffusion and specialized enzymes, pores, and channels at the cell membrane. Classical interpretation allows no role for the cytoplasmic components of the cells in regulation or control of transport. Scattered experimental evidence produced in the 1970's, however, suggests that the disruption of the microtubule and microfilament network of the cytoplasm can have an effect on the rate and direction of transport. This proposal is in the process of testing the idea that the cytoskeleton plays a role along with water to regulate transport. For this purpose the intestinal epithelium has been chosen as a membrane through which large amounts of sodium and water pass. This work is part of a coordinated effort among three labs to study this phenomenon in vivo in perfused rat intestine, in vitro on stripped intestinal mucosa in Ussing chambers, and in isolated intestinal enterocytes. Preliminary data produced by this team has already resulted in the acceptance of a role for cytoplasmic proteins in the electrical properties of the intestinal epithelium.

YEAR 1

In Year 1 the proposal is scheduled to address the proper procedures for the isolation of intestinal enterocytes in a viable state for subsequent transport studies in which the effects of cytoskeletal disrupting drugs will be tested.

1. Progress on perfusion of rat intestine in vivo: Before experiments to isolate enterocytes can begin base line parameters for the normal operation of the cells in the intact intestine must be determined. In collaboration with Dr. Cassidy, I have performed a sample of these experiments proving that cytochalasin B which disrupts actin filaments in cells does indeed reduce water and salt transport across the epithelium. I am continuing these experiments.


Methodology

- (a) L-15 medium: Because of its ability to sustain isolated epithelial cells in tissue culture and our preliminary experiments on isolation (section 3) we wish to use L-15 culture medium for the cell experiments. We must know how this effects transport in vivo.


1. It has been determined that the magnitude of water transport in normal intestine is greater with L-15 than with a balanced buffered salt solution.
2. It is possible that the variety of amino acids in L-15 stimulate transport.

Data

- (b) Effects of Cytochalasin B on transport in vivo: Experiments were done in which the intestine is first perfused with L-15 containing ^{14}C -PE6 as a volume marker. Absorption at normal levels was demonstrated. Then L-15, (^{14}C -PE6) was perfused through the intestine containing cytochalasin B at doses of 10 to 100 mg/ml. All doses caused secretion as predicted by the hypothesis:




- (c) Effects of Colchicine on transport in vivo: Experiments were done in which the intestine was perfused for control absorption and then with L-15 containing colchicine, a drug which disrupts microtubule polymerization.



A dose dependent response was seen for 0.5-2.5 mM colchicine. Secretion occurred as predicted by the hypothesis but to a lesser extent than with much lower doses of CB. Return to normal L-15 was able to slowly reactivate absorption over 1 hour.

- (d) Combination of CB and colchicine: Experiments were done in which intestine was perfused with L-15 to establish control absorption levels, and then with L-15 containing 20 mg/ml CB and 2.4 mM colchicine. Initially secretion was seen and then after $\approx$ 30 minutes, secretion ceased.



No return to absorption was noted upon perfusion with L-15 containing no drugs. The combined effects of the drugs are not reversible.

Results

- (e) All of the above experiments meet the predictions of the hypothesis that cytoskeletal proteins may play a role in transport in the intestine. These experiments are proceeding and will probably be published in 1982.

2. Progress on the Development of a Method for the Isolation of Viable Rat Enterocytes:

The experiments proposed in this project require large amounts of viable functioning cells from the surface of the small intestine. While methods had been developed for the isolation of enterocytes, these procedures produced cells that showed signs of morphological and biochemical damage. Experiments to determine the proper temperature the best isolation medium, and the oxygen tension requirements of the cells. These results are summarized in Table II.

TABLE II

Viability of Rat Enterocytes

Conditions of Isolation	% Cells Excluding Trypan Blue
L-15 Tissue Culture Medium bubbled with 95% oxygen- 5% CO ₂ (with or without serum)	90%
L-15 plus oxygen (preincubated with trypsin enzyme 1 hour)	80-90%
L-15 plus oxygen (preincubated with 0.05% collagenase for 1 1/2 hours)	70-80%
L-15 <u>NO oxygen</u>	50-70%
Buffer and 0.05% Hyaluronidase	20%
Balanced salt solution and EDTA	10-20%
Ringers Lactate	10-20%

- Conclusions:
1. Balanced salt solutions previously used for the isolation of chick enterocytes are not good for mammalian enterocytes.
 2. L-15 tissue culture medium which was designed to support colon and breast epithelial cells provides additional nutrients needed by these mammalian cells.
 3. The inclusion of oxygen dramatically increases short term survival.

These results were presented at the FASEB spring meetings 1981. In the same session were several people who had been attempting to isolate mamalian enterocytes with little success. All are going to try L-15, all agreed that salt solutions are not sufficient. In addition after these results were published another paper came out in Epithelial Transport of Water and Ions, W. Armstrong ed. 1981, extolling the virtues of L-15 culture medium for transport studies.

4. I will use L-15, plus oxygen at 25°C for my isolations.

3. Examination of Human Colon Cell Lines as Potential Model Systems:

Because a great deal of water and salt are readsorbed in the colon, the cells that line this organ might be of interest in the study also. In collaboration with Dr. L.P. Rutzky, of the University of Texas Medical School, Houston, I have examined the water and ion content and the NMR relaxation times of a number of normal and cancerous human colon cell lines. The results of the study are summarized in Table I.

TABLE I

Cell Line	(n)	Passage No.	Division Time-hrs	T ₁ (msec)	T ₂ (msec)	% Water
NORMAL COLON CELLS						
Adult NBV	(2)	12	24	1214±46	207±7	92.3±0.4
Adult NBM	(3)	13	24	1009±7	191±3	91.4±0.4
Fetal HSO677	(2)	20	20	1058±13	221±3	90.6±0.4
Fetal Hs0074	(2)	27	22	1106±24	163±23	
CANCER COLON CELLS						
LS 180	(5)	41	24	643±60	119±12	85.5±2.4
LS174T	(6)	37	20	744±16	121±3	86.6±0.5
Clone 3-5	(6)	93	19	663±65	127±11	89.7±1.9
Clone 6-6	(7)	60	20	716±39	114±10	86.4±1.2
HT 29	(7)	140	12	686±21	108±7	
SW 480	(4)	614	39	982±9	176±6	90.1±1.4
SW 1345	(6)	55	51	460±45	83±6	83.6±1.8

Conclusions: The normal cells will not continue to grow after a limited number of cell divisions in culture. This will make it difficult to obtain sufficient numbers of cells to test the hypothesis. However enough cells could be obtain to test the effects of the drugs in a complementary experiment to the freshly isolated small intestine cells. The morphological differences among the cancer cells make it unlikely, that they can be used as a model for transporting cells. However some investigators at UT Medical School are using an isolated epithelial cell from the kidney grown on millipore filters to undertake transport studies. These cultured cells may be a better model system.

Significant Achievement of Year 1

- 1.) The laboratory for the contract was set up in new department at Baylor College of Medicine after return from Washington.
- 2.) Methods for in vivo perfusion and isolation of intestinal enterocytes were developed.
- 3.) Experiments in the in vivo perfusion system showed that cytochalasin B and colchicine as cytoskeletal disrupting agents both caused secretion as predicted by the main hypothesis.
- 4.) Model systems of human colon cells were tested and may be used in experiments.
- 5.) Five major publications were published in Year 1.

Plans for Year 2

- 1.) Experiments on in vivo perfusion with cytochalasin and colchicine will be completed and published as background for conclusions on isolated enterocytes. All predictions of the hypothesis have held up so far, so some additional experiments may be conducted on other cytoskeletal disrupting agents.
- 2.) The in vivo model may be used to test drugs which will prevent secretion and restore normal function in secreting intestine. One such drug is "taxol" a newly discovered agent that causes rapid driving of polymerization of microtubule proteins.
- 3.) Sodium and water fluxes in isolated enterocytes will be measured and conditions for short term maintenance determined.
- 4.) Sodium and water fluxes in isolated enterocytes will be determined in the presence of cytochalasin B and colchicine.

CURRENT REPORTS AND PUBLICATIONS

Publications:

1. P.T. Beall, "Contribution of Cytoskeleton to Cellular Water Properties in Diverse Functional States". Fed. Proc. 40(2): 206-213, (1981).
2. P.T. Beall, "The Water of Life". In The Sciences Jan. 1981: 6-10 (1981).
3. P.T. Beall, S.P. Bowen, M.M. Cassidy and M. Dinno, "Effect of Structural Perturbation by Cytochalasin B on Ion and Water Transport in Tight and Leaky Epithelia. An Electrical Model". J. Physiology (London) 308: 90-91 (1980).
4. P.T. Beall, C.F. Hazlewood and L.P. Rutzky, "NMR Water Relaxation Times of Human Colon Cancer Cell Lines and Clones". Can. Biochem. Biophys (accepted for publication Dec. (1981)).
5. D.C. Chang, L.K. Misra, P.T. Beall, R.C. Fanguy and C.F. Hazlewood, "Nuclear Magnetic Resonance of Muscle Water Protons in Muscular Dystrophy of Chickens". J. Cell. Physiology 107:139-145 (1981).
6. P.T. Beall, "NMR Relaxation Times of Water Protons in Cultured Cells During Freezing and Under Osmotic Stress". In Biophysics of Water, Felix Frank, Editor, Elsevier Holland, (1982).

Chapters:

1. P.T. Beall, B.B. Asch, D. Medina and C.F. Hazlewood, "Distinction of Normal, Preneoplastic, and Neoplastic Mouse Mammary Cells and Tissues by Nuclear Magnetic Resonance Techniques". In The Transformed Cell, Ivan Cameron and Thomas Pool, Editors, Academic Press, N.Y. pp. 293-325, (1981).
2. P.T. Beall, D. Medina and C.F. Hazlewood, "The Systemic Effect of Elevated Tissue and Serum Relaxation Times for Water in Animals and Humans with Cancers". In NMR Basic Principles and Progress, P. Diehl, E. Fluck and R. Kosfeld, Editors, Springer-Verlag, Berlin, Vol. 19, pp. 38-57, (1981).

Abstracts:

1. P.T. Beall, "Effects of Cytochalasin B and Colchicine on Secretion in the Small Intestine". FASEB, (1982).
2. P.T. Beall, L.K. Misra, C.F. Hazlewood and R.L. Young, "Hormone Dependent Changes in NMR Relaxation Times in Tissues of Aged Female Rats". Biophysical Society, (1982).
3. P.T. Beall, "Isolation of Rat Intestinal Enterocytes in L-15 Medium". Fed. Proc. 40(3): 420a, (1981).
4. P.T. Beall, D.C. Chang and C.F. Hazlewood, "Proton NMR Relaxation Times of Water in Contact and Non Contact Inhibited Cells". Biophysical Journal (1981).
5. M.A. Dinno, S.P. Bowen, P.T. Beall and M.M. Cassidy, "Effects of Cytochalasin B on the Properties of Frog Gastric Mucosa". FASEB (1980).

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO. AD-A208 884	3. REPORT'S CATALOG NUMBER
4. TITLE (and Subtitle) A Role for Cytoplasmic Structural Proteins in the Transport of Water and Salts in the Intestine		5. TYPE OF REPORT & PERIOD COVERED Annual Feb. 1, 1981- Dec. 8, 1982 (Jan. 31, 1982)
7. AUTHOR(s) Paula T. Beall, Ph.D.		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Physiology Baylor College of Medicine Houston, Texas 77030		8. CONTRACT OR GRANT NUMBER(s) N00014-81-K-0167
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research-Biological Science Div. 800 North Quincy Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 207-275 Distribution of this report is unlimited
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE Dec. 8, 1981
		13. NUMBER OF PAGES 7
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Unlimited		
DISTRIBUTION STATEMENT A Approved for public release; Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Unlimited		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Cytoskeleton, water, transport, intestine		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This contract supports research to test the hypothesis - that the movement of water and salts across the epithelial membrane of the intestine depends not only on the properties of the cellular membranes but also on the structural properties of the cytoplasm of the cells. Current theories ascribe all controls of cellular transport of water and salts to passive diffusion and specialized enzymes, pores, and channels at the cell		

membrane.

→ Classical interpretation allows no role for the cytoplasmic components of the cells in regulation or control of transport. Scattered experimental evidence produced in the 1970's, however, suggests that the disruption of the microtubule and microfilament network of the cytoplasm can have an effect on the rate and direction of transport. This proposal is in the process of testing the idea that the cytoskeleton plays a role along with water to regulate transport. For this purpose the intestinal epithelium has been chosen as a membrane through which large amounts of sodium and water pass. This work is part of a coordinated effort among three labs to study this phenomenon in vivo in perfused rat intestine, in vitro on stripped intestinal mucosa in Ussing chambers, and in isolated intestinal enterocytes. Preliminary data produced by this team has already resulted in the acceptance of a role for cytoplasmic proteins in the electrical properties of the intestinal epithelium.

In Year 1, of this contract the following significant findings have been made: 1) cytochalasin B, an actin depolymerizing drug, causes the onset of a secretory state in rat intestine at doses from 10-100 mg/ml, 2) colchicine, a microtubule disrupting drug, caused secretion at doses of 0.5-2.5 mM, but to a lesser extent than CB, 3) effects of CB and colchicine are reversible, 4) combinations of CB and colchicine initiate secretion but then stop all secretion and absorption within 1 hr. These findings are consistent with the hypothesis that cytoskeletal proteins play a role in transport.

In Year 2, experiments are planned to test the hypothesis in isolated intestinal enterocytes.

OFFICE OF NAVAL RESEARCH
BIOLOGICAL SCIENCES DIVISION
BIOPHYSICS PROGRAM, Code 444
DISTRIBUTION LIST FOR TECHNICAL, ANNUAL AND FINAL REPORTS

Number of Copies

- (13) Administrator, Defense Documentation Center
Cameron Station
Alexandria, Virginia 22314
- (6) Director, Naval Research Laboratory
Attention: Technical Information Division
Code 2627
Washington, D. C. 20375
- (6) Office of Naval Research
Attention: Code 102IP (ONRL DOC)
800 N. Quincy Street
Arlington, Virginia 22217
- (3) Office of Naval Research
Biophysics Program
Code 444
Arlington, Virginia 22217
- (1) Commanding Officer
Naval Medical Research and Development Command
National Naval Medical Center
Bethesda, Maryland 20014
- (1) Chief, Bureau of Medicine and Surgery
Department of the Navy
Washington, D. C. 20375
- (2) Technical Reference Library
Naval Medical Research Institute
National Naval Medical Center
Bethesda, Maryland 20014
- (1) Office of Naval Research Branch Office
Building 114, Section D
666 Summer Street
Boston, Massachusetts 02210
- (1) Office of Naval Research Branch Office
536 South Clark Street
Chicago, Illinois 60605

- (1) Office of Naval Research Branch Office
1030 East Green Street
Pasadena, California 91106
- (1) Commanding Officer
Naval Medical Research Unit No. 2
Box 14
APO San Francisco 96263
- (1) Commanding Officer
Naval Medical Research Unit No. 3
FPO New York 09527
- (1) Officer in Charge
Submarine Medical Research Laboratory
Naval Submarine Base, New London
Groton, Connecticut 06342
- (1) Scientific Library
Naval Aerospace Medical Research Institute
Naval Aerospace Medical Center
Pensacola, Florida 32512
- (1) Commanding Officer
Naval Air Development Center
Attn: Aerospace Medical Research Department
Warminster, Pennsylvania 18974
- (1) DIRECTOR
Naval Biosciences Laboratory
Building 844
Naval Supply Center
Oakland, California 94625
- (1) Commander, Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709
- (1) DIRECTORATE OF LIFE SCIENCES
Air Force Office of Scientific Research
Bolling Air Force Base
Washington, D. C. 20332
- (1) Commanding General
Army Medical Research and Development Command
Forrestal Building
Washington, D. C. 20314

(1)

Department of the Army
U. S. Army Science and
Technology Center - Far East
APO San Francisco 96328

(1)

Assistant Chief for Technology
Office of Naval Research, Code 200
800 N. Quincy Street
Arlington, Virginia 22217