

AD-A110 483

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/O 12/1

NUMERICAL SOLUTION OF DEGENERATE VARIATIONAL INEQUALITY ARISING--ETC(U)

SEP 81 C W CRYER, S Z ZHOU

DAA629-80-C-0041

UNCLASSIFIED

MRC-TSR-2274

NL

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1

1 - 1


END

DATE

FILED

3 42

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963-A.

AD A110483

LEVEL

2

MRC Technical Summary Report #2274

NUMERICAL SOLUTION OF DEGENERATE
VARIATIONAL INEQUALITY ARISING
IN THE FLUID FLOW THROUGH POROUS MEDIA

C. W. Cryer and S. Z. Zhou

MRC-TR-11

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

12 30

DTIC
SELECTED
FEB 4 1982
H

September 1981

(Received July 28, 1981)

DTIC FILE COPY

Approved for public release
Distribution unlimited

221200

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

and

National Science Foundation
Washington, DC 20550

82 02 03 063

UNIVERSITY OF WISCONSIN - MADISON
MATHEMATICS RESEARCH CENTER

NUMERICAL SOLUTION OF DEGENERATE VARIATIONAL INEQUALITY
ARISING IN THE FLUID FLOW THROUGH POROUS MEDIA

C. W. Cryer* and S. Z. Zhou**

Technical Summary Report #2274

September 1981

ABSTRACT

In this paper we propose a numerical method for a degenerate variational inequality arising in the axisymmetric porous flow well problems which have been studied in Cryer and Zhou [1981]. We use the finite element method to discretize the problem, and we establish the convergence of the solution of the discrete problem to the solution of the degenerate variational inequality. The solution of the physical problem depends upon the unknown discharge q . A rapidly convergent numerical method for finding q is obtained.

AMS (MOS) Subject Classifications: 35J70, 35R35, 65N30, 76S05

Key Words: Porous flow; numerical methods; degenerate variational inequalities; convergence; finite elements.

Work Unit Number 3 - Numerical Analysis and Computer Science

* Computer Science Department and Mathematics Research Center, University of Wisconsin-Madison.

** Hunan University (Changsha, China) and Mathematics Research Center, University of Wisconsin-Madison.

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041 and the National Science Foundation under Grant No. MCS77-26732.

SIGNIFICANCE AND EXPLANATION

Most steady-state porous flow free boundary problems may be reduced to elliptic, variational or quasi-variational inequalities for which the numerical solutions have been studied by many authors. Some axisymmetric problems lead to another kind of variational inequality, namely degenerate variational inequalities. We give a numerical method for a degenerate variational inequality arising in the axisymmetric porous flow well problems, and study the convergence of the solution of the discrete problem. Numerical examples show that the method is efficient.

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Avail and/or	
Dist	Special
A	



The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the authors of this report.

NUMERICAL SOLUTION OF DEGENERATE VARIATIONAL INEQUALITY
ARISING IN THE FLUID FLOW THROUGH POROUS MEDIA

C. W. Cryer* and S. Z. Zhou**

1. Introduction

Most steady-state porous flow free boundary problems may be reduced to elliptic, variational or quasi-variational inequalities for which the numerical solutions have been studied by many authors (see, for instance, Baiocchi and Capelo [1978], Oden and Kikuchi [1979], and their references). In some axisymmetric problems there appears another kind of variational inequalities - degenerate variational inequalities. In this paper we propose a numerical method for a type of degenerate variational inequality arising in the axisymmetric porous flow well problems which have been studied in Cryer and Zhou [1981]. We use the finite element method to obtain the discrete problem. The convergence of the solution of the discrete problem to the solution of the degenerate variational inequality is proved. The solution of the physical problem depends upon the unknown discharge q . We give a numerical method for finding q which has been found to converge rapidly.

Here we recall some notations and results about weighted Sobolev spaces V^1 and V^2 . (See Chang and Jiang [1978], Zhou [1980].)

A - a bounded domain in (r,z) -plane with a locally Lipschitz boundary Γ , and with $r > 0$.

$C^\infty(R^2)$ - the space of functions infinitely differentiable in (r,z) -plane.

$C^\infty(\bar{A}) = \{v : v \text{ is defined in } A \text{ and has extension in } C^\infty(R^2)\}$

$C_0^\infty(A; \Gamma^*) = \{v \in C^\infty(\bar{A}) : v = 0 \text{ in some neighborhood of } \Gamma^*\}$, where $\Gamma^* \subset \Gamma$. If

$\Gamma^* = \Gamma$ then denote $C_0^\infty(A; \Gamma^*)$ by $C_0^\infty(A)$.

* Computer Science Department and Mathematics Research Center, University of Wisconsin-Madison.

** Hunan University (Changsha, China) and Mathematics Research Center, University of Wisconsin-Madison.

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041 and the National Science Foundation under Grant No. MCS77-26732.

$L^2(A, r) = \{v : v \text{ real measurable in } A, \|v\|_{L^2(A, r)}^2 < \infty\}$, where $\|v\|_{L^2(A, r)}^2 = \int_A r |v|^2 dr dz$.

$$V^0(A) = L^2(A, r), \|v\|_{V^0(A)} = \|v\|_{L^2(A, r)}.$$

$$V^1(A) = \{v : \partial^\alpha v \in L^2(A, r), |\alpha| \leq 1\}, \|v\|_{V^1(A)} = \sum_{|\alpha| \leq 1} \|\partial^\alpha v\|_{L^2(A, r)}.$$

$$V^2(A) = \{v : \partial^\alpha v \in L^2(A, r), |\alpha| \leq 2, \frac{1}{r} \frac{\partial v}{\partial r} \in L^2(A, r)\}.$$

$$\|v\|_{V^2(A)} = \sum_{|\alpha| \leq 2} \|\partial^\alpha v\|_{L^2(A, r)} + \left\| \frac{1}{r} \frac{\partial v}{\partial r} \right\|_{L^2(A, r)}.$$

$V_0^1(A)$ - the closure of $C_0^\infty(A)$ in $V^1(A)$.

$V_0^1(A, \Gamma^*)$ - the closure of $C_0^\infty(A, \Gamma^*)$ in $V^1(A)$.

The following propositions can be easily shown.

Proposition 1.1. If $\Gamma^* \cap \{r = 0\} = \emptyset$ and $\text{meas}(\Gamma^*) > 0$ then there exists a constant C such that

$$\|v\|_{V^1(A)}^2 \leq C \int_A \left[\left(\frac{\partial v}{\partial r} \right)^2 + \left(\frac{\partial v}{\partial z} \right)^2 \right] r dr dz, \quad v \in V_0^1(A, \Gamma^*).$$

Proposition 1.2. $V^1(A)$, $V^2(A)$ and $V_0^1(A, \Gamma^*)$ are Banach spaces, where we take

$$\|v\|_{V_0^1(A, \Gamma^*)} = \|v\|_{V^1(A)}.$$

Proposition 1.3. $C^\infty(\bar{A})$, $C_0^\infty(A, \Gamma^*)$ are respectively dense in $V^1(A)$, $V_0^1(A, \Gamma^*)$.

Proposition 1.4. If $\Gamma^* \cap \{r = 0\} = \emptyset$ then there exists a unique linear continuous operator $\text{tr} : V^1(A) \rightarrow L^2(\Gamma^*)$ such that $\text{tr } v = v$ on Γ^* for any $v \in C^\infty(\bar{A})$. Moreover, tr is a compact operator.

2. Continuous Problem

Let D be a L -shaped domain such as that shown in Figure 1. Set

$$\Omega_1 = \{(r, z) : 0 < r \leq R_0, 0 < z < h_0\}.$$

The original variational formulation of the physical problem is as follows (Cryer and Zhou [1981]).

Problem (PPW). Let R_0, R_1, h_0, h_w, H be numbers such that $R_1 > R_0 > 0, H > h_w > h_0 > 0$.

Find functions $\varphi(r)$ and $u(r, z)$ such that

$$\varphi \in C^0([R_0, R_1]), \varphi(R_0) \geq h_w, \varphi(R_1) = H \quad (2.1)$$

φ is strictly increasing

(2.2)

$$u \in V^1(\Omega) \cap C^0(\bar{\Omega}) .$$

(2.3)




Figure 1

$$\begin{aligned} u &= H & \text{on } \Gamma_2 \\ &= z & \text{on } \Gamma_0 \cup (\Gamma_4 \cap \partial\Omega) \\ &= h_w & \text{on } \Gamma_5 \cup \Gamma_6 \end{aligned}$$

(2.4)

$$\int_{\Omega} r \nabla u \cdot \nabla v \, dx \, dy = 0 \quad \text{for all } v \in K_1$$

(2.5)

where

$$\begin{aligned}
\Gamma_0 &= \{(r, z) : z = \varphi(r), R_0 < r < R_1\}, \\
\Omega &= \{(r, z) : 0 < z < \varphi(r), R_0 < r < R_1\}, \\
K_1 &= \{v \in V^1(\Omega) : v = 0 \text{ on } \Gamma_2 \cup \Gamma_3 \cup (\Gamma_4 \cap \partial\Omega) \cup \Gamma_5 \cup \Gamma_6\}.
\end{aligned} \tag{2.6}$$

By using a kind of Baiocchi transform

$$\begin{aligned}
\bar{u} &= u(r, z) & \text{in } \Omega \\
&= z & \text{in } \bar{\Omega} \setminus \bar{\Omega}
\end{aligned} \tag{2.7}$$

$$w(r, z) = \int_0^z [\bar{u}(r, t) - t] dt \text{ in } \bar{\Omega}. \tag{2.8}$$

Cryer and Zhou [1981] have derived the following degenerate variational inequality with a real parameter q .

Problem (PPW1). Find $w_q \in K_q^{**}$ such that

$$\begin{aligned}
&\int_D r \nabla w_q \cdot \nabla (v - w_q) dx dz \\
&> (h_w - h_0) \int_0^{R_0} (v - w_q)|_{z=h_0} r dx + \int_D (v - w_q) r dx dz, \quad \forall v \in K_q^{**}
\end{aligned} \tag{2.9}$$

where

$$K_q^{**} = \{v \in V^1(D) : v > 0 \text{ in } D, v < g_q(r, H) \text{ in } D \setminus \Omega_1, v = g_q \text{ on } \Gamma_D\}, \tag{2.10}$$

$$\Gamma_D = \bigcup_{i=1}^5 \Gamma_i, \tag{2.11}$$

$$\begin{aligned}
g_q(r, z) &= 0 & \text{on } \Gamma_1 \\
&= Hz - \frac{z^2}{2} & \text{on } \Gamma_2 \\
&= \frac{H^2}{2} + q \ln \frac{r}{R_1} & \text{on } \Gamma_3 \cup \Gamma_4 \\
&= \frac{H^2}{2} + q \ln \frac{R_0}{R_1} - \frac{(h_w - z)^2}{2} & \text{on } \Gamma_5.
\end{aligned} \tag{2.12}$$

For Problem (PPW1) there is a equivalent form which is more convenient with regard to the numerical solution.

Problem (PPW2). Find $w_q \in K_q^{**}$ such that

$$J(w_q) = \min_{v \in K_q^{**}} J(v) \quad (2.13)$$

where

$$J(v) = \int_D r |\nabla v|^2 dr dz - 2(h_w - h_0) \int_0^{R_0} v|_{z=h_0} r dx - 2 \int_D v r dr dz. \quad (2.14)$$

Cryer and Zhou [1981] have proved the following results.

Proposition 2.1. For any q with $0 < q < q_0$, where

$$q_0 = \frac{H^2 - (h_w - h_0)^2}{2 \ln(R_1/R_0)}, \quad (2.15)$$

(PPW1) has a unique solution.

Proposition 2.2. Let w_q be the solution of (PPW1). Then there exist α, β and $F^*(q)$ with α, β real numbers, $\beta \neq 0$, F^* a continuous bounded function, such that the following two conditions are equivalent:

$$w_q \in C^1(\bar{D}) \cap V^2(D) \quad (2.16)$$

$$F^*(q) - \beta q - \alpha = 0. \quad (2.16')$$

We call w_q a regular solution of (PPW1) if (2.16) is valid.

Proposition 2.3. If w_q is a regular solution of (PPW1) then $0 < \bar{q} < q_0$.

Proposition 2.4. There exists at least one regular solution w_q for Problem (PPW1).

Proposition 2.5. For the regular solution w_q of (PPW1) we have

$$\frac{\partial w}{\partial r} > \frac{\partial \bar{q}}{\partial r} > 0, \quad \frac{\partial w}{\partial z} > 0 \quad \text{in } D \quad (2.17)$$

$$\begin{aligned} \frac{\partial w}{\partial n} &= h_w - h_0 & \text{on } \bar{\Gamma}_6 \\ &= 0 & \text{on } \bar{\Gamma}_7. \end{aligned} \quad (2.18)$$

Proposition 2.6. If u is a solution of (PPW) then w defined by (2.8) is a regular solution of (PPW1) corresponding

$$q = \bar{q} = r \int_0^H \frac{\partial \bar{u}(r,t)}{\partial r} dt.$$

Conversely, if $w_{\bar{q}}$ is a regular solution of (PPW1) then the functions $\varphi_{\bar{q}}, u_{\bar{q}}$ defined as follows, is the solution of (PP'')

$$\bar{\Omega}_{\bar{q}} = \bar{\Omega}_1 \cup \{(r,z) \in D : r > R_0, w_{\bar{q}} < g_{\bar{q}}(r,H)\}$$

$$\varphi_{\bar{q}}(r) = \sup\{z : (r,z) \in \bar{\Omega}_{\bar{q}}\}, \quad R_0 < r < R_1$$

$$\varphi_{\bar{q}}(R_0) = \lim_{r \rightarrow R_0+0} \varphi_{\bar{q}}(r), \quad \varphi_{\bar{q}}(R_1) = \lim_{r \rightarrow R_1-0} \varphi_{\bar{q}}(r)$$

$$\bar{u}_{\bar{q}} = \frac{\partial w_{\bar{q}}}{\partial z} + z \quad \text{in } \bar{D}$$

$$u_{\bar{q}} = \bar{u}_{\bar{q}}|_{\bar{\Omega}_{\bar{q}}}.$$

Remark 2.1. Physically, $2\pi\bar{q}$ is the "discharge" while u is the "hydraulic head".

3. Numerical Approximation of (PPW2)

In this section we consider the approximate problem of Problem (PPW2): Given q and h with $0 < q < q_0, h > 0$, find $w_q^h \in K_q^h$ such that

$$J(w_q^h) = \min_{v \in K_q^h} J(v) \quad (3.1)$$

where K_q^h is a convex closed nonempty subset of V^h , and $\{V^h\}$ is a family of finite dimensional subspaces of $V^1(D)$. It is not required that $K_q^h \subset K_q^{**}$. So far, for space V^2 there is no approximation theorem similar to that for usual Sobolev space H^2 . Hence we can not prove the convergence theorem for our problem by using usual methods such as those in Falk [1974], Brezi and Sacchi [1976], Cryer and Fetter [1977].

Let $\{T_h\}$ be a family of triangulations of the domain D . The set of interior gridpoints will be denoted by D_h , and the set of boundary gridpoints by ∂D_h . Set $\Gamma_{Dh} = \partial D_h \cap \Gamma_D$. For each T_h and for each triangle $T \in T_h$ we set

$$\rho(T) = \text{diameter of } T$$

$$\theta(T) = \text{minimum angle of } T$$

$$\theta_h = \min_{T \in T_h} \theta(T).$$

We assume that

$$\lim_{h \rightarrow 0} \max_{T \in T_h} \rho(T) = 0, \quad (3.2)$$

and that there exists a positive constant θ_0 independent of h such that (Zlamal [1968])

$$\theta_h > \theta_0. \quad (3.3)$$

As v^h we take the space of linear finite elements corresponding to T_h in $V_0^1(D; \Gamma_1)$. Set

$$K_q^h = \{v \in V^h : v \geq 0 \text{ on } D_h, v \leq g_q(x, H) \text{ on } D_h - \partial_1, v = g_q \text{ on } \Gamma_{Dh}\}. \quad (3.4)$$

We need the following basic theorem (see, for instance, Glowinski [1980, Th. 5.2 in Ch. I]).

Theorem 3.1. Let V be a real Hilbert space, $a(\cdot, \cdot)$ - a bilinear, continuous, symmetric and coercive form on $V \times V$, $f(\cdot)$ - a continuous, linear functional on V , K - a closed, convex, nonempty subset of V , $\{K^h\}$ - a family of closed, convex, nonempty subsets of V with $K^h \subset V^h$. Assume that

(i) If $\{v^h\}$ is bounded in V and $v^h \in K^h$ then the weak cluster points of $\{v^h\}$ belong to K ;

(ii) There exists a set $X \subset V$ with $\bar{X} = K$ such that $\forall v \in X$ there exists $\{v^h\}$ satisfying that $v^h \in K^h$ and that $\lim_{h \rightarrow 0} v^h = v$ strongly in V .

Then

$$\lim_{h \rightarrow 0} \|u^h - u\|_V = 0$$

where u^h and u are respectively the solutions of the problems

$$J(u) = \min_{v \in K} J(v)$$

and

$$J(u^h) = \min_{v \in K^h} J(v)$$

provided that $J(v) = a(v, v) - 2f(v)$.

In our case we will take $C^1(\bar{D}) \cap K_q^{**}$ as the set X in the above theorem.

From now on we assume that $0 < q < q_0$.

Lemma 3.2. $\overline{C^1(\bar{D}) \cap K_q^{**}} = K_q^{**}$ in $V^1(D)$.

Proof. Set $Y = C^1(\bar{D}) \cap K_q^{**}$. By Cryer and Zhou [1981, (3.28)] there exists a function $v_q \in Y$, hence Y is nonempty.

Set

$$Y_1 = \{v \in V^1(D) : v \geq -v_q \text{ in } D, v \leq g_q(x, H) - v_q \text{ in } D \setminus \Omega_1, v = 0 \text{ on } \Gamma_D\}$$

$$Y_2 = Y_1 \cap C_0^\infty(D; \Gamma_D).$$

Then by the argument similar to that of Lemma 2.4 in Glowinski [1980, Ch. II] we can prove that

$$\overline{Y_2} = Y_1 \quad (3.5)$$

$\forall v \in K_q^{**}$ we have $v^* = v - v_q \in Y_1$. By (3.5) there exists a sequence $v_n^* \in Y_2$ such that $v_n^* \rightarrow v^*$ strongly in V^1 . Thus $v_n = v_n^* + v_q \in Y$ and $v_n \rightarrow v$ strongly in V^1 . So we obtain that

$$\overline{C^1(\bar{D}) \cap K_q^{**}} \supset K_q^{**}.$$

Using the well-known subsequence argument and proposition 1.4 we can easily obtain that

$$\overline{C^1(\bar{D}) \cap K_q^{**}} \supset K_q^{**}.$$

Q.E.D.

The following lemma may be easily proved.

Lemma 3.3. If $T \in T_h$ and $v^h \in V^h$ then

$$\int_T v^h \, dx = \frac{\text{meas}(T)}{3} \sum_{i=1}^3 v^h(M_{iT}).$$

where M_{iT} ($i = 1, 2, 3$) are the vertices of T .

Now we can prove the convergence theorem.

Theorem 3.4. Problem (3.1) has a unique solution w_q^h which converges to the solution w_q of Problem (PPW2) in $V^1(D)$ as $h \rightarrow 0$.

Proof. Let $V = V^h$, $a(v_1, v_2) = \int_D r \nabla v_1 \cdot \nabla v_2 \, dr dz$, and $f(v) = \int_D r v \, dr dz + (h_w - h_0) \int_0^{R_0} v|_{z=h_0} r \, dr$. Then it is easy to see that V is a Hilbert space with inner product

$$(v_1, v_2) = \int_D r (v_1 v_2 + \nabla v_1 \cdot \nabla v_2) \, dr dz,$$

that K_q^h is a closed, convex, nonempty subset of V , that $a(\cdot, \cdot)$ is a symmetric, continuous, coercive (by proposition 1.1), bilinear form on $V \times V$, and that $f(\cdot)$ is a linear, continuous (by proposition 1.4) functional on V . By the well-known theorem (Stampacchia [1964], or Lions and Stampacchia [1967]) we know that Problem (3.1) has a unique solution w_q^h .

Now we prove the convergence of w_q^h by using Theorem 3.1. Let $V = V_0^1(D; \Gamma_1)$, $K = K_q^{**}$. It is sufficient to verify the conditions (i) and (ii) because it is obvious that the rest of the conditions are satisfied.

Verification of (i): Let $\{v^h\}$ be a sequence such that $v^h \in K_q^h$, and

$$v^h \rightarrow v \text{ weakly in } V_0^1(D; \Gamma_1). \quad (3.6)$$

We prove that $v \in K_q^{**}$. First we prove that

$$v \leq g_q(r, H) \text{ in } D \setminus \Omega_1. \quad (3.7)$$

Define $g_q(r, z) = g_q(R_0, z)$ in Ω_1 . Let g^h be the piecewisely linear interpolation of $g_q(r, H)$. Then it is easy to see that

$$g^h \rightarrow g_q(r, H) \text{ in } C^0(\bar{D}). \quad (3.8)$$

For any $\psi \in C_0^\infty(D \setminus \Omega_1)$ with $\psi \geq 0$ we define $\psi = 0$ in Ω_1 and

$$\psi^h = \sum_{T \in \mathcal{T}_h} \psi(G_T) \phi(T)$$

where $\phi(T)$ is the characteristic function of T , and G_T is the centroid of T . It can be shown that

$$\psi^h \rightarrow \psi \text{ in } C^0(\bar{D}). \quad (3.9)$$

Denote by T_h^* The union of triangles T which are contained in $\bar{D} \setminus \Omega_1$. Then we have

$$\begin{aligned}
 \int_{D \setminus \Omega_1} (v^h - g^h) \psi^h dx dz &= \int_{T_h^*} + \int_{D \setminus (\Omega_1 \cup T_h^*)} \\
 &= \sum_{T \in T_h^*} \int_T + \int_{D \setminus (\Omega_1 \cup T_h^*)} \\
 &= \sum_{T \in T_h^*} \psi(G_T) \int_T (v^h - g^h) dx dz + \int_{D \setminus (\Omega_1 \cup T_h^*)} (v^h - g^h) \psi^h dx dz \\
 &= \sum_{T \in T_h^*} \frac{\psi(G_T) \text{meas}(T)}{3} \sum_{i=1}^3 (v^h(M_{iT}) - g^h(M_{iT})) + \int_{D \setminus (\Omega_1 \cup T_h^*)} \quad (\text{by lemma 3.3}) \\
 &< \int_{D \setminus (\Omega_1 \cup T_h^*)} (v^h - g^h) \psi^h dx dz \quad (\text{since } v^h < g^h \text{ in } D^h - \Omega_1) \quad (3.10)
 \end{aligned}$$

Noting that $\text{meas}(D \setminus (\Omega_1 \cup T_h^*)) \rightarrow 0$ as $h \rightarrow 0$ we obtain by (3.6), (3.8), (3.9) and (3.10) that

$$\int_{D \setminus \Omega_1} (v - g_q(r, H)) \psi dx dz < 0, \quad \forall \psi \in C_0^\infty(D \setminus \Omega_1) \text{ with } \psi > 0$$

which implies that (3.7) is valid.

By similar argument we obtain that

$$v > 0 \text{ in } D. \quad (3.11)$$

Finally, it is easy to see that

$$v^h + g_q(r, z) \text{ in } C^0(\bar{\Gamma}_D).$$

On the other hand, it follows from proposition 1.4 that $v^h \rightarrow v$ strongly in $L^2(\Gamma_D)$. Thus we have

$$v = g_q \text{ on } \Gamma_D. \quad (3.12)$$

This equation as well as (3.11) and (3.7) means that $v \in K_q^{**}$.

Verification of (ii): Let $X = C^1(\bar{D}) \cap K_q^{**}$. By lemma 3.2 we have $\bar{X} = K_q^{**}$. For any $v \in X$ we take piecewisely linear interpolation v^h of v . Then $v^h \in K_q^h$. By a result of Feng [1965] we have (note that (3.2) and (3.3))

$$v^h \rightarrow v, \quad \frac{\partial v^h}{\partial r} \rightarrow \frac{\partial v}{\partial r}, \quad \frac{\partial v^h}{\partial z} \rightarrow \frac{\partial v}{\partial z} \text{ in } L^\infty(D)$$

as $h \rightarrow 0$. Therefore we have that $v^h \rightarrow v$ strongly in V^1 .

Q.E.D.

Remark 3.1. The condition (3.3) may be replaced by a weaker condition $\theta_h^* \leq \pi - \theta_0$, where $\theta_h^* = \max_{T \in T_h} \theta^*(T)$. $\theta(T)$ = maximum angle of T . (Cf. Feng [1965].)

The Problem (3.1) is a quadratic programming problem which can be computed using S.O.R. with projection (Cryer [1971], Glowinski [1971]). The iterative process is convergent (see, for instance, Glowinski et al. [1976, p. 70]). We have used Carré's scheme (Carré [1961]) to choose the relaxation factor ω .

4. Numerical Method for Regular Solution

Numerical experiments indicate that $w_q^h \rightarrow w_q$ as $h \rightarrow 0$ for q with $0 < q < q_0$, just as theorem 3.4 claims. But w_q does not satisfy (2.17) if q does not correspond to the regular solution. Here we give a method for searching for regular solution and corresponding values of q .

The basic idea is as follows. For the solutions (PPW1), their derivatives are continuous everywhere in $\bar{D}$ except at the reentrant point P . The character that a regular solution possesses is that its derivatives are continuous even at the point P . Hence we may use (2.18) at P to find regular solutions. By the way, equation (2.40) in Baiocchi et al. [1973] may also be derived by the same idea. Now we turn to the concrete computation.

We choose a subset S of the family $\{T_h\}$ such that $\forall T_h \in S$ there is a element $T^* \in T_h$ which has a vertical edge and P is an endpoint of this edge. Denote by h^* and P' respectively the length and the other endpoint of the edge. Thus we have the discrete form of (2.18) at the point P :

$$f(q) = w_q^h(P) - w_q^h(P') - h^*(h_w - h_0) = 0. \quad (4.1)$$

This equation can be numerically solved by, for example, the secant method which is given by

$$q_{n+1} = q_n - \frac{q_n - q_{n-1}}{f(q_n) - f(q_{n-1})} f(q_n), \quad n = 2, 3, \dots \quad (4.2)$$

The initial values q_1, q_2 must be subject to the condition $0 < q < q_0$. Then we compute $w_{q_1}^h, w_{q_2}^h$ by using S.O.R. with projection, and $f(q_1), f(q_2)$ by using (4.1), and q_3 by using (4.2). This process is repeated until prescribed accuracy is reached.

5. Numerical Examples

We adopt the triangulation of D used by Cryer and Fetter [1979] (see Figure 2). Suppose that m denotes the number of subdivisions of D in z -direction, n the number of subdivisions of $D \setminus \Omega_1$ in r -direction. Then the coordinates of the gridpoints are given by

$$\begin{aligned} z_j &= (j-1)H/m, \quad 1 \leq j \leq m+1 \\ r_i &= R_0 \exp[(i-k-1)/n \cdot \ln(R_1/R_0)], \quad i = k+1, \dots, n+k+1 \\ r_i &= (i-1)R_0/k, \quad i = 1, \dots, k \end{aligned}$$

where

$$\begin{aligned} k &= [R_0/\Delta], \\ \Delta &= R_0[(R_1/R_0)^{1/n} - 1]. \end{aligned}$$

For $v^h \in K_q^h$ let $U_{ij} = v^h(r_i, z_j)$, vector $U = \{U_{ij}\}$. Then

$$J(v^h) = \sum_{R(i,j)} J_{R(i,j)}(v^h)$$

where $R(i,j)$ are the rectangles

$$R(i,j) = \{(r,z) : r_i \leq r \leq r_{i+1}, z_j \leq z \leq z_{j+1}\}.$$

It is easy to compute

$$J_{R(i,j)}(v^h) = U^T A_{R(i,j)} U + 2b_{R(i,j)}^T U.$$

The matrix $A_{R(i,j)}$ and the vector $b_{R(i,j)}$ are almost same as that in (7.11) and (7.12) of Cryer and Fetter [1977]. The only difference is that for $j = k_1 - 1$ we must add respectively $(h_0 - h_w)\Delta x \cdot x_1/2$ and $(h_0 - h_w)\Delta x \cdot x_u/2$, which correspond to the line integral in $J(v)$ now, to $b_{R(i,j)}(i, j+1)$ and $b_{R(i,j)}(i+1, j+1)$, where k_1 is the value of j corresponding $z = h_0$.

For the S.O.R. with projection we only note that there are two constraints in our problem - $v^h > 0$ in D_h and $v^h \leq q_q(r, H)$ in $D_h - \Omega_1$.

The equation (4.1) now becomes

$$f(q) = U(kk, k_1) - U(kk, k_1 - 1) - (h_w - h_0)H/m = 0$$

where $kk = k+1$.




Figure 2

Example 1. $R_0 = 4.8$, $R_1 = 76.8$, $h_0 = 9$, $h_w = 12$, $H = 48$;

discretization: (1) $m = 8$, $n = 12$; $kk = 4$

(2) $m = 16$, $n = 24$; $kk = 9$

(3) $m = 32$, $n = 48$; $kk = 17$

(4) $m = 64$, $n = 96$; $kk = 35$

stopping test:

$$\max_{i,j} |U^{(l+1)}(i,j) - U^{(l)}(i,j)| < 10^{-6} \text{ for inner iteration}$$

$$|f(q_g)| < 10^{-6} \text{ for outer iteration}$$

The results are shown in the following table.

discretization	ω (Carré's Scheme)	number of outer iterations	number of inner iterations	q	h_g
(1)	1.4993	2	40	399.26	30
(2)	1.7291	4	90	371.83	33
(3)	1.8419	4	160	362.75	31.5
(4)	1.9336	4	380	358.66	30.75

where h_g is the approximate value of $\varphi_{\frac{1}{q}}(R_0)$.

Example 2. $R_0 = 10$, $R_1 = 1130$, $h_0 = 120$, $h_w = 200$, $H = 460$;

discretization: $m = 50$, $n = 100$; $kk = 21$

$\omega = 1.939$

stopping test: same as that in example 1.

The following table shows the convergence of the outer iteration process.

outer iteration	q	f(q)	number of inner iterations
0	100	25380.14648	450
1	200	25206.47851	450
2	14714.1875	43.04459	440
3	14739.0156	0.089335	440
4	14739.0673	0.00000035	440

$h_L = 349.6$ (compared with $h_g = 350$ in Borelli [1955]).

The exact value of q is 14218.462.

The theoretical proof that $w_q^h + w_{\frac{1}{q}}$ as $h \rightarrow 0$ is still an open problem.

Appendix A: An approximation theorem (Feng Kang [1965]).

Theorem. Let T be a triangle such as in Figure 3, P_0, P_1 and P_2 its vertices, P_0P_1 the largest edge with length ρ . Assume that $u \in C^1(\bar{T})$, v is the linear interpolation of u with $v(P_i) = u(P_i)$, $i = 0, 1, 2$.

Let

$$\omega' = \sup_{\substack{P, P' \in \bar{T} \\ 0 < \theta < \pi}} |u_\theta(P) - u_\theta(P')|$$

where u_θ is the directional derivative of u . Then




Figure 3

$$\left| \frac{\partial v}{\partial x} - \frac{\partial u}{\partial x} \right| < \omega' \quad (A.1)$$

$$\left| \frac{\partial v}{\partial y} - \frac{\partial u}{\partial y} \right| < (1 + \operatorname{ctg} \alpha_0) \omega' \quad (A.2)$$

$$|v - u| < (2 + \operatorname{ctg} \alpha_0) \rho \omega' \quad (A.3)$$

where α_0 is the inner angle with vertex P_0 .

Proof: Let P_3 be a point such that $P_3 \in P_0P_1$ and $P_2P_3 \perp P_0P_1$. Denote by ρ_1 the length of P_2P_3 . It is easy to see that for any $P \in \bar{T}$ we have

$$v(P) = u(P_0) + \frac{u(P_1) - u(P_0)}{\rho} x + \frac{u(P_2) - \bar{u}(P_3)}{\rho_1} y \quad (A.4)$$

where

$$\begin{aligned} \bar{u}(P_3) &= \sigma u(P_1) + (1-\sigma)u(P_0) \\ \sigma &= P_0P_3/\rho \end{aligned}$$

By the mean value theorem there exist $Q_0 \in P_0P_3$ and $Q_1 \in P_1P_3$ such that

$$\begin{aligned} u(P_0) &= u(P_3) + \int_{P_3}^{P_0} \frac{\partial u}{\partial x} dx = u(P_3) - \sigma \rho \frac{\partial u(Q_0)}{\partial x} \\ u(P_1) &= u(P_3) + \int_{P_3}^{P_1} \frac{\partial u}{\partial x} dx = u(P_3) + (1-\sigma) \rho \frac{\partial u(Q_1)}{\partial x} \end{aligned}$$

Hence

$$\bar{u}(P_3) = u(P_3) + \sigma(1-\sigma)\rho \left[\frac{\partial u(Q_1)}{\partial x} - \frac{\partial u(Q_0)}{\partial x} \right] . \quad (A.5)$$

By (A.4) and (A.5) we obtain that

$$\begin{aligned} \frac{\partial v(P)}{\partial x} &= \frac{u(P_1) - u(P_0)}{\rho} = \frac{\partial u(Q)}{\partial x} \quad \text{for some } Q \in P_0 P_1 \\ \frac{\partial v(P)}{\partial y} &= \frac{u(P_2) - u(P_3)}{\rho_1} = \frac{\rho\sigma(1-\sigma)}{\rho_1} \left[\frac{\partial u(Q_1)}{\partial x} - \frac{\partial u(Q_0)}{\partial x} \right] \\ &= \frac{\partial u(Q^*)}{\partial y} - (1-\sigma) \operatorname{ctg} \alpha_0 \left[\frac{\partial u(Q_1)}{\partial x} - \frac{\partial u(Q_0)}{\partial x} \right] \quad \text{for some } Q^* \in P_2 P_3 . \end{aligned}$$

Therefore

$$\begin{aligned} \left| \frac{\partial v(P)}{\partial x} - \frac{\partial u(P)}{\partial x} \right| &= \left| \frac{\partial u(Q)}{\partial x} - \frac{\partial u(P)}{\partial x} \right| < \omega' \\ \left| \frac{\partial v(P)}{\partial y} - \frac{\partial u(P)}{\partial y} \right| &< \left| \frac{\partial u(Q^*)}{\partial y} - \frac{\partial u(P)}{\partial y} \right| + \operatorname{ctg} \alpha_0 \left| \frac{\partial u(Q_1)}{\partial x} - \frac{\partial u(Q_0)}{\partial x} \right| < (1 + \operatorname{ctg} \alpha_0) \omega' \\ |v(P) - u(P)| &< |v(P_0) - u(P_0)| + \left| \int_{P_0}^P \left(\frac{\partial v}{\partial x} - \frac{\partial u}{\partial x} \right) dx + \left(\frac{\partial v}{\partial y} - \frac{\partial u}{\partial y} \right) dy \right| \\ &< (2 + \operatorname{ctg} \alpha_0) \omega' \rho . \end{aligned}$$

Q.E.D.

Appendix B: The Computer Program

```
implicit double precision (a,b,c,d,e,f,h,i,o,p,q,r,s,t,u,y)
integer ub
common/rx/x(150)
common/sy/s(65)
common/coefo/c0(150,65)
common/coef1/c1(150,65)
common/coef2/c2(150,65)
common/coef3/c3(150,65)
common/unk/u(150,65)
common/par/imax,n,m,modit,omega,test,m1,n1,k1,k2,kk,q
common/par1/y1,y2,y3,b,f2,h2,eps,iter,testa
data m/8/,n/12/,a/4.8d0/,ab/72.0d0/,y1/48.0d0/,
q y2/12.0d0/,y3/9.0d0/,eps/1.0d-6/
c calculate the stepsize for the discretization, uniformly for y,
c nonuniformly for x
  a1=log(a/a)
  b=a+ab
  b1=log(b/a)
  a1b1=b1-a1
  do 114 ii=1,3
c make the meshes finer
  h1=a1b1/n
  h2=y1/m
  d=a*(exp(h1)-1)
c d is the first dx in the outside of the well
  k=ifix(a/d)
  kk=k+1
  d1=a/k
c d1 is the dx for uniformly dividing the underneath of
c the well
  n1=n+kk
  m1=m+1
  do 5 i=1,k
5    r(i)=(i-1)*d1
  do 6 i=kk,n1
6    r(i)=a*exp((i-kk)*h1)
  do 10 j=1,m1
```

```

10      s(j)=(j-1)*h2
c  r(i),s(j) are the coordinates of the mesh points
      k1=dint(y3/h2)+1
      k2=dint(y2/h2)+1
c  k1 corresponds the bottom of the well
c  k2 correspond the water level
      print 15,a,b,y3,y2,y1,n,m,kk,k1,k2,eps
15      format(2x,3hrw=,f12.4,2x,3hre=,f12.4,2x,2hh=,f12.4,
q          2x,3hhw=,f12.4,2x,3hhe=,f12.4/
q          2x,3hnx=,15,2x,3hny=,15,2x,3hkk=,15,2x,3hk1=,15,2x,
q          3hk2=,15,2x,4heps=,f12.8/)
c  d is now another constant (for saving storage)
      d=d1*(y3-y2)/2
c  calculate the term in the right side corresponding the line
c  integral
      do 50 i=1,k
      c3(i,k1)= c3(i,k1)+(r(i)+d1/3)*d
50      c3(i+1,k1)=c3(i+1,k1)+(r(i)+2/3)*d
c  calculate the coef's c0,c1,c2 and the right side term corres-
c  ponding the multiple integral
      do 65 i=1,n1-1
      do 60 j=1,m
      if (i .lt. kk .and. j .gt. k1-1) go to 65
      dx=r(i+1)-r(i)
      dy=s(j+1)-s(j)
      t1=(r(i)+dx/3.0d0)*dx*dy/2
      t2=dx**2*dy/36.0d0
      r2=dx*t2+(r(i)+dx/3.0d0)*t1
      r3=-dy*t2/2+(s(j)+dy/3.0d0)*t1
      c0(i,j)=c0(i,j)+t1/dx**2+t1/dy**2
      c1(i,j)=c1(i,j)-t1/dx**2
      c2(i,j)=c2(i,j)-t1/dy**2
      c3(i,j)=c3(i,j)-t1+(r2-r(i)*t1)/dx+(r3-s(j)*t1)dy
      c0(i+1,j)=c0(i+1,j)+t1/dx**2
      c3(i+1,j)=c3(i+1,j)-(r2-r(i)*t1)/dx
      c0(i,j+1)=c0(i,j+1)+t1/dy**2
      c3(i,j+1)=c3(i,j+1)-(r3-s(j)*t1)/dy
      t1p=(r(i)+2*dx/3.0d0)*dx*dy/2

```

```

r2p=dx*t2+(r(i)+2*dx/3.0d0)*t1p
r3p=-dy*t2/2+(s(j)+2*dy/3.0d0)*t1p
c0(i+1,j+1)=c0(i+1,j+1)+t1p/dx**2+t1p/dy**2
c3(i+1,j+1)=c3(i+1,j+1)-t1p*(r2p-r(i+1)*t1p)/dx
c3(i+1,j+1)=c3(i+1,j+1)-(r3p-s(j+1)*t1p)/dy
c0(i,j+1)=c0(i,j+1)+t1p/dx**2
c1(i,j+1)=c1(i,j+1)-t1p/dx**2
c3(i,j+1)=c3(i,j+1)+(r2p-r(i+1)*t1p)/dx
c0(i+1,j)=c0(i+1,j)+t1p/dy**2
c2(i+1,j)=c2(i+1,j)-t1p/dy**2
60  c3(i+1,j)=c3(i+1,j)+(r3p-s(j+1)*t1p)/dy
65  continue
c give the first initial value for q. calculate the optimum
c acceleration factor omega using carre's method
q=100.0d0
call init
imax=150
omega=1.0d0
call itera
omega=1.4d0
do 19 iter=1,20
call itera
if (iter .eq. 17) test17=testa
if (iter .eq. 18) test18=testa
if (iter .eq. 19) test19=testa
19  if (iter .eq. 20) test10=testa
p18=test18/test17
p19=test19/test18
p20=test20/test19
if ((p18-p19)*(p19-p20) .lt. 0.0d0) go to 20
if (dabs(p18-p19) .le. dabs(p19-p20)) go to 20
lamdag=p18-(p19-p18)**2/(p18+p20-2*p19)
print 17,lamdag
17  format (20x,7haiken=,f8.4)
go to 25
20  lamdag=p20

```



```

25  sq=sqrt(1.0d0-(lamdag+omega-1.0d0)**2/(lamdag*omega**2))
    omega1=omega0
    omega0=2.0d0/(1.0d0+sq)
    print 26,omega0
26  format (/20x,7homega0=,f8.4)
    domega=dabs(omega1-omega0)
    if (domega/(2.0d0-omega0) .lt. 0.01d0) go to 45
    omegam=omega0-(2.0d0-omega0)/4
    print 30,omegam
30  format (20x,7homegam=,f8.4)
    omega=omegam
    do 40 iter=1,20
    call itera
    if (iter .eq. 19) test19=testa
40  if (iter .eq. 20) test20=testa
    p20=test20/test19
    go to 20
45  omega=omega0
    print 46,omega
46  format (/2x,6homega=,f8.4)
    test=1.0d0
    call iterat
    f1=f2
    q2=q

c  give the second initial value for q
    q=200.0d0

c  outer iteration. secant method for computing q
    do 79 iter1=1,10
    call init
    test=1.0d0
    call iterat
    q1=q2
    q2=q
    print 68,iter1,q1,q2,f1,f2
68  format (2x,6hiter1=,i3,2x,3hq1=,f14.8,2x,3hq2=,f14.8,2x,
q 3hf1=,f14.8,2x,3hf2=,f14.8)
    if (dabs(f2) .lt. eps) go to 80
    q=q2-(q2-q1)*f2/(f2-f1)

```

```

79      f1=f2
c  print the final results
80      ub=0
          lb=1
          f=f2
          q=q2
          do 110 n0=1,n1-1,7
              if(n1-1-n0)115,85,85
85      np=min(n1-n0+1,7)
          ub=ub+np
          print 90,(r(i),i=lb,ub)
90      format(///15x,f15.8,2x,f15.8,2x,f15.8,2x,f15.8,
q 2x,f15.8,2x,f15.8,2x,f15.8/)
          do 100 j=1, m1,1
              j1=m1-j+1
              print 95, s(j1), (u(i,j1), i=lb,ub)
95      format(2x,f11.4,2x,f15.8,2x,f15.8,2x,f15.8,2x,f15.8,
q 2x,f15.8,2x,f15.8,2x,f15.8)
100     continue
          lb=ub + 1
110     continue
115     print 120, iter1,q,f,test
120     format(///5x,6hiter1=,i3,2x,2hq=,f15.8,
q 2x,2hf=,f15.8,2x,5htest=,f15.8///)
          do 113 i=1,n1
              do 113 j=1,m1
                  u(i,j)=0.0d0
                  c0(i,j)=0.0d0
                  c1(i,j)=0.0d0
                  c2(i,j)=0.0d0
113      c3(i,j)=0.0d0
          m=2*m
114      n=2*n
          stop
          end
c  inner iteration. s.o.r. method for computing u and f
          subroutine iterat
              implicit double precision(c,o,t,u,v,q,y,b,f,h,e)

```

```

common/coef0/c0(9750)
common/coef1/c1(9750)
common/coef2/c2(9750)
common/coef3/c3(9750)
common/unk/u(9750)
common/par/imax,n,m,modit,omega,test,m1,n1,k1,k2,kk,q
common/par1/y1,y2,y3,b,f2,h2,eps,iter
iter=0
70  iter=iter+1
    modit=mod(iter,10)
    if (modit .eq. 0) test=0.0d0
    do 7 j=2,m
      do 7 i=1,n1-1
        if (i .le. kk .and. j .gt. k1) go to 7
        if (i .eq. kk .and. j .eq. k1) go to 7
        ij=i+imax * (j-1)
        im1j=ij-1
        ip1j=ij+1
        ijm1=ij-imax
        ipj1=ij+imax
c on boundary segment gama 7, the mesh points do not have
c neighbor mesh points at their left side. We use then u(1)=0.0d0
c instead of u(im1j) in the equations
        if (i .eq. 1) im1j=1
        uold=u(ij)
        unew=-(c3(ij)+c1(ij)*u(ip1j)+c2(ij)*u(ijp1)
q +c1(im1j)*u(im1j)+c2(ijm1)*u(ijm1))/c0(ij)
        vint=(1.0d0-omega)*uold+omega*unew
        u(ij)=dmax1(vint,0.0d0)
        if (i .gt. kk) u(ij)=dmin1(u(ij),u(i+imax*m))
        if (modit.ne.0) go to 7
        vabs=dabs(u(ij)-uold)
        test=dmax1(test,vabs)
7    continue
    if (iter .ge. 500) go to 74
    if (test .gt. eps) go to 70
74  print 75,iter,test
75  format('//20x,5hiter=,i3,2x,5htest=,f15.8)

```

```

      f2=u(kk+imax*(k1-1))-u(kk+imax*(k1-2))-(y2-y3)*h2
      return
    end

c initialize u. order: gama 2, gama 3, gama 4, gama 5, else-
c where (linear interpolation, and constant on gama 6)
    subroutine init
      implicit double precision (b,i,q,r,s,u,y,o,t,f,h)
      common/unk/u(150,65)
      common/rx/r(150)
      common/sy/s(65)
      common/par/imax,n,m,modit,omega,test,m1,n1,k1,k2,kk,q
      common/par1/y1,y2,y3,b,f2,h2
      do 201 j=1,m1
201    u(n1,j)=y1*s(j)-s(j)**2/2
      do 202 i=kk,n1
202    u(i,m1)=y1**2/2+q*log(r(i)/b)
      do 203 j=k2,m
203    u(kk,j)=u(kk,m1)
      do 204 j=k1,k2
204    u(kk,j)=u(kk,m1)-(y2-s(j))**2/2
      do 206 j=2,m
      do 205 i=kk+1,n1-1
        lamda=s(j)/y1
205    u(i,j)=u(i,m1)*lamda
206    continue
      do 208 j=2,k1
      do 207 i=1,kk
        lamda=s(j)/y3
207    u(i,j)=u(kk,k1)*lamda
208    continue
      return
    end

c iteration for carre's method
    subroutine itera
      implicit double precision (a,b,c,d,e,f,h,o,p,q,r,s,t,u,y)
      common/coefo/c0(9750)
      common/coef1/c1(9750)
      common/coef2/c2(9750)

```

```

non/coef3/c3(9750)
non/unk/u(9750)
non/par/imax,n,m,modit,omega,test,m1,n1,k1,k2,kk
non/par1/y1,y2,y2,b,f1,h2,eps,iter,testa
:a=0.0d0
!1 j=2,m
!1 i=1,n1-1
! .le. kk .and. j .gt. k1) go to 71
! .eq. kk .and. j .eq. k1) go to 71
imax*(j-1)+1
j=ij-1
j=ij+1
l=ij-imax
l=ij+imax
(i .eq. 1) im1j=1
!u(ij)
!=(c3(ij)+c1(ij)*u(ip1j)+c2(ij)*u(ijp1)
!m1j)*u(im1j)+c2(ijm1)*u(ijm1))/c0(ij)
:=(1.0d0-omega)*uold+omega*unew
j)=dmax1(vint,0.0d0)
(i .gt. kk) u(ij)=dmin1(u(ij),u(i+imax*m))
!dabs(u(ij)-uold)
:a=testa+vabs
on that we choose this norm for error may be found
[1961]
irn

```

REFERENCES

- Baiocchi, C. and Capelo, A.: Desequazioni variazional e quasivariazionali; applicazioni a problemi di frontiera libera, Pitagora Editrice, Bologna (1978).
- Baiocchi, C., Comincioli, V., Guerri, L., and Volpi, G.: Free boundary problems in the theory of fluid flow through porous media: A numerical approach, *Calcolo* 10, 1-85 (1973).
- Boreli, M.: Free-surface flow toward partially penetrating wells, *Trans. Amer. Geophysical Union* 36, 664-672 (1955).
- Brezzi, F. and Sacchi, G.: A finite element approximation of variational inequalities related to hydraulics, *Calcolo*, 13 (1976), 259-273.
- Carré, B. A.: The determination of the optimum accelerating factor for successive over-relaxation, *Comp. J.* 4, 73-78 (1961).
- Chang, K. C. and Jiang, L. S.: The free boundary problem of the stationary water cone, *Acta Sci Natur. Univ. Pekin*, 1-25 (1978).
- Cryer, C. W.: The solution of a quadratic programming problem using systematic over-relaxation, *SIAM J. Control* 9, 385-392 (1971).
- Cryer, C. W. and Fetter, H.: The numerical solution of axisymmetric free boundary porous flow well problems using variational inequalities, Technical Summary Report #1761, MRC, University of Wisconsin-Madison (1977).
- Cryer, C. W. and Fetter, H.: The numerical solution of axisymmetric free boundary porous flow well problems using variational inequalities, *Constructive Method for Nonlinear Boundary Value Problems and Nonlinear Oscillations*, Birkhäuser Verlag, Basel (1979).
- Cryer, C. W. and Zhou, S. Z.: The solution of the free boundary problem for an axisymmetric partially penetrating well, Technical Summary Report #2245, MRC, University of Wisconsin-Madison (1981).
- Falk, R. S.: Error estimates for the approximation of a class of variational inequalities, *Math. Computation*, 18 (1974), 963-971.

- Feng, Kang: Differencing scheme based on variational principle, Applied and Computational Math. (Chinese), 1965:4, 238-262 (1965).
- Glowinski, R.: La methode de relaxation, Rendiconti di Matematica, 14 Universita di Roma (1971).
- Glowinski, R.: Lectures on numerical methods for nonlinear variational problems, Springer-Verlag, New York (1980).
- Glowinski, R., Lions, J. L., and Tremolieres, R.: Analyse numerique des inéquations variationnelles, Paris : Dunad, (1976).
- Lions, J. L. and Stampacchia, G.: Variational inequalities, Comm. Pure Appl. Math. 20, 493-519 (1967).
- Oden, J. T. and Kikuchi, N.: Theory of variational inequalities with applications to problems of flow through porous media, Pergamon Press, Oxford, U.K. (1979).
- Stampacchia, G.: Formes bilinéaires coercitives sur les ensembles convexes, Comptes Rendus Acad. Sci. Paris 158, 4413-4416 (1964).
- Zhou, S. Z.: Functional Spaces $W_{p,2}^n$, J. of Hunan University, 1980 : 4, (1980).

CWC/822/jvs

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER #2274	2. GOVT ACCESSION NO. AD-A110483	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Numerical Solution of Degenerate Variational Inequality Arising in the Fluid Through Porous Media		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) C. W. Cryer and S. Z. Zhou		8. CONTRACT OR GRANT NUMBER(s) MCS77-26732 DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Madison, Wisconsin 53706 Wisconsin		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 3 - Numerical Analysis and Computer Science
11. CONTROLLING OFFICE NAME AND ADDRESS (see Item 18 below)		12. REPORT DATE September 1981
		13. NUMBER OF PAGES 26
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES U. S. Army Research Office and National Science Foundation P. O. Box 12211 Washington, DC 20550 Research Triangle Park North Carolina 27709		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Porous flow; numerical methods; degenerate variational inequalities; convergence; finite elements.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this paper we propose a numerical method for a degenerate variational inequality arising in the axisymmetric porous flow well problems which have been studied in Cryer and Zhou [1981]. We use the finite element method to discretize the problem, and we establish the convergence of the solution of the discrete problem to the solution of the degenerate variational inequality. The solution of the physical problem depends upon the unknown discharge q . A rapidly convergent numerical method for finding q is obtained.		

DD FORM 1473 EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

END

DATE
FILMED

3-82

DTIC