

AD-A111 199

NEW YORK UNIV NY DEPT OF CHEMISTRY

F/S 20/8

CORRELATION BETWEEN THERMAL ELECTRON TRANSFER IN SOLUTION AND P-ETC(U)

FEB 82 P DELAHAY

N00014-75-C-0397

ML

UNCLASSIFIED

TR-18

[or]
20
21 9H

END
DATE
FILMED
DTIC

AD A111199

(12)

OFFICE OF NAVAL RESEARCH

Contract N00014-75-C-0397

Task No. NR 051-258

TECHNICAL REPORT NO. 18 (New Series)

Correlation Between Thermal Electron Transfer
in Solution and Photoelectron Emission

by

Paul Delahay

Accepted for Publication

in

Chemical Physics Letters

New York University
Department of Chemistry
New York, NY

February 1982

A

Reproduction in whole or in part is permitted for
any purpose of the United States Government

This document has been approved for public release
and sale; its distribution is unlimited

DTIC FILE COPY

82 02 22 033

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 18 (New Series)	2. GOVT ACCESSION NO. HD-1111 189	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) CORRELATION BETWEEN THERMAL ELECTRON TRANSFER IN SOLUTION AND PHOTOELECTRON EMISSION		5. TYPE OF REPORT & PERIOD COVERED Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Paul Delahay		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0397
9. PERFORMING ORGANIZATION NAME AND ADDRESS New York University New York, NY 10003		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 051-258
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217		12. REPORT DATE February 1982
		13. NUMBER OF PAGES 19
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Accepted for publication in Chemical Physics Letters.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Activation free energy Photoelectron emission Electrode reactions Reorganization free energy Electron transfer, optical Electron transfer, thermal		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The activation free energy for electron transfer in solution or at electrodes is correlated to the corresponding Franck-Condon determined reorganization free energy R_m for photoelectron emission. Excellent to fair agreement is obtained between the activation free energies predicted from R_m and experimental values. Data are given for V^{2+} , Cr^{2+} , Mn^{2+} , Fe^{2+} , Co^{2+} and $Fe(CN)_6^{4-}$ in aqueous solution. $\leftarrow$ (C)6(4)		

DD FORM 1473

JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

✓
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

CORRELATION BETWEEN THERMAL ELECTRON TRANSFER IN SOLUTION AND PHOTOELECTRON EMISSION

PAUL DELAHAY

Department of Chemistry, New York University, New York, New York 10003, U.S.A.

The activation free energy for electron transfer in solution or at electrodes is correlated to the corresponding Franck-Condon determined reorganization free energy R_m for photoelectron emission. Excellent to fair agreement is obtained between the activation free energies predicted from R_m and experimental values. Data are given for V^{2+} , Cr^{2+} , Mn^{2+} , Fe^{2+} , Co^{2+} and $Fe(CN)_6^{4-}$ in aqueous solution.

1. Introduction

Electron transfer between species M^{z+} and $M^{(z+1)+}$ ($z \geq 0$) in solution can be studied either as a homogeneous process such as electron exchange between isotopes of M or as a heterogeneous process at a metal or semiconductor electrode. The kinetics of these processes is characterized by a free energy of activation $\Delta G_x^\ddagger$ for electron transfer between isotopes and $\Delta G_e^\ddagger$ for electrode processes. These thermal electron transfer processes have their optical counterparts, namely intervalence-transfer absorption and photoelectron emission by solutions.

Correlations can be established between the preceding four types of electron transfer. The theories of electron transfer [1,2] of Marcus [3] and Hush [4] provide the key to such correlations. (See [1,2] for other theories.) The free energies of activation $\Delta G_x^\ddagger$ and $\Delta G_e^\ddagger$ in these theories are expressed as functions of reorganization free energies R_x and R_e , respectively. The latter are related to each other. Marcus [5] correlated in this fashion electron transfer in solution and at electrodes. Hush [6]

A

established the correlation between thermal electron transfer and intervalence-transfer absorption spectra. We examine in the present paper the relationship between thermal electron transfer and the energetics of photoelectron emission by solutions. This type of optical electron transfer is characterized by a reorganization free energy R_m which can be determined experimentally as was recently shown [7-9]. The correlation between the experimental quantities $\Delta G_x^\ddagger$ or $\Delta G_e^\ddagger$ and R_m thus obtained provides a very direct test of theories of electron transfer in solution. Such a test is also provided, albeit in a different way, by the widely obeyed Marcus cross relation [2,10]. The present test, just as the cross relation, is less dependent on model calculations of the reorganization free energies R_x and R_e than previous comparisons of experimental and predicted values of $\Delta G_x^\ddagger$ [2] and $\Delta G_e^\ddagger$ [11].

2. Reorganization free energy for photoelectron emission by solutions

Photoelectron emission by a species M^{z+} ($z \geq 0$) into the vapor phase of the solution of M^{z+} is represented by the two equations

$$M^{z+}(z,z) = M^{(z+1)+}(z+1,z) + e^-(g), \quad (1)$$

$$M^{(z+1)+}(z+1,z) = M^{(z+1)+}(z+1,z+1). \quad (2)$$

The symbol (z,z) denotes that both electronic and nuclear configurations of the ligand (if any) and solvent correspond to the ionic charge $z+$. The same definition applies to $(z+1,z+1)$ for the ionic charge $(z+1)+$. Conversely, $(z+1,z)$ indicates that the electronic configuration pertains to the charge $(z+1)+$ whereas the nuclear configuration corresponds to the charge $z+$ (Franck-Condon principle). Thus, the species produced by (1) reorganizes according to process (2).

The free energy of emission ΔG_m for (1) in the case of aqueous solutions is given by [7-9]

$$\Delta G_m = \Delta G_H + \Delta G - R_m. \quad (3)$$

There, ΔG_H (≈ 4.50 eV) correlates the free energy level of the normal hydrogen electrode to the electron vacuum level on the assumption of a negligible surface potential; ΔG is the change of standard free energy for the oxidation of $M^{z+}(z,z)$ to $M^{(z+1)+}(z+1,z+1)$; R_m (< 0) is the reorganization free energy for (2). The free energy ΔG_m was shown [7-9] to be equal (within ca. ± 0.1 eV) to the threshold energy determined from experimental emission spectra. Thus, R_m can be determined experimentally by application of eq. (3) provided that ΔG is known.

The reorganization free energy R_m can be subdivided into inner- and outer-sphere contributions as is done in electron transfer theories for the quantities R_x (chemical) and R_e (electrochemical). Thus,

$$R_m = R_m^{\text{in}} + R_m^{\text{out}}. \quad (4)$$

The outer-sphere contribution is interpreted by treating the solvent macroscopically as a continuous medium, and the inner-sphere contribution is treated microscopically [1,2].

The outer-sphere contribution R_m^{out} is calculated by application of the Marcus theory [12] of non-equilibrium polarization. By starting with eqs. (6.59) and (6.62) in [2] or eq. (25) in [12], one obtains (rationalized units)

$$R_m^{\text{out}} = -(\epsilon_0/2)(\epsilon_{\text{op}}^{-1} - \epsilon_s^{-1}) \int_V (\vec{E}_{z+1} - \vec{E}_z)^2 dV, \quad (5)$$

where ϵ_0 is the permittivity of free space; ϵ_{op} and ϵ_s are the optical and static dielectric constants of the solvent, respectively; $\vec{E}_{z+1}$ and $\vec{E}_z$ are the electric fields in vacuo of the charges $(z+1)|e|$ and $z|e|$, respectively; and V is the integration volume. It will be assumed to simplify matters that the ions involved have the same size. This is a minor approximation. The volume V extending to infinity is supposed to surround a sphere of radius a for both fields in (5), and the Coulombic potential is integrated from $r = a$ to ∞ . Thus,

$$R_m^{\text{out}} = - (1/4\pi\epsilon_0)(\epsilon_{\text{op}}^{-1} - \epsilon_s^{-1})e^2/2a. \quad (6)$$

The contribution R_m^{out} given by (6) is independent of the ionic charge of the species emitting photoelectrons. Thus, R_m^{out} according to this model should be the same for emission by $M^{z+}(z,z)$ and $M^{(z+1)+}(z+1,z+1)$ ions provided the radius a is the same. Conversely, the inner-sphere contribution R_m^{in} is expected to depend on the ionic charge z of the emitter because the force constants determining its magnitude [2,11] vary with z . The dependence of R_m on z nevertheless should not be a strong one because R_m^{out} is markedly larger than R_m^{in} (sec. 3). The application of the Born equation with the factor $(\epsilon_{\text{op}}^{-1} - \epsilon_s^{-1})$ to the calculation of the reorganization free energy in [8] leads to a $(z+1)^2 - z^2$ dependence and is not justified.

3. Correlation between $\Delta G_x^\ddagger$ for electron exchange and R_m for photoelectron emission

Electron exchange between cations of ionic charge z^+ and $(z+1)^+$ (isotope labeling) will be considered in this section. The change of free energy is equal to zero in that case. Moreover, the work w_r required to bring from infinity the two reactants together in the activated complex is equal to the work w_p needed to perform the same operation for the products. Hence, $w_p - w_r = 0$, and one has [1-4]

$$\Delta G_x^\ddagger = w_r + R_x/4. \quad (7)$$

A relationship between R_x of (7) and R_m of (3) will be obtained. The method of calculating R_x is briefly recalled for that purpose. The outer-sphere contribution R_x^{out} is calculated [1-4] by assuming that the separation between the two ions of the activated complex is sufficient to preserve the spherical symmetry of the field of each ion but not sufficient to allow the neglect of Coulombic interaction. R_x^{out} is then given by an equation identical to (6) (with a plus sign) and the factor $(1/2a_1 + 1/2a_2 - 1/r_{1,2})$ instead of $1/2a$. The change of sign arises because R_x is

defined in terms of the difference of free energies between the activated complex and the product of electron transfer (eq. (6.73) in [2]) whereas R_m is the change of free energy for process (2). The quantities a_1 and a_2 are the radii of the ions and $r_{1,2}$ is the distance between the centers of the reactants in the activated complex. In general, the radius a_i is set equal to the sum of the crystallographic radius of the ion and the diameter ($2 \times 1.38 \text{ \AA}$) of the water molecule [13]. Moreover, one sets $r_{1,2} = a_1 + a_2$. If one assumes $a_1 = a_2 = a$, the final equation for R_x^{out} is the same as eq. (6) except for the sign. Thus, $R_x^{\text{out}} = |R_m^{\text{out}}|$ according to this model. Two ions are involved as reactants in the exchange reaction whereas photons are absorbed by single ions. Hence, $R_x^{\text{in}} = 2|R_m^{\text{in}}|$. In view of eq. (4) and a similar relationship for R_x , one has

$$R_x = |R_m| + |R_m^{\text{in}}|. \quad (8)$$

This is the relationship needed to test eq. (7). The quantity R_m is experimental (cf. eq. (3)) and R_m^{in} is calculated for a bond-stretching model from eq. (6) in [10].

The computation of R_x by application of (8) does not require the use of the continuous medium model and the attending selection of the radius a and the usual assumption of dielectric unsaturation. The use of eq. (8) therefore removes the uncertainty resulting from the model computation of the outer-sphere contribution R_x^{out} in the verification of eq. (7). Equation (8), it should be noted, is not rigorous because it was obtained by assuming that the fields of the two reactants in the activated complex are spherically symmetrical. This is an approximation for which somewhat tentative corrections have been proposed [2]. Such corrections will not be attempted here to preserve the simplicity of the treatment.

Equations (7) and (8) were applied to the calculation of $\Delta G_x^\ddagger$ and comparison of the results with experiment. The work w_r in (7) was

calculated from [3]

$$w_r = z(z+1)(1/4\pi\epsilon_0\epsilon_s)(e^2/2a)F(\mu), \quad (9)$$

where the function $F(\mu)$ of the ionic strength μ corrects the Coulombic term in (9) for finite dilution. The linearized expression [14], $F(\mu) = (1 + 2a\kappa)^{-1}$, was used where κ is the reciprocal Debye length. This expression for $F(\mu)$ and the corresponding exponential form $\exp(-2a\kappa)$ are very crude at high ionic strength. Fortunately this does not matter much because w_r is small in comparison with $\Delta G_x^\ddagger$ for the systems studied here.

Values of $\Delta G_x^\ddagger$ calculated from (7) and (8) are compared with the experimental results in Table 1. The agreement is excellent for Mn^{2+} , Fe^{2+} , Co^{2+} but not as good for the other ions. The discrepancy may have several possible causes: e.g., kinetic complications in the experimental determination of $\Delta G_x^\ddagger$, transmission coefficient appreciably smaller than unity, departure from the medium-overlap case assumed in [3,4]. Errors in the determination of R_m from threshold energy measurements could hardly account for the discrepancies in Table 1.

The value of $|R_m^{out}| = |R_m| - |R_m^{in}|$ computed from the data in Table 1 will be compared with the values of $|R_m^{out}|$ calculated for the continuous-medium model. The latter is 1.14 ± 0.01 eV for the five cations [11] whereas the values of $|R_m^{out}|$ calculated from the data in Table 1 are 1.72, 1.47, 1.28, 1.65, 1.85 eV from V^{2+} to Co^{2+} . These values hold within ca. ± 0.3 eV because of the uncertainty on R_m (surface potential, extrapolation method) and the approximations and estimates in the computation of R_m^{in} . The minimum for Mn^{2+} matches perhaps not fortuitously the minimum in the corresponding sequence of the absolute values of the free energies of solvation [15], 19.12, 19.28, 18.92, 19.58, 20.85 eV. It seems that the continuous-medium model yields values of $|R_m^{out}|$ which are too low for the values of a in Table 1 for the five cations examined here.

The opposite holds for $\text{Fe}(\text{CN})_6^{4-}$, but the difference between the value of 0.86 eV from the model [11] and 0.29 eV from Table 1 appears too great and probably reflects errors, at least in part (preliminary value of R_m in [9] from analysis of literature data). The foregoing conclusion about the relative inadequacy of the continuous-medium model is not surprising since this model is not very satisfactory either in the treatment of ionic solvation [16,17].

4. Correlation between $\Delta G_e^\ddagger$ for electron transfer at electrodes and R_m for photoelectron emission

The counterpart of (7) for electron transfer at electrodes at zero overvoltage is [5,10]

$$\Delta G_e^\ddagger = w_r + (R_e/4)[1 + (w_p - w_r)/R_e]^2. \quad (10)$$

There w_r is the work done to bring the reactant from the bulk of the solution to the site near the electrode at which electron transfer occurs. The same definition applies to w_p for the product of the reaction. The work terms in (10) are of the form $w_r = z|e|\phi$ and $w_p = (z+1)|e|\phi$ for the oxidation of a species with ionic charge $z|e|$ to one with charge $(z+1)|e|$. The potential ϕ can, in principle, be calculated from double layer theory.

Thus, $w_p - w_r = w_r/z$, and eq. (10) becomes, after neglecting the term $(1/z^2)(w_r/R_e)^2$,

$$\Delta G_e^\ddagger = (1 + 1/2z)w_r + R_e/4. \quad (11)$$

The reorganization free energy R_e is given according to Marcus [5,10] by an equation similar to (6) except that the factor $1/2a$ is replaced by $(1/2)(a^{-1} - r_f^{-1})$, where a is now the distance from the center of the reactant to the electrode and $r_f = 2a$ pertains to the image forces. The value of a of sec. 3 is used in practice. Thus, the factor $1/2a$ of eq. (6) is now replaced by $1/4a$, and therefore $R_e^{\text{out}} = |R_m^{\text{out}}/2|$. Taking $R_e^{\text{in}} = R_m^{\text{in}}$, one obtains

$$R_e = (1/2)[|R_m| + |R_m^{in}|]. \quad (12)$$

By comparing (8) and (12) one verifies that $R_e = R_x/2$ in agreement with [5,10]. This relationship is not rigorous because it corresponds to a simplified treatment of the image force problem.

Equations (11) and (12) are the counterparts of (7) and (8). The quantity $\Delta G_e^\ddagger$ can be calculated from R_m provided that R_m^{in} and w_r can be computed. The term R_m^{in} is treated as in sec. 3, but the calculation of w_r from double layer theory is tentative at best (except for the mercury electrode [11]). We therefore use R_m and experimental values of $\Delta G_e^\ddagger$ to compute w_r . The electrochemical work term w_r can also be computed as follows from the experimental values of $\Delta G_x^\ddagger$ and $\Delta G_e^\ddagger$ of Tables 1 and 2. Thus, R_x is computed from (7) using the experimental $\Delta G_x^\ddagger$'s and calculated chemical w_r 's from Table 1. The quantity R_e follows from $R_e = R_x/2$ according to Marcus [5,10]. The electrochemical work w_r is then computed from (11) using the experimental $\Delta G_e^\ddagger$'s from Table 2. The agreement between the two sets of electrochemical w_r 's in Table 2 is very good especially for a comparison involving three different experimental methods ($\Delta G_x^\ddagger$, $\Delta G_e^\ddagger$, R_m).

5. Conclusion

Excellent to fair agreement was obtained between experimental activation free energies for chemical and electrochemical electron transfer in solution and the values predicted from the corresponding Franck-Condon determined reorganization free energies for photoelectron emission. The results reported here support the basic theoretical model [3-5] for electron transfer in solution used in this work.

Acknowledgment

This work was supported by the Office of Naval Research and the National Science Foundation. The author thanks Professor R. A. Marcus for pointing out the need for a finite dilution correction in (9) and the model-free character of cross relation tests (sec. 1).

References

- [1] J. Ulstrup, Charge transfer processes in condensed media (Springer, Berlin, 1979).
- [2] R. D. Cannon, Electron transfer reactions (Butterworths, London, 1980) pp. 175-222, 267-297.
- [3] R. A. Marcus, J. Chem. Phys. 24 (1956) 966; 43 (1965) 679.
- [4] N. S. Hush, Trans. Faraday Soc. 57 (1961) 557.
- [5] R. A. Marcus, Can. J. Chem. 37 (1959) 155.
- [6] N. S. Hush, Electrochim. Acta 13 (1968) 1005.
- [7] K. von Burg and P. Delahay, Chem. Phys. Letters 78 (1981) 287.
- [8] P. Delahay, K. von Burg and A. Dziedzic, Chem. Phys. Letters 79 (1981) 157.
- [9] P. Delahay, Accounts Chem. Research, in press.
- [10] R. A. Marcus, J. Phys. Chem. 67 (1963) 853.
- [11] J. M. Hale in Reactions of molecules at electrodes, N. S. Hush, ed. (Wiley-Interscience, New York, 1971) pp. 229-257.
- [12] R. A. Marcus, J. Chem. Phys. 24 (1956) 979.
- [13] R. A. Marcus, J. Chem. Phys. 26 (1957) 867.
- [14] G. M. Brown and N. Sutin, J. Am. Chem. Soc. 101 (1979) 883.
- [15] R. M. Noyes, J. Am. Chem. Soc. 84 (1962) 513.
- [16] B. Case in ref. [11], pp. 45-134.
- [17] B. E. Conway, Ionic hydration in chemistry and biophysics (Elsevier, Amsterdam, 1981) pp. 312-315.

Table 1

Correlation between $\Delta G_x^\ddagger$ for electron exchange and R_m for photoelectron emission

	$-R_m$ a)	$-R_m^{\text{in}}$ b)	a b)	w_r c)	$\Delta G_x^\ddagger$ (calculated)	$\Delta G_x^\ddagger$ d) (experimental)
	(eV)	(eV)	(Å)	(eV)	(eV)	(eV)
V^{2+}	2.13	0.41	3.48	0.04	0.68	0.87
Cr^{2+}	2.05	0.58	3.49	0.05	0.71	1.03
Mn^{2+}	2.02	0.74	3.56	0.03	0.72	0.75
Fe^{2+}	2.11	0.46	3.51	0.06	0.70	0.69
Co^{2+}	2.26	0.41	3.48	0.05	0.72	0.72
$Fe(CN)_6^{4-}$	0.67	0.38	4.65	0.04	0.30	0.47

a) From [8] except for $Fe(CN)_6^{4-}$ (preliminary result [9] from analysis of literature data)).

b) From [11].

c) From (9) with $F(\mu)$ calculated for the prevailing μ 's in the experimental determination of the $\Delta G_x^\ddagger$'s.

d) From [2] and references cited therein.

Table 2

Correlation between $\Delta G_e^\ddagger$ for electron transfer at electrodes and R_m for photoelectron emission

	R_e ^{a)}	$\Delta G_e^\ddagger$ ^{b)} (experimental)	w_r ^{c)}	w_r ^{d)}
	(eV)	(eV)	(eV)	(eV)
V^{2+}	1.27	0.38	0.05	-0.03
Cr^{2+}	1.32	0.54	0.17	0.04
Mn^{2+}	1.38	0.43	0.07	0.06
Fe^{2+}	1.29	0.37	0.04	0.04
Co^{2+}	1.34	0.59	0.20	0.20
$Fe(CN)_6^{4-}$	0.53	0.30	0.15	0.08

a) From values of R_m and R_m^{in} in Table 1 and eq. (12).

b) From [11] and references therein.

c) From eqs. (11) and (12).

d) From experimental values of $\Delta G_x^\ddagger$ and $\Delta G_e^\ddagger$ (see text).

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 472. 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Area Office Attn: Scientific Dept. 715 Broadway New York, New York 10003	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
ONR Western Regional Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
ONR Eastern/Central Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (RE&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, GENNo.
Copies

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco 96503

1

Mr. James Kelley
DTNSRDC Code 2803
Annapolis, Maryland 21402

1

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. Paul Delahay Department of Chemistry New York University New York, New York 10003	1	Dr. P. J. Hendra Department of Chemistry University of Southampton Southampton SO9 5NH United Kingdom	1
Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 41106	1	Dr. Sam Perone Department of Chemistry Purdue University West Lafayette, Indiana 47907	1
Dr. D. N. Bennion Department of Chemical Engineering Brigham Young University Provo, Utah 84602	1	Dr. Royce W. Murray Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514	1
Dr. R. A. Marcus Department of Chemistry California Institute of Technology Pasadena, California 91125	1	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
Dr. J. J. Auburn Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. C. E. Mueller The Electrochemistry Branch Materials Division, Research & Technology Department Naval Surface Weapons Center White Oak Laboratory Silver Spring, Maryland 20910	1
Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. G. Goodman Globe-Union Incorporated 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201	1
Dr. T. Katan Lockheed Missiles & Space Co, Inc. P.O. Box 504 Sunnyvale, California 94088	1	Dr. J. Boechler Electrochimica Corporation Attention: Technical Library 2485 Charleston Road Mountain View, California 94040	1
Dr. Joseph Singer, Code 302-1 NASA-Lewis 21000 Brookpark Road Cleveland, Ohio 44135	1	Dr. P. P. Schmidt Department of Chemistry Oakland University Rochester, Michigan 48063	1
Dr. B. Brummer EIC Incorporated 55 Chapel Street Newton, Massachusetts 02158	1	Dr. F. Richtol Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181	1
Library P. R. Mallory and Company, Inc. Northwest Industrial Park Burlington, Massachusetts 01803	1		

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. A. B. Ellis Chemistry Department University of Wisconsin Madison, Wisconsin 53706	1	Dr. R. P. Van Duyne Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. M. Wrighton Chemistry Department Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Dr. B. Stanley Pons Department of Chemistry University of Alberta Edmonton, Alberta CANADA T6G 2G2	1
Larry E. Plew Naval Weapons Support Center Code 30736, Building 2906 Crane, Indiana 47522	1	Dr. Michael J. Weaver Department of Chemistry Michigan State University East Lansing, Michigan 48824	1
S. Rubv DOE (STOR) 600 E Street Washington, D.C. 20545	1	Dr. R. David Rauh EIC Corporation 55 Chabel Street Newton, Massachusetts 02158	1
Dr. Aaron Wold Brown University Department of Chemistry Providence, Rhode Island 02192	1	Dr. J. David Margerum Research Laboratories Division Hughes Aircraft Company 3011 Malibu Canyon Road Malibu, California 90265	1
Dr. R. C. Chudacek McGraw-Edison Company Edison Battery Division Post Office Box 28 Bloomfield, New Jersey 07003	1	Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton 509 5NH England	1
Dr. A. J. Bard University of Texas Department of Chemistry Austin, Texas 78712	1	Dr. Janet Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. M. M. Nicholson Electronics Research Center Rockwell International 3370 Miraloma Avenue Anaheim, California	1	Dr. R. A. Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. Donald W. Ernst Naval Surface Weapons Center Code R-33 White Oak Laboratory Silver Spring, Maryland 20910	1	Mr. James R. Moden Naval Underwater Systems Center Code 3632 Newport, Rhode Island 02840	1

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. R. Nowak Naval Research Laboratory Code 6130 Washington, D.C. 20375	1	Dr. John Kincaid Department of the Navy Strategic Systems Project Office Room 901 Washington, DC 20376	1
Dr. John F. Houlihan Shenango Valley Campus Pennsylvania State University Sharon, Pennsylvania 16146	1	M. L. Robertson Manager, Electrochemical Power Sonices Division Naval Weapons Support Center Crane, Indiana 47522	1
Dr. M. G. Sceats Department of Chemistry University of Rochester Rochester, New York 14627	1	Dr. Elton Cairns Energy & Environment Division Lawrence Berkeley Laboratory University of California Berkeley, California 94720	1
Dr. D. F. Shriver Department of Chemistry Northwestern University Evanston, Illinois 60201	1	Dr. Bernard Spielvogel U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709	1
Dr. D. H. Whitmore Department of Materials Science Northwestern University Evanston, Illinois 60201	1	Dr. Denton Elliott Air Force Office of Scientific Research Bldg. 104 Bolling AFB Washington, DC 20332	1
Dr. Alan Bewick Department of Chemistry The University Southampton, SO9 5NH England	1		
Dr. A. Himy NAVSEA-5433 NC #4 2541 Jefferson Davis Highway Arlington, Virginia 20362	1		

TECHNICAL REPORT DISTRIBUTION LIST, 051B

	<u>No. Copies</u>		<u>No. Copies</u>
Professor K. Wilson Department of Chemistry, B-014 University of California, San Diego La Jolla, California 92093	1	Dr. B. Vonnegut State University of New York Earth Sciences Building 1400 Washington Avenue Albany, New York 12203	1
Professor C. A. Angell Department of Chemistry Purdue University West Lafayette, Indiana 47907	1	Dr. Hank Loos Laguna Research Laboratory 21421 Stans Lane Laguna Beach, California 92651	1
Professor P. Meijer Department of Physics Catholic University of America Washington, D.C. 20064	1	Dr. John Latham University of Manchester Institute of Science & Technology P.O. Box 88 Manchester, England M601QD	1
Dr. S. Greer Chemistry Department University of Maryland College Park, Maryland 20742	1		
Professor P. Delahay New York University 100 Washington Square East New York, New York 10003	1		
Dr. T. Ashworth Department of Physics South Dakota School of Mines & Technology Rapid City, South Dakota 57701	1		
Dr. G. Gross New Mexico Institute of Mining & Technology Socorro, New Mexico 87801	1		
Dr. J. Kassner Space Science Research Center University of Missouri - Rolla Rolla, Missouri 65401	1		
Dr. J. Telford University of Nevada System Desert Research Institute Lab of Atmospheric Physics Reno, Nevada 89507	1		

END

DATE
FILMED

3-82

DTIC