

AD-A111 394


PURDUE UNIV LAFAYETTE IN DEPT OF CHEMISTRY F/8 20/13
HEAT CAPACITY OF WATER AT EXTREMES OF SUPERCOOLING AND SUPERHEAT--ETC(U)
DEC 81 C A ANSELL, W J SICHINA, M OGUNI N00014-78-C-0035

UNCLASSIFIED

ML

1-1
2-1

END
DATE
FILMED
8-82
DTIC



AD A111394

Report No. N00014-78-C-0035

Heat Capacity of Water at Extremes of
Supercooling and Superheating

C. A. Angell
Purdue University
Department of Chemistry
West Lafayette, Indiana 47907

December 1981

Technical Report, December 1981

Distribution Statement

Prepared for

Office of Naval Research
800 N. Quincy Street
Arlington, VA 22217

Division of Sponsored Programs
Purdue Research Foundation
Purdue University
West Lafayette, Indiana 47907

DTIC
ELECTRONIC
FEB 26 1982
A

This document has been approved
for public release and sale; its
distribution is unlimited.

01 19 82 125

DTIC FILE COPY

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER N00014-78-C-0035	2. GOVT ACCESSION NO. 11-A114	3. RECIPIENT'S CATALOG NUMBER 394
4. TITLE (and Subtitle) Heat Capacity of Water at Extremes of Supercooling and Superheating		5. TYPE OF REPORT & PERIOD COVERED Technical Report, Dec. 1981
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) C. A. Angell, W. J. Sichina and M. Oguni		8. CONTRACT OR GRANT NUMBER(s) N78C0035
9. PERFORMING ORGANIZATION NAME AND ADDRESS Purdue University Department of Chemistry West Lafayette, IN 47907		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research 800 N. Quincy Street Arlington, VA 22217		12. REPORT DATE December 1981
		13. NUMBER OF PAGES 23
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Water; metastable; supercooled; superheated; heat capacities; homogeneous nucleation		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) New and more accurate measurements of the heat capacity of supercooled water and D ₂ O are reported. The behavior of the heat capacity in the other metastable region, the superheated liquid, is compared with that in the supercooled region, using calculated values from the sophisticated equation of state of Haar and colleagues at the Bureau of Standards. An impressive similarity exists, homogeneous nucleation in both regions occurring at approximately the same value of heat capacity.		

DD FORM 1 JAN 79 1473

EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-LF-014-6601

Unclassified 29
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

TABLE OF CONTENTS

Abstract	i
Introduction	1
Experimental	2
Results	4
Discussion	4
Acknowledgments	10
References	
Table 1	
Table 2	
Figure Captions	
Distribution List	



Title		Date	
Author		Date	
Subject		Date	
Abstract		Date	
Distribution		Date	
Availability		Date	
Dist		Date	
A		Date	

Heat Capacity of Water at Extremes of Supercooling and Superheating

C. A. Angell, W. J. Sichina and M. Oguni
Department of Chemistry
Purdue University
West Lafayette, Indiana 47907

Abstract

deuterium oxide
Low temperature heat capacities for water and D_2O have been remeasured using different sample preparation and data analysis procedures from those used in the original studies of this anomalous region. Self-consistent results have been obtained with reproducibilities of 1%. Comparisons are made with the behavior in the superheated region at 1 atm pressure calculated from a refined equation of state by Haar et al. It appears that the kinetic metastability limit of the liquid (nucleation terminated) at both hot and cold extremes may be reached at the same value of the liquid heat capacity.

Heat Capacity of Water at Extremes of Supercooling and Superheating

C. A. Angell, W. J. Sichina and M. Oguni
Department of Chemistry
Purdue University
West Lafayette, Indiana 47907

In 1972 Rasmussen et al.⁽¹⁾ reported that the heat capacity of water increases very rapidly on supercooling below $\sim -15^{\circ}\text{C}$ and reaches values of $\sim 1.5 \text{ cal gm}^{-1}\text{deg}^{-1}$ ($113 \text{ J mol}^{-1}\text{deg}^{-1}$) at -38°C at which temperature crystallization of ice I terminated the measurements. At the other extreme, equally striking increases in heat capacity of water near the superheating limit have been determined by extrapolations to low pressure (based on corresponding states arguments) of high pressure stable range data.^(2,3) It had earlier been suggested by Anasimov et al.⁽⁴⁾ that water would show causally related anomalies in C_v at high and low temperature extremes.

The purpose of this report is to present new and improved data for the low temperature range and to point out that the phase change which terminates the metastable region at each extreme occurs in the region of sharply increasing heat capacity.

There is a particular need for improved data at low temperatures. The results of Rasmussen et al.,⁽¹⁾ which have helped generate much of the current interest in supercooled water, were essentially of exploratory character. They revealed an interesting phenomenon but exhibited differences of up to 10% between data sets from different methods of measurement, and different instruments. Nevertheless, for want of better, these data have been used as a basis for predicting and analyzing the behavior of other properties, such as sound velocities,^(5,6) possibly to the detriment of comparisons with experiment. For instance, if the values of C_p used were to contain an error of + 5% growing in between -25 and -33°C in the region of fast-increasing C_p , then a predicted

velocity minimum based on

$$v = \left(\frac{1}{\rho \kappa_g} \right)^{1/2} = \left(\frac{C_p}{C_v} \times \frac{1}{\rho \kappa_T} \right)^{1/2} \quad (1)$$

would disappear, if C_v were assumed constant. If C_v were (more correctly) subjected to the same correction, then the predicted velocity minimum, and the associated κ_g maximum, would be displaced to lower temperature, and a phenomenological relationship to the maximum in C_v ,^(7,8) which would now be revealed more clearly at $\sim -20^\circ\text{C}$, would become more obvious. Likewise, the thermodynamic absurdity

$$0 > C_p/C_v = [1 - T\alpha^2/\rho C_p \kappa_T]^{-1} \quad (2)$$

obtained⁽⁶⁾ by boldly extrapolating, to $T < -40^\circ\text{C}$, power law equations fitting α , C_p , and κ_T data measured at higher temperatures, will not appear.

We have therefore carefully repeated the low temperature measurements a number of times using differential scanning calorimetry of dispersed droplet samples, prepared and measured by different investigators, using different sample preparation techniques and somewhat different data analysis methods. We present here sets of emulsion data which are self-consistent to $\pm 1\%$, and agree to 0.5% with bulk sample data to the low temperature limit of the latter, -13°C . At the low temperature extreme, the new data differ by almost 5% from the average values previously reported.

EXPERIMENTAL

In the low temperature emulsion measurements, the water contribution to the total heat capacity of the two phase sample has been separated not only by the technique of comparing frozen and unfrozen sample data used previously⁽¹⁾

but also by the technique of off-setting the non-aqueous component with an equal mass of dispersant-surfactant solution in the reference pan. Comparison of results with literature data in the temperature range above 0°C suggest that these techniques are of approximately the same overall reliability. The former suffers from the addition of the random errors from two scans, while the latter suffers from a combination of errors from two weighings of small quantities, and additional uncertainties from possible changes in emulsion volume fraction of water during transfer to the sample pan.

Measurements were performed using the Perkin Elmer DSC-2 instrument equipped with the P.E. Scanning Auto Zero Accessory. Samples of mass ≈ 15 mgm were down-scanned from $\sim 20^\circ\text{C}$ at 10 deg min^{-1} on a sensitivity setting of $2 \geq \text{mcal/s}$. In the case of investigator I's determinations, the pen displacement with the instrument in the isothermal hold condition was carefully checked, for both empty and loaded pans, at short intervals between the initial and final temperatures of the scanning run. Under favorable instrument conditions this is unnecessary. However, in practice, variations in these relative isothermal hold pen displacements which would imply heat capacity errors of up to 5% for small samples studied in uncalibrated wide temperature range scans, can at times be observed, and this practice is necessary to obtain the best performance from the instrument.⁽⁹⁾

The question of whether the lowest temperature C_p results are influenced by partial crystallization during cooling cannot be satisfactorily answered by the obvious procedure of making measurements during temperature increase rather than decrease, though data above -33°C were confirmed by this method in earlier work.^(1c) The reason is that any measurements must start from an "isothermal hold" position, and a modest time, hence temperature, interval ($\sim 2^\circ\text{C}$) must be allowed for the instrument-sample ensemble to reach its correct steady-state condition for measurement. The isothermal hold temperature cannot be

set too close to T_H because the total elapsed time near T_H determines the probability of nucleation. Thus the temperature is already above the region, $T < 35^\circ\text{C}$, in which the threat of incipient crystallization influence is real, before the measurement can commence. A better confirmation of the absence of such artifacts is perhaps the absence of any pre-crystallization upswings in the case of emulsions of binary solutions in which the anomalous character of water has been "quenched" by the second component. The behavior of such solutions is systematically presented for the case of $\text{H}_2\text{O}_2 + \text{H}_2\text{O}$, in a previous paper.⁽⁹⁾

The most serious source of error in these measurements at low temperature would appear to be in the temperature calibration. This must be assumed the same for downscans as it is for upscans because due, to supercooling of standards, calibrations are always conducted on upscans.

RESULTS

The new data for water are collected in Table 1, and displayed graphically in Fig. 1. Discussion of the form of this heat capacity function, and the manner in which it is affected by changes of external pressure, and additions of second components, is given elsewhere.⁽⁹⁻¹¹⁾

A redetermination of C_p for D_2O has reproduced earlier reported DSC data⁽¹⁾ to better than the experimental uncertainty of $\pm 2\%$. The new data for bulk and emulsion samples are compared with the old, and with literature data for bulk samples, in Table 2.

DISCUSSION

Anomalous increases in liquid heat capacity at low temperature of the form of Fig. 1 are not found in other molecular liquids, even those with otherwise water-like properties, such as H_2O_2 and N_2H_4 , nor in mixtures of water with more than 20% of these liquids (even though some of these mixtures can be studied during continuous cooling down to -140°C where they vitrify).⁽⁹⁾

On the other hand, all molecular liquids which are internally stable against decomposition must exhibit anomalies of this character on sufficient superheating. This follows from the general form of the fluid equation of state which shows that a region of mechanical instability must exist in the isobaric V-T diagram for all pressures below the critical pressure. Such mechanical instabilities are generally illustrated by reference to the impossible negative compressibility region in the ordinary van der Waals equation P-V isotherm for $T < T_c$. For qualitative purposes, we illustrate the corresponding T-V behavior in Fig. 2 for a series of isobars at $P < P_c$ calculated for water using a van der Waals equation with parameters obtained, as usual, from the T_c and P_c values. The maxima correspond to minima in the corresponding P-V isotherms, hence to points at which the superheated liquid becomes mechanically unstable, and must spontaneously rupture. Of course, the simple van der Waals equation is inadequate to describe fluid water in detail, and it is therefore not surprising that T-V maximum, hence superheating limit, of 277°C predicted by this equation for 1 atm pressure, is less than the experimental superheating limit, T_H , of 279.5°C, observed for small droplets by different authors,^(12,13) or 302°C obtained by Skripov and coworkers from explosive boiling nucleation rate studies.⁽¹⁴⁾ The latter value is close to that calculated by Apfel for nucleation in bulk.⁽¹²⁾ The theoretical limit (the spinodal temperature T_g), should actually lie somewhat above the homogeneous nucleation temperature (which is the kinetically limited reflection of the impending catastrophe) and has been estimated at 305°C by Eberhart and Schnyders⁽¹⁵⁾ from equation of state calculations, and at 324°C at 1 atm by Skripov.⁽¹⁴⁾ In this region the impending instability must be anticipated by large fluctuations in density and entropy and these will be reflected in the magnitudes of the related thermodynamic properties, compressibility and heat capacity, in which we are interested here.

Although direct measurements of heat capacity or expansivity in this region have not yet been made they may be calculated using the equation of state described by Haar et al.⁽¹⁶⁾ The compressibility calculations up to 240°C from the same equation are in very good agreement with values obtained recently by Hareng and Leblond⁽¹⁷⁾ by combining direct light scattering measurements of adiabatic compressibility with extrapolated density data. Values of C_p calculated by Haar⁽¹⁸⁾ at normal pressure are plotted in Fig. 3 together with the low temperature data. It is notable that even the upswing in C_p at low temperatures is predicted by this equation. The increases at high temperature are more gradual than those predicted using a scaling equation suggested by Lienhard.^(2,3)

The compressibility data of Hareng and Leblond,⁽¹⁷⁾ which were in good agreement with earlier data to 180°C of Trinh and Apfel⁽⁵⁾ and more recent data to 240°C of Skripov and coworkers,⁽¹⁴⁾ were shown to fit very well an equation of the familiar critical law form

$$\kappa_T = A(T/T_s - 1)^{-\gamma} \quad (3)$$

with $T_s = 588$ K and $\gamma = 1$. This form had also been tested by one of the present authors⁽³⁾ on calculated heat capacities based on Lienhard's equation, under the impression that a value of $\gamma = 1$ was predicted by mean field theory. However, it now seems that $\gamma = 1$ is only applicable at the critical density. Away from the critical density, i.e. approaching a spinodal point, it appears from theoretical analyses by Compagne⁽¹⁹⁾ and Speedy⁽²⁰⁾ that the exponent should be 1/2 for κ_T , and also for α and C_p .

If we take the calculated C_p values of Haar covering the same C_p range as has been measured at low temperatures, we find that after subtracting a constant "background" contribution of $75 \text{ J K}^{-1} \text{ mol}^{-1}$ the excess C_p can be fitted to Eq. (1) with $T_g = 600.0 \text{ K}$ and $\gamma = 1.0$ - values similar to those found by Hareng and Leblond - as shown in the insert to Fig. 3. On the other hand, extension of the equation of state calculations (into regions where no validity would be claimed by its authors) shows $C_p \rightarrow \infty$ at 595.9 K , cf 597 K by Skripov⁽¹⁴⁾ and the limit is approached with a smaller Eq. (1) slope, $\gamma \approx 1/2$, as the new theoretical treatments suggest. It is unfortunate that, because of metastability and nucleation kinetics, it will not be possible to resolve the problem of the appropriate power law by experiment, as in the case of critical phenomena.

It has been shown elsewhere⁽⁹⁾ that the excess heat capacity of supercooled water obtained by subtracting, from the Table 1 values, a background C_p (determined from binary molecular and ionic solution data described elsewhere⁽⁷⁾) is also well described by Eq. (1) with a T_g value of 226 K and $\gamma \approx 1.0$. Interestingly enough, this singular temperature is just 5 degrees below the lowest freezing point ever achieved for pure water samples (cloud chamber experiments).⁽²¹⁾ The implied possibility that the low temperature phenomenon might be due to an instability was discussed in some detail in an earlier paper⁽²²⁾ and has now been incorporated into broad unified treatment of high temperature, low temperature and negative pressure instabilities in liquid water, by Speedy.⁽²³⁾

For the emulsion droplets studied here, which freeze commencing at -38.5°C , $\epsilon (= T/T_g - 1)$ at the phase change is 0.038 . The corresponding value for ϵ at the superheating limit of liquid-suspended droplets is 0.078 if we use $T_g = 600 \text{ K}$ from Haar's equation and $T_h = 553$ from droplet superheating limit studies.^(9,10)

However, this value of T_H may be too low since it corresponds with nucleation at the silicone oil + water surface.⁽¹²⁾ For bulk nucleation T_h is estimated to be about 298°C (571 K) by Apfel, and explosive nucleation measured by Skripov et al⁽¹¹⁾ yields $T_h = 573$ K, for which case $\epsilon = 0.045$, very close to the value at freezing. On the other hand the liquid heat capacities at the droplet superheating and supercooling limits are remarkably similar. This could be rationalized by arguing that, since C_p is proportional to the mean square entropy fluctuations, and since a critical nucleus is essentially a special entropy fluctuation, a correlation of nucleation probability and C_p might be expected.

A quantitative theory for the low temperature heat capacity behavior, in particular the apparent accord with a critical law, is currently lacking. Stanley and Teixeira⁽²¹⁾ have recently shown that the singularity, if real, could be associated with a percolation threshold for clusters of "four-bonded" water molecules, hence amorphous ice-like density fluctuations, in the liquid. On the other hand, Speedy⁽²⁰⁾ has now shown that all the volumetric observations on superheated, stretched (negative pressure), normal and supercooled water can be self-consistently explained by postulating the existence of a spinodal ($\kappa_T \rightarrow \infty$) line bounding the mechanically stable liquid regime, and deriving, by a Landau expansion, an equation of state which yields P_s on fitting to the available volumetric data. The form of this P_s, T_s boundary line is that of a normal liquid at high temperature, but at lower temperature and negative pressures, it departs from the normal van der Waals behavior (which is to constantly increasing negative P_s with decreasing T_s) and at $T_s = 35^\circ\text{C}$ $P_s = -2100$ bar, it reverses slope. The reversal point corresponds with the intersection of the spinodal with the extrapolation, to negative pressures, of the observed

temperatures of density maxima.⁽²⁰⁾ The spinodal (instability) thus reappears at positive pressures at temperature below the freezing point. The peculiar reversal would be a feature common to liquids in which the lowest energy state, which is approached with decreasing temperature, is correlated with the largest volume state. Unfortunately, there are very few such liquids (SiO_2 and some tetrahedrally coordinated antimonides and tellurides) so the phenomenon cannot be broadly studied.

Both the percolation and the spinodal models have attractive and challenging features, and will doubtless stimulate increasing experimental and theoretical work in this area. However, a tendency evident in some of the current literature to see these two approaches as somehow mutually exclusive should be discouraged. The language used in ref. 22 to describe the molecular nature of the increasing density fluctuations which the power law for the compressibility required to be present, ("density fluctuations presumably arise and decay by the migration of large $\text{O}\cdots\text{H}-\text{O}$ angle ($\rightarrow 180^\circ$) hydrogen bonds which 'prop open' the expanded structure when present in high local concentration but allow collapsed dense regions to form when local concentrations are smaller",) accurately previews the topological picture provided by the percolation model.⁽²⁴⁾ Future refinements of the current simplified percolation model to incorporate the actual continuous distribution of hydrogen bond energies and other interaction terms may well lead to an equation of state containing an instability at low temperatures, as indeed is contained in the optimized empirical equation of state of Haar and colleagues (which we have only used in this paper to discuss the high temperature anomalies⁽²⁵⁾). In this case the two approaches would obviously be mutually reinforcing.

Both approaches are, however, antagonistic to heterophase fluctuation interpretations of water anomalies since the latter contain no reason for excluding any liquids susceptible to crystallization from showing qualitatively similar anomalies.

REFERENCES

1. (a) Rasmussen, D. H.; MacKenzie, A. P.; Tucker, J. C.; Angell, C. A. Science 1973, 181, 4079.
(b) Rasmussen, D. H.; MacKenzie, A. P. J. Chem. Phys. 1973, 59, 5003.
(c) Angell, C. A.; Shuppert, J.; Tucker, J. C. J. Phys. Chem. 1973, 77, 3092.
2. Lienhard, J. H. Nucl. Sci. Eng. 1977, 62, 302.
3. Angell, C. A. "Proc. 9th Intern. Conf. on Properties of Steam", Eds. Straub and Grigull, Pergammon Press: London, 1980.
4. Anasimov, M. A.; Voronel, A. V., Zaugol'nikova, N. S.; Ovodov, G. I. JETP Letters 1972, 15, 317.
5. Trinh, E.; Apfel, R. E. J. Chem. Phys. 1978, 69, 4245.
6. D'Arrigo, G. Il Nuovo Cimento 1981, 61B, 123.
7. Chen, C. -H.; Frank, H. S., presented at the 167th Meeting of the Amer. Chem. Soc., Los Angeles, March 31 - April 5, 1974.
8. Angell, C. A. "Supercooled Water", in "Water: A Comprehensive Treatise", Vol. 7, ed. F. Franks, Plenum Press (in press).
9. Oguni, M.; Angell, C. A. J. Chem. Phys. 1980, 73, 1948.
10. Angell, C. A.; Shuppert, J.; Tucker, J. C. J. Phys. Chem. 1973, 77, 3092.
11. Kanno, H.; Angell, C. A. J. Chem. Phys. 1980, 73, 1940.
12. Apfel, R. E. Nature, Physical Science 1972, 238, 63; J. Chem. Phys. 1971, 54, 62.
13. Blander, M.; Hengstenberg D.; Katz, J. L. J. Phys. Chem. 1971, 75, 3613.
14. (a) Skripov, V. P. "Metastable Liquids", New York - Jerusalem, Wiley, Israel, Progr. Sci. Transl., 1974.
(b) Pavlov, P. A.; Skripov, V. P. Teplofizika Vishokikh Tempetarur 1970, 8, 579-585.
(c) Skripov, V. P.; Shuravenko, N. A.; Isaev, O. A. Teplofizika Visokikh Temperatur 1978, 16, 563-568.
15. Eberhard, J. G.; Schnyders, H. C. J. Phys. Chem. 1973, 77, 2730.
16. Haar, L.; Gallagher, J.; Kell, G. S. "Proc. 9th Intern. Conf. on Properties of Steam", Eds. Straub and Grigull, Pergammon Press: London, 1980.
17. Hareng, M.; Leblond, J. J. Chem. Phys. 1980, 73, 622.
18. Haar, L. (private communication).
19. Compagne, A. Physica 1973, 72, 115.

ACKNOWLEDGMENTS

The authors are indebted to the Office of Naval Research and the National Science Foundation for support of this research under Agreement and Grant Nos. N00014-78-C-0035 and DMR 77-04318A1, respectively, and to Lester Haar of the National Bureau of Standards for providing the equation of state data plotted in Fig. 3.

20. Speedy, R. J. (to be published).
21. Mason, B. J. Clouds, Rain and Rainmaking, University Press: Cambridge, 1962.
22. Speedy, R. J.; Angell, C. A. J. Chem. Phys. 1976, 65, 851.
23. Speedy, R. J. J. Phys. Chem. (in press).
24. Stanley, H. E.; Teixeira, J. J. Chem. Phys. 1980, 73, 3404.
25. The instability is actually predicted to occur at too high a temperature, -34°C ,⁽¹⁸⁾ though its pressure dependence at low pressures⁽¹¹⁾ is approximately the same as that of T_s and T_h displayed in ref. 11.

Table 1

Heat Capacity of Water in $\text{J K}^{-1} \text{mol}^{-1}$

Temperature T/K		Investigator I		Investigator II		Suggested Value									
							Emulsion Samples (sample freeze technique)		Bulk Sample	lit. data					
											n-octane	methyl cyclohexane carrier	Average	This work	lit. data
290	17	75.4			75.4	75.4 ^a									
285	12	75.4			75.2	75.5 ^a									
280	7	75.5			75.5	$\begin{cases} 75.6^a \\ 75.5^b \\ 75.9^a \end{cases}$									
275	2	75.4			75.5	75.9 ^a									
270	- 3	76.2	75.9	76.0	75.7	76.2 ^b	76.1								
265	- 8	76.6	75.7	76.2	76.6	76.8 ^b	76.4								
260	-13	77.5	77.3	77.4	77.6		77.6								
255	-18	78.3	79.5	78.9			78.8								
250	-23	80.6	82.2	81.4			81.4								
245	-28	84.4	86.4	85.4			85.2								
242	-31														
240	-33	92.7	92.7	92.7			92.3								
239	-34														
238	-35	96.6	96.8	96.7			96.5								
237	-36						99.0								
236	-37					102.2	102.7								

(a) Handbook of Physics and Chemistry, CRC press, 58th Edition.

(b) Anasimov et al, ref. 15.

Table 2

Heat Capacity of D₂O in J K⁻¹ mol⁻¹

Temperature		Investigator I					ref. 1
		Bulk		lit.	emulsion		emulsion
T/K	T/°C	Run I	Run II		Run I	Run II	
298.15	25			84.3 ⁺			
295				82.75 ⁺			
290	17	83.6		82.4 [*]			
289	16			83.58 ⁺			
285	12	83.75			85.4	83.5	
284	11			83.62 ⁺			
280	7	84.1	84.9		84.9	83.7	
279	6			84.36 ⁺			
275	2	84.7	85.2		85.1	84.1	
270	-3	86.1	86.0		86.0	85.0	84
265	-8	87.4	87.2		87.2	86.5	85.8
263	-10						87.7
260	-13				89.6	88.7	
255	-18				93.3	92.3	92.1
253	-20						95.5
250	-23				99.9	99.2	101.3
245	-28				109.3	109.5	110.0
242	-31				119.6		
240	-33				129.3		129.2

⁺ Termicheskie Konstanty Veshchestv (Thermal Constants of Substances) Academy of Sciences, USSR Moscow, 1965, Part I, p. 33.

^{*} NBS Circular 500

⁺ Kemp and Roth, J. Am. Chem. Soc. 1936, 58, 1829.

FIGURE CAPTIONS


Figure 1 Heat capacity of water in normal and supercooled temperature ranges at 1 atm pressure. ● literature data for bulk sample ○, □, △, data obtained in this study by investigator I for bulk sample (○) and emulsion samples using MCH carrier (□) and n-octane carrier (△): ■ data obtained in this study by investigator II for emulsion sample in n-octane carrier.


Figure 2 Isobars for water at subcritical pressures according to the Van der Waals equation showing the metastability limits at various pressures.

Figure 3 Combination of low temperature range data of this study with metastable superheated range data calculated from the equation of state of Haar et al for superheated range. The metastable state low and high temperature limits observed for small droplets are -42°C and 280°C , respectively (marked T_h).

Insert: Test of critical type law for high temperature equation-of-state C_p values after subtracting a constant "background" of $75 \text{ cal mol}^{-1} \text{ deg}^{-1}$. The plot uses $T_s = 600.0 \text{ K}$, and yields a slope of unity as indicated by the solid line in the inset.

Use of (calculated) C_p values beyond the C_p range of those measured in low temperature experiments, yields slightly lower T_s ($T_s \rightarrow 596 \text{ K}$) and smaller slopes ($\gamma \rightarrow 0.4$).








Fig. 3

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Area Office Attn: Scientific Dept. 715 Broadway New York, New York 10003	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
ONR Western Regional Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
ONR Eastern/Central Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (RE&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, GENNo.
Copies

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco 96503

1

Mr. James Kelley
DTNSRDC Code 2803
Annapolis, Maryland 21402

1

TECHNICAL REPORT DISTRIBUTION LIST, 051B

	<u>No. Copies</u>		<u>No. Copies</u>
Professor K. Wilson Department of Chemistry, B-014 University of California, San Diego La Jolla, California 92093	1	Dr. B. Vonnegut State University of New York Earth Sciences Building 1400 Washington Avenue Albany, New York 12203	1
Professor C. A. Annett Department of Chemistry Purdue University West Lafayette, Indiana 47907	1	Dr. Hank Loos Laguna Research Laboratory 21421 Stans Lane Laguna Beach, California 92651	1
Professor P. Meijer Department of Physics Catholic University of America Washington, D.C. 20064	1	Dr. John Latham University of Manchester Institute of Science & Technology P.O. Box 88 Manchester, England M601QD	1
Dr. S. Greer Chemistry Department University of Maryland College Park, Maryland 20742	1		
Professor P. Delahay New York University 100 Washington Square East New York, New York 10003	1		
Dr. T. Ashworth Department of Physics South Dakota School of Mines & Technology Rapid City, South Dakota 57701	1		
Dr. G. Gross New Mexico Institute of Mining & Technology Socorro, New Mexico 87801	1		
Dr. J. Kassner Space Science Research Center University of Missouri - Rolla Rolla, Missouri 65401	1		
Dr. J. Telford University of Nevada System Desert Research Institute Lab of Atmospheric Physics Reno, Nevada 89507	1		

