

AD-A111 966

FOREIGN TECHNOLOGY DIV WRIGHT-PATTERSON AFB OH
'RYTERSKI' AND 'RYMANOWSKI' WINDS, (U)

F/G 4/2

UNCLASSIFIED


FEB 82 J LEWINSKA
FTD-ID(RS)T-1715-81

NL

1-1
2-1



END
DATE
FEB 82
DTIC




MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A111966

DTIC FILE COPY

DTIC-1000000-1000000


FOREIGN TECHNOLOGY DIVISION



"RYTERSKI" AND "RYLANOWSKI" WINDS

by

Janina Lewinska



DTIC does not
production

DTIC
ELLITE
MAR 12 1982
A

Approved for public release;
distribution unlimited.

82 03 11 162

DISCLAIMER NOTICE

**THIS DOCUMENT IS BEST QUALITY
PRACTICABLE. THE COPY FURNISHED
TO DTIC CONTAINED A SIGNIFICANT
NUMBER OF PAGES WHICH DO NOT
REPRODUCE LEGIBLY.**

EDITED TRANSLATION

FTD-1D(RS)T-1715-81

12 February 1982

MICROFILME NR: FT 82-C-000194

"RYTERSKI" AND "RYMANOWSKI" WINDS

By: Janina Lewinska

English pages: 11

Source: Przegląd Geofizyczny, Vol. 3(11), Nr. 1, 1958, pp. 17-26

Country of origin: Poland

Translated by: SCITKAN

F33657-81-D-0263

Requester: USAF/ETAC/MAC

Approved for public release; distribution unlimited.



A 23 CP

THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:

TRANSLATION DIVISION
FOREIGN TECHNOLOGY DIVISION
WP-AFB, OHIO.

FTD-1D(RS)T-1715-81

Date 12 Feb 19 82

"RYTERSKI" AND "RYMANOWSKI" WINDS

Jadwiga Lewinska

Winds, as one of the weather-forming factors, undoubtedly exert a profound effect on the formation of weather in valleys. The present work deals with description of winds blowing in the valleys of the Poprad River on the Stary Sacz, the Leluchow section and the Tabor River. Periodic winds appear along the axis of these valleys which is close to the S-N direction. Because of their force, frequency and effect on local living conditions, these winds are known to the inhabitants of these valleys under the local names of ryterski and rymanowski. For the study of these winds, I utilized results of meteorological observations obtained at the stations in Stary Sacz and Rymanow-Zdroj and for comparative purposes also at the Nowy Sacz and Kasprowy Wierch stations in 1954-55. Because of the short time from the data input, the obtained conclusions in this work should be considered as of orientation type only, particularly those of a quantitative character.

The first of the mentioned valleys (Poprad River) on the S-N axis in the Piwniczna--Stary Sacz section is located in Beskid Sadecki and cuts the massive Carpathians in the area of Leluchow. The second valley (Tabor River) also lies on an axis close to S-N, and is located in Beskid Niski. The sizes of these two valleys are not really comparable (the length of Poprad Valley in Polish territory is about 55 km, and the Tabor Valley merely 27 km); nevertheless, anemometric conditions reigning in these valleys are of similar character. When treating the material, I observed the following rules: winds from sectors SE to SW with velocities 5 m and above were considered, and in more detailed treatment, winds with velocities 10 m/sec and above were considered if they were observed at least once during the day. By following this rule, I could use the criterium adopted by Gellert [1]. Moreover, when treating materials, I took into consideration the air temperature, humidity at 13:00 hr, and atmospheric pressure. At the same time, I also paid attention to the synoptic

situation on the basis of the map of Europe at 00 hr EST. A situation which could cause air advection from the south (with particular interest in a situation which could lead to strong winds from the south direction). Preliminary analysis of the material led us to assume that the ryterski and ryanowski winds are winds of foehn type (known in Tatras as halny).

The values of air temperature and atmospheric pressure (monthly averages) and their average daily values the highest (or lowest) during the wind duration, and differences between these values, are presented in Tables 1 and 2. In all the observed cases, one can notice a very distinct temperature rise as well as fall of pressure, as compared to air covering the valley in the period preceding the southern wind. We have not included the data on the loss of humidity because of considerable gaps in observational material; nevertheless, even here one can see considerable losses of humidity (reaching 35%). Already on the basis of these data we can assume that we are dealing with falling winds, causing clear thermo-hygro-baric effects both in the Poprad Valley and in the Tabor Valley.

The frequency of appearance of winds from the southern quadrant at the above mentioned stations is shown graphically in Figures 1 and 2 (the fact that in specific cases we included in the graphs winds with velocity on orographic form, hence in the narrow valley these velocities will be usually much higher than, for instance, in the broad and flat Sadecki Valley). In agreement with generally accepted views, we assumed that winds of foehn type appear here when there are centers of low pressure north of the Carpathian chain and centers of high pressure south of it. Examples of such weather conditions are provided by situations shown on synoptic maps on May 18, 1955 (Figure 3) and on January 31, 1955 (Figure 4). In these cases, centers of low pressure were moving from the Atlantic to the area of Great Britain and the North Sea, and centers of high pressure--from the Mediterranean Sea to the Black Sea area, Ukraine and Asia Minor. Such situations lead to strong winds of foehn type in the Carpathians. The isobars on synoptic maps run from SW to NE. Low baric centers moving along the Atlantic axis from Great Britain cause a drop of pressure over the




Figure 1

Key: 1--winds from S quadrant with velocities < 5 m/sec
 2--winds from S quadrant with velocities > 5 m/sec
 3--days with synoptic situations typical for halny winds

Carpathians which, in turn, results in southern gradient winds.

Comparing baric situations with results of meteorological observations at the above mentioned stations, we arrive at the following conclusions (compare Figures 1 and 2):

- 1) winds from the S quadrant appear simultaneously at meteorological stations Kaszowy Wierch, Nowy Sącz, Rymanów and Starey Sącz, hence at stations located in different orographic conditions;
- 2) this appearance of winds from the S quadrant simultaneously in situations described above takes place in 70-90% of cases;
- 3) the duration of the above mentioned winds is from 2-7 days;
- 4) the highest frequency of these winds occurs in the autumn-winter period and the lowest in summer.




Figure 2

Key: 1--winds from S quadrant with velocities < 5 m/sec
 2--winds from S quadrant with velocities > 5 m/sec
 3--days with synoptic situations typical for balny winds

The above conclusions justify us to maintain that the ryterski and rymanowski winds do not have the character of local winds caused by thermal gradients between mountain peaks and depths of valleys, but in the majority of cases arise as the result of atmospheric processes resulting from the general baric situation.

Tables 3, 4 and 5 show frequencies of winds with foehn character, and periods of their duration for Kasprowy Wierch, Rymanow and Stary Sącz stations. It follows from these tables that the average number of days with winds of foehn type for Kasprowy Wierch is 72 days, and for Stary Sącz, 73 days ($v > 10$ m/sec). If, following H. Fikler, we accept the average frequency for the Alps of the appearance of foehn winds to be 30-40 days per year, and the maximal frequency--60 days; for Tatra, 45 days and for Tatras on the average 80 days, according to W. Kilian and M. Orlica, then the above given values for ryterski wind obtained on the basis of observations in two years do not

TABLE 1. Differences between the monthly (I-IV) and daily (V-XII) differences in the air temperature according to the occurrence of the wind (for 1951-55).

GPR. year.			I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII
Nowy Sącz	1951	t_m	-9.2	-9.0	4.0	1.6	12.1	17.2	16.3	17.8	.	8.9	3.0	2.6
		$t_{max} R$	2.8	1.2	11.3	11.7	16.5	21.3	20.9	21.5	.	15.6	10.9	8.1
		Δ	12.0	13.1	7.3	7.1	4.1	4.1	4.1	6.7	.	8.7	7.9	5.3
	1955	t_m	-3.1	-2.1	-0.1	4.8	11.1	11.6	17.1	.	13.8	8.9	3.2	1.1
		$t_{max} R$	4.4	6.8	10.7	12.0	18.2	21.5	19.2	.	19.3	11.3	9.3	1.8
		Δ	7.5	9.2	11.1	7.2	6.8	6.9	1.8	.	5.5	5.1	6.1	3.1
Rymanów	1951	t_m	-10.1	-7.9	3.2	3.6	11.5	16.9	15.6	16.3	13.5	7.3	2.6	1.5
		$t_{max} R$	1.2	2.2	8.7	11.6	11.1	21.2	17.5	22.7	19.2	13.8	10.8	9.1
		Δ	9.2	10.1	5.1	8.0	2.6	4.5	1.9	6.1	5.7	6.5	8.1	7.6
	1955	t_m	-3.2	-2.6	-0.5	3.6	10.5	14.2	17.1	.	12.6	8.3	2.5	0.2
		$t_{max} R$	3.3	3.8	12.0	9.9	19.9	21.1	24.1	.	17.5	11.2	10.5	3.6
		Δ	6.5	6.1	12.5	6.3	9.1	6.9	6.3	.	4.9	5.9	7.8	3.1

t_m -- average monthly air temperature; $t_{max} R$ -- average daily temperature the highest during the occurrence of ryterski or rymanowski wind; $\Delta = t_{max} R - t_m$

TABLE 2. Differences between the average monthly atmospheric pressure and the average daily lowest pressure during the occurrence of ryterski wind (for Nowy Sącz)

yr.		I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII
1951	P_m	79.9	81.5	78.2	80.1	78.3	80.1	76.6	78.1	81.6	82.6	82.9	77.3
	$P_{min} R$	80.1	79.8	83.3	71.6	69.0	71.9	73.1	71.6	79.5	75.3	71.9	59.5
	Δ	9.8	10.7	19.9	8.8	9.3	8.5	3.5	6.3	2.1	7.3	11.0	17.8
1955	P_m	78.7	70.3	19.5	81.5	80.1	80.2	78.3	.	82.7	80.7	83.7	75.3
	$P_{min} R$	81.6	62.6	10.9	78.7	71.1	69.7	73.2	.	65.1	70.8	80.2	69.9
	Δ	23.1	8.5	8.6	7.8	6.9	13.5	5.1	.	17.6	10.1	3.5	11.1

P_m -- average monthly atmospheric pressure; $P_{min} R$ -- average daily atmospheric pressure in the period of occurrence of ryterski wind; $\Delta = P_m - P_{min} R$

TABLE 3. Frequency of occurrence of ryterski wind (Stary Sącz), ryterski wind compared with halny wind (Kasprowy Wierch) on the basis of observations in the years 1954-55, and of halny wind (Kasprowy Wierch) in the years 1938-51 [Pctny%]

		I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII	Rok
$\frac{v}{m/sec}$	Ryt.	10.5	11.0	9.0	4.5	7.0	6.5	3.5	4.0	9.0	11.5	10.5	11.5	102.5
	Rym.	10.5	9.0	5.5	7.0	4.5	6.0	6.5	3.0	5.0	9.0	12.0	12.0	90.0
	Halny	15.0	15.0	16.5	9.0	13.0	12.0	11.0	7.0	12.0	15.0	15.0	14.5	135.0
$\frac{v}{m/sec}$	Ryt.	10.5	10.5	7.5	2.0	4.5	2.5	0.0	2.0	4.5	7.5	6.5	12.5	73.0
	Rym.	2.5	2.0	0.0	0.0	1.0	0.5	0.0	0.0	1.0	0.5	2.5	3.5	13.5
	Halny	7.5	9.0	9.5	2.5	4.5	5.0	3.0	1.0	6.5	7.5	8.0	8.5	72.5
	Halny*	7.6	7.9	7.1	8.0	5.5	3.9	2.9	5.0	6.1	7.2	10.1	9.2	81.0

TABLE 4. Average time of duration (in days) of ryterski winds with velocities $v \geq 5$ m/sec, on the basis of the results of observations in the years 1954-55

I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII
5.0	4.5	3.5	2.0	2.5	2.5	1.0	1.5	4.5	4.5	5.5	4.0

TABLE 5. Maximal periods of duration (in days) of ryterski winds with velocities $v \geq 10$ m/sec on the basis of the results of observations at Stary Sącz (1954-55), compared with the periods of duration of halny winds at Kasprowy Wierch (1938-51).

	I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII
Stary Sącz	5	5	1	2	3	3	1	2	5	5	6	4
Kasprowy Wierch	3	11	5	9	5	6	2	1	6	9	9	7



Figure 3. Synoptic map at 0.00 hrs on May 18, 1955
W = wysoki = high; N = niski = low

deviate from average values and should be considered as reliable. According to S. Leszczycki, winds of foehn type occur on the average 24 days per year at the Tatras foothills in Zakopane. Hence, it seems right to point out large velocities of these winds in the Poprad Valley and that these velocities are close to those observed in the Tatras. We have to assume that in the Rytro locality where the wind reaches maximum velocity, the number of days of southern wind velocity of the order of 10 m/sec and above is higher than in the Tatras. The analysis of wind directions for Rymanow and Stary Sącz (Figure 5) shows their large similarity.

The frequency of appearance of rymanowski winds with velocities 5 m/sec and above agrees with analogical data for the remaining stations, but the frequency of winds with large velocities of the order of 10 m/sec and above is only several days per year and in this connection does not cause any strong weather contrasts.




Figure 4. Synoptic map at 0.00 hrs on January 31, 1955

We checked the above obtained results by the field examination in Poprad Valley at the section of Nowy Sącz, Muszyna (of length about 40 km), whereas in the Taber Valley we limited ourselves to only a local inspection in the Rymanów-Zdrój area. The results of this checking are as follows: Strong periodic southern winds are observed in the Poprad Valley at the section below Piwniczna to Stary Sącz, particularly in the autumn-winter season, and less frequently in summer. These winds cause the warming up. Their strength frequently exceeds 10 m/sec (and not infrequently even 20 m/sec). This strength increases considerably in the vicinity of Rytro town where the width of the valley does not exceed 100 m. In the same locality, according to descriptions of local inhabitants, there often occur winds of destructive power which causes tearing of roofs, snapping of trees, etc. On the basis of analysis of damages caused by the wind, and of descriptively local people, one can assume that the power of that wind reaches and sometimes exceeds 35 m/sec. After the period of occurrence




Figure 5.

of the wind, usually not a period of calm, which seldomly occurs. In the middle of winter, the Carpat Valley turns to the east and then, due to its direction in small sections because of wind shift of the Carpat River, hence the absence of the ryterski-type wind along it. The mesometeoric conditions in the Tatra River Valley take rising shape although because of relatively small and "shaded" valleys, we do not observe strong winds in this area.

Summarizing the above results, we can state that in valleys perpendicular to the main axis of mountain ridges one periodically observes winds blowing along the axis of the valley. These winds, called ryterski and rymonowski, have all the features of falling winds. They are formed in synoptic situations when centers of low pressure lie NW from the Carpathian chain over the Atlantic and North Sea (particularly over Great Britain) and move toward the east. Then one can observe centers of high pressure SE from the Carpathians, characteristically localized over the Black Sea, the Bosphorus and Asia Minor. According to the criterion proposed by W. Milata, the ryterski and rymonowski winds should be considered in the majority of cases as winds of halny-lowland type. The frequency of appearance of ryterski and rymonowski winds is of the same order as the frequency of halny wind in Tatras. In the yearly course one can see the maximum of frequency of the appearance of wind in the autumn-winter period and the minimum in summer.

Manuscript received by the editor on January 29, 1957.

REFERENCES

1. Bartolicki, G. The balny wind in Tatras. Czas. Geogr., 2 (1944).
2. von Ficker, L. and von Radtke, R. Föhn und Föhnwindung. Der geographische Stand der Frage. Leipzig 1943.
3. Guenther, Paul. Die Lora der Adria.
4. Chumakov, S. F. Principles of synoptic meteorology [in Russian] Leningrad, 1943.
5. Kosinska-Larinińska, S. Föhnerscheinungen im Tatragebirge, Met. Zeitschr., 46 (1932).
6. Kosinska-Larinińska, S. The balny wind on July 31, 1931.
7. Leczynski, S. Region of Föhnwind. Biul. Kom. Mgi Pop. Turystyki 2 (1935).
8. Lewinski, J. Distribution of wind directions in Polish Tatras and at Föhnwind (manuscript).
9. Migonies, W. Z. Consideration of anemological problems in Tatras. Przegl. Geogr., 10 (1936).
10. Milata, W. The balny winds. Wierchy. 20 (1950-51)
11. Milata, W. The balny winds in the Carpathians. Kalendarz IKC 10 (1937).
12. Milata, W. The balny winds in the Sudeten and Carpathians. Taternik, 29 (1947).
13. Orlicki, M. Anemometric conditions at Tatra peaks. Wiad. Sluszy Hydrol. i Meteor., III, 2, 4 (1954).
14. Wigilew, B. A few words about the balny winds. Pam. Tow. Tatr., 33 (1912).
15. Wigilew, B. Studies of the climate at Zakopane. Spraw. Kom. Fizjogr. AU, 51 (1917).

**DATA
FILM**