

MICROCOPY RESOLUTION TEST CHART
NBS 1963-A

ADA 112122

12

AD

Report DELNV-TR-0024

QUANTUM MECHANICS APPLIED TO DYE LASERS

by
Carol K. Pearce

September 1981

Approved for public release; distribution unlimited.

MAR 18 1982

E

DTIC FILE COPY

U.S. ARMY ELECTRONICS R&D COMMAND
NIGHT VISION & ELECTRO-OPTICS LABORATORY
FT. BELVOIR, VIRGINIA 22060




104

**Destroy this report when it is no longer needed.
Do not return it to the originator.**

**The citation in this report of trade names of
commercially available products does not constitute
official endorsement or approval of the use of such
products.**

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)



UNCLASSIFIED

ii

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

CONTENTS

Section	Title	Page
I	INTRODUCTION AND BACKGROUND	1
II	APPLICATION OF THE LAMB LASER THEORY TO DYE LASERS	6
III	PROPAGATION OF LIGHT PULSES	10
IV	SPECTRAL CROSS-RELAXATION AND COLLISIONAL DEPHASING EFFECTS	12
V	A QUANTUM-MECHANICAL DYE LASER THEORY INVOLVING EXACT TREATMENT OF TRIPLET STATES	16
VI	CONCLUSIONS	24



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Dist _____	
Avail _____	
Dist _____	
A	

QUANTUM MECHANICS APPLIED TO DYE LASERS

I. INTRODUCTION AND BACKGROUND

Successful lasing in an organic dye was not reported until 1966 although the first successful laser was developed in 1961. Use of dyes as saturable absorber laser Q-switches preceded the laser dye success and continues to the present. NV&EOL has successfully applied the incorporation of a dye Q-switch in low-energy laser rangefinding systems. High-energy applications were explored. A concerted effort was made to extend the long wavelength applicability of the dye Q-switch through the creation of new dyes. A close relationship exists between dye lasers and dye Q-switches. One of the first instances of lasing in a dye occurred inadvertently during a search for new dye Q-switches. Developments in the more widely studied phenomenon of dye lasers support our mission in dye Q-switches and have provided a stimulus for this report.

Both pulsed and continuous wave (CW) dye lasers are operational in configurations much like those used for other lasers (except for modifications required to offset losses due to the triplet state). Numerous organic dyes have been found to lase; the cyanine, coumarin, xanthene, and oxazine classes are of major importance.¹ The tunability inherent in dye lasers is probably their most useful asset. Depending upon the choice of dye, tunable radiation can be generated from the near ultraviolet to the visible and near infrared regions of light. Linewidths as narrow as 10^{-4} nanometers are available, as are pulse widths of 0.01 nanosecond duration.² When saturable absorber mode locking dyes are incorporated in the laser cavity, picosecond pulsewidths become attainable. Both CW and pulsed (0.1 to over 300 p/s) operation can be obtained commercially, with peak power up to 2×10^6 watts being cited.² Energies between several microjoules and up to over 10 joules per pulse have been reported.³ The potential of the dye laser in integrated optical elements has been discussed with possible applications in communications and data transmission.³ The precise tunability of dye lasers mandate their use in spectroscopic studies and in such practical applications as LIDAR. LIDAR (Light Detection and Ranging) enables one to detect and measure trace atmospheric pollutants by a radar type of fluorescence technique.³ Dye lasers have been investigated for optical warning/detections systems and as human optical countermeasure sources.^{4 5} More extensive application of

- ¹ K. H. Drexhage, "Structure and Properties of Laser Dyes," in "Dye Lasers," Chap. 4, ed. by F. P. Schaefer, Springer-Verlag, New York (1973).
- ² "Laser Focus Buyer's Guide," pp 143-144, Advanced Technology Publications, Inc., Newton, MA (1979).
- ³ T. W. Hänsch, "Applications of Dye Lasers," in "Dye Lasers," Chap. 5, ed. by F. P. Schaefer, Springer-Verlag, New York (1973).
- ⁴ L. Gampel, S. Guch, and H. Wang, "Dye Laser Module," U.S. Army Research and Development Report, ECOM 76-1972-1 (Feb 77).
- ⁵ D. H. Sliney and T. L. Lyon, "Evaluation of a Phase-R Dye Laser in Phase II of an ECOM Countermeasure Source Evaluation," AD-917 016L, Defense Documentation Center, Alexandria, VA (December 1973).

the dye laser can be expected in all of these areas, particularly as the state-of-the-art advances. Some of the improvements that should develop are: dyes and host materials designed specifically for dye laser systems, vapor dye lasers operable in the visible and near infrared (and pumped by electrical discharge or electron beam sources, and modifications in the laser module itself.

As dye laser technology requires increasingly unique responses, rather than merely borrowing from the existing technologies of dyes, photochemistry, and lasers, guidance will be needed from theoretical treatments. Much of the theoretical work done on dye lasers is based upon energy balance and rate studies. Quantum mechanical treatments of dye laser systems are, at present, much less numerous than for other laser systems. Among the few quantum mechanical dye laser studies are the semiclassical theories of Eberly, Chin and Zardecki, and Vahey and Yariv.^{6 7 8} A fully quantum-mechanical study was reported recently by Schaefer and Willis.⁹ The intent of this report is to discuss dye lasers as a case of light interacting with matter, the case being treated quantum mechanically. How quantum mechanics is applied, and the kinds of problems solved, will be of major interest.

Aspects of dye lasers covered in these quantum mechanical treatments to be discussed more fully below, include:

- a. Application of the Lamb laser theory to dye lasers.⁶
- b. The propagation and superposition of light pulses in a dye-solvent medium.
- c. The absorption of light in organic dyes (both in dye lasers and as saturable absorbers) as it is affected by spectral cross-relaxation and collisional dephasing.⁸
- d. Development of a completely quantum mechanical theory of dye lasers, including an exact calculation for the dye triplet state.⁹

Basic to all of these studies is the fact that organic dye molecules, upon absorption of light quanta, undergo an optical transition from a ground (singlet) state energy level to a higher excited singlet energy level. Decay to the ground state results in fluorescent emission, comparable to the laser emission when the dye is in a laser cavity. Intersystem crossing (where k_{ST} is the intersystem crossing rate) can easily occur, however, so that the excited singlet may decay instead to a triplet level (Figure 1). Thus the efficiency of the (singlet) emission process is reduced.

⁶ J. H. Eberly, "Toward a Theory of the Dye Laser," "Atomic and Molecular Optics," Proc. Rochester Symp., ed. by J. H. Eberly, Univ of Rochester (March 1971); also US Gov't Publication AD725218 (Defense Documentation Ctr), Alexandria, VA.

⁷ A. L. Chin and A. Zardecki, *Phys. Rev. A*, *13*, 1528 (1976).

⁸ D. W. Vahey and A. Yariv, *Phys. Rev. A*, *10* (6) 1578 (1974).

⁹ R. B. Schaefer and C. R. Willis, *Phys. Rev. A*, *13* (5) 1874 (1976).




Figure 1. Typical Jablonski energy level diagram for dye molecules.

The distribution of molecules upon interaction with light will be among a minimum of four energy levels, so that the organic dye laser must be considered as a four-level system at least, in contrast to the two-level laser of Lamb's theory.⁶ Unlike other laser media, furthermore, organic dyes are characterized by broad absorption and emission bands. These broad bands afford the tunability for which dye lasers are known. They indicate also that many vibrational/rotational levels are populated for a particular electronic transition.

In the discussions to follow, the use of density matrices will be found to be most useful in formulating the equations of motion for dye molecules moving through an electric field (the field due to the electric component of light waves).^{10 11} The density matrix is one whose elements are bilinear combinations of the probability amplitudes, C_A and C_B (if $\psi(r, t)$ is a known wave function):¹⁰

$$\psi(r, t) = C_A(t)\psi_A(r) + C_B(t)\psi_B(r), \quad (1)$$

where $\psi(r)$ is the stationary, or time-independent, wave function for molecules A or B.

Sargent, Scully, and Lamb (hereafter SS&L) have discussed the above "pure case" as well as the more generally useful "mixed case" when the state vector (in Dirac notation) $|\psi(t)\rangle$ is unknown.¹⁰ The wave function $\psi(r, t)$ is an expansion coefficient of the state vector.¹⁰ Thus, ρ is the general density operator

$$\rho = \sum_j P_j |\psi_j\rangle \langle \psi_j|. \quad (2)$$

P_j is a probability factor for systems which have a state vector $|\psi(t)\rangle$ and where two-level state vectors are:

$$|\psi_j(t)\rangle = C_{aj}(t)|a\rangle + C_{bj}(t)|b\rangle, \quad (3)$$

and so

$$\rho = \sum_j P_j \begin{pmatrix} |C_{aj}|^2 & C_{aj}C_{bj}^* \\ C_{bj}C_{aj}^* & |C_{bj}|^2 \end{pmatrix}. \quad (4)$$

⁶ J. H. Eberly, "Toward a Theory of the Dye Laser," "Atomic and Molecular Optics," Proc. Rochester Symp., ed. by J. H. Eberly, Univ of Rochester (March 1971); also US Gov't Publication AD725218 (Defense Documentation Ctr), Alexandria, VA.

¹⁰ M. Sargent III, M. O. Scully, and W. E. Lamb, Jr., "Laser Physics," Chaps 7, 14, and 16, Addison-Wesley, New York (1974).

¹¹ W. E. Lamb, Jr., Phys. Rev., 134 A1429 (1964).

The equation of motion, utilizing Schrodinger's time-dependent equation

$$\hat{H} \psi(r, t) = i\hbar \frac{\partial \psi}{\partial t}(r, t) \quad (5)$$

is:

$$\dot{\rho} = -\frac{i}{\hbar} [\hat{H}, \rho]. \quad (6)$$

SS&L describe the use of the density matrix equations of motion in various aspects of laser quantum mechanical problems.¹⁰

SS&L showed that these matrix equations of motion could be used to derive equations for the Bloch model. In the Bloch model, a two-level system of magnetic dipoles of spin $\pm \frac{1}{2}$, is used to represent another two-level system (as a two-level laser system). This approach was used by Vahey and Yariv⁸ for the singlet-triplet dye system. The Bloch equations are based upon the equations of motion through a magnetic field and include two relaxation terms of equations.^{9 10 12}

Reservoir theory can be used with the density operator, ρ , in laser calculations. The quantum mechanical treatment of dye lasers by Schaefer and Willis⁹ combines reservoir theory with density matrices. As described by SS&L, the *reservoir approach facilitates the division of a system into portions of primary (A) and secondary (B) interest. The "secondary" part becomes the reservoir, which can be eliminated by use of a "reduced density operator,"* $\rho_{AB}(t)$.¹⁰

The "rotating wave approximation" has been used below.^{6 7 9} This approximation allows one to eliminate terms where $(\omega + \nu)$ is in the denominator, or $e^{i(\omega + \nu)t}$ is present.^{6 10} Since ω , the "matter" wave frequency in

$$\psi(r, t) = C_a \psi_a(r) e^{-i\omega_a t} + C_b \psi_b(r) e^{-i\omega_b t}. \quad (7)$$

⁶ J. H. Eberly, "Toward a Theory of the Dye Laser," "Atomic and Molecular Optics," Proc. Rochester Symp., ed. by J. H. Eberly, Univ of Rochester (March 1971); also US Gov't Publication AD725218 (Defense Documentation Ctr), Alexandria, VA.

⁷ S. L. Chin and A. Zardecki, Phys. Rev. A, 13, 1528 (1976).

⁸ D. W. Vahey and A. Yariv, Phys. Rev. A, 10 (6) 1578 (1974).

⁹ R. B. Schaefer and C. R. Willis, Phys. Rev. A, 13 (5) 1874 (1976).

¹⁰ M. Sargent III, M. O. Scully, and W. E. Lamb, Jr., "Laser Physics," Chaps 7, 14, and 16, Addison-Wesley, New York (1974).

¹² A. Yariv, "Quantum Electronics," Chap. 8, John Wiley and Sons, Inc., New York (1967).

and ν , the "field" frequency of

$$E(t) \propto \cos(\nu t), \quad (8)$$

are equal under near resonance conditions, and $\omega \gg 1$. ($\omega + \nu$) is relatively large. For $e^{i(\omega + \nu)t}$, rapid oscillation occurs, so that such terms average zero.⁶

In each of the discussions below, the symbolism as used by the authors, respectively, has been maintained in the section discussing their work. Kauzmann has discussed the fundamentals of the quantum mechanics of light interacting with matter and time-dependent processes.¹³

II. APPLICATION OF THE LAMB LASER THEORY TO DYE LASERS

Eberly sought to derive a dye laser theory from the jointly interacting laser electric field and the polarization density of the dye.⁶ He applied the Lamb laser theory to organic dye lasers, noting two important modifications would be necessary.¹¹ The dye laser must be treated as a four-level system at least, not a two-level one, and Lamb's use of the extreme Doppler limit would be inappropriate since dyes are conventionally used in solution. (One should note, however, that more recently lasing has been achieved with organic dye vapors.)^{14 15}

The polarization density $P(t, z)$ was derived, based upon the Lamb theory, for dye molecules in the presence of the perturbing field $E(t, z)$. For a single mode of oscillation of an electromagnetic (light) wave in the laser cavity, given by $U(z)$, the natural frequency was $\Omega = \nu + \dot{\phi}$ (where $\phi(t)$ is the function describing the slow variation of ϕ with time, and ν is the frequency of the light wave.)

Further, assumptions were made that discrete modes were present in the laser cavity, that the vector nature of the field could be ignored (as well as its polarization), and that all parameters are functions of time, t , and cavity axial position, z , only. For the perturbing electric field,

$$E(t, z) = U(z) \epsilon(t) \cos(\nu t + \phi(t)).$$

⁶ J. H. Eberly, "Toward a Theory of the Dye Laser," "Atomic and Molecular Optics," Proc. Rochester Symp., ed. by J. H. Eberly, Univ of Rochester (March 1971); also US Gov't Publication AD725218 (Defense Documentation Ctr), Alexandria, VA.

¹¹ W. E. Lamb, Jr., *Phys. Rev.*, **134** A1429 (1964).

¹³ W. Kauzmann, "Quantum Chemistry," Part V, Academic Press, New York (1957).

¹⁴ P. W. Smith, P. F. Liao, et al., *Appl. Phys. Ltrs.*, **25** 144 (1974).

¹⁵ B. Steyer and F. P. Schaefer, *Opt. Comm.*, **10** 219 (1974).

Here ϵ is the amplitude and is assumed to vary slowly with time. Only the first derivatives of $\epsilon(t)$ and $\phi(t)$ were used in Eberly's derivation. Since both $\phi(t)$ and $\epsilon(t)$ have been assumed to exhibit little change with time, this must be true for $E(t, z)$ also, so that the treatment is semiclassical.

The time-dependent projection of the polarization density $P(t, z)$ can be written as a function composed of oscillations in and out of phase with $\cos(\nu t + \phi(t))$ of $E(t, z)$:

$$P(t) = \int_{\text{cavity}} dz U(z) P(t, z) \quad (9)$$

$$= C(t) \cos(\nu t + \phi) + S(t) \sin(\nu t + \phi).$$

Using Maxwell's wave equation modified with an arbitrary conductivity constant σ as a cavity loss correction factor, coupled differential equations for $\epsilon(t)$ and $\phi(t)$ were obtained (where η is an arbitrary constant undefined further by Eberly):

$$(\nu + \dot{\phi} - \Omega) \epsilon(t) = -\frac{1}{2} (\nu/\eta^2) C(t) \quad (10)$$

$$\dot{\epsilon}(t) + \frac{1}{2} (\sigma/\eta^2) \epsilon(t) = -\frac{1}{2} (\nu/\eta^2) S(t). \quad (11)$$

These two equations, along with that for the polarization density, $P(t, z)$, are those equations subject to self-consistent (field) solution in Eberly's theory.

Eberly determined the polarization density according to Lamb's laser theory but adapted it to the unique characteristics of a dye system. The optical transitions involved in a lasing dye are shown in Figure 1. The assumptions basic to Eberly's treatment are:

a. None but optical transitions contribute to the polarization density, and

b.
$$P^2(t, z) = \sum_{i=1}^n p_i^2(t_i, z) \quad (12)$$

where $n = 3$ in the present case and p_i is the partial polarization associated with the i 'th optical transition.

Addressing oneself to one of the transitions involved in the above summation, where $n = (\text{level B} - \text{level a})$, and specifying $\lambda_B(t) = \text{rate-time}^{-1}\text{-volume}^{-1}$ for the excitation of molecules into level/state "B," $P_i(t)$ becomes $P_B(t)$ or

$$P_B(t) = \int_{-\infty}^t dt_0 \lambda_B(t_0) \bar{\mu}_B(t, t_0). \quad (13)$$

For the dye molecules undergoing the transition B-a, there will be more than one final state possible, as noted above, so that:

$$\bar{\mu}(t, t_0) = \sum_a (\mu_{Ba} \rho_{aB}^B + \rho_{Ba}^B \mu_{aB}). \quad (14)$$

where μ_{Ba} and μ_{aB} are the dipole moments corresponding to the different final states, $\bar{\mu}_B(t, t_0)$ is the average or expectation value of μ_{molecule} in state B at time t_B and ρ^B is the density matrix of that same molecule.^{6 11}

$$\rho^B = \begin{pmatrix} |B|^2 & Ba^* \\ B^*a & |a|^2 \end{pmatrix} = \begin{pmatrix} \rho_{BB}^B & \rho_{Ba}^B \\ \rho_{aB}^B & \rho_{aa}^B \end{pmatrix} \quad (15)$$

From the equation of motion for this density matrix,

$$i \hbar \dot{\rho} = [H, \rho] + \frac{\hbar}{2} (\gamma \rho + \rho \gamma), \quad (16)$$

γ being the diagonal matrix:¹⁰

$$\gamma = \begin{pmatrix} \gamma_B & 0 \\ 0 & \gamma_a \end{pmatrix} \quad (17)$$

where γ_B and γ_a are the radiative decay constants for states B and a, and following the development of Eberly and Lamb, and invoking the rotating wave approximation, the polarization density for the B-a optical transition will be:^{6 11}

$$P_B(t, z) = \frac{\lambda_B(t)}{2\gamma_D} \sum_a \left(-\frac{i}{\hbar} \right) \frac{\mu_{Ba} \mu_{aB}}{\gamma_{Ba} + i(\omega_{Ba} - \nu)} U(z) \epsilon(t) e^{-i(\nu t + \phi)} \quad (18)$$

+ complex conjugate.

⁶ J. H. Eberly, "Toward a Theory of the Dye Laser," "Atomic and Molecular Optics," Proc. Rochester Symp., ed. by J. H. Eberly, Univ of Rochester (March 1971); also US Gov't Publication AD725218 (Defense Documentation Ctr), Alexandria, VA.

¹⁰ M. Sargent III, M. O. Scully, and W. E. Lamb, Jr., "Laser Physics," Chaps 7, 14, and 16, Addison-Wesley, New York (1974).

¹¹ W. E. Lamb, Jr., Phys. Rev., 134 A1429 (1964).

Expressions for $C_B(t)$ and $S_B(t)$ were obtained by Eberly from the P(t) equation (9):

$$C_B(t) = \frac{\eta^2}{\nu} \left(\frac{\lambda_B}{\gamma_B} \right) \beta \epsilon(t) \quad (19)$$

$$S_B(t) = \frac{\eta^2}{\nu} \left(\frac{\lambda_B}{\gamma_B} \right) \alpha \epsilon(t), \quad (20)$$

where

$$\alpha + i\beta = \int_{\pi} \frac{M_{Ba} D_a(\omega_{Ba}) d\omega_{Ba}}{\gamma_{Ba} + i(\omega_{Ba} - \nu)} \quad (21)$$

here, $M = 1$, since restricted to single mode, and $D_a(\omega_{Ba}) =$ energy density of state a with energy $\hbar\omega_{Ba}$.

Eberly obtained one last significant result from the theory outlined above, for the zero order. Using the above expressions for $C_B(t)$ and $S_B(t)$, the photon gain from the $B \leftarrow a$ transitions, $G_{B \leftarrow a}$ is:

$$G_{B \leftarrow a} = (2\dot{\epsilon}/\epsilon)_{B \leftarrow a} = \frac{\nu}{\eta^2} S_B(t)/\epsilon(t) \quad (22)$$

$$= \left(\frac{\lambda_B}{\gamma_B} \right) \frac{\gamma_{Ba}}{\pi} \int \frac{M_{Ba} D_a(\omega) d\omega}{[(\omega - \nu)^2 + \gamma_{Ba}^2]} \quad (23)$$

This expression was similar to one Eberly had obtained earlier using rate equations and applied to problems associated with the threshold for organic dye lasers.^{6 16}

⁶ J. H. Eberly, "Toward a Theory of the Dye Laser," "Atomic and Molecular Optics," Proc. Rochester Symp., ed. by J. H. Eberly, Univ of Rochester (March 1971); also US Gov't Publication AD725218 (Defense Documentation Ctr), Alexandria, VA.

¹⁶ O. G. Peterson, J. P. Webb, W. C. McColgin, and J. H. Eberly, J. Appl. Phys., 42 (5) 1917 (1971).

III. PROPAGATION OF LIGHT PULSES

The semiclassical self-consistent approach was also used by Chin and Zardecki to describe the propagation of light pulse through a dye medium and also the superposition of short light pulses in that medium.⁷ A multi-level model for a dye molecule in solution was presented. Triplet states are ignored since the duration of the pulse trains is too short for intersystem crossing to occur. Essentially, the "supermolecule" (the dye molecule in its solvent cage) when perturbed by the absorption of a light pulse undergoes several transitions ($A \rightarrow b$, $b \rightarrow B$, $B \rightarrow a$, and $a \rightarrow A$ of Figure 1) between equilibrium (A and B of Figure 1) and non-equilibrium states (b and a of Figure 1).

Equations of motion were derived using a population matrix formulation, $\rho(z, t)$, for the interaction of the radiation field and the dye "supermolecules." A basic assumption is made that:

$$\gamma_{ab}^{-1} \ll T_p \ll \gamma_a^{-1}, \quad (24)$$

where T_p is the pulse duration, γ_{ab}^{-1} is the atomic memory time, and γ_a^{-1} is the non-radiative decay time from b to B in Figure 1. The field is regarded as a linearly polarized running wave for which the amplitude and phase vary slowly. The amplitude envelope function, ξ , used was:

$$\frac{\partial \xi}{\partial z} + \frac{1}{c} \frac{\partial \xi}{\partial t} = -\kappa \xi - \frac{1}{2\epsilon_0} k S_{AB}, \quad (25)$$

where $k = \frac{\omega_L}{c} = \frac{\text{mean frequency of incident pulse}}{\text{velocity of light}}$, κ represents nonresonant

losses in the medium, and ϵ_0 is the vacuum permittivity. S_{AB} is the quadrature component of the total polarization, P_{AB} , and is computed using the rotating wave approximation. The phase function, ϕ , for the electric field is:

$$\phi = \omega_L t - kz + \phi(z, t). \quad (26)$$

As in Eberly's treatment, a perturbation method was used, the perturbation being $\frac{\mu E}{\hbar}$ where μ is the dipole matrix element between states and E is the energy of the transition. Expressions for the perturbed and unperturbed Hamiltonians were given to be used in generating the equations of motion.

⁷ S. L. Chin and A. Zardecki, Phys. Rev. A, 13, 1528 (1976).

The total polarization P_{AB} can be expressed as the sum of the partial polarizations, P_{ab} :

$$P_{AB} = \sum_a \sum_b P_{ab} \quad (27)$$

$$= C_{AB}(z, t)\cos\phi + S_{AB}(z, t)\sin\phi. \quad (28)$$

$$P_{ab}(z, t) = \mu_{ba} \rho_{ab}(z, t) + \mu_{ab} \rho_{ab}(z, t). \quad (29)$$

The population matrix used to express the equations of motion for the "supermolecule" in any of the states, α , in levels A or B was:

$$\rho(z, t) = \sum_{\alpha} \int_{-\infty}^t dt_0 \lambda_{\alpha}(z, t_0) \rho(\alpha, z, t_0, t). \quad (30)$$

Here, $\rho(\alpha, z, t_0, t)$ is the density matrix for a molecule (at place z , time t_0) in state α , and $\lambda_{\alpha}(z, t_0)$ is the molecular concentration per unit time in state α . Four equations of motion,

$\dot{\rho} = \frac{-i}{\hbar} (\hat{H}, \rho)$, were given, $\dot{\rho}_{aa}$, $\dot{\rho}_{bb}$, $\dot{\rho}_{ba}$, and $\dot{\rho}_{ab}$. Using zeroth order elements of the population matrix, Chin and Zardecki define the population difference in a and b, $N_{ba}^{(0)}$:

$$N_{ba}^{(0)} = \rho_{bb}^{(0)} - \rho_{aa}^{(0)}. \quad (31)$$

If the photon flux is defined as $I = \frac{\epsilon_0 \xi^2 c}{2\pi \omega_L}$, and I_0 is the input pulse at $z = 0$, then the flux modified by multiple transitions is:

$$I(z, t) = \frac{I_0 (t - z/c) e^{(a_1 - 2\kappa)z}}{1 - [a_2/(a_1 - 2\kappa)] I_0 (t - z/c) \{1 - e^{(a_1 - 2\kappa)z}\}} \quad (32)$$

Resonant absorption to the first and second order in the photon flux is given by the coefficients a_1 and a_2 .

Chin and Zardecki applied their theory to the spectral hole burning phenomena, obtaining results consistent with those of Mouro¹⁷ for dye molecules in a solvent cage. They also addressed the case of two adjacent pulses being superposed in a dye medium. The first pulse obeys the condition of equation (24). The second pulse is similar to the first and occurs after a time delay, T . This delay is much shorter than the fluorescence time of the dye molecule in level B and is much shorter than the solvent-solute relaxation time. As in the propagation study, equations of motion were derived for the time evolution of the population matrix elements, and expressions for the photon flux were found. The pulse superposition effect is defined by the authors as the storage of the energy of a first pulse as a population inversion, and stimulated emission occurring subsequently upon irradiation with a second pulse. Their derivation involved separating the absorption and amplification processes, so that in the direct-transition approximation the photon flux becomes:

$$\frac{\partial I}{\partial z} + \frac{1}{c} \frac{\partial I}{\partial t} = \left[\rho_{BB} (T) \sum_d \sigma_{Bd} - 2\kappa - \sum_{a,b} \sigma_{ba} \rho_{aa}^{(o)} \left(1 + \frac{2\sigma_{ba}}{\hat{\gamma}_{ba}} \right)^{-1} \right] I. \quad (33)$$

Qualitative experimental support for the superposition theory was presented using oxazine perchlorate in glycerine.

IV. SPECTRAL CROSS-RELAXATION AND COLLISIONAL DEPHASING EFFECTS

That the absorption of light by organic dye molecules is far from a simple process was shown in the semiclassical theory of Vahey and Yariv.⁸ They treated dye solutions interacting with light at high-incident flux, as an ensemble of optical dipoles. Absorption of light in this case may exhibit inhomogeneous broadening or absorption occurring for the ensemble over a wider frequency band than would be anticipated for a lone optical dipole.⁸⁻¹⁰ Spectral cross-relaxation results when the configuration of the solvent cage or environment surrounding the optical dipole-dye molecule is altered in some way. Another effect which may occur during the interaction of light and an optical dipole-dye molecule, is that of a random shift in the phase of the radiating dipole due to collisions in the dye system.⁸⁻¹⁰

⁸ D. W. Vahey and A. Yariv, *Phys. Rev. A*, **10** (6) 1578 (1974).

¹⁰ M. Sargent III, M. O. Scully, and W. E. Lamb, Jr., "Laser Physics," Chaps 7, 14, and 16, Addison-Wesley, New York (1974).

¹⁷ G. Mouro, *IEEE J. Quantum Electron.*, **QE-11** (1) 1 (1975).

Vahey and Yariv point out that earlier treatments of the interaction between light and matter have included neither spectral cross-relaxation effects nor collisional dephasing. They presented a semiclassical study which did consider these effects.⁸ Application of their theory to experimental data for certain well-known cyanine saturable absorber dyes accounted for observed "hole-burning" in the absorption spectrum due to cross-relaxation (see Figure 3 of Mouro¹⁷). They determined that if:

T_1 = time constant for deactivation or decay of excited molecules

T_2 = time constant for collisional dephasing

T_3 = time constant for spectral cross-relaxation (and if T_A = time constant for absorption, and $\frac{1}{T} \propto$ rate of optical process), then distortion in the absorption spectrum will occur when $\frac{1}{T_A} > \frac{1}{T_3}$, and coherent interaction effects will be significant

when $\frac{1}{T_A} \sim \frac{1}{T_2}$. Vahey and Yariv are concerned here with systems in which

$T_2 \leq \frac{1}{10} T_1$. Rate equations were derived by Vahey and Yariv, based upon concepts evolving from likening the dye absorbers to an ensemble of liquid-free spins interacting with radiation. The Bloch equations were applied to such systems, affording a vector model of the density matrix equations of motion.^{8 10 12} The model used was that of a single dye molecule (state vector (u', v', r')) which underwent only one collision, a strong collision, at time t_0 . The undamped Bloch equation,

$$\frac{d\vec{r}}{dt} = \vec{\omega} \times \vec{r}' \quad (34)$$

was applied to the molecule with the conditions that:

$$\vec{r}'(t = t_0) = (0, 0, \pm 1); \vec{r}'(t) = (u', v', r')$$

$$\vec{\omega} = (\omega_1(x, t), 0, \omega_0 - \omega + \delta(t))$$

$(\omega_0 - \omega)$ is the detuning frequency,

ω_1 = a frequency proportional to the field amplitude,

$\delta(t)$ = the temporal detuning generated by cross-relaxation.

⁸ D. W. Vahey and A. Yariv, *Phys. Rev. A*, **10** (6) 1578 (1974).

¹⁰ M. Sargent III, M. O. Scully, and W. E. Lamb, Jr., "Laser Physics," Chaps 7, 14, and 16, Addison-Wesley, New York (1974).

¹² A. Yariv, "Quantum Electronics," Chap. 8, John Wiley and Sons, Inc., New York (1967).

¹⁷ G. Mouro, *IEEE J. Quantum Electron.*, *QE-11* (1) 1 (1975).

A solution of the Bloch equation is $v'(\omega_1, \omega_0 - \omega, t - t_0, \pm 1)$. Since the undamped equation was used, Vahey and Yariv note that u' , v' , and r' are not ensemble averages, but are averages determined by a summation of microscopic state vectors produced since times t_0 by random collisions at interval T_2 .

Using this model to address the question of optical absorption in dye molecules, where $\vec{p}(x, t, \omega_0)$ is the induced dipole moment (due to a polarized field, $\vec{E}(x, t)$) and \vec{p}_0 is the dipole matrix element due to the initial and final states, the authors obtain:

$$\vec{p}(x, t, \omega_0) = \vec{p}_0 [-u(x, t, \omega_0) \cos(\omega t - kx) + v(x, t, \omega_0) \sin(\omega t - kx)] \quad (35)$$

since

$$\vec{E}(x, t) = \hat{e}_z \xi(x, t) \cos(\omega t - kx). \quad (36)$$

The out-of-phase component of polarization, $v(x, t, \omega_0)$, is:

$$v(x, t, \omega_0) = \int_{-\infty}^t \frac{dt_0}{T_2} e^{-(t-t_0)/T_2} [n_b(x, t_0 +, \omega_0) v'(\omega_1, \omega, t - t_0 - 1) + n_a(x, t_0 +, \omega_0) v'(\omega_1, \omega_0 - \omega, t - t_0 + 1)]. \quad (37)$$

This is the instantaneous polarization. The probability that the dye ground state remained occupied after collision occurred (at time t_0) was given by $n_b(x, t_0 +, \omega_0)$. The probability that dye molecules would be found in an excited state at time $(t_0 +)$ was $n_a(x, t_0 +, \omega_0)$; to eliminate systems showing stimulated emissions, $n_a = 0$. Development of an expression for $n_b(x, t_0 +, \omega_0)$ leads to expressions for the induced polarization:

$$v(x, t, \omega_0) = \frac{\omega_1(x, t) T_2}{1 + \omega_1^2(x, t) T_2^2 + (\omega_0 - \omega)^2 T_2^2} \left\{ n_b(x, t, \omega_0) [1 - (T_2/T_3)] + \int_{-\infty}^{\infty} d\omega'_0 g(\omega'_0 - \bar{\omega}_0) m_b(x, t, \omega'_0) (T_2/T_3) + n_c(x, t) (T_2/T_1) \right\} \quad (38)$$

In phase $u(x, t, \omega_0)$:

$$u(x, t, \omega_0) = -(\omega_0 - \omega) T_2 v(x, t, \omega_0). \quad (39)$$

To relate the induced polarization and light absorption, the electromagnetic wave equation was separated into two equations, one for phase and one for amplitude. The wave amplitude, $\xi(x, t)$ is

$$\frac{\partial \xi}{\partial x} + \frac{n_o}{c} \frac{\partial \xi}{\partial t} = -\frac{1}{2} N_o \frac{\omega P_o}{\epsilon_o n_o c} \int_{-\infty}^{\infty} d\omega_o g(\omega_o - \bar{\omega}_o) v(x, t, \omega_o). \quad (40)$$

Here, N_o is the absorber concentration, while the refractive index of the solvent is given by n_o . One of the basic consequences obtained by Vahey and Yariv, the photon-transport equation, evolved from the wave amplitude equation. The transport equation is:

$$\left(\frac{\partial}{\partial x} + \frac{n_o}{c} \frac{\partial}{\partial t} \right) F(x, t) = -N_o \sigma_h F(x, t) \langle n_{\text{eff}}(x, t, \omega_o) \mathcal{L}(F, \omega_o - \omega) \rangle_{\text{av.}} \quad (41)$$

where the flux density is:

$$F(x, t) \equiv \frac{1}{2} \epsilon_o c n_o \xi^2(x, t) / \hbar \omega, \quad (42)$$

the homogeneous cross section for interaction of molecules in exact resonance with the wave is

$$\sigma_h = \omega p_o^2 T_2 / \epsilon_o c n_o \hbar, \quad (43)$$

The effective ground-state population is

$$\begin{aligned} n_{\text{eff}}(x, t, \omega_o) &= n_b(x, t, \omega_o) [1 - (T_2/T_3)] \\ &+ \langle n_b(x, t, \omega_o) \rangle_{\text{av.}} (T_2/T_3) \end{aligned} \quad (44)$$

and $\mathcal{L}(F, \omega_o - \omega)$ is a modified Lorentzian function.

The other significant theoretical result obtained was the population rate equation:

$$\begin{aligned} \frac{\partial}{\partial t} n_b(x, t, \omega_o) &= -\sigma_n F(x, t) n_{\text{eff}}(x, t, \omega_o) \mathcal{L}(F, \omega_o - \omega) - \\ &\{ [n_b(x, t, \omega_o) - \langle n_b(x, t, \omega_o) \rangle_{\text{av.}}] / T_3 \} + n_c(x, t) T_1. \end{aligned} \quad (45)$$

Vahey and Yariv applied their theory to two classes of absorbers, according to:

- (1) Steady-state limit, where the pulse duration is long with respect to T_1 and T_2 .
- (2) Short-pulse limit, where the pulse duration is very short compared to T_1 and T_2 .

For the steady-state case, they found that the experimental parameter in giant-pulse studies with organic dyes is the steady-state transmission:

$$\frac{F(x)}{F(0)} = \frac{Q^2(x) - 1}{Q^2(0) - 1} \quad (46)$$

where

$$Q = [1 + F\sigma_n(T_2 + T_3)]^2 \quad (47)$$

and $F(0)$ is a known input flux density. In the short-pulse case, the theory was extended to such problems occurring in real-world absorbing systems as:

- (a) Excited state absorption.
- (b) The molecular anisotropy dependence.
- (c) The multiplicity of interacting excited states.

V. A QUANTUM-MECHANICAL DYE LASER THEORY INVOLVING EXACT TREATMENT OF TRIPLET STATES

Schaefer and Willis were the first to present a dye laser theory which was not only fully quantum-mechanical, including radiation fluctuations, but also treated the triplet states exactly.⁹ As noted above, dye lasers are unique among lasers because both singlet and triplet levels can be populated during lasing.

The model upon which this quantum-mechanical treatment was based, consisted of two main parts—matter and radiation. Reservoir theory was invoked, creating a matter or molecule reservoir (singlet and triplet level decay) and a radiation reservoir (accounting for radiative losses in the laser cavity).¹⁰

⁹ R. B. Schaefer and C. R. Willis, *Phys. Rev. A*, **13** (5) 1874 (1976).

¹⁰ M. Sargent III, M. O. Scully, and W. E. Lamb, Jr., "Laser Physics," Chaps 7, 14, and 16. Addison-Wesley, New York (1974).

The Hamiltonian, \hat{H} , for this model became:

$$\hat{H} = [\hat{H}_R + \hat{H}_M + V_{R-M}] + [\hat{H}_{R-rs} + V_{R-rs}] + [\hat{H}_{M-rs} + V_{M-rs}] \quad (48)$$

where V represents the interaction Hamiltonian potential. The terms are defined by Schaefer and Willis as:

\hat{H}_R = radiation Hamiltonian

\hat{H}_M = molecule Hamiltonian

\hat{H}_{R-rs} = radiation-reservoir Hamiltonian

\hat{H}_{M-rs} = molecule-reservoir Hamiltonian

V_{R-M} = radiation-molecule interaction

V_{R-rs} = radiation-radiation reservoir interaction

V_{M-rs} = molecule-molecule reservoir interaction.

The reservoir terms were eliminated in Schaefer and Willis because the only reservoir attributes of interest were those of pumping and decay and not specific details about the reservoirs.

The radiation field Hamiltonian was:⁹

$$\hat{H} = \sum_{\vec{k}} \sum_{\alpha} \hbar \Omega_{\vec{k}} (a_{\vec{k},\alpha}^{\dagger} a_{\vec{k},\alpha} + 1/2) \quad (49)$$

where $\sum_{\vec{k}}$ represents summation over all cavity modes (in the present case $\vec{k} = 1$, since only a single-mode was treated), and \sum_{α} represents summation over all α polarizations. The frequency associated with \vec{k} is $\Omega_{\vec{k}}$. The "a and a^{\dagger} " operators are annihilation and creation photon operators, respectively.¹⁰

⁹ R. B. Schaefer and C. R. Willis, Phys. Rev. A, 13 (5) 1874 (1976).

¹⁰ M. Sargent III, M. O. Scully, and W. E. Lamb, Jr., "Laser Physics," Chaps 7, 14, and 16, Addison-Wesley, New York (1974).

Turning to the Hamiltonian for matter, \hat{H}_M , the singlet-triplet states associated with dye molecule energy levels require that:

$$\hat{H}_M = \hat{H}_S + \hat{H}_T \quad (50)$$

where

$$\hat{H}_S = \sum_{\sigma} \sum_i (s_{\sigma,i}^{\dagger} s_{\sigma,i} \epsilon_i)$$

$$\hat{H}_T = \sum_{\sigma} \sum_j (t_{\sigma,j}^{\dagger} t_{\sigma,j} \epsilon_j)$$

and σ = total dye-molecules active in laser system

i = singlet states

j = triplet states

ϵ_i & ϵ_j = energies corresponding to i or j levels, respectively

$s_{\sigma,i}^{\dagger}$ & $t_{\sigma,j}^{\dagger}$ = creation operators for singlets, triplets, respectively

$s_{\sigma,i}$ & $t_{\sigma,j}$ = annihilation operators for singlets and triplets, respectively.

The relationship between the molecular annihilation and creation operators were given:

$$[s_{\sigma,i}^{\dagger}, s_{\sigma,j}] = \delta_{ij} \quad [t_{\sigma,i}^{\dagger}, t_{\sigma,j}] = \delta_{ij} \quad (51)$$

$$[s_{\sigma,i}^{\dagger}, s_{\sigma,j}^{\dagger}] = [s_{\sigma,i}, s_{\sigma,j}] = [t_{\sigma,i}^{\dagger}, t_{\sigma,j}^{\dagger}] = [t_{\sigma,i}, t_{\sigma,j}] = 0 \quad (52)$$

$$[s_{\sigma,i}^{\dagger}, t_{\sigma,j}^{\dagger}] = [s_{\sigma,i}^{\dagger}, t_{\sigma,j}] = [s_{\sigma,i}, t_{\sigma,j}^{\dagger}] = [s_{\sigma,i}, t_{\sigma,j}] = 0 \quad (53)$$

That the molecule or matter terms must be treated as singlets and triplets was shown also in the radiation-matter interaction Hamiltonians:

$$V_{R-M} = V_{R-S} + V_{R-T} \quad (54)$$

From the matter-radiation interaction (dipole approximation), Schaefer and Willis found:

$$(55)$$

$$V_{R-S} = i \sum_{\sigma} \sum_{i,j} \sum_k \gamma_{\sigma ijk} (s_{\sigma,i}^{\dagger} s_{\sigma,j} a_k^{\dagger} + s_{\sigma,j}^{\dagger} s_{\sigma,i} a_k), \quad (55)$$

where $\epsilon_j > \epsilon_i$,

$$\gamma_{\sigma ijk} = [4 \pi e^2 (\epsilon_j - \epsilon_i)^2 h \Omega_k V]^{1/2} \times (\vec{\epsilon} \cdot \vec{\chi}) \Gamma(\chi);$$

$$V_{R-T} = i \sum_{\sigma} \sum_{l,m} \sum_k \gamma_{\sigma lmk} (t_{\sigma,l}^{\dagger} t_{\sigma,m} a_k^{\dagger} + t_{\sigma,m}^{\dagger} t_{\sigma,l} a_k) \quad (56)$$

where $\epsilon_l > \epsilon_m$,

$$\gamma_{\sigma lmk} = [4 \pi e^2 (\epsilon_m - \epsilon_l)^2 h \Omega_k V]^{1/2} \times (\chi_{lm}) \Gamma(\chi).$$

m = electron mass

χ_{lm} = dipole matrix element

e = electronic charge

V = volume of laser cavity

$\Gamma(\chi)$ = eigenfunction for the spacial cavity

\sum_{ij} for $j > i$

$\sum_{l,m}$ for $m > l$

$\sum_k = 1$ for single-mode operation.

The rotating wave approximation was applied to the interaction Hamiltonians by the authors. (Thus transitions were eliminated where photon *absorption* might produce a *lower energy level* and photon *emission* might yield a *higher energy level*). Thus, from the commutation relations cited, $s^{\dagger} s a^{\dagger}$ (or $t^{\dagger} t a^{\dagger}$) terms correspond to the creation (emission) of a photon and decrease in molecular energy level. Similarly, $s^{\dagger} s a$ (or $t^{\dagger} t a$) terms indicate photon annihilation (absorption) and an increase in molecular energy levels.

Having defined the system Hamiltonian, Schaefer and Willis utilized the Born-Markoff¹⁸ approximation for the Liouville equation.

$$\frac{d\hat{\rho}(t)}{dt} = \frac{i}{\hbar} [\hat{H}, \hat{\rho}(t)], \hat{\rho}(t) \text{ being the density operator of the system} \quad (57)$$

To demonstrate the validity of such an approach, they calculated that experimental data for Rhodamine 6G, one of the best-studied laser dyes, did indeed satisfy the criterion for validity that its (continuous-wave lasing) interaction time is minimal compared to the relaxation time, or $(t_{int}/t_{rel}) \ll 1$.

Thus,

$$0.025 \leq \left(\frac{S N}{\nu_{S,1}} \right) \leq 0.04 \quad (58)$$

and

$$0.01 \leq \left(\frac{T N}{\nu_{T,u}} \right) \leq 0.03 \quad (59)$$

where

N = number of lasing organic molecules

S = stimulated emission rate of singlets

T = stimulated absorption rate for triplets

$\nu_{S,1}$ = lower singlet decay rate

$\nu_{T,u}$ = upper triplet decay rate.

Clearly, both the singlet and triplet contributions satisfy the $(t_{int}/t_{rel}) \ll 1$ criterion.

¹⁸ SS&L, in discussing the Markoff approximation, note that the average value of a random shift in an energy difference is zero, so that if the variations of this shift are rapid compared to other decay times (relaxation times) in the system, then, $\frac{\text{(average shift)}}{\text{decay time}} \sim 0$.

If $\hat{H} = \hat{H}_0 + \lambda V$ and λ is dimensionless, the above approximation was given as:

$$\frac{d\hat{p}(t)}{dt} + \frac{i}{h} [\hat{H}_0, \hat{p}(t)] = -\lambda \int_0^{\infty} dT [V, (V(T), \hat{p})], \quad (60)$$

$$\text{where } V(T) = e^{i\hat{H}_0 T/h} V e^{-i\hat{H}_0 T/h}. \quad (61)$$

This equation was shown by Willis to describe laser photon statistics if one does not separate matter-radiation interactions into factorable matrices.¹⁹

Equations of motion were derived next. As derived by Schaeffer and Willis, using the Born approximation master equation where the diagonal matrix is coupled only to the diagonal elements, two matrices result:

Radiation Density Matrix

$$R_n = \langle n | R | n \rangle \quad (62)$$

Diagonal Single-particle Radiation Density Matrix

$$D_n^j(t) \equiv \langle n | \langle j | D(1, t) | j \rangle | n \rangle; \quad (63)$$

here n is the photon number, and j refers to the state of the j th molecule. Two sets of equations of motion were formulated from these matrices. One set, the equation of motion for the radiation density operator $\hat{R}(t)$ is:

$$\begin{aligned} \frac{d R_n(t)}{dt} &= \nu_R [n \overset{\textcircled{1}}{R}_n - (n+1) \overset{\textcircled{2}}{R}_{n+1}] \\ &- S N [(n+1) (D_n^{S,u} \overset{\textcircled{3}}{-} D_{n-1}^{S,q}) + n (D_n^{S,q} \overset{\textcircled{4}}{-} D_{n-1}^{S,u})] \\ &- T N [(n+1) (D_n^{T,u} \overset{\textcircled{5}}{-} D_{n+1}^{T,q}) + n (D_n^{T,q} \overset{\textcircled{6}}{-} D_{n-1}^{T,u})]. \end{aligned} \quad (64)$$

¹⁹ C. R. Willis, Phys. Rev., 165 420 (1968).

The terms labeled (1) and (2) give the radiation decay associated with the radiation reservoir, (3) is emission (stimulated and spontaneous) in the singlet state, (4) is stimulated absorption for the ground singlet state, and (5) and (6) are the terms corresponding to (3) and (4) respectively, for the triplet state. The other set for the singlet-molecule-radiation density operator $\hat{D}(\ell, t)$ was composed of four equations to satisfy the possibility that the molecule could be in a single (S) or triplet (T) state, and in an upper (u) or lower (ℓ) level of state:

$$\frac{dD_n^{S,u}(t)}{dt} = -(\nu_{S,u} + K_{ST}) D_n^{S,u} + R_n(t) \nu_e - S(n+1) (D_n^{S,u} - D_{n+1}^{S,\ell}), \quad (65)$$

$$\frac{dD_n^{S,\ell}(t)}{dt} = -\nu_{S,\ell} D_n^{S,\ell} - S_n (D_n^{S,\ell} - D_{n+1}^{S,u}) (n+1) \quad (66)$$

$$\frac{dD_n^{T,u}(t)}{dt} = -\nu_{T,u} D_n^{T,u} - T(D_n^{T,u} - D_{n+1}^{T,\ell}) (n+1) \quad (67)$$

$$\frac{dD_n^{T,\ell}(t)}{dt} = K_{ST} D_n^{S,u} + \beta \nu_{T,u} D_{n-1}^{T,u} - \nu_{\ell} D_n^{T,\ell} - nT(D_n^{T,\ell} - D_{n-1}^{T,u}). \quad (68)$$

These two sets of equations are the fundamental ones for the Schaefer-Willis dye laser theory. Several solutions were obtained, incorporating the parameters α and β :

α = a triplet parameter:

the value of α increases with triplet concentration.

$$\alpha \equiv (T K_{ST} / S \nu_{T,\ell}).$$

β , where $0 \leq \beta \leq 1$, and

where

T = triplet stimulated absorption rate

S = singlet stimulated emission rate

K_{ST} = intersystem crossing rate

β = fraction return of excited triplets, $0 \leq \beta \leq 1$

V_e = molecular excitation rate due to external pump

n = number of photons

V_R = radiation decay rate.

The dye laser exact solution for the steady state was obtained:

$$R_{n+1} = \prod_{p=0}^n G(p) R_0 \quad (69)$$

where $G(p) \equiv \left(\frac{S\eta_p^{S,u} + T\eta_p^{T,u}}{S\eta_{p+1}^{S,\ell} + T\eta_{p+1}^{T,\ell} + \nu_R/N} \right)$

p is an integer.

$\prod_{p=0}^n$ is the product of terms for $0 \leq p \leq n$.

and $\eta_n^{S,u}$ is obtained from $D_n^{S,t}(t) = \eta_n^{S,t}(t) R_n(t)$ when

$$\frac{dD_n^{S,u}(t)}{dt} = Q \text{ in equations (65) - (68).}$$

Simplified solutions of this equation were obtained by (1) eliminating the triplets, giving the "singlet-state case," and (2) including the triplets, but neglecting terms for triplet-state emission and singlet-state absorption to obtain the "approximate solution." Finally, graphical solutions of the exact equation were obtained by computer methods.

Results of the various solutions were expressed as the extrema of R_n , the threshold value of pumping, and variations in R_n when pumping values are above threshold. One interesting result of this theory is that, instead of one (maximum) extremum for the radiation distribution, R_n , as is found for a gas laser, for the dye laser it is possible to expect both a maximum and a minimum for R_n , depending upon other laser parameters.

VI. CONCLUSIONS

The basic quantum mechanical studies of dye lasers have been reviewed. Despite the complexity of dye laser systems, compared to other laser systems, successful results have been achieved. An acceptable understanding exists of dye lasers and their potential prospects in the areas of spectral hole burning, gain, propagation of light pulses in the dye medium (and the superposition of two light pulses), light absorption and emission processes for states of different multiplicities, and pumping thresholds. Although much remains to be done, these studies demonstrate the unique contribution quantum mechanics can make upon an examination of a time-dependent interaction of matter with the electrical field component of light in a dye medium.

DISTRIBUTION FOR NV&EOL REPORT DELNV-TR-0024

No. Copies	Addressee	No. Copies	Addressee
1	Commander ERADCOM ATTN: DRDEL-CP-C 2800 Powder Mill Road Adelphi, MD 20783	1	Project Manager M60 Tank System Warren, MI 48090
1	Director Atmospheric Sciences Lab ATTN: DELAS-D White Sands Missile Range, NM 88802	1	Project Manager TOW/DRAGON Redstone Arsenal, AL 35809
1	Director CS&TA Laboratory ATTN: DELCS-D Fort Monmouth, NJ 07703	1	Project Manager VIPER/AHAMS Redstone Arsenal, AL 35809
1	Director Electronic Warfare Lab ATTN: DELEW-D Fort Monmouth, NJ 07703	1	Project Manager AAH AVARADCOM St. Louis, MO 63166
1	Director Electronics Technology & Devices Laboratory ATTN: DELET-D Fort Monmouth, NJ 07703	1	Project Manager TADS/PNVS AVARADCOM St. Louis, MO 63166
1	Commander Harry Diamond Labs ATTN: DELHD-AC Adelphi, MD 20783	1	Commander ARRCOM ATTN: DRSAR-CPE Rock Island, IL 61299
1	Director Signal Warfare Lab ATTN: DELSW-D Vint Hill Station, VA 22186	1	Project Manager RPV AVARADCOM St. Louis, MO 63166
1	Project Manager XM-1 Tank System Warren, MI 48090	1	Commander ARRADCOM ATTN: DRDAR-SEC Dover, NJ 07801

No. Copies	Addressee
1	Commander CERCOM ATTN: DRSEL-CP-CR Fort Monmouth, NJ 07703
1	Project Manager MICV Warren, MI 48090
1	Commander MIRADCOM ATTN: DRDMI-DC Redstone Arsenal, AL 35809
1	Commander MIRCOM ATTN: DRSMI-FO Redstone Arsenal, AL 35809
1	Commander AVARADCOM ATTN: DRCAV-BC St. Louis, MO 63166
1	Commander TSARCOM ATTN: DRSTS-CO St. Louis, MO 63166
1	Commander TARADCOM ATTN: DRCTA-VC Warren, MI 48090
1	Commander TARCOM ATTN: DRSTA-NC Warren, MI 48090
1	Commander NARADCOM ATTN: DRDNA-O Natick, MA 01760

No. Copies	Addressee
1	Commander AMMRC Watertown, MA 01272
1	Commandant US Army Infantry School Fort Benning, GA 31905
1	Director Defense Advanced Research Project Agency Rosslyn, VA 22209
1	Commander MERADCOM ATTN: DRDME-CA 3 DRDME-WP Fort Belvoir, VA 22060
1	Commandant Defense Systems Management School Fort Belvoir, VA 22060
1	Commander US Army Engineer School Fort Belvoir, VA 22060
1	Commander US Operational Test and Evaluation Agency 5600 Columbia Pike Falls Church, VA 22041
1	Commander USACSC Fort Belvoir, VA 22060
1	HQDA ATTN: DACA-CA Washington, DC 20310

No. Copies	Addressee	No. Copies	Addressee
12	Defense Technology Information Center ATTN: DDC-TCA Cameron Station (Bldg 5) Alexandria, VA 22314	1	Department of Defense Production Engineering Spt Ofc (PESO) ATTN: D. Anderson Cameron Station Alexandria, VA 22314
1	Commander US Army Training & Doctrine Command ATTN: ATCD-AN Fort Monroe, VA 23651	1	Project Manager CAC ATTN: DRCPM-CAC Vint Hill Station, VA 22186
2	Commander US Army Logistics Center Fort Lee, VA 23801	1	Project Manager SOTAS ATTN: DRCPM-STA Fort Monmouth, NJ 07703
1	Commander US Army Systems Analysis Agency Aberdeen Proving Ground, MD 21005	1	Commandant US Army Armor School Fort Knox, KY 40121
1	Commander US Naval Research Lab Washington, DC 20375	1	Commandant US Army Field Artillery School Fort Bliss, TX 79916
1	Commander HQ DARCOM ATTN: DRCDE-DE Alexandria, VA 22333	3	Commander AFSC/ACC Andrews AFB, MD 20334
1	Director ATTN: DELNV-TMS/SEMCO Fort Belvoir, VA 22060	1	Study Center National Maritime Research Center King's Point, NY 11024
1	Project Manager REMBASS ATTN: DRCPM-RBS Fort Monmouth, NJ 07703	1	Commander USAAVNC Fort Rucker, AL 36362
1	Project Manager FIREFINDER ATTN: DRCPM-FF Fort Monmouth, NJ 07703	1	Commander ESD/ACC Hanscomb AFB, MA 01730

No. Copies	Addressee
1	Commander US Naval Ordnance Lab/ White Oak ATTN: Technical Library Silver Spring, MD 20910
1	Commander Naval Electronics Lab Ctr ATTN: Library San Diego, CA 92152
1	Armament Development and Test Center ATTN: DLOSL, Tech Library Eglin Air Force Base, FL 32542
2	Director NV&EOL ATTN: DELNV-L (R. Buser)
1	DELNV-D (L. Cameron)
1	DELNV-L (T. Paul)
	Fort Belvoir, VA 22060
1	Commander Sacramento Army Depot Sacramento, CA 95813
1	Commander New Cumberland Army Depot New Cumberland, PA 17070
1	Commander Anniston Army Depot Anniston, AL 36201
1	Commandant US Army Air Mobility R&D Ctr Ames Research Center Moffett Field, CA 94035
20	Commander US Army Communications and Electronics Material Readiness Command ATTN: DRSEL-CP-CR (C. Pearce) Fort Monmouth, NJ 07703

DATE
ILME