

AD-A115 817

OHIO STATE UNIV COLUMBUS DEPT OF CHEMISTRY
STUDIES ON THE SYNTHESIS OF BORON HYDRIDE SYSTEMS. (U)
NOV 81 S G SHORE

F/G 7/3

DAA629-78-6-0112

UNCLASSIFIED

ARO-15611.9-C

NL

1 of 1

4-1A

08/7

END

DATE

FORMED

07-82

DTIC

15611.9-C

RF Project 761119/711144
Final Report

(12)

**the
ohio
state
university**

research foundation

1314 kinnear road
columbus, ohio
43212

STUDIES ON THE SYNTHESIS OF
BORON HYDRIDE SYSTEMS

Sheldon Shore
Department of Chemistry

For the Period
July 1, 1978 - September 30, 1981

U.S. DEPARTMENT OF THE ARMY
Army Research Office
Research Triangle Park, North Carolina 27709

Grant No. DAAG29-78-G-0112

November, 1981

DTIC
ELECTE
JUN 21 1982
S D
E

This document has been approved
for public release and sale; its
distribution is unlimited.

AD A115817

DTIC FILE COPY

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO. AD-A115817	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Studies on the Syntheses of Boron Hydride Systems.		5. TYPE OF REPORT & PERIOD COVERED Final 7-1-78 to 9-30-81
		6. PERFORMING ORG. REPORT NUMBER 761119/711144
7. AUTHOR(s) Dr. Sheldon, G. Shore		8. CONTRACT OR GRANT NUMBER(s) DAAG 29-78-G-0112
9. PERFORMING ORGANIZATION NAME AND ADDRESS The Ohio State University Research Foundation 1314 Kinnear Rd Columbus, Ohio 43214		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS P-15611-C
11. CONTROLLING OFFICE NAME AND ADDRESS U.S. Army Research Office Post Office Box 12211 Research Triangle Park, NC 27709		12. REPORT DATE 11-20-81
		13. NUMBER OF PAGES 11
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE NA
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) NA		
18. SUPPLEMENTARY NOTES The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Boron hydrides, carboranes, preparation of		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The boron hydrides B_2H_6 , B_4H_{10} , B_5H_{11} , and $B_{10}H_{14}$ are prepared in good yields through hydride ion abstraction reactions when the borane anions BH_4^- , $B_3H_8^-$, $B_4H_9^-$, and $B_9H_{14}^-$, respectively, are treated with a molar equivalent of a Lewis acid BX_3 ($X=F, Cl$, or Br), generally in the absence of a solvent, for reaction periods of 1-4 hrs. A high yield (up to 90%) method for the conversion of B_5H_9 to $B_9H_{14}^-$ is presented as the precursor to the practical		

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20.

conversion of B_5H_9 to $B_{10}H_{14}$ (up to (50%)). Additionally a good conversion of B_5H_9 to $n-B_{18}H_{22}$ (up to (50%)) and the preparation of the nido-carborane $\phi_2C_2B_8H_{10}$ have been achieved. The hydride ion abstraction reactions by BBr_3 and BCl_3 lead to the new anions $HBBr_3^-$ and $HBCl_3^-$.

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	



Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

Forward

In the past three years under ARO support, we have developed a new approach to boron hydride syntheses¹ which has led to convenient high yield preparations of B_4H_{10} and B_5H_{11} by procedures which are safer, cleaner, and more easily scaled-up than the standard pyrolytic and chemical methods. Additionally, this new approach has led to a practical and simple conversion of B_5H_9 to $B_{10}H_{14}$ in a "one-pot" reaction. This latter result is of special significance in that 200,000 pounds of B_5H_9 is in storage, a legacy from the high energy fuels program from the 1950's. Thus there is the potential for putting this B_5H_9 to good use in preparing $B_{10}H_{14}$ which is currently in short supply. The U.S. Army has applied for a patent for this procedure. Furthermore, in the course of our work under ARO support we have developed additional chemistry of B_5H_9 which leads to a high yield preparation of $B_9H_{14}^-$ (90% yield), a good conversion of B_5H_9 to $B_{18}H_{22}$ (ca 50%), and most recently the preparation of the nido-carborane $\phi_2C_2B_8H_{10}$. In effect, B_5H_9 can be considered to be a raw material which can be the starting point for preparing materials which are normally prepared from $B_{10}H_{14}$ from B_5H_9 or proceeding directly to materials which are normally synthesized from $B_{10}H_{14}$. Principles which have been developed for the preparation of $B_{10}H_{14}$, B_5H_{11} , and B_4H_{10} have also been applied to the development of a high yield synthesis (up to 90%) of B_2H_6 in a procedure which does not require a solvent. The Ohio State University has applied for a patent on this process.


The above statements briefly summarize background information to the proposed program. Details are provided below.

Results and Discussion

1. Hydride Ion Abstraction Reactions: New, Systematic Good Yield Syntheses of B_4H_{10} , B_5H_{11} , and $B_{10}H_4$


We have developed a new systematic approach to boron hydride syntheses which is based upon the principle of hydride ion abstraction.

The systematic nature of these syntheses relates to our observation that hydride ion can be abstracted from certain boron hydride anions to give as one of the final products a neutral boron hydride which contains one more boron atom than the anionic starting material. The simplest reaction observed involves $[N(\underline{n}\text{-C}_4\text{H}_9)_4][BH_4]$ and is quantitative



where $BX_3 = BCl_3, BBr_3$. The tetra- $\underline{n}$ -butyl ammonium salts of the previously unreported anions $HBBR_3^-$ and $HBCl_3^-$ are stable, free flowing solids under a dry atmosphere at room temperature. Very recently we have discovered that $NaBH_4$ can react quantitatively with BF_3 in the absence of a solvent to give high purity B_2H_6 . This "dry" preparation of B_2H_6 is a potentially practical synthesis.

Tetraborane(10) and pentaborane(11) are prepared by the following reactions in which 1:1 molar ratios of reactants are stirred vigorously in the absence of a solvent.




Tetraborane(10) and pentaborane(11) are obtained in 65% and 60% yields, respectively. These yields are based upon the amount of boron in the borane anion. In both reactions the borane anion is completely consumed. The starting material $[N(\underline{n}\text{-C}_4\text{H}_9)_4][\text{B}_3\text{H}_8^-]$ is well known and of course KB_4H_9 is easily obtained from the B_4H_{10} prepared in the reaction cited on page 4. We routinely prepare B_4H_{10} and B_5H_{11} in 10 to 20 millimole quantities in the time frames cited above. Scale-up to larger quantities is practical.

One of the principal handicaps to investigating the chemistry of the intermediate boron hydrides B_4H_{10} and B_5H_{11} has been the absence of simple preparative procedures which would provide these materials in relatively large quantities in good yield. Traditionally, B_4H_{10} and B_5H_{11} have been prepared by hot-cold reactor techniques^{2,3} and more recently from the protonation of B_3H_6^- salts.⁴⁻⁶ Additionally B_5H_{11} has been prepared from the protonation^{5b} of $\text{B}_5\text{H}_{12}^-$. The procedures outlined here are much safer and simpler than the classical hot cold reactor techniques. Additionally, requirements for product purification are minimal for the present method compared to the hot-cold reactor methods and the method of protonation of B_3H_8^- salts. When carried out under conditions indicated, the presence of volatile impurities (trace quantities of B_2H_6 and B_5H_9 from reaction 2 and trace quantities of $\underline{n}\text{-B}_9\text{H}_{15}$ from reaction 3) present no problems in purifying B_4H_{10} and B_5H_{11} . Our present method for preparing B_5H_{11} is also superior to the earlier reported protonation of $\text{B}_5\text{H}_{12}^-$ since it gives comparable yields but requires one less step in the preparative procedure.



Reaction 1 can be viewed as hydride abstraction from BH_4^- ions to

give BH_3 units which combine to form B_2H_6 .




This reaction differs from the traditional syntheses of B_2H_6 in which diborane(6) is generated through hydride-halide exchange in reactions of metal borohydrides with solvents.⁷⁻⁹

For reactions 2 and 3 hydride abstraction would yield the boranes B_3H_7 and B_4H_8 respectively. In view of the products obtained, it is reasonable to assume that subsequent reactions involve, effectively, transfer of BH_3 , for example from one B_4H_8 to another B_4H_8 to produce B_5H_{11} . Indeed very recent results we have obtained which are described in the Proposed Investigation Section of this proposal support this reaction scheme. Thus, viewing reactions 2 and 3 in this light suggests the following stoichiometries.




In these reactions 67% of the available boron in B_3H_8^- is converted to B_4H_{10} and 63% of the available boron in B_4H_9^- is converted to B_5H_{11} . The close correspondence of experimental yields to these proposed stoichiometries (2' and 3') suggests that within experimental error reactions 2 and 3 are quantitative with respect to yields of B_4H_{10} and B_5H_{11} . Residues of empirical compositions $(\text{BH}_2)_x$ and $(\text{B}_3\text{H}_5)_x$ decompose at room temperature to give small amounts of B_5H_9 and $n\text{-B}_9\text{H}_{15}$ respectively.

The systematic nature of this procedure was further demonstrated in an extension of reactions 2 and 3. Treatment of $[\text{N}(\text{CH}_3)_4]\text{B}_9\text{H}_{14}$ with BCl_3 gave $\text{B}_{10}\text{H}_{14}$ in a yield of 50% based on $\text{B}_9\text{H}_{14}^-$ (Reaction 4).



A reaction stoichiometry analogous to 2' and 3' is suggested
(Reaction 4')



In this stoichiometry 56% of the boron in $B_9H_{14}^-$ is converted to $B_{10}H_{14}$, which corresponds well with our results and is consistent with reaction 4 being close to quantitative.

2. The Use of Pentaborane(9) as a Starting Material for the Preparation of Higher Boron Hydrides and Carborane Systems

Described here are our recent studies which have led to the practical conversion of B_5H_9 to $B_9H_{14}^-$, $B_{10}H_{14}$, $B_{18}H_{22}$, $B_9H_{13}L$, and most recently the nido-carborane system $R_2C_2B_8H_{10}$. The key to this work is our development of a good high yield conversion of B_5H_9 to $B_9H_{14}^-$ which is described below.

a. $B_9H_{14}^-$. Although $B_9H_{14}^-$ is generally prepared through the degradation of $B_{10}H_{14}$ by base,¹⁰ it is also possible to prepare this ion through the thermal decomposition of $B_5H_8^-$ which is generated by deprotonating B_5H_9 . However, the yields of $B_9H_{14}^-$ prepared this way from B_5H_9 do not exceed 60%.¹¹⁻¹⁴ By allowing $B_5H_8^-$ to react with an equimolar amount of B_5H_9 in THF at room temperature we have been able to obtain good quality $B_9H_{14}^-$ in 90% yield.¹⁵ This is achieved experimentally by treating B_5H_9 with NaH in a 2:1 molar ratio. Although not described here, we have examined in detail the reaction of $B_5H_8^-$ with B_5H_9 not only to optimize the yield and purity of $B_9H_{14}^-$, but also to identify intermediates formed in the overall reaction scheme. We have prepared, according to our particular needs the salts $[N(n-C_4H_9)_4][B_9H_{14}]$, $Na[B_9H_{14}]$ and $K[B_9H_{14}]$.

b. $B_{10}H_{14}$. Our high yield preparation of $[N(\underline{n-C_4H_9})_4][B_9H_{14}]$ coupled with our preparation of $B_{10}H_{14}$ from the reaction of $[N(\underline{n-C_4H_9})_4][B_9H_{14}]$ with BCl_3 provides a practical route to $B_{10}H_{14}$ from B_5H_9 employing a single reactor for the entire procedure.

In a typical preparation of $B_{10}H_{14}$ from B_5H_9 , 21.6 millimoles of NaH, 43.2 millimoles of B_5H_9 and 22 millimoles of $[N(CH_3)_4]Cl$ are stirred for 12 hrs. in 16 ml of THF at room temperature. Hydrogen gas and THF are pumped away, leaving behind a dry solid which is good quality $[N(CH_3)_4][B_9H_{14}]$ and NaCl. Then 22 millimoles of BCl_3 are condensed onto the solid reaction products and this mixture is stirred vigorously for 6 hrs at 25° . The $B_{10}H_{14}$ is then sublimed from the flask under dynamic vacuum. A 9.57 mmole quantity of $B_{10}H_{14}$ representing a 45% conversion of B_5H_9 to $B_{10}H_{14}$ is obtained. This percent conversion of starting material to $B_{10}H_{14}$ is comparable to that reported for the conversion of $NaBH_4$ to $B_{10}H_{14}$ by a nonpyrolytic method.¹⁶ The present procedure, however, requires fewer steps, less solvent, and it also can be scaled-up.



Very recently we have been able to modify the preparation of $B_{10}H_{14}$ to give a more effective conversion of B_5H_9 to $B_{10}H_{14}$. Details of this modification and the principles behind it are discussed in the Proposed Program Section of this proposal since we hope to use this modification in a general way in our systematic synthetic procedures which are based upon hydride ion abstraction reactions.

c. $B_9H_{13}L$ and $\underline{n-B_{18}H_{22}}$. By taking advantage of the simple high-yield conversion of B_5H_9 to $B_9H_{14}^-$ described above, we found that it is possible to convert B_5H_9 to $B_9H_{13}L$, where $L = Et_2S, P\phi_3$ and $\underline{n-B_{18}H_{22}}$ in practical "one-pot" procedures which are modifications of an earlier method¹⁷. For example, $B_9H_{14}^-$ is converted to $B_9H_{13}OBu_2$ according to the reaction.



Pyrolysis of $\text{B}_9\text{H}_{13}\text{OBu}_2$ gave $n\text{-B}_{18}\text{H}_{22}$ in a yield which represented a 45% conversion of B_5H_9 . This procedure for the preparation of $n\text{-B}_{18}\text{H}_{22}$ not only frees us of the requirement of $\text{B}_{10}\text{H}_{14}$, but it also rivals in terms of simplicity and yield other procedures¹⁸ for the preparation of this compound.

d. 5,6- $\phi_2\text{C}_2\text{B}_8\text{H}_{10}$. We have succeeded in preparing from B_5H_9 the the nido carborane 5,6- $\phi_2\text{C}_2\text{B}_8\text{H}_{10}$ in a "one-pot" synthesis.



Normally this nido carborane is prepared¹⁹ from a higher carborane system through a degradation process or from the degradation of B_9H_{15} or B_8H_{12} . These earlier methods were dependent upon $\text{B}_{10}\text{H}_{14}$ since it is the common starting point not only for the higher carboranes but has also been used¹⁰ to prepare $\text{B}_9\text{H}_{14}^-$ which is then converted to B_9H_{15} and B_8H_{12} .²⁰

References

1. Leach, J. B.; Toft, M.; Himpsl, F. L.; and Shore, S. G.; J. Am. Chem. Soc.; 1981, 103, 988.
2. Stock, A. and Mathing, W.; Ber. Deut. Chem. Ges.; 1936, 69, 1456.
3. Klein, M. J.; Harrison, B. C. and Solomon, I. J.; J. Am. Chem. Soc.; 1958, 80, 4149.
4. Jeffers, W.; Brit 853727, 1960; Chem. Abstr.; 1961, 26835.
5. Schaeffer, R. and Tebbe, F.; J. Am. Chem. Soc.; 1962, 84, 3974.
6. Gaines, D. F. and Schaeffer, R.; Inorg. Chem.; 1964, 3, 438.
7. Brown, H. C. and Tierney, P. A.; J. Am. Chem. Soc.; 1958, 80, 1552.
8. Shapiro, I.; Weiss, H. G.; Schmich, M.; Skolnik, S.; and Smith, G. B. L.; J. Am. Chem. Soc.; 1952, 74, 901.
9. Brown, H. C.; Murray, K. J.; Murray, L. J.; Snover, J. A.; and Zweifel, G.; J. Am. Chem. Soc.; 1960, 82, 4233.
10. Benjamin, L. E.; Stafiej, S. F. and Tackacs, E. A.; J. Am. Chem. Soc.; 1963, 85, 2674.
11. Brice, V. T.; Johnson, H. D.; Denton, D. L. and Shore, S. G.; Inorg. Chem.; 1974, 11, 1135.
12. Savory, C. G. and Wallbridge, M. G. H.; J. Chem. Soc. Dalton Trans.; 1973, 179.
13. Denton, D. L.; Clayton, W. R.; Mangion, J.; Shore, S. G. and Meyers, E. A. Inorg. Chem.; 1976, 15, 541.
14. Barton, L.; Onak, T. and Shore, S. G.; "Gmelin Handbuch der Anorganischen Chemie"; Springer-Verlag: Berlin, 1979; Vol. 54, Chapter 4. pp. 113-114.
15. Keller, P. C.; Himpsl, F. L.; Toft, M. A.; Boocock, S. K. and Shore, S. G.; "Abstracts of Papers"; 177th National Meeting of the American Chemical Society.
16. Dunks, G. B.; Barker, K.; Hedaya, E.; Hefner, D.; Ordonez, K.; Remec, P.; Inorg. Chem.; 1981, 20, 1962.
17. Dobson, J.; Keller, P. C. and Schaeffer, R.; Inorg. Chem.; 1968, 7, 399.
18. Barton, L.; Onak, T. and Shore, S. G.; "Gmelin Handbuch der Anorganischen Chemie"; Springer-Verlag: Berlin 1979, Vol. 54, Chapter 4. pp. 253-254.

19. a) Stibr, B.; Plesek, J. and Hermanek, S.; Coll. Czech Chem. Commun.; 1973, 38, 338. b) Plesek, J. and Hermanek, S.; ibid.; 1974. 39, 821 c) Rietz, R. R. and Schaeffer, R.; J. Am. Chem. Soc.; 1971, 93, 1263 d) Reitz, R. R. and Schaeffer, R.; ibid.; 1973, 95, 6254.
20. J. Dobson, P. C. Keller, and R. Schaeffer, Inorg. Chem., 1968, 7, 399.

Patents Applications

1. "Preparation of Decaborane 14 Through Hydride Ion Abstraction Reactions". Filed by U.S. Army Missile Command. Redstone Arsenal.
2. "A Dry Process for Producing Diborane. Filed by Ohio State University".

Special Technical Reports

Submitted to ARO

1. "Studies on the Conversion of Pentaborane(9) to Decaborane(14)" Part I.
2. "Studies on the Conversion of Pentaborane(9) to Decaborane(14)" Part II.
3. "Studies on the Conversion of Pentaborane(9) to Decaborane(14)" Part III.

Appendix

Publications

1. "New Systematic, Good Yield Syntheses of Boron Hydrides: Preparation of B_4H_{10} and B_5H_{11} . A Practical Conversion of B_5H_9 to $B_{10}H_{14}$ ", J. Am. Chem. Soc. 1981, 103, 988.
2. "New Systematic Syntheses of Boron Hydrides Via Hydride Ion Abstraction Reactions: Preparation of B_2H_6 , B_4H_{10} , B_5H_{11} and $B_{10}H_{14}$ ". Inorg. Chem. In press.
3. "Hexaborane(10) Derivatives: Relative Acidities of $2-CH_3B_6H_9$ and $2-BrB_6H_9$ and NMR Spectra of $2-CH_3B_6H_8^-$, $2-BrB_6H_8^-$, and $(THF)_2Mg(2-CH_3B_6H_8)_2$ ". Inorg. Chem. 1981, 20, 1270.
4. "Neutron and X-ray Diffraction Studies of Tris(methyl-diphenylphosphine) [tetrahydroborato(1-)]copper, $Cu[P(C_6H_5)_2(CH_3)_3(BH_4)]$. The First Accurate Characterization of an Unsupported Metal-Hydrogen-Boron Bridge Bond". J. Am. Chem. Soc., 1981, 103, 5165.

FILMED
7-8