


AD-A116 340


NATIONAL MARINE FISHERIES SERVICE PANAMA CITY BEACH F--ETC F/G 8/1
BENTHIC COMMUNITY RESPONSE TO DREDGING BORROW PITS, PANAMA CITY--ETC(U)
MAR 82 C H SALOMAN, S P NAUGHTON, J L TAYLOR DACW72-81-M-0198
CERC-MR-82-3 NL

UNCLASSIFIED

1-2

0000





12
MR 82-3


Benthic Community Response to Dredging Borrow Pits, Panama City Beach, Florida

by

Carl H. Saloman, Steven P. Naughton, and John L. Taylor

MISCELLANEOUS REPORT NO. 82-3

MARCH 1982



DTIC
SELECTED
JUL 1 1982
H

Approved for public release;
distribution unlimited.

Prepared for

**U.S. ARMY, CORPS OF ENGINEERS
COASTAL ENGINEERING
RESEARCH CENTER**

Kingman Building
Fort Belvoir, Va. 22060

82 07 01 012

AD A116340

DTIC FILE COPY

Reprint or republication of any of this material shall give appropriate credit to the U.S. Army Coastal Engineering Research Center.

Limited free distribution within the United States of single copies of this publication has been made by this Center. Additional copies are available from:

*National Technical Information Service
ATTN: Operations Division
5285 Port Royal Road
Springfield, Virginia 22161*

Contents of this report are not to be used for advertising, publication, or promotional purposes. Citation of trade names does not constitute an official endorsement or approval of the use of such commercial products.

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER MR 82-3	2. DTIC ACCESSION NUMBER AD-A116340	3. REPORT NUMBER'S CATALOG NUMBER
4. TITLE (and Subtitle) BENTHIC COMMUNITY RESPONSE TO DREDGING BORROW PITS, PANAMA CITY BEACH, FLORIDA	5. TYPE OF REPORT & PERIOD COVERED Miscellaneous Report	
7. AUTHOR(s) Carl H. Saloman Steven P. Naughton John L. Taylor	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS National Marine Fisheries Service, Southeast Fisheries Center, 3500 Delwood Beach Drive, Panama City Beach, Florida 32407	8. CONTRACT OR GRANT NUMBER(s) DACW72-81-M-0198	
11. CONTROLLING OFFICE NAME AND ADDRESS Department of the Army Coastal Engineering Research Center Kingman Building, Fort Belvoir, Virginia 22060	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS G31266	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	12. REPORT DATE March 1982	
	13. NUMBER OF PAGES 138	
	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release, distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Beach nourishment Dredging Benthos Panama City Beach, Florida		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) > This report gives biological and physical oceanographic data from base-line work, and studies of dredged and undredged sediments before and after dredging (9-meter contour) for beach nourishment at Panama City Beach, Florida. These studies were designed to show major short-term environmental effects of offshore dredging and included analyses of hydrology, sediments, and benthos. (continued)		

DD FORM 1 JAN 79 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

Hydrological measurements were limited to water temperature and salinity. Analysis of surface sediments included particle-size distribution, carbon chemistry, and statistical properties of mean grain size, sorting, skewness, and kurtosis. Average and extreme periods of water temperature and salinity were recorded. Regional nearshore sediments proved to be fine sand, containing less than 1 percent silt-clay, that was moderately well to well sorted, symmetrical to coarsely skewed, and leptokurtic. Total carbon content averaged less than 0.30 percent, and most of that occurred in the form of carbonate deposits. Over a postdredging study period of 1 year, sediment samples from borrow pits showed little variation from these general features.

In studies of the benthos, 362 species and 58,068 individuals were recorded among 14 invertebrate phyla and bony fishes. Dominant groups by species and abundance included annelida, mollusca, and arthropoda (crustacea). Faunal comparisons between dredged and undredged areas were made on the basis of species richness and abundance, the Shannon-Weaver index of diversity (H'), Pielou's index of equitability (J'), Morisita's index of faunal similarity (together with matrices and classification diagrams derived from that index), and two statistical derivations, based on diversity and abundance data, that were designed to show sample-to-sample faunal variations and the time period required for faunal recovery in borrow pits. Information obtained from these procedures showed that recovery began soon after dredging and was complete, or nearly so, within 1 year.

These results were similar in most respects to those from study of offshore dredging elsewhere in comparable geographic settings. Even so, the need for close association between ecological research and coastal engineering programs is emphasized.

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special



UNCLASSIFIED

PREFACE

This report gives preconstruction and postconstruction environmental data related to short-term effects of beach nourishment at Panama City Beach, Florida. Areas of study included water quality, sediments, and benthic invertebrates. Dredging and beach restoration were done by the U.S. Army Engineer District, Mobile, and research was sponsored by the U.S. Army Coastal Engineering Research Center (CERC), and by the National Marine Fisheries Service (NMFS), Gulf Fisheries Center, Panama City Beach, Florida. The work was carried out under the coastal ecology research program.

The report is based on data collected and compiled by Carl H. Saloman and Steven P. Naughton, NMFS, who assisted Dr. John L. Taylor, Taylor Biological Company, Inc., in preparing the report under CERC Contract No. DACW72-81-M-0198. Invaluable assistance with statistical programs and data processing was provided by Dr. S.A. Bloom, Department of Zoology, University of Florida, Gainesville. Editorial reviews were provided by E. Nakamura, NMFS, and by B. Hall, CERC.

The authors acknowledge the assistance of their colleagues for identification of the following faunal groups: Dr. R.W. Heard, Jr., Gulf Coast Research Laboratory, Ocean Springs, Mississippi (crustacea); and J.R. Hall, National Marine Fisheries Service, Washington, D.C. (mollusca). Identification of species in other groups was done by the authors with the aid of reference material available from NMFS.

E.J. Pullen, Chief, Coastal Ecology Branch, served as contract monitor for this report, under the general supervision of R.P. Savage, Chief, Research Division; he also assisted in the editorial review process and made arrangements for several technical aspects of manuscript preparation and publication.

Comments on this publication are invited.

Approved for publication in accordance with Public Law 166, 79th Congress, approved 31 July 1945, as supplemented by Public Law 172, 88th Congress, approved 7 November 1963.


TED E. BISHOP

Colonel, Corps of Engineers
Commander and Director

CONTENTS

	Page
CONVERSION FACTORS, U.S. CUSTOMARY TO METRIC (SI)	6
I INTRODUCTION.	7
1. Background	7
2. Purpose.	8
II STUDY AREA.	9
III SAMPLING STATIONS AND RATIONALE	9
IV SAMPLING AND ANALYTICAL PROCEDURES.	12
1. Hydrology.	12
2. Sedimentology.	12
3. Benthos.	13
V RESULTS	14
1. General.	14
2. Hydrology.	15
3. Sedimentology.	15
4. Benthos.	18
VI CONCLUSIONS AND DISCUSSION.	27
LITERATURE CITED.	30
APPENDIX	
A HYDROLOGICAL AND SEDIMENT DATA BY STATION	31
B CHECKLIST OF ORGANISMS.	42
C BIOLOGICAL AND BIOSTATISTICAL DATA BY STATION	46
D FAUNAL SIMILARITY MATRICES.	131
E FAUNAL CLASSIFICATION ANALYSES.	134
F STABILITY ANALYSES.	137

TABLES

1 Water temperature and salinity at stations A and B before the 1974-75 dredging, and at station 1 before and after the 1976 dredging for beach nourishment at Panama City Beach, Florida	16
2 Textural and statistical properties of sediments in control (undredged bottom) and experimental (borrow pit) samples taken 1 year after dredging at stations 1 to 6 along the 9-meter depth contour off Panama City Beach, Florida, July 1977	16
3 Species in dominant phyla (listed alphabetically) that were numerically abundant at one or more base-line or control stations offshore Panama City Beach, Florida, November 1974 to November 1977	19

CONTENTS

TABLES--Continued

	Page
4 Species richness, abundance, diversity (H'), and equitability (J') and base-line stations offshore Panama City Beach, Florida, November 1974 to July 1976.	19
5 Species richness, abundance, diversity (H'), and equitability (J') at control stations offshore Panama City Beach, Florida, August 1976 to November 1977.	20
6 Species richness, abundance, diversity (H'), and equitability (J') at experimental stations offshore Panama City Beach, Florida, August 1976 to November 1977	21
7 Species and their frequency of occurrence in the first 3 weeks after dredging at station 1 offshore Panama City Beach, Florida, August 1976	23

FIGURES

1 Study area at Panama City Beach, Florida, showing stations 1 to 6, July 1977	10
2 Schematic representation of sampling plan, Panama City, Florida.	11

CONVERSION FACTORS, U.S. CUSTOMARY TO METRIC (SI) UNITS OF MEASUREMENT

U.S. customary units of measurement used in this report can be converted to metric (SI) units as follows:

Multiply	by	To obtain
inches	25.4	millimeters
	2.54	centimeters
square inches	6.452	square centimeters
cubic inches	16.39	cubic centimeters
feet	30.48	centimeters
	0.3048	meters
square feet	0.0929	square meters
cubic feet	0.0283	cubic meters
yards	0.9144	meters
square yards	0.836	square meters
cubic yards	0.7646	cubic meters
miles	1.6093	kilometers
square miles	259.0	hectares
knots	1.852	kilometers per hour
acres	0.4047	hectares
foot-pounds	1.3558	newton meters
millibars	1.0197×10^{-3}	kilograms per square centimeter
ounces	28.35	grams
pounds	453.6	grams
	0.4536	kilograms
ton, long	1.0160	metric tons
ton, short	0.9072	metric tons
degrees (angle)	0.01745	radians
Fahrenheit degrees	5/9	Celsius degrees or Kelvins ¹

¹To obtain Celsius (C) temperature readings from Fahrenheit (F) readings, use formula: $C = (5/9) (F - 32)$.

To obtain Kelvin (K) readings, use formula: $K = (5/9) (F - 32) + 273.15$.

BENTHIC COMMUNITY RESPONSE

TO DREDGING BORROW PITS,

PANAMA CITY BEACH, FLORIDA

by

Carl H. Saloman, Steven P. Naughton,

and

John L. Taylor

I. INTRODUCTION

1. Background.

On the gulf coast of northwestern Florida, at Panama City Beach, major environmental alterations over the past 10 years have provided an exceptional opportunity to determine the degree and duration of these alterations associated with the practice of dredging and beach nourishment. Historically, these events have included the development of several engineering plans, the intervention of a major hurricane, an emergency dredging and beach restoration program, and several ecological studies related to disturbances caused by both the hurricane and the dredging.

In 1970, the Senate Committee on Public Works acknowledged an urgent need for beach erosion control and hurricane protection at Panama City Beach. This critical situation was referred to the U.S. Army Engineer District, Mobile, for study. In 1975, the Mobile District completed a feasibility report that contained recommendations for beach nourishment and maintenance along 29.8 kilometers of shoreline from the entrance to St. Andrew Bay, west to Philips Inlet (Wilson, 1975). During preparation of the report, the U.S. Army Coastal Engineering Research Center (CERC) sponsored a research program to determine ecological changes that could be expected from the dredging and coastal construction work. This investigation, which was conducted by the National Marine Fisheries Service (NMFS) between November 1974 and October 1975, involved the study of hydrology, sediments, and benthic fauna at two offshore stations, and at five stations on each of nine nearshore transects. Emphasis was placed on diversity, abundance, and distribution of bottom-dwelling invertebrates which are directly affected by dredging and redistribution of sediments (Saloman, 1976).

Before this investigation was completed, Hurricane Eloise struck Panama City Beach (25 September 1975). Winds up to 185 kilometers per hour and seas estimated at 9 meters caused severe erosion and extensive property damage (Saloman, 1976; Salsman and Ciesluk, 1978). In winter months that followed, high wind and waves associated with periodic cold fronts caused further shoreline erosion.

In anticipation of the storm, and realizing the opportunity to measure large-scale environmental changes alongshore, NMFS conducted an intertidal benthic survey that consisted of faunal sampling before the storm and during a 1-month period after the storm. The pattern of faunal disruption and recovery recorded in this unique study provided considerable insight into the sequence of population changes to be expected in the proposed beach nourishment program (Saloman and Naughton, 1977).

In the next year (July-August 1976), the Corps of Engineers funded an emergency dredging operation to restore the most ravaged beach areas and established berms to provide temporary protection against storms normally occurring during fall and winter seasons. Numerous borrow areas, 305 to 610 meters offshore (6- to 9-meter depth) were dredged and about 306,000 cubic meters of sand was pumped ashore at 23 distribution sites (U.S. Army Engineer District, Mobile, 1976).

At the same time, NMFS again conducted studies of the nearshore environment over a 3-month period prior to dredging, during dredging, and for about 6 months after dredging was completed. Benthic sampling sites were selected in nourishment areas and in unrestored areas. The location of the three nourishment areas coincided with the location of benthic base-line data collected in 1974 (Saloman and Naughton, unpublished data).

Based on emergency nourishment experience and the analysis of the Hurricane Eloise data collected, the Mobile District revised original plans for shoreline protection and maintenance at Panama City Beach. The revised plan included berm enlargement on the beach front and additions to height and width of backbeach dunes. Consequently, the volume of sand estimated for original construction was increased from 4 to 8 million cubic meters; and borrow areas formerly selected at 9-meter depths were relocated seaward along the 18-meter bottom contour (Wilson, 1976).

Onshore, the environmental impact of this latest plan can probably be predicted to a high degree of accuracy on the basis of findings in NMFS beach surveys in 1974 and 1976. Briefly stated, the results of these investigations showed that shallow, subtidal and intertidal faunas recover rapidly following major disturbances (natural or man-induced). A more recent study funded by CERC provides additional information on the long-term environmental effects of dredging in offshore borrow areas at Panama City Beach (Culter and Mahadevan, 1982). A study of short-term environmental effects of dredging in offshore borrow areas at Panama City Beach is the subject of the present report.

2. Purpose.

This report provides a comprehensive analysis of benthic data from studies designed to show short-term environmental effects of offshore dredging during the emergency restoration project at Panama City Beach in July-August 1976.

It is based on comparisons of hydrological, sedimentological, and biological data from collections at stations A and B in base-line studies that began in 1974 (Saloman, 1976), and from control and experimental samples taken by NMFS in undredged bottom and borrow areas over a 20-month period between April 1976 and November 1977.

II. STUDY AREA

Panama City Beach is located on the northwestern gulf coast of Florida about 145 kilometers east of Pensacola. The study area covers 35 kilometers and extends from West Pass at the entrance to St. Andrew Bay, to Philips Inlet (Fig. 1). The beach's sugarlike sand and exceptionally clear water are major attractions for about 2 million visitors annually. Tourism is a great economic asset and most of the beach has been developed to accommodate tourists and provide various types of recreation.

Regional meteorological and oceanographic conditions were described by Salsman and Ciesluk (1978). Climate is humid and subtropical. Average summer and winter air temperatures are 28° and 12° Celsius, with about the same water temperatures at respective seasons. Winds are 20 kilometers per hour or less at most times, and rarely exceed 37 kilometers per hour. From spring through late summer, the net wind direction is southerly, but between September and January, the direction shifts to northerly. Waves are usually about 0.9 meter; tides are diurnal, and tidal amplitude is normally about 0.6 meter; and tidal currents are generally below 4 kilometers per hour. However, during tropical storms and ahead of cold fronts, strong winds off the gulf produce waves, tides, and currents far greater than average. Even in less severe weather, beach sand is easily eroded because of its fine texture (0.1- to 0.2-millimeter median diameter). Seaward, a series of parallel sandbars protects the beach to some extent, but beyond, the featureless bottom slopes rather quickly to a 15-meter depth at 1.6 kilometers from shore. At greater depths, sediments are somewhat coarser and widely scattered limestone reefs appear in low relief.

III. SAMPLING STATIONS AND RATIONALE

The sampling data in this report were collected in about 9 meters of water at stations located offshore of Panama City Beach. As a matter of convenience, and for clarity, these stations have been separated into three groups since there were differences in their locations, sampling procedures, and objectives.

The first group includes stations A and B (Fig. 2) of the preconstruction investigation of 1974-75. Station A was located seaward of the Fiesta Motel about midway between West Pass and Philips Inlet. Station B was seaward of the Roundtowner Motel, which is just east of Philips Inlet. The sampling schedule at these stations consisted of an initial collection in November 1974, and subsequent quarterly collections in February, May, and August 1975. Both were sampled before beach nourishment to determine seasonal environmental conditions (base-line data) in the zone designated for dredging (Saloman, 1976).




Figure 1. Study area at Panama City Beach, Florida, showing stations 1 to 6, July 1977.




Figure 2. Schematic representation of sampling plan, Panama City, Florida.

The second group contains station 1 (Fig. 2), located seaward of Treasure Island Motel (near the eastern end of the study area), which had two collecting areas--one at the borrow site and the other a short distance away on undredged bottom. Samples were taken from the designated borrow site before dredging in April, June, and July 1976. Then 2 days after dredging (10 August 1976), concurrent sampling was started inside and outside the borrow pit. Sampling in both the pit (experimental samples) and adjacent to it (control samples) continued on a weekly schedule for 1 month. Samples were taken twice the next month, and then monthly thereafter until the study was concluded in November 1977. These samples were collected to record diversity and abundance of benthic fauna at a specific dredge site before dredging started, and then, over time, to compare population characteristics of control samples with experimental samples.

The third group includes stations 1, 2, 3, 4, 5, and 6 for one-time sampling only inside and outside borrow pits during July 1977--about 12 months after dredging (Fig. 2). The six stations were located seaward of the following landmarks: station 1, Treasure Island Motel; station 2, Sun and Swim Motel; station 3, Hilton Holiday Inn; station 4, Sandpiper Motel; station 5, Peppertree Condominium (now Landmark Condominium); and station 6, Blue Dolphin Motel. These collections provided a comparison of fauna in control and experimental samples from a number of borrow pits for an evaluation of short-term recovery within a period of 1 year. Throughout this report, samples from stations A and B, and preconstruction samples from station 1, are referred to as baseline or control samples; all other samples from outside borrow pits are called control samples, and all samples from within borrow pits are designated experimental samples.

IV. SAMPLING AND ANALYTICAL PROCEDURES

1. Hydrology.

Surface water temperature and salinity measurements were recorded in each sampling period at stations A and B, and on a monthly schedule over the duration of sampling at station 1. Temperature was taken using a hand-held, mercury bulb thermometer graduated in Celsius degrees. Salinity, in parts per thousand, was determined with a Goldberg temperature-compensated refractometer (American Optical Co., Model No. 10419).

2. Sedimentology.

Sediment samples were collected to determine textural features, statistical properties, and carbon chemistry. Textural parameters included weight percentages of granules, sand, and silt-clay. Mean grain size, standard deviation (as a measure of sorting), skewness, and kurtosis were calculated and interpreted according to the system described by Folk (1974). The carbon analyses included total carbon, total organic carbon, and total carbonate carbon.

Collections were limited to surface samples that included the upper 10 centimeters of sediment. Sediments were collected in standard 8-ounce, screw-cap jars; all samples were stored frozen prior to analyses. Detailed analytical methods are described by Saloman (1976).

For textural analyses, sediment samples were sieved at 1-phi intervals in nested screens placed on a mechanical shaker. Fraction weights were recorded to the closest milligram and tabulated as weight percentages. No hydrometer or pipette determinations were required because silt-clay percentages were quite low. Based on grain-size distribution curves, formulas introduced by Folk (1974) were used to calculate statistical properties. Carbon analyses were made using a Leco 750-100, 90-second carbon analyzer.

Additionally, divers recorded observations of sediment inside and outside the borrow pit at station 1. These observations were made on a regular basis during the first postconstruction collection, and in subsequent collections, until the study ended.

3. Benthos.

At all collecting points, infauna was sampled with a hand-operated plug sampler (box corer) that covered a surface area of 1/64 square meter and penetrated the bottom to a depth of 23 centimeters (Saloman, 1976). Replicate samples were taken at each site, but the number was not always the same for each of the three station groups. At stations A and B, four replicates composed a sample (preconstruction base-line study of 1974-75). At station 1, the first collection contained 32 replicates (19 April 1976), while second and third preconstruction samples each consisted of 36 replicates. After dredging, however, both control and experimental samples from station 1 each included 16 replicates. Finally, in the one-time collection at stations 1 to 6, 1 year after dredging, control and experimental samples were each composed of 40 replicates. The decision to take more than 4 replicates in most samples was somewhat arbitrary, since sampling to develop a species rarefaction curve showed that 4 plugs comprised an adequate qualitative and quantitative sample of the nearshore benthos (Saloman, 1976). For reference, a schematic representation of the overall sampling plan was prepared to show geographic relationships among stations within the study area, landmarks along the shore, pertinent transect locations from studies started in 1974, and the sampling locations of borrow pits and undredged bottom studied between April 1976 and July 1977 (Fig. 2).

All benthic samples were taken by scuba divers and sieved on shipboard in a 0.3-meter diameter screen of 0.7 millimeter mesh. Material remaining on the screen was preserved with 10-percent seawater formalin in standard 2-quart, screwcap jars. Rose bengal dye was added to the formalin to stain organisms and facilitate their subsequent separation from debris. In the laboratory,

each collection was resieved under tapwater and all specimens from respective samples were stored in 70-percent isopropanol for final sorting, taxonomic determinations, and species counts. The 0.7 millimeter screen was used instead of a conventional 0.5 millimeter one because the former facilitated sieving operations and retained a percentage of infauna that was shown to be very nearly equivalent to that sampled by the smaller mesh size.

As in Saloman's (1976) work, biological data presented here include a species checklist and individual station listings that show species occurrence and frequency, together with calculations for number of individuals per square meter and the Shannon-Weaver index of faunal diversity (H'). Also, as a measure of relative species dominance, equitability (J') was computed for each station (Pielou, 1975). Two other statistical procedures were also employed. The first, Morisita's Index (Morisita, 1959; Bloom, 1981), provided a numerical method of comparing faunal similarity between comparable sets of control and experimental samples, and was used to develop similarity matrices and classification diagrams that graphically show faunal relationships based on station data for diversity and abundance.

The second procedure, a stability analysis (Bloom, 1980), is a multivariate, nonparametric statistical and geometric procedure that converts biotic data from control and experimental samples into communities that can be represented mathematically. For one representation all base-line and control data were used to define numerical characteristics of a preconstruction community cluster that has a central point, or centroid, and certain specific spatial limits. In the first stability analysis, the distance from the centroid to control and experimental samples was used to determine variability among samples from undredged and dredged bottoms. In the second analysis, community clusters calculated for experimental samples were compared to the preconstruction cluster, in postconstruction sequence. When a boundary or an experimental cluster met the limit of the preconstruction cluster, faunal recovery was accepted. Experimental collections from station 1, where sampling over time was done, were the only borrow pit samples used in this analysis.

V. RESULTS

1. General.

The findings in this section are based on the detailed information given in Appendixes A to F. Appendix A lists abiotic parameters by station. Appendix B is a checklist of all organisms collected at offshore stations from November 1974 to November 1977. Appendix C contains all biological station data and indices of diversity (H') and equitability (J'). Appendix D (Similarity Matrices) and E (Classification Analyses and Dendrograms) are both based on Morisita's index of faunal similarity. Appendix F is a graphic representation of the two stability analyses. The first graph shows comparative variability among control and experimental samples when compared with the centroid of a community cluster calculated from all base-line and control samples. The second

is a stability plot for experimental samples from station 1 showing the post-construction time lapse before faunal recovery appears evident.

2. Hydrology.

Water temperature and salinity data from the 1974-75 sampling at stations A and B were compared to data from station 1 sampled during similar months in 1976-77 (Table 1). Both sets of data show normal seasonal trends in water temperature, except for one abnormally low value of 9° Celsius recorded in February 1977.

Salinity was low at stations A and B in August 1975, but salinity during other months was 32 parts per thousand or higher, and similar to station 1 records (Table 1). Appreciable declines in salinity apparently coincide with periods of seasonally heavy rainfall.

3. Sedimentology.

The influence of dredging on sediment composition was determined by analyses of base-line and control samples, compared to samples taken from borrow pits. Base-line data came from seasonal sediment collections at stations A and B, and from those taken before dredging at station 1 in April, June, and July 1976. Control data were available from samples outside the borrow pit at station 1, and from samples collected in an undredged bottom at stations 1 to 6 in July 1977. Data from experimental samples also came from periodic collections at station 1, and from borrow pit collections in the single survey in July at stations 1 to 6.

Textural, statistical, and chemical properties of base-line samples (Table 2) were used to describe natural features of offshore sediments, since these samples were collected in all seasons prior to dredging at eastern, central, and western locations within the study area (see App. A).

a. Texture. Sediment composition was about 99-percent sand, and both granules and silt-clay size particles contributed less than 1 percent.

b. Statistical Properties. Values for mean grain size, standard deviation, skewness, and kurtosis classified these sediments as fine sand that is moderately well to well sorted, symmetrical to coarsely skewed, and leptokurtic (sorted better in the center than at the ends of grain size distribution curves).

c. Carbon Chemistry. Total carbon content of base-line samples was less than 0.30 percent. Carbonate carbon contributed somewhat more to this total than organic carbon, indicating that most carbon occurred in the form of shell fragments rather than as organic deposits.

For station 1, when these features were compared to control and experimental samples, noteworthy differences appeared only in experimental samples.

Table 1. Water temperature and salinity at stations A and B before the 1974-75 dredging, and at station 1 before and after the 1976 dredging for beach nourishment at Panama City Beach, Florida.

Station	Date	Water Temp. (°C)	Salinity (ppt)
<u>1974</u>			
A	18 Nov.	21.0	34.5
B	18 Nov.	20.8	34.3
<u>1975</u>			
A	20 Feb.	17.4	34.4
B	20 Feb.	17.5	33.9
A	20 May	26.2	32.2
B	20 May	26.0	32.2
A	12 Aug.	28.3	26.2
B	12 Aug.	28.5	26.1
<u>1976</u>			
1 (before)	Apr.	20.2	33.3
	May	20.2	34.9
	June	25.7	32.3
	July	28.0	33.3
	Aug.	27.0	35.3
	Sept.	27.8	32.6
	Oct.	24.9	33.1
	Nov.	18.0	33.2
	Dec.	12.5	34.1
<u>1977</u>			
1 (after)	Jan.	12.4	33.3
	Feb.	9.0	34.3
	Mar.	14.3	34.4
	Apr.	22.4	33.5
	May	21.8	34.3
	June	25.7	32.1
	July	27.5	33.6
	Aug.	29.0	35.3
	Sept.	27.7	32.6
	Oct.	25.0	33.1
	Nov.	-	-

Table 2. Textural and statistical properties of sediments in control (undredged bottom) and experimental (borrow pit) samples taken 1 year after dredging at stations 1 to 6 along the 9-meter depth contour off Panama City Beach, Florida, July 1977.

Station	Textural			Statistical			
	Granule (pct)	Sand (pct)	Silt-clay (pct)	Mean grain size (phi)	Std. dev. (phi)	Skewness	Kurtosis
1							
Control		99.70	0.30	2.45	0.45	-0.19	1.18
Experimental		98.64	1.36	2.50	0.53	-0.00	1.39
2							
Control		99.65	0.35	2.45	0.44	-0.18	1.15
Experimental		99.80	0.20	2.43	0.48	-0.19	1.21
3							
Control		99.88	0.12	2.21	0.62	-0.32	1.11
Experimental	0.92	98.96	0.11	1.75	1.06	-0.46	0.82
4							
Control		99.86	0.14	2.24	0.61	-0.31	1.16
Experimental	0.08	99.81	0.11	2.01	0.83	-0.41	0.95
5							
Control		99.86	0.14	2.31	0.59	-0.33	1.34
Experimental		99.86	0.14	2.26	0.58	-0.28	1.11
6							
Control	0.34	99.52	0.14	2.11	0.76	-0.40	1.09
Experimental	0.14	99.76	0.11	2.31	0.61	-0.34	1.39

The particle-size distribution of sand was below 99 percent in experimental samples from September, October, and November 1976, and from January, June, July, August, and September 1977. The lowest level (92 percent) was recorded in September 1976. Other low values were only in the 97- to 98-percent range. Granule-size particles were consistently under 1 percent, but 11 experimental samples contained more than 0.30-percent silt-clay. The highest value for the silt-clay fraction was 8.1 percent in a sample collected on 21 September 1976. Values of more than 1-percent silt-clay were also recorded in another September sample as well as in October and November 1976, and again in January, June, July, August, and September 1977.

Mean grain size for experimental samples did not range below fine sand. Sorting categories changed for two experimental samples. In the September 1976 sample, sorting was only moderate; in the May 1977 sample, it proved to be extremely poor. For skewness, five experimental samples exhibited an uncharacteristic trend that placed them in classifications of fine skewed to strongly fine skewed. The single sample classified as strongly fine skewed was obtained in September 1976; the others were collected in September and October 1976, and August and September 1977. Deviation from the normal leptokurtic condition was recorded for five experimental samples. Values corresponding to mesokurtic were recorded in August 1976, and April and May 1977. Values in the very leptokurtic range were recorded in January and June 1977.

A carbon content percentage greater than the base-line average was recorded in 12 experimental samples; however, this number of samples may be low since no carbon analyses were made after the June 1977 sampling. The highest recorded value was 2.32 percent for the September 1976 sample. Other slightly elevated values ranged between 0.31 and 1.21 percent. Among these 12 samples, the proportion of organic carbon to carbonate carbon was higher for carbonate in 6 samples, higher for organic in 5, and in 1, the ratio was nearly even.

Sediment data for control and experimental samples collected at the six stations in July 1977 has been tabulated for comparison (Table 2). These analyses include only textural and statistical properties; no information on carbon chemistry was available.

At the six stations, granule-size particles were present in only four samples, and three of these came from borrow pits at stations 3, 4, and 6. The single control sample containing granules also came from station 6, and the overall granule distribution was under 1 percent. Sand content was about 99 percent in all collections. For the silt-clay fraction, only one value was considered abnormally high and that was recorded for the experimental sample from station 1 (1.36 percent).

With the one exception of medium sand (station 3, experimental), all samples fell into the classification of fine sand. Calculations for sorting showed that 9 of 12 samples were well to moderately well sorted. Other classifications included moderately sorted (station 4, experimental and station 6, control) and

poorly sorted (station 3, experimental). Skewness values were characteristic for five samples (symmetrical to coarsely skewed), and the other seven samples fit the strongly coarse-skewed classification and were about equally divided between the control and experimental samples. The normal, or leptokurtic condition, was found in nine samples. Of the remaining three, the experimental sample from station 4 and the control sample from station 6 were mesokurtic, while the experimental sample from station 3 was platykurtic.

Although sedimentological conditions in some experimental samples varied from the base-line criteria until late 1977, large variations were confined to borrow pit sediments at station 1 within 2 months after dredging. During that period, properties which may have been limiting to benthos were high silt-clay and organic carbon content.

Diver reports between 18 August (10 days after dredging) and 4 October 1976, stated that the station 1 borrow pit was 3 to 5 meters deep and had very dark surface sediments of an extremely soft, silty texture. Initially no surface signs of benthic life (burrows, mounds, or trails) were reported. Within the next month, sediments had become firmer and sandier; signs of infauna activity were conspicuous, crabs and other epibenthos were numerous, and a variety of fishes was observed. After 12 months, and on the last dive at station 1 in November 1977, divers concluded that borrow pits had filled to within a meter of surrounding bottom, and that sediments inside were still finer, darker, and less compact than sediments outside, but marine life appeared similar in control and experimental areas.

4. Benthos.

The checklist of organisms in Appendix B contains about 362 organisms at the species level, representing 14 invertebrate phyla and the vertebrate class, Osteichthyes (bony fishes). Of this number, Annelida had 152 species (42 percent), Arthropoda had 108 (30 percent), and there were 69 mollusks (19 percent). The remaining 33 species (9 percent) were divided among 11 groups: Cnidaria, Platyhelminthes, Nemertinea, Nematoda, Phoronida, Brachiopoda, Sipunculida, Echiurida, Echinodermata, Hemichordata, and Cephalochordata.

Species counts from each station showed a total of 58,068 individuals collected. On a percentage basis, more than half were annelids (55 percent), 19 percent were mollusks, 18 percent were arthropods, Cnidaria and Cephalochordata each accounted for 2 percent, Nematoda and Echinodermata both had 1 percent, and the other seven groups contained 2 percent, collectively. For the three major phyla, species that were numerically dominant in one or more of the base-line or control site collections are given in Table 3.

All station data for richness, quantitative abundance, diversity (H'), and equitability (J') were tabulated by base-line, control, and experimental sample categories (Tables 4, 5, and 6). Graphic analyses of Morisita's Index and stability are given in Appendixes D, E, and F.

Table 3. Species in dominant phyla (listed alphabetically) that were numerically abundant at one or more base-line or control stations offshore Panama City Beach, Florida, November 1974 to November 1977.

MOLLUSCA

Acteocina candel
Cylichnella bidentata
Diastrum varium
Ervilla concentrica
Lepton sp.
Lucina multilineata

Natica pusilla
Periploma margaritaceum
Pitar simpsoni
Strigilla mirabilis
Tellina texana
Tellina versicolor

ANNELIDA

Armandia agilis
Armandia maculata
Brania wellfleetensis
Ceratonereis irritabilis
Chone sp.
Dispio uncinata
Eteone lactea
Glycera americana
Goniada littorea
Haploscoloplos foliatus
Lumbrineris cruzensis
Lumbrineris tenuis
Lumbrineris tetraodon
Magelona riojai
Magelona sp.
Mesochaetopterus longitarsus

Nephtys buccera
Nephtys picta
Onuphis eremita oculata
Onuphis nebulosa
Owenia fusiformis
Paranides lyra
Paranides fulgens
Paraprionospio pinnata
Prionospio cristata
Rullierineris mexicana
Scolelepis texana
Scoloplos armiger
Spio pettiboneae
Spiophanes bombyx
Unidentified Oligochaete

ARTHROPODA

Acanthohaustorius sp.
Albunea paretii
Ampelisca abdita
Ampelisca verrilli
Cyclaspis varians
Cyclaspis sp.
Erichthonius sp.
Lepidactylus sp.

Monoculodes sp.
Oxyurostylis smithi
Processa hemphilli
Protohaustorius sp.
Pseudohaustorius sp.
Pseudoplatyschnopus sp.
Syncheildium sp.
Unidentified Ostracod

Table 4. Species richness, abundance, diversity (H'), and equitability (J') and base-line stations offshore Panama City Beach, Florida, November 1974 to July 1976.

Station	Date	Replicates per sample (No.)	Species (No.)	Individuals per m ² (No.)	H'	J'
A	Nov. 1974	4	15	2,064	1.9	0.7
	Feb. 1975		27	3,008	2.2	0.7
	May 1975		41	4,784	2.8	0.8
	Aug. 1975		43	3,888	3.1	0.8
	Avg.		32	3,436	2.5	0.8
	Range		15 to 43	2,064 to 4,784	1.9 to 3.1	0.7 to 0.8
B	Nov. 1974	4	27	3,808	1.9	0.6
	Feb. 1975		26	3,984	2.3	0.7
	May 1975		28	5,344	2.3	0.7
	Aug. 1975		47	5,248	3.0	0.8
	Avg.		32	4,596	2.4	0.7
	Range		26 to 47	3,808 to 5,344	1.9 to 3.0	0.6 to 0.8
1	Apr. 1976	32	67	1,506	2.5	0.6
	June 1976	36	94	1,902	3.5	0.8
	July 1976	36	120	7,178	3.1	0.6
	Avg.		94	3,529	3.0	0.7
	Range		67 to 120	1,506 to 7,178	2.5 to 3.5	0.6 to 0.8
	Overall					
	Avg.		49	3,883	2.6	0.7
	Range		15 to 120	1,506 to 7,178	1.9 to 3.5	0.6 to 0.8

Table 5. Species richness, abundance, diversity (H'), and equitability (J') at control stations offshore Panama City Beach, Florida, August 1976 to November 1977.

Station	Date	Replicates per sample (No.)	Species (No.)	Individuals per m ² (No.)	H'	J'
1	10 Aug. 1976	16	72	5,576	2.4	0.6
	18 Aug. 1976		80	5,500	2.8	0.6
	24 Aug. 1976		84	4,836	2.9	0.6
	1 Sept. 1976		74	3,080	2.9	0.7
	8 Sept. 1976		83	2,260	3.4	0.8
	21 Sept. 1976		89	3,128	3.0	0.7
	4 Oct. 1976		87	3,116	3.3	0.7
	18 Oct. 1976		77	3,912	2.6	0.6
	1 Nov. 1976		67	3,020	2.6	0.6
	1 Dec. 1976		74	3,080	3.0	0.7
	5 Jan. 1977		56	1,724	3.0	0.8
	2 Feb. 1977		53	1,516	3.1	0.8
	1 Mar. 1977		64	2,360	3.1	0.7
	1 Apr. 1977		57	2,632	3.1	0.8
	2 May 1977		55	2,572	2.7	0.7
	1 June 1977		55	1,976	3.3	0.8
	5 July 1977		64	3,264	3.1	0.7
	2 Aug. 1977		80	5,168	3.0	0.7
	1 Sept. 1977		70	3,572	2.9	0.7
	3 Oct. 1977		64	2,112	2.8	0.7
	1 Nov. 1977		72	2,904	3.0	0.7
Avg.			70	3,205	3.0	0.7
Range			53 to 89	1,515 to 5,576	2.4 to 3.3	0.6 to 0.8
1	11 Jul. 1977	40	99	3,365	3.2	0.7
2	15 Jul. 1977	40	112	3,750	3.4	0.7
3	25 Jul. 1977	40	105	4,326	3.2	0.7
4	26 Jul. 1977	40	74	4,050	2.9	0.7
5	27 Jul. 1977	40	57	1,408	3.0	0.7
6	28 Jul. 1977	40	66	2,483	3.0	0.7
Avg.			86	2,817	3.1	0.7
Range			57 to 112	1,408 to 4,326	2.9 to 3.4	0.6 to 0.8
<u>Overall</u>						
Avg.			74	3,119	3.0	0.7
Range			53 to 112	1,408 to 5,576	2.4 to 3.4	0.6 to 0.8

Table 6. Species richness, abundance, diversity (H'), and equitability (J') at experimental stations offshore Panama City Beach, Florida, August 1976 to November 1977.

Station	Date	Replicates per sample (No.)	Species (No.)	Individuals per m ² (No.)	H'	J'
1	10 Aug. 1976	16	20	324	2.0	0.7
	18 Aug. 1976		38	976	2.2	0.6
	24 Aug. 1976		60	2,136	2.6	0.6
	1 Sept. 1976		38	1,612	2.1	0.6
	8 Sept. 1976		47	1,344	2.7	0.7
	21 Sept. 1976		45	924	2.9	0.8
	4 Oct. 1976		85	2,440	3.7	0.8
	18 Oct. 1976		46	1,124	2.9	0.8
	1 Nov. 1976		55	2,044	2.5	0.6
	1 Dec. 1976		54	3,540	2.3	0.6
	5 Jan. 1977		36	2,192	1.8	0.5
	2 Feb. 1977		44	2,212	1.9	0.5
	1 Mar. 1977		62	3,732	2.6	0.6
	1 Apr. 1977		52	3,144	2.2	0.6
	2 May 1977		54	1,656	2.8	0.7
	1 June 1977		69	3,256	3.2	0.8
	5 July 1977		49	1,964	2.7	0.7
	2 Aug. 1977		70	2,920	3.2	0.8
	1 Sept. 1977		32	440	2.9	0.8
	3 Oct. 1977		61	1,588	3.1	0.8
	1 Nov. 1977		54	1,220	2.9	0.7
Avg. Range		51 20 to 85	1,942 324 to 3,732	2.6 1.8 to 3.7	0.7 0.5 to 0.8	
1	11 July 1977	40	81	2,422	2.9	0.7
2	15 July 1977	40	114	3,862	3.5	0.7
3	25 July 1977	40	98	4,037	3.3	0.7
4	26 July 1977	40	94	2,587	3.4	0.8
5	27 July 1977	40	80	2,644	2.9	0.7
6	28 July 1977	40	83	3,034	3.4	0.8
Avg. Range		92 80 to 114	3,101 2,422 to 4,037	3.2 2.9 to 3.5	0.7 0.7 to 0.8	
<u>Overall</u>						
Avg. Range			60 20 to 114	2,200 324 to 4,037	2.8 1.8 to 3.7	0.7 0.5 to 0.8

a. Richness. The data from base-line and control samples indicate that species richness followed an irregular seasonal pattern. Generally, numbers of species were lowest in a period between late fall and spring, and showed one or more peaks sometime between midsummer and late fall.

For base-line collections (Table 4), the number of species per sample averaged 49 and ranged between 15 (November) and 120 (July). The average for control samples was 74 and ranged between 53 (February) and 112 (July). Intermediate values were recorded for experimental samples. In these collections, average number of species per sample was 60; the low, which was only 20, occurred in the first collection after dredging; the high was 114, recorded in July 1 year later.

On a date-to-date comparison at station 1 and stations 1 to 6, richness data for control and experimental samples (Tables 5 and 6) gave somewhat conflicting results. For time-sequence samples at station 1, richness data showed incomplete borrow pit recovery as numbers of species prove to be consistently higher for controls on every occasion except 1 June 1977. This was reflected in the average of 70 and the range between 53 and 89 for control samples, as opposed to an average of 51 and a range of 20 to 85 for experimental samples. Even so, a degree of recovery was evident at station 1 a few weeks after dredging, and richness data for control and experimental samples first approximated one another by October 1976. Species recorded in the early stages of recovery at station 1 are of special interest because they include survivors, migrators, and perhaps the first recruits (Table 7).

Contrary to indications of the incomplete recovery discussed above, results for richness in the one-time sampling at stations 1 to 6 showed that borrow pits generally supported more species than undredged bottom at 1 year. This was true for stations 2, 4, 5, and 6. Findings at station 1 were contradictory, and at station 3, species in experimental collections were outnumbered by those in control collections. The number of species in control samples averaged 86 and ranged between 57 and 112; the number for experimental samples was higher with an average of 92 and a range between 80 and 114.

Even though richness data are somewhat inconsistent, overall they indicate that faunal recovery began rapidly and was virtually complete throughout the study area in about 1 year. Data from the one-time sampling at six stations support this statement to a greater degree than those from regular time-sequence samples at station 1.

b. Abundance. Except for a few anomalies, seasonal cycles of faunal abundance coincided with periods of low and high species diversity, i.e., fewer animals were recorded in winter collections, and peak numbers generally occurred at various times between March and December. In base-line samples, numbers of individuals per square meter of bottom averaged 3,883 and ranged from 1,506 (April) to 7,178 (July). The average for control samples was 3,119, with a range between 1,408 (July) and 5,576 (August). Experimental samples had an

Table 7. Species and their frequency of occurrence in the first 3 weeks after dredging at station 1 offshore Panama City Beach, Florida, August 1976.

Species	No. of individuals (by date)			Species	No. of individuals (by date)		
	10 Aug.	18 Aug.	24 Aug.		10 Aug.	18 Aug.	24 Aug.
CHIDARIA				ANNELIDA (Cont'd)			
Unid. sp.			1	<i>Polydora tetrabranchia</i>	1		114
PLATYHELMINTHES				<i>Prionospio cristata</i>	43		1
Unid. sp.			2	<i>Rullierieris mexicana</i>	1		21
HEMERTHREA				<i>Scoloplos armiger</i>	5		1
Unid. sp.	1	3	9	<i>Scoloplos rubra</i>			19
HEMATOIDA				<i>Spio pettibonense</i>			1
Unid. sp.	2	3	1	<i>Spio phanias bombyx</i>			
BRANCHIOPODA				SIPUNCULIDA			
<i>Glyptotendipes pyramidalis</i>			1	<i>Golfingia trichoscephala</i>	1		
MOLLUSCA				ARTERPODA			
<i>Caecum faridensis</i>	1		1	<i>Acanthohauastorius</i> sp.			1
<i>Cardiomya costellata</i>			1	<i>Ampelisca abdita</i>	1		1
<i>Lepton</i> sp.			4	<i>Ampelisca verrilli</i>	3		27
<i>Lucina multilineda</i>			1	<i>Monoculodes</i> sp.	1		1
<i>Periphaea margaritaceum</i>			1	<i>Prothauastorius</i> sp.	4		1
<i>Pitar simplex</i>			3	<i>Pseudohauastorius</i> sp.	1		31
<i>Strigilla mirabilis</i>			2	<i>Synchelidium</i> sp.	1		1
<i>Tellina texana</i>	12	18	23	<i>Albunea paretii</i>	3		1
<i>Tellina varicolor</i>				<i>Callianassa jamaicensis</i>			1
ANNELIDA				<i>Petrochirus diogenes</i>			1
Unid. Oligochaete			2	<i>Petrochirus galenthus</i>			1
<i>Ampelisca pygmaea</i>			1	<i>Pinnixa retinens</i>			1
<i>Aranidia maculata</i>			1	<i>Processa vicina</i>	1		1
<i>Brania velifera</i>	1		9	<i>Cyclaspis</i> sp.			1
<i>Capitellidae</i> Jonesi			1	<i>Cyclaspis varians</i>	4		2
<i>Caulerella</i> sp.	1		1	<i>Oxyurostylis smithi</i>	5		3
<i>Ceratonereis irritabilis</i>			1	<i>Nebalia</i> sp.	1		1
<i>Diapatra cuprea</i>			1	Unid. Mysid	2		1
<i>Eteone lactea</i>	1		1	Unid. Ostracod	2		1
<i>Eulella sanguinea</i>			1	<i>Penaeus duorarum</i>			1
<i>Glycera americana</i>	1		1	<i>Stacyonia brevirostris</i>	1		
<i>Glycera dibranchiata</i>			1	<i>Stacyonia typica</i>			
<i>Glycera</i> sp.			1	<i>Acanthosquilla blumienalis</i>	1		
<i>Glycyde solitaria</i>			1	ECHINODERMATA			
<i>Goniada littorea</i>	1		1	<i>Leptosynapta</i> sp.			1
<i>Gyptis vittata</i>	1		1	<i>Metilia quinquesperforata</i>			
<i>Haploscoloplos foliosus</i>			170	Unid. Ophiuroid	2		1
<i>Lumbrineris cruzensis</i>			3	HEMICHORDATA			
<i>Meloboris indicus</i>	38	113	13	Unid. Enteropneust	1		
<i>Mesochestopterus sagittarius</i>			1	CEPHALOCHORDATA			
<i>Nephtys pecta</i>			1	<i>Branchiostoma floridae</i>	2		11
<i>Onuphis a. eculaza</i>			1	VERTEBRATA			
<i>Onuphis nebulosa</i>	4		1	<i>Lepophidium gracilis</i>			1
<i>Paranoides species</i>			2	<i>Symphurus</i> sp.	1		
<i>Paranoides lyra</i>			2				
<i>Paronis fulgens</i>			2				
<i>Paraprionospio pinnata</i>			2				
<i>Phyllodoce arenae</i>			6				
				TOTAL SPECIES/INDIVIDUALS	20/81	38/24	60/534

average of 2,200, with a range between 324 (immediately after dredging), and 4,037 1 year following dredging.

Results of periodic sampling at station 1 showed that numbers of individuals within the borrow pit first reached control sample abundance in December 1976, or about 3 months after dredging had been completed. From that time through the next four sampling periods, individuals in experimental samples were more numerous than in control samples. In May, abundance values were reversed, then again favored the experimental sample in June but remained higher in controls until collecting terminated in November 1977. Thus, a pattern of abundance indicative of faunal recovery within 3 months did not occur the following summer and fall seasons.

At stations 1 to 6, one-time sampling in July neither confirmed nor refuted evidence of recovery from time-sequence sampling at station 1. Numbers of individuals were higher in control samples at stations 1, 3, and 4, while abundance values were higher in experimental samples at stations 2, 5, and 6. A comparison of averages and ranges showed that the average number of individuals per square meter was higher for experimental samples. The low for experimental collections was well above that of control samples, and the high for experimental samples was comparable to the high for control samples. In summary, abundance values demonstrated rapid initial faunal recovery in the borrow pits that was practically complete after about 12 months.

c. Diversity (H') and Equitability (J'). For comparable pairs of control and experimental samples, species richness and abundance data were converted statistically to provide an index of diversity (H') that was used to numerically determine degrees of difference between faunal communities in undredged bottom and borrow pits. Observed differences were validated for each sample set by calculating equitability (J'), which is a mathematical measurement of how evenly organisms in a sample are divided among the various species represented (Pielou, 1975). Used in combination, values of H' and J' for base-line and control samples were regarded normal. For experimental samples, lesser values of H' and J' were attributed to dredging effects, and equal or higher values were considered evidence of faunal recovery. In base-line samples, values for both parameters were slightly higher in summer months, but control samples at station 1 showed no seasonal trend.

Average values for H' and J' in base-line samples were 2.6 and 0.7 respectively, with H' ranging from 1.9 to 3.5 and J' ranging from 0.6 to 0.8. Average H' in control samples was a little higher than base-line but J' was the same and ranges of both were within base-line limits. Among experimental collections, average H' was 2.8 and ranged between 1.8 and 3.7. The average for J' was the same as for base-line and control samples, but the low was 0.5 and the high was 0.8. Lowest values for H' and J' were recorded in January and February, and may have been a result of low water temperature as well as dredging.

When H' and J' values for control and experimental samples taken on the same data were compared, the results showed little regularity. In the series from

station 1, the first experimental sample to equal or surpass control values of H' and J' was collected in October, about 2 months after dredging. From that time until November of the next year, only 5 of 14 experimental samples showed evidence of faunal recovery. Recovery was demonstrated somewhat better by H' and J' data from the six stations sampled in July 1977. At four borrow pit stations, experimental samples had the same or higher diversity and equitability values than control samples. Also, average H' for experimental samples was higher than that for control samples, and averages of J' were the same inside and outside borrow pits.

A review of diversity and equitability results suggests the following: (1) the benthos off Panama City Beach exhibited an annual cycle in which species diversity and abundance were greater in warm water months than in winter; (2) faunal recovery in the borrow pit at station 1 was evident to a considerable degree within 2 to 3 months after dredging, and became nearly complete by the end of sampling in November 1977; and (3) faunal recovery also occurred within 1 year of dredging in at least half of the six borrow pits sampled. To further test these inferences, sets of biotic data from control and experimental samples were evaluated using Morisita's index of faunal similarity and stability analyses. Morisita's index was first used to develop similarity matrices (App. D), and then to perform a classification analysis that arranged control and experimental samples in the form of a dendrogram according to their various degrees of likeness (App. E). Two stability analyses were made (App. F). The first shows the amount of sample variation among the control and experimental samples when compared to the centroid of the statistical faunal cluster calculated from all base-line and control data. The second shows time to faunal recovery by plotting experimental sample data against the nearest mathematical edge of the same statistical cluster.

d. Morisita's Index. Similarity matrices were calculated and displayed for time-sequence samples from station 1, and for one-time collections at stations 1 to 6 (App. D). A regular pattern of light cells (no similarity) and dark cells (high similarity) was not evident because 45 percent or more of station-to-station comparisons in both values had faunal overlap of at least 50 percent. For additional clarification, the same data were used to generate a classification analysis for presentation as a cluster diagram (App. E). In performing the necessary calculations, a Q-mode (normal) analysis was made to show faunal relationships on a station-to-station basis; no data transformations were made because doing so would obscure the dominant ranking of any faunal elements in the samples; and group averaging was selected as the sorting strategy.

For time-sequence samples, the first five (1 September 1977-experimental to 10 August 1976-experimental) show very little similarity to any other samples and were therefore considered unrelated, or outliers. These outliers include two summer-fall experimental samples taken 1 year after dredging, two similar winter collections taken about 6 months after dredging, and the first experimental sample taken a few days after the dredging. The interpretation here is that the two experimental samples 1 year after dredging are as unrelated to other samples

as the one taken immediately after dredging and the two taken in winter during the presumed period of least faunal diversity and abundance.

The next group is the first cluster and has five samples (1 April 1977-experimental to 2 May 1977-experimental). These are related by season (spring), and consist of a base-line sample and control and experimental samples collected 8 to 9 months after dredging. This mixture, and close correspondence between control and experimental samples suggests that community recovery has occurred within the borrow pit at station 1.

Then there is a single, odd sample with no close associates (1 November 1977-experimental), followed by the second cluster which contains eight samples (4 October 1976-experimental to 3 October 1977-experimental). Except for the two control samples, this group represents the experimental samples in the fall during the first 3 months after dredging.

Cluster three is considered the opposite of cluster two. It has seven samples (10 August 1976-control to 2 August 1977-control); five are post-dredging late summer and fall control samples; one a preconstruction control sample from July; and one a winter experimental sample.

Cluster four is the largest grouping and contains the next 15 samples (1 September 1977-control to 1 November 1977-control); 8 of these are fall control samples and closely associated with experimental samples taken as soon as 2 weeks after dredging, as well as in various other months. Here, the indication is that recovery at station 1 began very quickly after dredging.

The fifth and last cluster contains six samples (1 June 1977-control to 11 July 1977-experimental), which are equally divided among summer control and experimental samples taken about 1 year after dredging. Similarities between clusters one and five provide substantial evidence of faunal recovery over a postconstruction period of 8 to 11 months.

For the one-time sampling at six stations, control and experimental collections all show a high level of faunal affinity and therefore support cluster data from station 1 showing a recovery time of 1 year or less. At the time these samples were taken, the diagram shows that station location east to west along the coast was a greater clustering factor than whether or not a sample came from a dredged or undredged bottom. This is not surprising considering the daily discharge of estuarine water through West Pass and into nearshore waters at the eastern end of the study area.

e. Stability Analyses. In the first analysis, control and experimental samples are represented along the x-axis according to the number of days before and after dredging (see App. F). The y-axis is a scale of increasing distance from a statistically determined centroid, or midpoint within a community cluster represented mathematically and calculated from all available base-line and

control data. This graph shows a large variation occurring in control and experimental samples, and at corresponding times, both appear about equally distant from the centroid--distance to maximum community stability. In other words, control samples did not show close connections to the centroid, nor did they follow a seasonal or any other discernible pattern in relation to that point. Likewise, experimental samples showed no definite postconstruction deviation from the centroid, and followed no subsequent trend that might have indicated recovery. In fact, when respective sample distances from the centroid were compared in a Mann-Whitney U-Test, it was found that variations among control and experimental samples were statistically indistinguishable. The point emphasized by this analysis is that faunal variation was a major feature of both control and experimental samples.

In the second graph, the y-axis scale (labeled distance to cluster edge) refers to the edge of the statistical community (to a 95-percent confidence level) that has the centroid as its midpoint (App. F). The zero point on the scale represents the nearest edge of the community, higher positive values are increasing distances from the edge, and negative values show that the experimental sample falls inside the cluster about the centroid and cannot be statistically separated from it. Experimental samples along the x-axis are arranged by day number in postdredging sequence. The x-y plots show that an experimental sample first touched the edge of the centroid cluster on day 332 (5 July 1977), about 11 months after dredging was completed at station 1. This intersection of an experimental sample with the zero line represents time to faunal recovery. However, in several later samples, the plot again falls outside the cluster edge, and does not return until October, 14 months after dredging and 1 month before sampling ended. This situation may be due to normal sample variation.

VI. CONCLUSIONS AND DISCUSSION

Study results indicate several general conclusions related to hydrology, sediments, and benthic fauna of borrow pits and undredged adjacent bottom. Hydrological measurements included temperature and salinity, recorded quarterly at stations A and B in 1974 and 1975, and monthly at station 1 during a 20-month period between April 1976 and November 1977. Temperature data showed that regular seasonal changes are subject to rather wide year-to-year variations. Summer temperature was the most consistent, but in spring, fall, and winter, observed yearly differences were on the order of 10° Celsius. In part, fluctuations of this magnitude could conceivably mediate events responsible for changes in benthic diversity and abundance recorded in base-line, control, and experimental samples.

Salinity was characteristically high (above 32 parts per thousand); however, a low value of 26 parts per thousand, recorded in August 1975, showed that the study area may at times be influenced by estuarine water masses from St. Andrew Bay and perhaps other areas as well (Salsman and Ciesluk, 1978). As with temperature, such periodic change could be translated into adjustments in community structure. In the case of salinity, however, the effects might be

more than physiological, as foreign water masses would undoubtedly introduce a variety of immigrant organisms and potential community recruits.

A comparison of sediments from undredged bottom and borrow pits showed that most deviations from normal properties appeared in experimental samples. Major sedimentological differences could be identified due to accumulation of loosely packed, darker, and siltier sediments in the pits shortly after dredging. These distinctions became more subtle with time, and by the following year, the surface samples (in nearly filled pits) were very similar to sediments on the adjacent undisturbed sea floor. When compared to base-line samples, specific differences included the following: (1) lower sand content, (2) higher silt-clay content, (3) poorer sorting, (4) more finely skewed, (5) more variation in both directions from a leptokurtic condition, and (6) higher content of organic carbon.

In the borrow pit at station 1, altered sediment texture was confirmed by divers, and bathymetric changes were recorded over time. Depth of the cut was 3 to 5 meters below the sea floor, and sediment at the bottom initially appeared dark, soft, and silty. Within a few months this material was covered by fine sand. By the end of sampling in November 1977, the pit had filled to within a meter of the surrounding bottom. A final visual impression was that sediments were still finer and darker, but no distinction could be made between epibenthic and pelagic marine life inside and outside the borrow pit.

Dredging caused an immediate decline in the bottom community followed by a rapid postconstruction recovery that was virtually complete after 1 year. This, or even a shorter recovery period of 8 to 9 months, was supported by analyses that included: (1) species richness, (2) abundance of individuals, (3) diversity and equitability indexes, (4) Morisita's index of faunal similarity, and (5) stability analyses. It is important to again note that sampling beyond 1 year indicated lack of complete faunal recovery. This may be true, or these samples may merely be representative of large natural environmental variations that were shown to be an inherent characteristic of the shallow coastal system off Panama City Beach.

On the basis of data presented here, and complementary studies by Saloman (1976) and Culter and Mahadevan (1982), it is evident that dredging done at Panama City Beach has had no adverse long-term effect on bottom dwelling invertebrates, sediments, or water quality either along the shore or in offshore borrow areas. Short-term ecological consequences of dredging were shown to last only about 1 year, and included only minor sedimentological changes and only a small decline in diversity and abundance among bottom dwelling invertebrates. This lack of evident protracted environmental alteration is due to factors related to physical and biological oceanography within the dredging and disposal areas, and to certain engineering features of the beach restoration project. The natural factors would include the following regional characteristics: (1) moderate to high wave energy capable of eroding and transporting large volumes of sediment annually, (2) tidal, longshore, offshore, and storm generated currents that have

the same, or greater, capability of transporting nearshore sediments, (3) a geographic location that is regularly influenced by water masses and marine life of estuarine, coastal, and oceanic origins, (4) a native infauna that is diversified, abundant, and well adapted to substrate disruption and movement, and (5) a fauna that is composed of subtropical and temperate species whose active reproductive periods are limited by low water temperatures normally recorded in only 1 or 2 winter months.

As for features of the dredging project, numerous small borrow areas were used, instead of fewer larger ones, and they were dredged only to a depth of about 5 meters or less. At this level, no strata of silt, clay, or rock were uncovered so that sediment type in dredged areas remained very much like sediment in undredged areas. Also, dredging occurred in fairly shallow water where sediment transport supplied the volume of sand required to rapidly fill the borrow pits. In this connection, it is important to mention that because of their fast filling rate, and the normally low concentration of suspended solids in overlying water, no biologically detrimental quantities of silt and clay size particles accumulated in borrow areas off Panama City Beach. If anything, during the recovery period, data support the theory that within borrow pits a relative decrease in turbulence and a slight increase in organic deposits may have been responsible for figures showing a higher diversity and abundance of infauna in some dredged areas compared to figures for bottom left undisturbed.

In general, results of coastal restoration studies at Panama City Beach agree with findings for similar projects in comparable surroundings (Thompson, 1973), and along with more recent work (Turbeville and Marsh, 1982), provide additional information that can be used both locally and elsewhere to more accurately predict and evaluate environmental effects of beach nourishment operations. Nevertheless, since each coastal and estuarine area has certain unique features, it is important to continue a close association between ecological research and coastal engineering. Ideally, the research should be conducted to collect base-line data, proceed during all phases of construction, and continue after project completion for a sufficient period of time to obtain short-term (1 year) and long-term data (2 years or longer). In all instances major research emphasis should at least include: (1) factors related to geographic and meteorological conditions, (2) sedimentology, (3) water quality, (4) hydrodynamics, (5) resident and migratory biota at the bottom and throughout the water column, (6) interactions between biotic and abiotic elements, and (7) socioeconomic circumstances. By using such a research-oriented approach in future engineering projects, many important coastal resources could be protected, or even enhanced, and most environmental problem areas would be identified and avoided.

LITERATURE CITED

- BLOOM, S.A., "Multivariate Quantification of Community Recovery," *The Recovery Process in Damaged Ecosystems*, J. Cairns, Jr., ed., 1st ed., Ann Arbor, Mich., 1980, pp. 141-151.
- BLOOM, S.A., "Similarity Indices in Community Studies: Potential Pitfalls," *Marine Ecology Program*, Ser. 5, 1981, pp. 125-128.
- CULTER, J.K., and MAHADEVAN, S., "Long-Term Effects of Beach Nourishment on the Benthic Fauna of Panama City Beach, Florida," MR 82-2, U.S. Army, Corps of Engineers, Coastal Engineering Research Center, Fort Belvoir, Va., Jan. 1982.
- FOLK, R.L., "Petrology of Sedimentary Rocks," Hemphill Publishing Co., Austin, Tex., 1974.
- MORISITA, M., "Measuring of Interspecific Association and Similarity Between Communities," *Memoirs of the Faculty of Science*, Kyushu University, Ser. E (Biology), Vol. 3, No. 1, 1959, pp. 65-80.
- PIELOU, E.C., "Ecological Diversity," John Wiley & Sons, New York, 1975.
- SALOMAN, C.H., "The Benthic Fauna and Sediments of the Nearshore Zone off Panama City Beach, Florida," MR 76-10, U.S. Army, Corps of Engineers, Coastal Engineering Research Center, Fort Belvoir, Va., Aug. 1976.
- SALOMAN, C.H., and NAUGHTON, S.P., "Effects of Hurricane Eloise on the Benthic Fauna of Panama City Beach, Florida, USA," *Marine Biology*, Vol. 42, 1977, pp. 357-363.
- SALSMAN, G.G., and CIESLUK, A.J., "Environmental Conditions in Coastal Waters Near Panama City, Florida," NCSC TR 337-78, U.S. Naval Coastal Systems Center, Panama City, Fla., 1978.
- THOMPSON, J.R., "Ecological Effects of Offshore Dredging and Beach Nourishment: A Review," MP 1-73, U.S. Army, Corps of Engineers, Coastal Engineering Research Center, Fort Belvoir, Va., Jan. 1973.
- TURBEVILLE, D.B., and MARSH, G.A., "Benthic Fauna of an Offshore Borrow Area in Broward County, Florida," MR 82-1, U.S. Army, Corps of Engineers, Coastal Engineering Research Center, Fort Belvoir, Va., Jan. 1982.
- U.S. ARMY ENGINEER DISTRICT, MOBILE, "Bay County, Florida, Construction Plans for Emergency Beach Protection," Mobile, Ala., 1976.
- WILSON, D., "Feasibility Report for Beach Erosion Control and Hurricane Protection, Panama City Beaches, Florida," U.S. Army Engineer District, Mobile, Mobile, Ala., 1975.
- WILSON, D., "Panama City Beaches, Florida, Interim Feasibility Report for Beach Erosion Control and Hurricane Protection," U.S. Army Engineer District, Mobile, Mobile, Ala., 1976.

APPENDIX A

HYDROLOGICAL AND SEDIMENT DATA BY STATION

Hydrological and sedimentological data, by station and date, for offshore stations (9-meter depth) before and after dredging--beach restoration project, Panama City Beach, Florida (November 1974 to November 1977).

HYDROLOGICAL AND SEDIMENTOLOGICAL DATA, BY STATION AND DATE, FOR
OFFSHORE STATIONS (30-FOOT DEPTH) BEFORE AND AFTER DREDGING -
BEACH RESTORATION PROJECT, PANAMA CITY BEACH, FLORIDA (NOVEMBER
1974 TO NOVEMBER 1977).

STATION A - CONTROL

PARAMETER	DATE				MEAN	RANGE	
	11/74	2/75	5/75	8/75			
HYDROLOGICAL							
SALINITY, ‰	34.500	34.390	32.220	26.220	31.832	26.22 TO	34.50
WATER TEMP., C	21.000	17.400	26.200	28.300	23.225	17.40 TO	28.30
SEDIMENT							
GRANULE, WT. %	99.861	99.892	99.826		99.860	99.83 TO	99.89
SAND	0.139	0.108	0.174		0.140	0.11 TO	0.17
SILT							
CLAY							
MEAN GRAIN SIZE, #	2.203	2.294	2.433		2.310	2.20 TO	2.43
ST. DEVIATION, #	0.715	0.595	0.499		0.603	0.50 TO	0.71
SKEWNESS	-0.156	-0.246	-0.199			-0.25 TO	-0.16
KURTOSIS	1.014	1.145	1.227		1.129	1.01 TO	1.23
T. CARBON, WT. %	0.113	0.144	0.080	0.070	0.102	0.07 TO	0.14
T. ORGANIC C	0.081	0.024	0.047	0.050	0.050	0.02 TO	0.08
T. CARBONATE C	0.032	0.120	0.033	0.020	0.051	0.02 TO	0.12

STATION B - CONTROL

PARAMETER	DATE				MEAN	RANGE	
	11/74	2/75	5/75	8/75			
HYDROLOGICAL							
SALINITY, ‰	34.330	33.890	32.170	26.110	31.625	26.11 TO	34.33
WATER TEMP., C	20.800	17.500	26.000	28.500	23.200	17.50 TO	28.50
SEDIMENT							
GRANULE, WT. %		0.502			0.502	0.50 TO	0.50
SAND	99.871	99.341	100.000	99.886	99.774	99.34 TO	100.00
SILT	0.129	0.157		0.114	0.133	0.11 TO	0.16
CLAY							
MEAN GRAIN SIZE, #	2.213	2.169	2.330	2.447	2.290	2.17 TO	2.45
ST. DEVIATION, #	0.802	0.744	0.562	0.554	0.665	0.55 TO	0.80
SKEWNESS	-0.236	-0.382	-0.234	-0.059		-0.38 TO	-0.09
KURTOSIS	1.262	1.177	1.134	1.376	1.237	1.13 TO	1.38
T. CARBON, WT. %	0.106	0.334	0.082		0.174	0.08 TO	0.33
T. ORGANIC C	0.084	0.114	0.008		0.069	0.01 TO	0.11
T. CARBONATE C	0.022	0.220	0.074		0.105	0.02 TO	0.22

TREASURE ISLAND MOTEL (STATION 1) - CONTROL

PARAMETER	DATE			MEAN	RANGE	
	4/76	6/76	7/76			
HYDROLOGICAL						
SALINITY, ‰	33.330	32.330	33.280	32.980	32.33 TO	33.33
WATER TEMP., C	20.200	25.700	29.000	24.633	20.20 TO	28.00
SEDIMENT						
GRANULE, WT. %	0.156			0.156	0.16 TO	0.16
SAND	99.836			99.836	99.84 TO	99.84
SILT	0.008			0.008	0.01 TO	0.01
CLAY						
MEAN GRAIN SIZE, #	2.407			2.407	2.41 TO	2.41
ST. DEVIATION, #	0.470			0.470	0.47 TO	0.47
SKEWNESS	0.020			0.020	0.02 TO	0.02
KURTOSIS	1.228			1.228	1.23 TO	1.23
T. CARBON, WT. %	0.269			0.269	0.27 TO	0.27
T. ORGANIC C	0.032			0.032	0.03 TO	0.03
T. CARBONATE C	0.237			0.237	0.24 TO	0.24

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 8/10/76	DATE - EXPERIMENTAL 8/10/76
HYDROLOGICAL		
SALINITY, ‰	35.280	35.280
WATER TEMP., °C	27.000	27.000
SEDIMENT		
GRANULE, WT. %		99.056
SAND		0.144
SILT		
CLAY		
MEAN GRAIN SIZE, #		2.481
ST. DEVIATION, #		0.411
SKEWNESS		-0.137
KURTOSIS		1.017
T. CARBON, WT. %		0.347
T. ORGANIC C		0.336
T. CARBONATE C		0.011

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 8/18/76	DATE - EXPERIMENTAL 8/18/76
HYDROLOGICAL		
SALINITY, ‰	35.280	35.280
WATER TEMP., °C	27.000	27.000
SEDIMENT		
GRANULE, WT. %		0.271
SAND		99.418
SILT		0.311
CLAY		
MEAN GRAIN SIZE, #		2.493
ST. DEVIATION, #		0.530
SKEWNESS		-0.067
KURTOSIS		1.436
T. CARBON, WT. %		0.308
T. ORGANIC C		0.300
T. CARBONATE C		0.008

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 8/24/76	DATE - EXPERIMENTAL 8/24/76
HYDROLOGICAL		
SALINITY, ‰	35.280	35.280
WATER TEMP., °C	27.000	27.000
SEDIMENT		
GRANULE, WT. %		0.063
SAND		99.634
SILT		0.303
CLAY		
MEAN GRAIN SIZE, #		2.501
ST. DEVIATION, #		0.458
SKEWNESS		0.024
KURTOSIS		1.209
T. CARBON, WT. %		0.361
T. ORGANIC C		0.177
T. CARBONATE C		0.184

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL		
PARAMETERS	DATE - CONTROL 9/1/76	DATE - EXPERIMENTAL 9/1/76
HYDROLOGICAL		
SALINITY, ‰	32.610	32.610
WATER TEMP., C	27.800	27.800
SEDIMENT		
GRANULE, WT. %	0.187	
SAND	99.672	97.108
SILT	0.141	2.892
CLAY		
MEAN GRAIN SIZE, ϕ	2.323	2.747
ST. DEVIATION, ϕ	0.558	0.587
SKEWNESS	-0.281	0.285
KURTOSIS	1.189	1.115
T. CARBON, WT. %	0.348	1.123
T. ORGANIC C	0.100	0.039
T. CARBONATE C	0.248	1.084

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL		
PARAMETERS	DATE - CONTROL 9/8/76	DATE - EXPERIMENTAL 9/8/76
HYDROLOGICAL		
SALINITY, ‰	32.610	32.610
WATER TEMP., C	27.800	27.809
SEDIMENT		
GRANULE, WT. %		
SAND		99.776
SILT		0.224
CLAY		
MEAN GRAIN SIZE, ϕ		2.508
ST. DEVIATION, ϕ		0.507
SKEWNESS		-0.015
KURTOSIS		1.348
T. CARBON, WT. %		0.302
T. ORGANIC C		0.257
T. CARBONATE C		0.045

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL		
PARAMETERS	DATE - CONTROL 9/21/76	DATE - EXPERIMENTAL 9/21/76
HYDROLOGICAL		
SALINITY, ‰	32.610	32.610
WATER TEMP., C	27.800	27.800
SEDIMENT		
GRANULE, WT. %		
SAND		91.896
SILT		8.104
CLAY		
MEAN GRAIN SIZE, ϕ		2.835
ST. DEVIATION, ϕ		0.736
SKEWNESS		0.340
KURTOSIS		1.070
T. CARBON, WT. %		2.318
T. ORGANIC C		0.462
T. CARBONATE C		1.856

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 10/24/76	DATE - EXPERIMENTAL 10/24/76
HYDROLOGICAL		
SALINITY, ‰/0	33.060	33.060
WATER TEMP., C	24.900	24.900
SEDIMENT		
GRANULE, WT. %		0.092
SAND		99.626
SILT		0.283
CLAY		
MEAN GRAIN SIZE, Ø		2.452
ST. DEVIATION, Ø		0.481
SKEWNESS		-0.165
KURTOSIS		1.202
T. CARBON, WT. %		0.281
T. ORGANIC C		0.187
T. CARBONATE C		0.094

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 10/18/76	DATE - EXPERIMENTAL 10/18/76
HYDROLOGICAL		
SALINITY, ‰/0	33.060	33.060
WATER TEMP., C	24.900	24.900
SEDIMENT		
GRANULE, WT. %		98.611
SAND		1.389
SILT		
CLAY		
MEAN GRAIN SIZE, Ø		2.536
ST. DEVIATION, Ø		0.411
SKEWNESS		0.155
KURTOSIS		1.068
T. CARBON, WT. %		0.722
T. ORGANIC C		0.700
T. CARBONATE C		0.072

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 11/1/76	DATE - EXPERIMENTAL 11/1/76
HYDROLOGICAL		
SALINITY, ‰/0	33.170	33.170
WATER TEMP., C	18.000	18.000
SEDIMENT		
GRANULE, WT. %		0.108
SAND		98.769
SILT		1.123
CLAY		
MEAN GRAIN SIZE, Ø		2.507
ST. DEVIATION, Ø		0.536
SKEWNESS		-0.042
KURTOSIS		1.492
T. CARBON, WT. %		0.519
T. ORGANIC C		0.316
T. CARBONATE C		0.203

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL		
PARAMETERS	DATE - CONTROL 12/1/76	DATE - EXPERIMENTAL 12/1/76
HYDROLOGICAL		
SALINITY, ‰	34.060	34.060
WATER TEMP., °C	12.500	12.500
SEDIMENT		
GRANULE, WT. %		0.052
SAND	99.876	99.086
SILT	0.124	0.862
CLAY		
MEAN GRAIN SIZE, φ	2.300	2.524
ST. DEVIATION, φ	0.577	0.471
SKEWNESS	-0.267	0.074
KURTOSIS	1.118	1.225
T. CARBON, WT. %	0.275	0.498
T. ORGANIC C	0.060	0.110
T. CARBONATE C	0.215	0.388

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL		
PARAMETERS	DATE - CONTROL 1/5/77	DATE - EXPERIMENTAL 1/5/77
HYDROLOGICAL		
SALINITY, ‰	33.280	33.280
WATER TEMP., °C	12.400	12.400
SEDIMENT		
GRANULE, WT. %		0.437
SAND		97.222
SILT		2.341
CLAY		
MEAN GRAIN SIZE, φ		2.518
ST. DEVIATION, φ		0.597
SKEWNESS		-0.037
KURTOSIS		1.664
T. CARBON, WT. %		0.919
T. ORGANIC C		0.327
T. CARBONATE C		0.592

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL		
PARAMETERS	DATE - CONTROL 2/2/77	DATE - EXPERIMENTAL 2/2/77
HYDROLOGICAL		
SALINITY, ‰	34.330	34.330
WATER TEMP., °C	9.000	9.000
SEDIMENT		
GRANULE, WT. %		0.085
SAND		99.701
SILT		0.215
CLAY		
MEAN GRAIN SIZE, φ		2.499
ST. DEVIATION, φ		0.486
SKEWNESS		-0.036
KURTOSIS		1.295
T. CARBON, WT. %		0.313
T. ORGANIC C		0.298
T. CARBONATE C		0.017

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 3/1/77	DATE - EXPERIMENTAL 3/1/77
HYDROLOGICAL		
SALINITY, ‰/0	34.440	34.440
WATER TEMP., C	14.300	14.300
SEDIMENT		
GRANULE, WT.%		0.652
SAND		99.265
SILT		0.084
CLAY		
MEAN GRAIN SIZE, ϕ		2.316
ST. DEVIATION, ϕ		0.571
SKEWNESS		-0.297
KURTOSIS		1.228
T. CARBON, WT.%		0.253
T. ORGANIC C		0.163
T. CARBONATE C		0.090

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 3/1/77	DATE - EXPERIMENTAL 3/1/77
HYDROLOGICAL		
SALINITY, ‰/0	33.500	33.500
WATER TEMP., C	22.400	22.400
SEDIMENT		
GRANULE, WT.%		0.201
SAND	99.829	99.214
SILT	0.171	0.585
CLAY		
MEAN GRAIN SIZE, ϕ	2.303	2.487
ST. DEVIATION, ϕ	0.560	0.414
SKEWNESS	-0.275	-0.103
KURTOSIS	1.140	1.031
T. CARBON, WT.%	0.214	0.339
T. ORGANIC C	0.202	0.328
T. CARBONATE C	0.012	0.011

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 3/2/77	DATE - EXPERIMENTAL 3/2/77
HYDROLOGICAL		
SALINITY, ‰/0	34.800	34.280
WATER TEMP., C	21.800	21.800
SEDIMENT		
GRANULE, WT.%		0.016
SAND		99.801
SILT		0.183
CLAY		
MEAN GRAIN SIZE, ϕ		2.491
ST. DEVIATION, ϕ		10.389
SKEWNESS		-0.100
KURTOSIS		0.937
T. CARBON, WT.%		0.244
T. ORGANIC C		0.097
T. CARBONATE C		0.147

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL		
PARAMETERS	DATE - CONTROL 6/1/77	DATE - EXPERIMENTAL 6/1/77
HYDROLOGICAL		
SALINITY, ‰	32.060	32.060
WATER TEMP., °C	25.700	25.700
SEDIMENT		
GRANULE, WT. %		0.085
SAND		97.964
SILT		1.951
CLAY		
MEAN GRAIN SIZE, Ø		2.356
ST. DEVIATION, Ø		0.677
SKEWNESS		-0.193
KURTOSIS		1.572
T. CARBON, WT. %		1.206
T. ORGANIC C		0.206
T. CARBONATE C		1.000

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL		
PARAMETERS	DATE - CONTROL 7/5/77	DATE - EXPERIMENTAL 7/5/77
HYDROLOGICAL		
SALINITY, ‰	33.560	33.560
WATER TEMP., °C	27.500	27.500
SEDIMENT		
GRANULE, WT. %	0.335	
SAND	99.422	98.705
SILT	0.244	1.295
CLAY		
MEAN GRAIN SIZE, Ø	2.456	2.507
ST. DEVIATION, Ø	0.453	0.483
SKEWNESS	-0.193	0.034
KURTOSIS	1.195	1.274
T. CARBON, WT. %		
T. ORGANIC C		
T. CARBONATE C		

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL		
PARAMETERS	DATE - CONTROL 8/2/77	DATE - EXPERIMENTAL 8/2/77
HYDROLOGICAL		
SALINITY, ‰	35.330	35.330
WATER TEMP., °C	29.000	29.000
SEDIMENT		
GRANULE, WT. %		
SAND		97.489
SILT		2.511
CLAY		
MEAN GRAIN SIZE, Ø		2.529
ST. DEVIATION, Ø		0.463
SKEWNESS		0.161
KURTOSIS		1.201
T. CARBON, WT. %		
T. ORGANIC C		
T. CARBONATE C		

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 9/1/77	DATE - EXPERIMENTAL 9/1/77
HYDROLOGICAL		
SALINITY, ‰	32.610	32.610
WATER TEMP., C	27.700	27.700
SEDIMENT		
GRANULE, WT. %		96.923
SAND		3.077
SILT		
CLAY		
MEAN GRAIN SIZE, ϕ		2.544
ST. DEVIATION, σ		0.465
SKEWNESS		0.197
KURTOSIS		1.219
T. CARBON, WT. %		
T. ORGANIC C		
T. CARBONATE C		

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 10/3/77	DATE - EXPERIMENTAL 10/3/77
HYDROLOGICAL		
SALINITY, ‰	33.060	33.060
WATER TEMP., C	25.000	25.000
SEDIMENT		
GRANULE, WT. %		0.092
SAND		99.597
SILT		0.311
CLAY		
MEAN GRAIN SIZE, ϕ		2.491
ST. DEVIATION, σ		0.505
SKEWNESS		-0.037
KURTOSIS		1.327
T. CARBON, WT. %		
T. ORGANIC C		
T. CARBONATE C		

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 11/1/77	DATE - EXPERIMENTAL 11/1/77
HYDROLOGICAL		
SALINITY, ‰		
WATER TEMP., C		
SEDIMENT		
GRANULE, WT. %		0.101
SAND		99.163
SILT		0.736
CLAY		
MEAN GRAIN SIZE, ϕ		2.551
ST. DEVIATION, σ		0.516
SKEWNESS		0.075
KURTOSIS		1.282
T. CARBON, WT. %		
T. ORGANIC C		
T. CARBONATE C		

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 7/11/77	DATE - EXPERIMENTAL 7/11/77
HYDROLOGICAL		
SALINITY, 00/0	33.560	33.560
WATER TEMP., C	27.500	27.500
SEDIMENT		
GRANULE, WT.%		
SAND	99.700	98.641
SILT	0.300	1.359
CLAY		
MEAN GRAIN SIZE, ϕ	2.445	2.499
ST. DEVIATION, σ	0.445	0.525
SKEWNESS	-0.187	-0.001
KURTOSIS	1.178	1.388
T. CARBON, WT.%		
T. ORGANIC C		
T. CARBONATE C		

SUN & SWIM MOTEL (STATION 2) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 7/11/77	DATE - EXPERIMENTAL 7/11/77
HYDROLOGICAL		
SALINITY, 00/0	33.560	33.560
WATER TEMP., C	27.500	27.500
SEDIMENT		
GRANULE, WT.%		
SAND	99.646	99.796
SILT	0.354	0.204
CLAY		
MEAN GRAIN SIZE, ϕ	2.452	2.425
ST. DEVIATION, σ	0.440	0.479
SKEWNESS	-0.179	-0.194
KURTOSIS	1.148	1.205
T. CARBON, WT.%		
T. ORGANIC C		
T. CARBONATE C		

MILTON HOLIDAY INN (STATION 3) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 7/11/77	DATE - EXPERIMENTAL 7/11/77
HYDROLOGICAL		
SALINITY, 00/0	33.330	33.330
WATER TEMP., C	26.800	26.600
SEDIMENT		
GRANULE, WT.%		
SAND	99.879	0.922
SILT	0.121	98.964
CLAY		0.114
MEAN GRAIN SIZE, ϕ	2.214	1.749
ST. DEVIATION, σ	0.615	1.064
SKEWNESS	-0.319	-0.460
KURTOSIS	1.109	0.824
T. CARBON, WT.%		
T. ORGANIC C		
T. CARBONATE C		

SANDPIPER MOTEL (STATION 4) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 7/11/77	DATE - EXPERIMENTAL 7/11/77
HYDROLOGICAL		
SALINITY, ‰	33.330	33.330
WATER TEMP., C	26.800	26.800
SEDIMENT		
GRANULE, WT. %		0.079
SAND	99.859	99.810
SILT	0.141	0.111
CLAY		
MEAN GRAIN SIZE, Ø	2.244	2.006
ST. DEVIATION, Ø	0.608	0.831
SKEWNESS	-0.307	-0.414
KURTOSIS	1.158	0.954
T. CARBON, WT. %		
T. ORGANIC C		
T. CARBONATE C		

PEPPERTREE CONDOMINIUM (STATION 5) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 7/11/77	DATE - EXPERIMENTAL 7/11/77
HYDROLOGICAL		
SALINITY, ‰	33.330	33.330
WATER TEMP., C	26.800	26.800
SEDIMENT		
GRANULE, WT. %		
SAND	99.864	99.863
SILT	0.136	0.137
CLAY		
MEAN GRAIN SIZE, Ø	2.305	2.257
ST. DEVIATION, Ø	0.593	0.575
SKEWNESS	-0.331	-0.280
KURTOSIS	1.344	1.111
T. CARBON, WT. %		
T. ORGANIC C		
T. CARBONATE C		

BLUE DOLPHIN MOTEL (STATION 6) - CONTROL & EXPERIMENTAL

PARAMETERS	DATE - CONTROL 7/11/77	DATE - EXPERIMENTAL 7/11/77
HYDROLOGICAL		
SALINITY, ‰	33.330	33.330
WATER TEMP., C	26.800	26.800
SEDIMENT		
GRANULE, WT. %		
SAND	0.340	0.137
SILT	99.520	99.757
CLAY	0.139	0.106
MEAN GRAIN SIZE, Ø	2.114	2.311
ST. DEVIATION, Ø	0.760	0.612
SKEWNESS	-0.397	-0.340
KURTOSIS	1.092	1.391
T. CARBON, WT. %		
T. ORGANIC C		
T. CARBONATE C		

APPENDIX B

CHECKLIST OF ORGANISMS

Checklist of organisms collected at offshore stations (9-meter depth)
before and after dredging--beach restoration project, Panama City Beach,
Florida (November 1974 to November 1977).

CHECKLIST OF ORGANISMS COLLECTED AT OFFSHORE STATIONS (30-FOOT DEPTH)
BEFORE AND AFTER DREDGING - BEACH RESTORATION PROJECT, PANAMA CITY
BEACH, FLORIDA (NOVEMBER 1974 TO NOVEMBER 1977).

CNIDARIA
ACTINIARIA (SEA ANEMONES)
UNIDENTIFIED SP.

PLATYHELMINTHES
TURBELLARIA (FLATWORMS)
UNIDENTIFIED SP.

NEMERTINEA (RIBBON WORMS)
UNIDENTIFIED SP.

NEMATODA (ROUNDWORMS)
UNIDENTIFIED SP.

PHORONIDA (PHORONIDS)
PHORONIS ARCHITECTA

BRACHIOPODA (LAMP SHELLS)
GLOTTIDIA PYRAMIDATA

MOLLUSCA (SHELLFISH)
GASTROPODA (SNAILS)

ACTEOCINA CANALICULATA
ACTECCINA CANDEI
ACTEON PUNCTOSTRIATUS
ANACHIS FLORICANA
BULLA STRIATA
CAECUM FLORIDANUM
CAECUM IMBRICATUM
CAECUM PULCHELLUM
CYLICHNELLA BIDENTATA
DIASIMA VARIUM
MELANELLA JAMAICENSIS
NASSARIUS ACUTUS
NATICA PUSILLA
OLIVA SAYANA
OLIVELLA BULLULA
OLIVELLA MINUTA
OLIVELLA MUTICA
OLIVELLA PUSILLA
PHILINE SAGRA
POLINICES DUPLICATUS
TEREBRA CONCAVA
TEREBRA DISLOCATA
TURBNILLA CONRADI
TURBNILLA ELEGANTULA
TURBNILLA SP.

PELECYPODA (CLAMS)
ANADARA FLORIDANA
ANATINA ANATINA
CARDIOMYA COSTELLATA
CHIONE CANCELLATA
CHIONE GRUS
CUMINGIA TELLINOIDES
CUMINGIA T. VANHYNINGI
CUNA CALLI
DIPLOCCNYA SEMIASPERA
DIPLOCCNYA SP.
ERVILIA CONCENTRICA
LAEVICARDIUM LAEVIGATUM
LAEVICARDIUM MORTONI
LAEVICARDIUM PICTUM

LEPTON SP.
LUCINA MULTILINEATA
LUCINA RADIANI
LYNSIA H. FLORICANA
MACOMA CONSTRICTA
MACRACALLISTA MACULATA
MACRACALLISTA NIMBOSA
MACRA SP.
MUSCULUS LATERALIS
NUCULANA ACUTA
PANDORA TRILINEATA
PARVIDEA SOLENIIFORMIS
PARVILUCINA BLANDA
PEPILOMA MARGARITACEUM
PITAE SIMPSONI
SEMLE PROFICUA
SOLEMYA SP.
SOLEMYA VELUM
SOLEMYA VIRIDIS
STRIGILLA MIRABILIS
TELLIDORA CRISTATA
TELLINA A. TAYLORIANA
TELLINA AEGUISTRIATA
TELLINA IRIS
TELLINA TAMPAENSIS
TELLINA TEXANA
TELLINA VERSICOLOR
TRACHYCARDIUM MURICATUM
VARICORBULA OPERCULATA
VENERIDAE UNIDENTIFIED SP.

ANNELIDA (SEGMENTED WORMS)

CLIGOCFAETA
UNIDENTIFIED SP.

POLYCHAETA
AGLAPHAMUS VERRILLI
AMERICONUPHIS MAGNA
AMPHARETE ACUTIFRONS
ANATIDES ERYTHROPHYLLUS
ANTINCE SP.
ARONICES MAYAQUEZENSIS
ARONICES NOBISIO PYGMAEA
ARENICOLA CRISTATA
ARICIDEA CERRUTI
ARICIDEA FAUVELI
ARICIDEA FRAGILIS
ARICIDEA PHILIPINAE
ARICIDEA SUECICA
ARICIDEA TAYLORI
ARICIDEA WASSI
ARICIDEA SP.
ARMANDIA AGILIS
ARMANDIA MACULATA
ASYCHIS CAROLINAE
AXIOPELLA MUCOSA
BRANCHIOASYCHIS AMERICANA
BRANIA CLAVATA
BRANIA WELFLEETENSIS
CASIRA INCERTA
CAPITELLA CAPITATA
CAPITELLIDES JONESI
CAPITELLIDAE UNIDENTIFIED SP.
CARAZZIELLA SP.
CAULLERIELLA SP.
CERATONEREIS TRITABILIS
CERATONEREIS MIRABILIS
CHAETONE GAYHEADIA

CHAETOCZENE SETCSA
 CHLOEIA VIRIDIS
 CHONE SP.
 CIRRIATULIDAE UNIDENTIFIED SP.
 CIRRORPHORUS LYRIFORMIS
 CISTERNICES GULOLI
 CASTERANCHUS LUMBRICOIDES
 CIPATRA CUPREA
 CISPIC UNGINATA
 COVILLEA SOCIABILIS
 COITESCHIA BELLUCIDA
 COPILOBRANCHUS SANGUINEUS
 CYDNE ALBA
 CYDNE LACTEA
 CYDNE SANGUINEA
 EUNICE ANTIENNATA
 EURYTHOE COMPLANATA
 EXOGONE DISPAR
 FLABELLIGERA SP.
 GLYCERA AMERICANA
 GLYCERA DIBRANCHIATA
 GLYCERA OXYCEPHALA
 GLYCERA SP.
 GLYCINE SOLITARIA
 GONIADA LITTOREA
 GRUBELEPIS MEXICANA
 GYPTIS BREVIPALPA
 GYPTIS VITTATA
 HAPLISCLOPLOS FOLIOSUS
 HAPLISCLOPLOS FRAGILIS
 HAPLISCLOPLOS ROBUSTUS
 HARMOTHOE IMBRICATA
 HARMOTHOE LUNULATA
 HENYCDUS ROSEUS
 HETEROMASTUS FILIFORMIS
 ISOLDA PULCHELLA
 LAEONEREIS CULVERI
 LOIMIA MEDUSA
 LOIMIA VIRIDIS
 LUMBRINERIS ACUTUS
 LUMBRINERIS CRUZENSIS
 LUMBRINERIS ERECTA
 LUMBRINERIS TENUIS
 LUMBRINERIS TETRAURA
 LYSIDICE NINETTA
 LYSILLA ALBA
 MACROCLYMENE ZONALIS
 MAGELONA LONGICORNIS
 MAGELONA PETTIBONEAE
 MAGELONA RIOJAY
 MAGELONA SP.
 MALACCERUS INDICUS
 MEDIDMASTUS CALIFORNIENSIS
 MEGALOMMA BIOCULATUM
 MESOCIAETOPTERUS SAGITTARIUS
 MICROPHTHALMUS ABERRANS
 MICROPHTHALMUS SCZELKOWII
 MICROPHTHALMUS SP.
 MICROCSPID PIGMENTATA
 MINUSPID CIRRIFERA
 MYRICHELE SP.
 NEANTHES ACUMINATA
 NEANTHES SP.
 NEANTHES SUCCINEA
 NEPHYS BUCERA
 NEPHYS PICTA
 NEREIS LAMELLOSA
 NEREIS PELAGICA
 NEREIS SP.
 NOTOMASTUS HENIPDOUS
 NOTOMASTUS LATERICEUS
 ONUPHIS EPEMITA OCULATA
 ONUPHIS NEBULOSA
 ONUPHIS PALLICA
 ONUPHIS SP.

ORHINIA RISERI
 OWENIA FUSIFORMIS
 PARANITES SPECIOSA
 PARACNIDES LYRA
 PARACNIDES SP.
 PARACNIS FULGENS
 PARACNIS SP.
 PARAPRICNOSPID PINNATA
 PARAPRICNOSPID LONGICARRATA
 PHERUS EMERSI
 PHYLLODICE ARENAE
 PHYLLODICE SP.
 PHYLLOCRINATUS
 PISTA CRISTATA
 PISTA PALMATA
 PODARKE OBSCURA
 POECILOCHASTUS JOHNSONI
 POLYCIPRUS EXIMIUS
 POLYOCRA SOCIALIS
 POLYOCRA TETRABRANCHIA
 POLYODONTES LUPINA
 POLYODONTES UNIDENTIFIED SP.
 PRIONOSPID CRISTATA
 PRIONOSPID STEENSTRUPI
 PSEUDEURYTHOE AMBIGUA
 RULLIERINEREIS MEXICANA
 SABELLA MICROPHTHALMA
 SCOLELEPIS SQUAMATA
 SCOLELEPIS TEXANA
 SCOLOPES ARMIGER
 SCOLICUS RUBRA
 SIGALION ARENICOLA
 SIGAMBRA BASSI
 SIGAMBRA TENTACULATA
 SPHAEROSYLIS SP.
 SPID PETTIBONEAE
 SPIDONTES UNIDENTIFIED SP.
 SPIDOCIAETOPTERUS OCULATUS
 SPIDONANES BOMBYX
 STHEKELAIS BOA
 STREETOSYLIS ARENAE
 THARYX ANNULOSUS
 TRAVISIA HORSNAE
 WEBSTERINEREIS TRIDENTATA

SIPUNCULICA (PEANUT WORMS)
 ASPIDOSIPHON SP.
 GOLFINGIA TRICHOCEPHALA
 SIPUNCULUS LONGIPAPILLOSUS
 UNIDENTIFIED SP.

ECHIURIDA (ECHIURIIDS)
 UNIDENTIFIED SP.

ARTHROPODA (CRUSTACEANS)

AMPHIPCCA
 ACANTHCHAUSTORIUS SP.
 AMPELISCA ABDITA
 AMPELISCA SP.
 AMPELISCA VADORUM
 AMPELISCA VERRILLI
 ARGISSA SP.
 CAPRELLIDAE UNIDENTIFIED SP.
 CARINCRATEA SP.
 COROPHIUM SP.
 CYNADLSA SP.
 ELASMOPUS SP.
 ERICHTHONYUS SP.
 GAMMAROBSIS SP.
 GITANOPSIS SP.
 HIPPOMEDON SP.
 HYPERIA SP.

LEMBOS SP.
LEPIDACTYLUS SP.
LISTERELLA SP.
LYSIANOPSIS SP.
MELITA APPENDICULATA
MICRODEUTOPUS SP.
MICROPROTOPUS SP.
MONOCULODES SP.
PARAPHOXUS SP.
PHOTIS SP.
PROTOMASTORIUS SP.
PSUDOMASTORIUS SP.
PSUDOPLATYSCNOPUS SP.
SYNCELIUM SP.
TIRCA BIOSCELLATUS
TIRON SP.
 UNIDENTIFIED SP.
 ANOMURA
ALBONEA PARETII
EUCERAMUS PRAELONGUS
LEPIDOPA WEBSTERI
PAGURUS LONGICARPUS
PAGURUS SP.
PETRECHIRUS DIOGENES
PETRELISTHES GALATHINUS
 BRACHYURA
CALLINECTES SAPIDUS
CALLINECTES SP.
DISSODACTYLUS MELLITAE
HEPATUS EPHELITICUS
LIGINIA DUBIA
METOPORHAPIS CALCARATA
OSACHILA TUBEROSA
OVALIPES OCELLATUS
PANOPAEUS HERBSTII
PERSEPHONA P. AQUILONARIS
PINNIXIA CHAETOPTERANA
PINNIXIA CYLINDRICA
PINNIXIA CRISTATA
PINNIXIA LEPTOSYNAPTAE
PINNIXIA LUNZI
PINNIXIA PEARSEI
PINNIXIA RETINENS
PINNIXIA SAYANA
PINNIXIA SP.
PINNOCHERES MACULATUS
PINNOCHERES OSTREUM
PINNOCHERES SP.
PORTUNUS GIBBESII
PORTUNUS SAYI
PORTUNUS SP.
PORTUNUS SPINIMANUS
 PORTUNIDAE UNIDENTIFIED SP.
RANILIA MURICATA
 CALLINANASSIDAE
CALLINANASSA JAMAICENSE
 CARIDEA
ALPHEUS HETEROCHAELIS
AMBICEPTER SYMMETRICUS
HIPPOLYTE PLEURACANTHA
LATREUTES PARVULUS
LEPTOCHELA SERRATORBITA
OGYRIDES ALPHAEOSTRIS
OGYRIDES LIMICOLA
BERGIMENES LONGICAUDATUS
PROCESSA HEMPHILLI
PROCESSA VICINA
SYNALPHEUS SP.
 UNIDENTIFIED SP.
 CUMACEA
CYCLAPSIS SP.
CYCLAPSIS VARIANS
OXYUROSSTYLIS SMITHI

SPILOCUMA SALOMANI
 UNIDENTIFIED SP.
 ISOPODA
ANCINA DEPRESSUS
APANTHURA MAGNIFICA
CHIROCTEA EXCAVATA
EDOTEA MONTOSA
 LEPTOSTRACA
NEBALIA SP.
 MYSIDACEA
BOWMANIELLA SP.
MYSTICOPSIS BIGELOWI
PRAUNUS FLEXUOSUS
 UNIDENTIFIED SP.
 OSTRACODA
HAPLOCYTHERIDEA SEPTIPUNCTATA
SARIELLA CHILDI
 UNIDENTIFIED SP.
 PENAIDEA
ACETES AMERICANUS
LUCIFER FAXONI
PENAEUS DUORARUM
SICYCNIA BREVIDSTRIS
SICYCNIA SP.
SICYCNIA TYPICA
TRACHYPENAEUS CONSTRICTUS
 STOMATOPODA
ACANTHOSQUILLA BIMINIENSIS
CORONIS EXCAVATRIX
 TANAIIDACEA
 UNIDENTIFIED SP.

ECHINODERMATA
 ASTEROIDEA (STARFISHES)
ASTROPECTEN ARTICULATUS
LUZIA ALTERNATA
 ECHINOIDEA (SAND DOLLARS; URCHINS)
LYTECHINUS VARIEGATUS
MOIRA ATROPS
MELLITA QUINQUESPERFORATA
 UNIDENTIFIED SP.
 HOLOTHURIDAE (SEA CUCUMBERS)
LEPTOSYNAPTA SP.
 UNIDENTIFIED SP.
 OPHIUROIDEA (BRITTLE STARS)
HEMIPHOLIS ELONGATA
MICROPHOLIS GRACILLIMA
OPHIOPHRAGMUS FIDIGRANEUS
OPHIOPHRAGMUS MOOREI
OPHIOPHRAGMUS WOODMANI
 UNIDENTIFIED SP.

HEMICHORDATA
 ENTEROPNEUSTA (ACORN WORMS)
 UNIDENTIFIED SP.

CEPHALOCHORDATA (LANCELETS)
BRANCHIOSTOMA FLORIDAE

VERTEBRATA
 PISCES (FISHES)
 GOBIIDAE, UNIDENTIFIED SP.
 HEMIPTERONCTUS NOVACULA
LEBOPHIDIUM GRAELSI
MICROGOBIUS CARRI
 OPHIDIIDAE, UNIDENTIFIED SP.
SYMPHURUS SP.

APPENDIX C

BIOLOGICAL AND BIOSTATISTICAL DATA BY STATION

Biological and biostatistical data, by station and date, for offshore stations (9-meter depth) before and after dredging--beach restoration project, Panama City Beach, Florida (November 1974 to November 1977).

BIOLOGICAL AND BIOSTATISTICAL DATA, BY STATION AND DATE, FOR OFFSHORE STATIONS (30-FOOT DEPTH) BEFORE AND AFTER DREDGING - BEACH RESTORATION PROJECT, PANAMA CITY BEACH, FLORIDA (NOVEMBER 1974 TO NOVEMBER 1977).

STATION A - CONTROL

SPECIES	NO. OF INDIVIDUALS				TOTAL	PCT.
	11/74	2/75	5/75	8/75		
PLATYHELMINTHES						
TURBELLARIA (FLATWORMS)						
UNIDENTIFIED SP.	0	0	4	1	5	0.58
NEMERTINEA (RIBBON WORMS)						
UNIDENTIFIED SP.	1	4	7	8	20	2.33
NEMATODA (ROUNDWORMS)						
UNIDENTIFIED SP.	0	19	7	18	44	5.12
MOLLUSCA (SHELLFISH)						
GASTROPODA (SNAILS)						
ACTECCINA CANDEI	0	0	0	1	1	0.12
ACTECCIN PUNCTOSTRIATUS	0	0	0	1	1	0.12
NATICA PLISILLA	0	0	0	2	2	0.23
OLIVELLA MLYICA	0	0	0	2	2	0.23
POLINICES DUPLICATUS	0	0	2	0	2	0.23
TEREBRA DISLOCATA	0	0	1	0	1	0.12
PELECYPODA (CLAMS)						
ERVILIA CONCENTRICA	0	0	1	1	2	0.23
LUCINA MULTILINEATA	0	2	2	4	8	0.93
PERIPLOMA MARGARITACEUM	0	1	0	0	1	0.12
STRIGILLA MIRABILIS	0	0	2	10	12	1.40
TELLINA VERSICOLOR	0	0	1	19	20	2.33
ANNELIDA (SEGMENTED WORMS)						
OLIGOCHAETA						
UNIDENTIFIED SP.	22	35	5	0	62	7.22
POLYCHAETA						
APCCEICNO SPIO PYGMAEA	0	1	1	3	5	0.58
ARICILEA SP.	2	2	0	0	4	0.47
ARMANCA MACULATA	5	2	18	4	29	3.38
ERANIA CLAVATA	0	0	0	1	1	0.12
ERANIA VELLFLEETENSIS	0	0	2	4	6	0.70
CAPITELLIDAE UNIDENTIFIED SP.	0	2	0	0	2	0.23
DIOPATRA CUPREA	0	1	0	0	1	0.12
DISPIO UNCINATA	0	0	0	1	1	0.12
ETEDNE LACIEA	0	0	1	7	8	0.93
GLYCERA AMERICANA	0	0	1	4	5	0.58
HAPLOSCOLOPLOS FOLIOLUS	0	0	0	1	1	0.12
HAPLOSCOLOPLOS ROBUSTUS	0	1	0	0	1	0.12
LUMBRINERIS CRUZENSIS	0	0	0	11	11	1.28
MAGELCNA RIOJAI	0	0	1	0	1	0.12
MAGELCNA SP.	0	0	1	0	1	0.12
MESOCIAETOPTERUS SAGITTARILIS	0	0	1	0	1	0.12
MINUSPIO CYRRIFERA	0	1	1	0	2	0.23
NEPHIYS BUCERA	0	0	0	2	2	0.23
NEPHIYS PICTA	0	2	7	6	15	1.75
ONUPHIS FRENITA OCULATA	1	0	0	0	1	0.12
PARANAIRES SPECIOSA	0	0	0	1	1	0.12
PARACNIDES LYRA	19	5	0	1	25	2.91
PARACNIDES SP.	2	0	0	0	2	0.23
PARAPRIONOSPIO PINNATA	17	1	3	0	21	2.44
PHYLLCDOCE ARENAE	0	0	4	0	4	0.47
PHYLLCDOCE SP.	0	0	5	0	5	0.58
POECILOCHAETUS JOHNSONI	0	0	1	0	1	0.12
PRIONOSPIO CRISTATA	47	76	4	5	132	15.37
SCOLELEPIS SOVAMATA	2	0	0	0	2	0.23
SCOLELEPIS TEXANA	0	4	5	0	9	1.05
SCOLOPLOS RUBRA	0	1	0	0	1	0.12

STATION A - CONTROL
(CONTINUED)

SPECIES	NO. OF INDIVIDUALS				TOTAL	PCT.
	11/74	2/75	5/75	8/75		
<u>SIGAMBRA BASSI</u>	0	1	1	2	4	0.47
<u>SPID. PETTIBONEAE</u>	7	5	9	1	22	2.56
<u>SPIDOCFAEIOPTERUS OCULATUS</u>	1	0	0	0	1	0.12
<u>SPIDOPHANE BOMBYX</u>	0	1	42	2	45	5.24
ARTHRACOPODA (CRUSTACEANS)						
AMPHIPODA						
<u>ACANTHOMAUSTORIUS SP.</u>	0	0	8	2	10	1.16
<u>LYSIANOPSIS SP.</u>	0	0	1	0	1	0.12
<u>PROTOFAUSTORIUS SP.</u>	0	12	58	15	85	9.90
<u>PSEUDOPHAUSTORIUS SP.</u>	0	3	4	3	10	1.16
<u>PSEUDOPHYTISCHNOPUS SP.</u>	1	2	2	16	21	2.44
<u>SYNCHELYDIUM SP.</u>	0	2	3	0	5	0.58
ANOMURA						
<u>ALBUEA PARETII</u>	1	0	0	0	1	0.12
BRACHYURA						
<u>PINNIZIA CRISTATA</u>	0	0	0	1	1	0.12
<u>PINNATHERES MACULATUS</u>	0	0	2	0	2	0.23
<u>PORTUNUS GIBBESII</u>	0	0	0	1	1	0.12
<u>PORTUNUS SPINIFRONS</u>	1	0	0	0	1	0.12
<u>RANILIA RUBICATA</u>	0	0	0	2	2	0.23
CARIDEA						
<u>PROCESSA MEMPHILLI</u>	0	0	7	0	7	0.81
<u>PROCESSA VICINA</u>	0	0	0	1	1	0.12
CLANACEA						
<u>CYCLAPSIS VARIANS</u>	0	0	0	1	1	0.12
<u>ORYZODONTIS SMITHI</u>	0	0	3	0	3	0.35
<u>UNIDENTIFIED SP.</u>	0	0	0	2	2	0.23
OSTRACCA						
<u>UNIDENTIFIED SP.</u>	0	0	0	8	8	0.93
PENAIDEA						
<u>SICYCINIA BREVIROSTRIS</u>	0	1	0	0	1	0.12
ECHINODERMATA						
ECHINOIDEA (SAND DOLLARS; URCHINS)						
<u>HELLITA QUINQUESPERFORATA</u>	0	0	0	45	45	5.24
HOLOTHURIDEA (SEA CUCUMBERS)						
<u>UNIDENTIFIED SP.</u>	0	0	0	3	3	0.35
OPHIURIDEA (BRITTLE STARS)						
<u>UNIDENTIFIED SP.</u>	0	0	11	0	11	1.28
CEPHALOCHORDATA (LANCELETS)						
<u>BRANCHIOSTOMA FLORIDAE</u>	0	1	59	19	79	9.20
VERTEBRATA						
PISCES (FISHES)						
<u>HEMITELEOSTEUS NOVACULA</u>	0	0	0	1	1	0.12
<u>OPHIOTIDAE, UNIDENTIFIED SP.</u>	0	0	1	0	1	0.12
TOTALS						
NC. SPECIES	125	188	299	243	855	
NC. IND. PER M2	15	27	41	43	75	
S-W INDEX - H'(LN)	2064	3008	4784	3888		
EVENNESS - J	1.923	2.154	2.801	3.113		
	0.710	0.654	0.754	0.828		
AV. NO. SPECIES	31.5	AV. S-W INDEX	2.498			
AV. NC. IND. PER M2	34.36.0	AV. EVENNESS	0.736			

STATION B - CONTROL

SPECIES	NO. OF INDIVIDUALS					PCT.
	11/74	2/75	5/75	8/75	TOTAL	
CNIDARIA						
ACTINIARIA (SEA ANEMONES)						
UNIDENTIFIED SP.	0	0	0	2	2	0.17
PLATYHELMINTHES						
TURBELLARIA (FLATWORMS)						
UNIDENTIFIED SP.	0	0	2	1	3	0.26
NEMERTINEA (RIBBON WORMS)						
UNIDENTIFIED SP.	0	3	6	7	16	1.39
NEMATODA (ROUNDWORMS)						
UNIDENTIFIED SP.	0	18	2	11	31	2.70
MOLLUSCA (SHELLFISH)						
GASTROSCCA (SNAILS)						
ACTECCIA CANDEI	0	0	0	6	6	0.52
PELECYFIDA (CLAMS)						
CHIONE CANCELLATA	0	1	0	0	1	0.09
ERVILIA CONCENTRICA	0	0	2	1	3	0.26
STRIGILLA MIRABILIS	0	1	7	74	82	7.14
TELLINA VERSICOLOR	0	0	0	28	28	2.44
ANNELIDA (SEGMENTED WORMS)						
CLIGGCHAETA						
UNIDENTIFIED SP.	18	26	1	10	55	4.79
POLYCHAETA						
AGLACPHAMUS VERRILLI	1	0	0	0	1	0.09
AMPHARETE ACUTIFRONS	1	0	0	0	1	0.09
ANATIDES ERYTHROPHYLLS	0	0	1	0	1	0.09
APOPHIONOSPION PYGMAEA	0	0	2	0	2	0.17
ARICICEA FRAGILIS	1	0	0	1	2	0.17
ARMANCIA MACULATA	11	13	24	1	49	4.26
BRANIA WELFLEETENSIS	4	1	1	3	9	0.78
CAPITELLIDAE UNIDENTIFIED SP.	1	0	0	0	1	0.09
CAULLERIELLA SP.	0	0	0	2	2	0.17
CERATONEREIS IRRITABILIS	0	0	0	2	2	0.17
CIRRATULIDAE UNIDENTIFIED SP.	0	1	0	0	1	0.09
DISPID UACINATA	0	0	0	1	1	0.09
EIECNE LACTEA	0	0	1	4	5	0.44
GLYCERA AMERICANA	0	0	0	3	3	0.26
GYPTIS VITTATA	4	1	0	0	5	0.44
HAPLOSCLIOPLIS FRAGILIS	0	1	0	0	1	0.09
HETERONASTIUS FILIPICRMIS	3	0	0	0	3	0.26
LUMBINERIS CROZENSIS	0	0	0	2	2	0.17
LAGELINA SP.	1	0	0	0	1	0.09
MEDICASTIUS CALIFCENIENSIS	0	0	0	1	1	0.09
MESOCAPTEPTERUS SAGITTARIUS	0	0	0	3	3	0.26
MINUSPIC CIRRIFERA	0	1	0	0	1	0.09
NEPHIYS BUCERA	0	0	1	0	1	0.09
NEPHIYS PICTA	0	0	6	4	10	0.87
NOTICASTIUS HEMIPEDUS	0	0	0	2	2	0.17
OPHELIA SP.	9	3	0	5	17	1.48
OWENIA FUSIFCRMIS	1	0	0	0	1	0.09
PARACNIDES LYRA	3	3	0	1	7	0.61
PARACNIDES FULGENS	0	3	0	0	3	0.26
PARAPHILINUS PINNATA	10	1	0	0	11	0.96
PHYLLIDICE ARENAE	0	0	2	0	2	0.17
PHYLLIDICE SP.	0	0	2	0	2	0.17
PLECILLICHAETUS JOHNSONI	0	1	0	0	1	0.09

STATION B - CONTROL
(CONTINUED)

SPECIES	NO. OF INDIVIDUALS				TOTAL	PCT.
	11/74	2/75	5/75	8/75		
<u>FRICNCSPID CRISTATA</u>	134	55	3	18	210	18.28
<u>SCOLELEPIS SQUAMATA</u>	1	1	0	0	2	0.17
<u>SCOLELEPIS TEXANA</u>	0	3	11	0	14	1.22
<u>SCOLEPLOS RUBRA</u>	0	2	0	0	2	0.17
<u>SPID PETTIBONEAE</u>	9	1	22	5	37	3.22
<u>SPIONIDAE UNIDENTIFIED SP.</u>	2	0	0	0	2	0.17
<u>SPIOPHANES BCMBYX</u>	0	0	29	7	36	3.13
<u>TRAVISIA HOBSCNAE</u>	0	0	0	3	3	0.26
 <u>SIPUNCULICA (PEANUT WORMS)</u>						
<u>SIPUNCULUS LONGIPAPILLCSUS</u>	0	1	0	1	2	0.17
 <u>ARTHROPODA (CRUSTACEANS)</u>						
<u>AMPHICDA</u>						
<u>ACANTHOHAUSTORIUS SP.</u>	0	6	16	7	29	2.52
<u>AMPELISCA SP.</u>	1	0	0	1	2	0.17
<u>LISTRIELLA SP.</u>	0	0	0	3	3	0.26
<u>MONOCLODES SP.</u>	0	0	0	1	1	0.09
<u>PROTOMALSTORIUS SP.</u>	0	29	100	8	137	11.92
<u>PSEUDOMALSTORIUS SP.</u>	0	0	1	1	2	0.17
<u>PSEUDOPLATYISCHNOPUS SP.</u>	1	4	3	11	19	1.65
<u>SYNCHELIDIUM SP.</u>	3	0	6	1	10	0.87
<u>BRACHYURA</u>						
<u>PINNIXIA CRISTATA</u>	0	0	1	0	1	0.09
<u>PINNIXIA SAYANA</u>	0	0	0	6	6	0.52
<u>RANILIA MURICATA</u>	0	0	0	2	2	0.17
<u>CARIDEA</u>						
<u>PROCESSA TEMPHILLI</u>	1	0	1	11	13	1.13
<u>PROCESSA VICINA</u>	0	0	0	1	1	0.09
<u>CUMACEA</u>						
<u>CYCLAPSID VARIANS</u>	0	0	0	2	2	0.17
<u>UNIDENTIFIED SP.</u>	0	0	0	3	3	0.26
<u>GSTRACCA</u>						
<u>UNIDENTIFIED SP.</u>	0	0	0	3	3	0.26
<u>PENAEIDAE</u>						
<u>TRACHYPENAEUS CONSTRICTUS</u>	1	0	0	0	1	0.09
 <u>ECHINODERMATA</u>						
<u>ECHINOCIDEA (SAND DOLLARS; URCHINS)</u>						
<u>MELLITA QUINQUE SPERFORATA</u>	5	0	0	6	11	0.96
<u>UNIDENTIFIED SP.</u>	0	0	7	0	7	0.61
<u>HOLOTHUROIDEA (SEA CUCUMBERS)</u>						
<u>LEPTOSYNAPTA SP.</u>	0	0	0	1	1	0.09
<u>OPHIUROIDEA (BRITTLE STARS)</u>						
<u>OPHIOBRANCHIUS FILOGRANEUS</u>	1	0	0	0	1	0.09
 <u>CEPHALOCORDATA (LANCELETS)</u>						
<u>BRANCHIOSTOMA FLORIDAE</u>	10	69	74	51	204	17.75
 <u>VERTEBRATA</u>						
<u>PISCES (FISHES)</u>						
<u>MICROGASTER CARRI</u>	1	0	0	0	1	0.09
 TOTALS	238	249	334	328	1149	
NO. SPECIES	27	26	28	47	75	
NO. IND. PER M2	3808	3984	5344	5248		
S-W INDEX - F'(LN)	1.898	2.247	2.320	3.000		
EVENNESS - J	0.576	0.690	0.696	0.779		
 AV. NO. SPECIES	32.0	AV. S-W INDEX	2.366			
AV. NO. IND. PER M2	4596.0	AV. EVENNESS	0.685			

TREASURE ISLAND MOTEL (STATION 1) - CONTROL

SPECIES	NO. OF INDIVIDUALS			TOTAL	PCT.
	4/76	6/76	7/76		
CNIDARIA					
ACTINIARIA (SEA ANEMONES)					
UNIDENTIFIED SP.	0	1	2	3	0.06
PLATYHELMINTHES					
TURBELLARIA (FLATWORMS)					
UNIDENTIFIED SP.	0	3	0	3	0.06
NEMERTINEA (RIBBON WORMS)					
UNIDENTIFIED SP.	25	37	62	124	2.34
NEMATODA (ROUNDWORMS)					
UNIDENTIFIED SP.	25	48	133	206	3.89
PHORONIDA (PHORONIDS)					
PHORONIS ARCHITECTA	2	2	1	5	0.09
MOLLUSCA (SHELLFISH)					
GASTROPODA (SNAILS)					
ACTECCINA CANALICULATA	1		0	2	0.04
ACTECCINA CANDEI	0	18	24	42	0.79
CAECUM FLORIDANUM	0	9	20	29	0.55
CAECUM IMBRICATUM	0	0	1	1	0.02
CYLICINELLA BICENTATA	0	3	11	14	0.26
DIASITOMA VARIUM	0	0	5	5	0.09
NATICA PUSILLA	0	0	16	16	0.30
OLIVELLA BULLULA	0	0	11	11	0.21
OLIVELLA MUTICA	1	0	2	3	0.06
OLIVELLA PUSILLA	0	3	0	3	0.06
TURBONILLA CONRADI	0	0	10	10	0.19
TURBONILLA ELEGANTULA	0	0	5	5	0.09
TURBONILLA SP.	0	0	1	1	0.02
PELECYPODA (CLAMS)					
ANADARA FLORIDANA	0	3	22	25	0.47
CHIONE GRUS	0	1	0	1	0.02
CUMINGIA T. VANHYNINGI	0	0	1	1	0.02
DIPLODONTA SP.	0	0	2	2	0.04
ERVILLA CONCENTRICA	1	15	223	239	4.52
LEPTA SP.	3	0	10	13	0.25
LUCINA MULTILINEATA	6	35	30	71	1.34
LUCINA FADIANI	1	0	0	1	0.02
LYONSIA H. FLORIDANA	0	2	4	6	0.11
MACRICALLISTA NIMBOSA	0	0	1	1	0.02
MACTEA SP.	0	0	1	1	0.02
PAPYRIDEA SOLENIIFORMIS	1	0	0	1	0.02
PARVILUCINA BLANCA	1	0	0	1	0.02
PERILICHA MARGARITACEUM	1	1	0	2	0.04
PITAE SIMPSONI	1	4	5	10	0.19
SEPIE PECTICUA	0	0	6	6	0.11
STRIGILLA MIRABILIS	1	1	22	24	0.45
TELLINA TEXANA	0	7	90	97	1.83
TELLINA VERSICOLOR	13	43	555	611	11.54
VENETIDAE UNIDENTIFIED SP.	0	9	3	12	0.23
ANNELIDA (SEGMENTED WORMS)					
CLIOCHÆTA					
UNIDENTIFIED SP.	46	20	31	97	1.83
POLYCHÆTA					
AMPHARETE ACUTIFECUS	5	0	0	5	0.09
APCERINUS PYGMAEA	2	5	6	13	0.25

TREASURE ISLAND MOTEL (STATION 1) - CONTROL
(CONTINUED)

SPECIES	NO. OF INDIVIDUALS			TOTAL	PCT.
	4/76	6/76	7/76		
ARMANCIA AGILIS	0	1	6	7	0.13
ARMANCIA MACULATA	11	7	29	47	0.89
ERANCIASUCHIS AMERICANA	1	0	0	1	0.02
ERANIA CLAVATA	0	0	1	1	0.02
ERANIA WELFLEETENSIS	1	3	10	14	0.26
CAPITELLA CAPITATA	0	0	1	1	0.02
CAULLERIELLA SP.	0	0	1	1	0.02
CERATONEREIS MIRABILIS	0	0	3	3	0.06
CHONE SP.	0	1	9	10	0.19
DIOPATRA CUPREA	0	0	2	2	0.04
CISPIC UNCINATA	3	1	0	4	0.08
ENOPLOERANCHUS SANGUINEUS	0	0	1	1	0.02
ETECNE LACTEA	2	5	7	14	0.26
EULALIA SANGUINEA	0	0	1	1	0.02
EXOGCNE CISPAR	0	0	1	1	0.02
GLYCERA AMERICANA	1	33	25	59	1.11
GLYCERA DIERANCHIATA	0	0	2	2	0.04
GLYCERA OXYCEPHALA	8	0	0	8	0.15
GLYCERA SP.	0	0	2	2	0.04
GONIATA LITTOREA	0	24	17	41	0.77
GRUEFULEPIS MEXICANA	0	1	0	1	0.02
GYPTIS VITTATA	0	0	1	1	0.02
HAPLOCYCLOPLOS FOLIOSUS	2	4	8	14	0.26
PARMOTICE LUNULATA	0	1	0	1	0.02
ISOLTA PULCHELLA	1	0	0	1	0.02
LUMBERINERIS CRUZENSIS	2	146	940	1088	20.56
LUMBERINERIS TETRAURA	0	5	0	5	0.09
LYSILLA ALBA	1	0	0	1	0.02
MAGELCNA RIOJAI	1	0	0	1	0.02
MAGELCNA SP.	0	1	6	7	0.13
MEDICMASTUS CALIFORNIENSIS	0	2	0	2	0.04
MESOCYCTOPTERUS SAGITTARIUS	0	0	35	35	0.66
MYRICICELE SP.	1	0	0	1	0.02
NEANTHES ACUMINATA	1	0	0	1	0.02
NEANTHES SUCCINEA	0	0	1	1	0.02
NEPHTYS BUCERA	2	2	11	15	0.28
NEPHTYS PICTA	48	37	56	141	2.66
NEREIS PELAGICA	1	0	6	7	0.13
NOTOMASTUS HEMIPODUS	0	2	2	4	0.08
NOTOMASTUS LATERICELUS	0	3	0	3	0.06
ONUPHIS EREMITA OCULATA	3	17	32	52	0.98
ONUPHIS NEBULOSA	2	1	0	3	0.06
OVENIA FUSIFORMIS	7	10	8	25	0.47
PARANATHES SPECIOSA	0	2	0	2	0.04
PARACNIDES LYRA	3	6	3	12	0.23
PARACNIS FULGENS	4	4	10	18	0.34
PARAPRIONOSPIO PINNATA	16	0	1	17	0.32
PHYLLODOCE ARENAE	5	3	24	32	0.60
PISTA CRISTATA	1	0	0	1	0.02
PISTA PALMATA	0	0	1	1	0.02
POECILOCHAETUS JOHNSONI	1	0	0	1	0.02
POLYDORA TETRABRANCHIA	0	4	0	4	0.08
PRIONOSPIO CRISTATA	16	105	205	326	6.16
PRIONOSPIO STEENSTRUPI	0	0	11	11	0.21
PSEUDOCURYTHOE AMBIGUA	1	0	0	1	0.02
RULLIERINEREIS MEXICANA	0	2	6	8	0.15
SABELLA MICROPHYLMA	0	0	1	1	0.02
SCOLELEPIS SOVAMATA	0	1	1	2	0.04
SCOLELEPIS TEXANA	4	2	3	9	0.17
SCOLICLOS ARMIGER	0	1	17	18	0.34
SYLICHN ARENICOLA	0	1	0	1	0.02

TREASURE ISLAND HOTEL (STATION 1) - CONTROL
(CONTINUED)

SPECIES	NO. OF INDIVIDUALS			TOTAL	PCT.
	4/76	6/76	7/76		
<u>SIGAMBRA BASSI</u>	1	1	5	7	0.13
<u>SPIC PELLIBONEAE</u>	12	1	15	28	0.53
<u>SPICCAETICTERUS OCULATUS</u>	4	0	2	6	0.11
<u>SPICPHANES BCMBYX</u>	336	40	21	397	7.50
<u>STHEBELAIS BCA</u>	1	0	1	2	0.04
 <u>SIPUNCULICA (PEANUT WORMS)</u>					
<u>GOLFINGIA TRICHOCEPHALA</u>	1	0	1	2	0.04
 <u>ARTHOPODA (CRUSTACEANS)</u>					
<u>AMPHIPODA</u>					
<u>ACANTHCHAUSTICEIUS SP.</u>	1	0	7	8	0.15
<u>AMPELISCA ABDITA</u>	0	1	0	1	0.02
<u>AMPELISCA VERNILLI</u>	2	7	89	98	1.85
<u>ARGYSSA SP.</u>	0	1	5	6	0.11
<u>COROPHIUM SP.</u>	0	0	1	1	0.02
<u>CYMAULSA SP.</u>	1	0	0	1	0.02
<u>ERICTHONIUS SP.</u>	0	0	1	1	0.02
<u>HYPERIA SP.</u>	0	0	1	1	0.02
<u>LEPTACTYLUS SP.</u>	0	1	0	1	0.02
<u>LISTERIELLA SP.</u>	0	1	6	7	0.13
<u>LYSIANOPSIS SP.</u>	0	1	0	1	0.02
<u>MICROPCTOPUS SP.</u>	0	3	0	3	0.06
<u>MONOCULODES SP.</u>	1	1	25	27	0.51
<u>PARAPLOXUS SP.</u>	0	1	1	2	0.04
<u>PROTIS SP.</u>	0	1	0	1	0.02
<u>PROTIFAUSTORIUS SP.</u>	2	0	27	29	0.55
<u>PSEUDOMALSTORIUS SP.</u>	1	1	7	9	0.17
<u>PSEUDOPLATYISCHNOPUS SP.</u>	56	19	209	284	5.37
<u>SYNCHELIDIUM SP.</u>	19	17	58	94	1.78
<u>TIRON BISCOCELLATUS</u>	0	0	2	2	0.04
<u>TIRON SP.</u>	1	0	0	1	0.02
<u>ANOMURA</u>					
<u>ALBUNEA PARETII</u>	0	1	0	1	0.02
<u>LEPIDODA WEBSTERI</u>	0	0	3	3	0.06
<u>PAGURUS LONGICARPUS</u>	0	0	8	8	0.15
<u>BRACHYURA</u>					
<u>CALLINECTES SP.</u>	0	0	5	5	0.09
<u>HEPATUS EPHELITICUS</u>	0	0	3	3	0.06
<u>LIBinia CUBIA</u>	0	0	1	1	0.02
<u>OVALIPES OCELLATUS</u>	0	0	1	1	0.02
<u>PERSPECTNA P. AQUILONARIS</u>	0	0	1	1	0.02
<u>PINNIXIA CRISTATA</u>	6	0	0	6	0.11
<u>PINNIXIA RETINENS</u>	0	3	8	11	0.21
<u>PINNIXIA SAYANA</u>	0	2	4	6	0.11
<u>CALLINASSIDAE</u>					
<u>CALLINASSA JAMAICENSE</u>	0	0	1	1	0.02
<u>CARIDEA</u>					
<u>ALPHEUS FETEROCHAEILIS</u>	0	1	0	1	0.02
<u>AMBICEPTER SYMMETRICUS</u>	0	1	0	1	0.02
<u>HIPPOLYTE PLEURACANTHA</u>	0	0	1	1	0.02
<u>LATHEUTES PARVULUS</u>	0	0	3	3	0.06
<u>PROCESSA FEMPHILLI</u>	0	3	3	6	0.11
<u>PROCESSA VICINA</u>	0	1	0	1	0.02
<u>CUMACEA</u>					
<u>CYCLIPSIS SP.</u>	0	1	6	7	0.13
<u>CYCLIPSIS VARIANS</u>	14	20	26	60	1.13
<u>OXYUEDOSTYLIS SMITHI</u>	4	11	13	28	0.53
<u>ISOPODA</u>					
<u>EGGTEA MCNIQUA</u>	0	1	4	5	0.09
<u>LEFTICSTRACA</u>					

TREASURE ISLAND HOTEL (STATION 1) - CONTROL
(CONTINUED)

SPECIES	NO. OF INDIVIDUALS			TOTAL	PCT.
	4/76	6/76	7/76		
<u>NEEALIA SP.</u>	0	0	6	6	0.11
<u>MYSTICACIA</u>					
<u>PRALLIS FLEXUCSUS</u>	0	1	0	1	0.02
<u>UNIDENTIFIED SP.</u>	3	1	2	6	0.11
<u>GSTRACCIA</u>					
<u>HAPLOCYTHERIDEA SEPTIFUNCIATA</u>	0	29	0	29	0.55
<u>UNIDENTIFIED SP.</u>	0	0	17	17	0.32
<u>PENAICEA</u>					
<u>SICYPTIA TYPICA</u>	0	1	0	1	0.02
<u>SICMATICCOA</u>					
<u>ACANTHOSQUILLA BIRMINIENSIS</u>	0	1	3	4	0.08
<u>ECHINODERMATA</u>					
<u>ASTEROCIDEA (STARFISHES)</u>					
<u>ASTROPECIEN ARTICULAILS</u>	0	0	1	1	0.02
<u>ECHINOIDEA (SAND DOLLARS; URCHINS)</u>					
<u>MELLITA QUINQUEPERFORATA</u>	1	50	123	174	3.29
<u>HOLOTHURACIDEA (SEA CUCUMBERS)</u>					
<u>UNIDENTIFIED SP.</u>	0	1	0	1	0.02
<u>OPHIURIDEA (BRITTLE STARS)</u>					
<u>OPHIOPHRAGMUS BURDEMANNI</u>	0	1	1	2	0.04
<u>UNIDENTIFIED SP.</u>	0	8	14	22	0.42
<u>PENICHOCEATA</u>					
<u>ENTEROPNEUSTA (ACORN WORMS)</u>					
<u>UNIDENTIFIED SP.</u>	0	3	0	3	0.06
<u>CEPHALOCYRCATA (LANCELETS)</u>					
<u>BRANCHIOSICMA FLORICAE</u>	0	4	23	27	0.51
<u>VERTEBRATA</u>					
<u>PISCES (FISHES)</u>					
<u>HEMIPYRROQUILS NOVACULA</u>	0	0	1	1	0.02
TOTALS	753	951	3589	5293	
NO. SPECIES	67	94	120	166	
NO. IND. PER M2	1566	1902	7178		
S-W INDEX - H' (LN)	2.516	3.482	3.084		
EVENNESS - J	0.598	0.766	0.644		
AV. NO. SPECIES	93.7	AV. S-W INDEX	3.027		
AV. NO. IND. PER M2	3528.7	AV. EVENNESS	0.670		

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
8/10/76

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNICARIA				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	4	0.287	0	0.0
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	1	0.072	0	0.0
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	23	1.650	1	1.235
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	33	2.367	2	2.469
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
CAECID FLORIDANUM	2	0.143	1	1.235
CYLICHELLA BIDENTATA	1	0.072	0	0.0
NASSARIUS ACUTUS	1	0.072	0	0.0
NATICA PUSILLA	2	0.143	0	0.0
OLIVELLA PULLULA	3	0.215	0	0.0
TEREBA DISLOCATA	2	0.143	0	0.0
TURBELLULA CONRADI	2	0.143	0	0.0
PELECYPODA (CLAMS)				
CUMINGIA TELLINOIDES	3	0.215	0	0.0
ERVILLIA CONCENTRICA	38	2.726	0	0.0
LEPTIS SP.	7	0.502	0	0.0
LUCINA MULTILINEATA	8	0.574	0	0.0
STRIGILLA MIRABILIS	13	0.933	0	0.0
TELLINA TAMPAENSIS	1	0.072	0	0.0
TELLINA TEXANA	93	6.671	0	0.0
TELLINA VERSICOLOR	123	8.824	12	14.815
ANNELIDA (SEGMENTED WORMS)				
CLIGOCFAETA				
UNIDENTIFIED SP.	7	0.502	0	0.0
POLYCHAETA				
AMPHARETE ACUTIFRONS	1	0.072	0	0.0
ARMANDIA MACULATA	1	0.072	0	0.0
AXIOHELLA MUCOSA	1	0.072	0	0.0
BRANIA BELLEFLEETENSIS	14	1.004	1	1.235
CAULIERIELLA SP.	1	0.072	1	1.235
CHONE SP.	1	0.072	0	0.0
ETECE LACTEA	4	0.287	1	1.235
GLYCERA AMERICANA	6	0.430	1	1.235
GLYCERA SP.	2	0.143	0	0.0
GONATA LITTOREA	0	0.0	1	1.235
GYPTIS VITTATA	0	0.0	1	1.235
HAPLOSCLOPUS FOLIOSUS	2	0.143	0	0.0
PARALICHOE LUNULATA	1	0.072	0	0.0
LOINIA MUCOSA	1	0.072	0	0.0
LUMEFINERIS CRUZENSIS	669	47.991	38	46.914
MAGELCNA SP.	1	0.072	0	0.0
MESOCALOPTERUS SAGITTARIUS	2	0.143	0	0.0
NEANITES ACUMINATA	2	0.143	0	0.0
NEPHYS BUCERA	1	0.072	0	0.0
NEPHYS PICTA	5	0.359	0	0.0
ONUPHIS EREMITA OCULATA	9	0.646	0	0.0
ONUPHIS NEBULOSA	11	0.789	4	4.938
PARALICHOE SPECIOSA	5	0.359	0	0.0

TREASURE ISLAND MOTEL (STATICA 1) - CONTROL AND EXPERIMENTAL
8/10/76
(CONTINUED)

SPECIES	NO. CF IND. (C.)		NO. CF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
PHYLLODICE ARENAE	5	0.359	0	0.0
PRIONOSPION CRISTATA	69	4.950	0	0.0
NULLITERINEREYS MEXICANA	3	0.215	1	1.235
SCOLOPLOS ARMIGER	14	1.004	5	6.173
SYGAMERA BASSI	5	0.359	0	0.0
SPIONA PETTIBONEAE	2	0.143	0	0.0
SPIONOPHANES BOHMYX	7	0.502	0	0.0
ARTROPODA (CRUSTACEANS)				
AMPHIPODA				
ACANTHOCHAUSTORIUS SP.	1	0.072	0	0.0
AMPELISCA VERRILLI	3	0.215	0	0.0
COROPHIUM SP.	1	0.072	0	0.0
MICROPROTOPUS SP.	1	0.072	0	0.0
MONOCLODES SP.	5	0.359	0	0.0
PROTCHALSTICRIS SP.	15	1.076	4	4.938
PSEUDOCOAUSTORIUS SP.	8	0.574	1	1.235
PSEUDOCALATYISCHNEPUS SP.	74	5.308	1	1.235
SYNCHELIDIUM SP.	10	0.717	0	0.0
BRACHYURA				
OVALIPES OCELLATUS	1	0.072	0	0.0
PINNIPEDIA RETINENS	3	0.215	0	0.0
CARIDEA				
PROCESSA FEMPHILLI	2	0.143	0	0.0
UNIDENTIFIED SP.	2	0.143	0	0.0
CLMACEA				
CYCLAPSIS SP.	4	0.287	0	0.0
CYCLAPSIS VARIANS	5	0.359	0	0.0
CYXIDOSTYLIS SMITHI	1	0.072	0	0.0
ISOPODA				
EDOTEA MONTOSA	3	0.215	0	0.0
MYSIDACEA				
UNIDENTIFIED SP.	1	0.072	0	0.0
OSTRACCA				
UNIDENTIFIED SP.	12	0.861	0	0.0
PENAEIDAE				
SICYPTIA TYPICA	0	0.0	1	1.235
ECHINODERMATA				
ASTEROIDEA (STARFISHES)				
ASTROPECTEN ARTICULATUS	1	0.072	0	0.0
ECHINOIDEA (SAND DOLLARS; URCHINS)				
MELLITA QUINQUESPERFORATA	14	1.004	2	2.469
OPHIURIDEA (BRITTLE STARS)				
OPHIOURUS WUEGENANI	1	0.072	0	0.0
HEMICHORCATA				
ENTEROPNEUSTA (ACORN WORMS)				
UNIDENTIFIED SP.	2	0.143	0	0.0
CEPHALOCHORDATA (LANCELETS)				
FRANCISCOTOMA FLORIDAE	12	0.861	2	2.469
TOTALS				
NO. SPECIES	1394	72	81	20
NO. IND. PER M2		5576		324
S-W INDEX - H'(LN)		2.3604		2.0322
EVENNESS - J		0.5519		0.6784

TREASURE ISLAND MCTEL (STATION 1) - CONTROL AND EXPERIMENTAL
8/18/76

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNIDARIA				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	3	0.218	0	0.0
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	8	0.582	0	0.0
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	33	2.400	3	1.230
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	59	4.291	3	1.230
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
ACTECCINA CANDEI	1	0.073	0	0.0
NATICA PUSILLA	3	0.218	0	0.0
CLIVELLA PULCILLA	5	0.364	0	0.0
TURBINILLA CONRADI	1	0.073	0	0.0
PELECYPODA (CLAMS)				
ERVILIA CONCENTRICA	17	1.236	0	0.0
LEPTICA SP.	4	0.291	0	0.0
LUCINA MULTILINEATA	11	0.800	0	0.0
PAPYRICEA SOLENIIFORMIS	1	0.073	0	0.0
PERIPLOMA MARGARITACEUM	2	0.145	0	0.0
PITAE SIMPSONI	2	0.145	3	1.230
STRIGILLA MIRABILIS	13	0.945	0	0.0
TELLINA TEXANA	55	4.000	2	0.820
TELLINA VERSICOLOR	79	5.745	18	7.377
ANNELIDA (SEGMENTED WORMS)				
OLIGOCHAETA				
UNIDENTIFIED SP.	15	1.091	0	0.0
POLYCHAETA				
AMPHAEJE ACUTIFRONS	1	0.073	0	0.0
APOEICHNESEIC PYGMAEA	0	0.0	1	0.410
ARMANDIA MACULATA	12	0.873	1	0.410
BRANIA WELFLEETENSIS	9	0.655	0	0.0
CAPITELLA CAPITATA	1	0.073	0	0.0
CAPITELLIDES JONESI	1	0.073	1	0.410
CAULLETTIELLA SP.	1	0.073	0	0.0
CERATONEREIS IRREITABILIS	7	0.509	1	0.410
CHCNE SP.	3	0.218	0	0.0
CIOPATRA CUPREA	1	0.073	1	0.410
ETECNE LACTEA	2	0.145	1	0.410
EXOCNE DISPAR	1	0.073	0	0.0
FLABELLIGERA SP.	1	0.073	0	0.0
GLYCERA AMERICANA	3	0.218	1	0.410
GLYCERA DIBRANCHIATA	6	0.436	1	0.410
GLYCERA SP.	4	0.291	0	0.0
GONIATA LITTOREA	1	0.073	0	0.0
LAPLOSCHLOPLOS FOLIOLIS	0	0.0	3	1.230
LARNOTHOE CUNULATA	1	0.073	0	0.0
LUMBERINERIS CRUZENSIS	499	36.291	113	46.311
MAGELONA SP.	1	0.073	0	0.0
MESOCYCTOPTERUS SAGITTABILIS	4	0.291	3	1.230
NEPHIYS BUCERA	1	0.073	0	0.0
NEPHIYS PICTA	6	0.655	0	0.0
ONUPHIS EREMYIA OCULATA	0	0.0	1	0.410

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
8/18/76
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>GNUPHIS NEBULOSA</u>	7	0.509	0	0.0
<u>PARANAITES SPECIOSA</u>	1	0.073	0	0.0
<u>PARACNICES LYRA</u>	0	0.0	2	C.820
<u>PARACNICES FULGENS</u>	1	0.073	0	0.0
<u>PARAPRIONOSPION PINNATA</u>	1	0.073	2	C.820
<u>PHYLLOCOCE ARENAE</u>	3	0.218	2	C.820
<u>POLYDORA TETRABRANCHIA</u>	1	0.073	1	C.410
<u>PRIONOSPION CRISTATA</u>	200	14.545	43	17.623
<u>RULLIERINEREIS MEXICANA</u>	4	0.291	C	C.0
<u>SCOLOPLCS ARMIGER</u>	30	2.182	7	2.869
<u>SIGALICN ARENICOLA</u>	2	0.145	C	C.0
<u>SIGAMERA BASSI</u>	9	0.655	C	C.0
<u>SPIO PETTIBONEAE</u>	6	0.436	C	C.0
<u>SPIOPHANES BOMBYX</u>	5	0.364	C	C.0
 <u>SIPLUNCULICA (PEANUT WORMS)</u>				
<u>GOLFINGIA TRICHOCEPHALA</u>	1	0.073	1	C.410
 <u>ARTHROPODA (CRUSTACEANS)</u>				
<u>AMPHIPODA</u>				
<u>ACANTHOHAUSTORIUS SP.</u>	12	0.873	0	0.0
<u>AMPELISCA ABDITA</u>	3	0.218	1	C.410
<u>AMPELISCA VERRILLI</u>	15	1.091	3	1.230
<u>ARGISSA SP.</u>	2	0.145	0	0.0
<u>MONOCILCOES SP.</u>	C	0.0	1	0.410
<u>PROIONHALSTORILS SP.</u>	15	1.091	0	0.0
<u>PSEUDOHALSTORILS SP.</u>	6	0.436	0	0.0
<u>PSEUDOPALATIUSCHNITZUS SP.</u>	45	3.564	4	1.639
<u>SYNCELIDILM SP.</u>	13	0.945	1	0.410
<u>BRACHYURA</u>				
<u>CALLINECTES SP.</u>	1	0.073	0	C.0
<u>PINNIXIA RETINENS</u>	5	0.364	C	0.0
<u>CALLINANASSIDAE</u>				
<u>CALLINANASSA JAMAIGENSE</u>	0	0.0	3	1.230
<u>CARIDEA</u>				
<u>ERCESSA LEMPELLI</u>	7	0.509	0	0.0
<u>ERCESSA VICINA</u>	3	0.218	1	C.410
<u>CUMACEA</u>				
<u>CYCLAPSIS SP.</u>	5	0.364	0	0.0
<u>CYCLAPSIS VARIANS</u>	12	0.873	4	1.639
<u>ORYZOPSIS STYLI SMITHI</u>	7	0.509	5	2.049
<u>LEPTOSTRACA</u>				
<u>NEBALIA SE.</u>	1	0.073	1	0.410
<u>MYSTACEA</u>				
<u>UNIDENTIFIED SP.</u>	2	0.145	1	C.410
<u>OSTRACCA</u>				
<u>UNIDENTIFIED SP.</u>	10	0.727	2	C.820
<u>PENAEIDAE</u>				
<u>SICYCNIA BREVIROSTRIS</u>	C	0.0	1	C.410
<u>STOMATOPODA</u>				
<u>ACANTHOSQUILLA BIMINIENSIS</u>	0	0.0	1	C.410
 <u>ECHINODERMATA</u>				
<u>ASTEROIDEA (STARFISHES)</u>				
<u>ASTROPECTEN ARTICULATUS</u>	1	0.073	0	0.0
<u>ECPHINOIDEA (SAND DOLLARS; URCHINS)</u>				
<u>MELLIA CLINGUESPERECRATA</u>	15	1.091	C	0.0
<u>HOLOTHURIDEA (SEA CUCUMBERS)</u>				
<u>LEPTOSYNAPIA SP.</u>	2	0.145	C	C.0
<u>OPHYURIDEA (BRITTLE STARS)</u>				
<u>OPHIOURAGMUS MURDEMANNI</u>	1	0.073	0	0.0

APPENDIX B (CONTINUED)

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
8/18/76
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
UNIDENTIFIED SP.	2	0.145	0	0.0
HEMICHORDATA				
ENTEROPNEUSTA (ACORN WORMS)				
UNIDENTIFIED SP.	1	0.073	0	0.0
CEPHALOCORDATA (LANCELETS)				
BRANCHIOSTOMA FLORIDAE	32	2.327	0	0.0
VERTEBRATA				
PISCES (FISHES)				
SYMPHYRUS SP.	1	0.073	0	0.0
TOTALS	1375		244	
NO. SPECIES		80		38
NO. IND. PER M2		5500		576
S-W INDEX - H'(LN)		2.7517		2.1746
EVENNESS - J		0.6280		0.5678

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
8/24/76

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNICARIA				
ACTINIAEIA (SEA ANEMONES)				
UNIDENTIFIED SP.	3	0.248	1	0.187
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	6	0.496	2	0.375
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	25	2.068	9	1.625
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	16	1.323	1	0.187
BRACHIOPODA (LAMP SHELLS)				
GLYPTIDIA PYRAMICATA	0	0.0	1	0.187
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
ACTECCINA CANDEI	3	0.248	0	0.0
NATICA PLISILLA	1	0.083	0	0.0
OLIVELLA BULLULA	1	0.083	0	0.0
PELECYPODA (CLAMS)				
ANADARA FLORIDANA	1	0.083	0	0.0
CARDICMYA COSTELLATA	0	0.0	1	0.187
ERVILIA CONCENTRICA	42	3.474	0	0.0
LEPTON SP.	1	0.083	4	0.749
LUCINA MULLINEATA	6	0.744	1	0.187
BAPYDEIA SOLENIIFORMIS	2	0.165	0	0.0
PERILLINA MARGARITACEUM	1	0.083	1	0.187
PITAE SIMPSONI	27	2.233	0	0.0
STRIGILLA MIRABILIS	14	1.158	1	0.187
TELLINA TAMPAENSIS	2	0.165	0	0.0
TELLINA TEXANA	21	1.737	9	1.685
TELLINA VERSICOLOR	78	6.452	23	4.307
VENERIDAE UNIDENTIFIED SP.	6	0.496	0	0.0
ANNELIDA (SEGMENTED WORMS)				
OLIGOCHAETA				
UNIDENTIFIED SP.	17	1.406	2	0.375
POLYCHAETA				
APOEICNCSILC PYGMAEA	1	0.083	0	0.0
ABERICCLA CRISTATA	1	0.083	0	0.0
ARMADIA MACULATA	9	0.744	9	1.685
AXIDIMELLA MUCOSA	2	0.165	0	0.0
BRANIA WELFLEETENSIS	7	0.579	0	0.0
CERATINEFIS INRIIABILIS	6	0.496	7	1.311
CHONE SP.	9	0.744	0	0.0
DIOPATRA CUPREA	1	0.083	2	0.375
ETECNE LACIEA	7	0.579	4	0.749
EULALIA SANGUINEA	1	0.083	1	0.187
GLYCERA AMERICANA	4	0.331	3	0.562
GLYCERA DIERANCHIATA	3	0.248	1	0.187
GLYCERA SP.	3	0.248	2	0.375
GLYCINE SOLITARIA	0	0.0	1	0.187
GONIATA LITTOREA	7	0.579	1	0.187
GYPTIS VITIATA	1	0.083	0	0.0
HAPLOSCHOLOPUS FOLIOSUS	1	0.083	0	0.0
ISOLLA PULCHELLA	1	0.083	0	0.0
LUMBERINERIS CRUZENSIS	476	39.371	170	21.835

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
8/24/76
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>MALACCCEUS INDICUS</u>	0	0.0	1	0.187
<u>MESOCHAETOPTERUS SAGITTARIUS</u>	11	0.910	13	2.434
<u>NEPHYS PICTA</u>	12	0.993	1	0.187
<u>NOTICASTIUS LATERICUS</u>	2	0.165	0	0.0
<u>ONUPHIS EREMITA CULATA</u>	34	2.812	3	0.562
<u>ONUPHIS NEBULOSA</u>	2	0.165	2	0.375
<u>OWENIA FUSIFORMIS</u>	4	0.331	0	0.0
<u>PARANITES SPECIOSA</u>	3	0.248	1	0.187
<u>PARACNICES LYRA</u>	2	0.165	0	0.0
<u>PARACNIS FULGENS</u>	2	0.165	1	0.187
<u>PARAPRIONOSPIO PINNATA</u>	1	0.083	2	0.375
<u>PHYLLODOCE ARENAE</u>	4	0.331	6	1.124
<u>POLYDORA TETRABRANCHIA</u>	1	0.083	0	0.0
<u>PRIONOSPIO CRISTATA</u>	101	8.354	114	21.348
<u>RULLIERINEREIS MEXICANA</u>	15	1.241	1	0.187
<u>SCOLOPLOS ARMIGER</u>	39	3.226	21	3.933
<u>SCOLOPLOS RUBRA</u>	0	0.0	1	0.187
<u>SIGALION ARENICOLA</u>	1	0.083	0	0.0
<u>SIGAMERA BASSI</u>	2	0.165	0	0.0
<u>SPIO PETTIBONEAE</u>	12	0.993	19	3.558
<u>SPIOCHAETOPTERUS OCULATUS</u>	2	0.165	0	0.0
<u>SPIOPHANES BOMBYX</u>	5	0.414	1	0.187
ARTHROPODA (CRUSTACEANS)				
AMPHIPODA				
<u>ACANTHOHAUSTORIUS SP.</u>	0	0.0	1	0.187
<u>AMPELISCA AEDITA</u>	1	0.083	1	0.187
<u>AMPELISCA VERRILLI</u>	31	2.564	27	5.056
<u>LISTRIELLA SP.</u>	1	0.083	0	0.0
<u>MONOCULODES SP.</u>	1	0.083	0	0.0
<u>PROTOKHAUSTORIUS SP.</u>	4	0.331	1	0.187
<u>PSEUDOKHAUSTORIUS SP.</u>	1	0.083	1	0.187
<u>PSEUDOPLATYISCHNOPUS SP.</u>	8	0.662	31	5.805
<u>SYNCELIDILM SP.</u>	4	0.331	1	0.187
ANCMURA				
<u>ALBINEA PARETII</u>	1	0.083	1	0.187
<u>PETROCHIRUS OTTOGENES</u>	0	0.0	1	0.187
<u>PETROLISTHES GALATHINUS</u>	0	0.0	1	0.187
BRACHYURA				
<u>CALLINECTES SP.</u>	2	0.165	0	0.0
<u>HEPATILUS EPHELIICUS</u>	1	0.083	0	0.0
<u>PINNIXIA RETINENS</u>	0	0.0	1	0.187
CARIDEA				
<u>OGYRIDES LIMICOLA</u>	1	0.083	0	0.0
CUMACEA				
<u>CYCLAPSIS SP.</u>	10	0.827	1	0.187
<u>CYCLAESIS VARIANS</u>	3	0.248	2	0.375
<u>XYUPESTYLIS SMITHI</u>	16	1.323	3	0.562
ISCOPODA				
<u>EDOLIA MONIOSA</u>	3	0.248	0	0.0
LEPTOSTRACA				
<u>NEBALIA SP.</u>	2	0.165	1	0.187
OSTRACCEA				
<u>UNICENTIFIED SP.</u>	13	1.075	1	0.187
PENAEIDAE				
<u>PENAEUS CUORARUM</u>	0	0.0	1	0.187
STENACROPODA				
<u>ACANTHOSQUILLA BIMINIENSIS</u>	1	0.083	0	0.0

ECHINODERMATA
ASTEROIDEA (STARFISHES)

TREASURE ISLAND MCTEL (STATION 1) - CONTROL AND EXPERIMENTAL
8/24/76
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
~~ASTROPECTEN ARTICULATUS~~	1	0.083	0	0.0
~~ECHINOIDEA (SAND DOLLARS; URCHINS)~~				
~~MELLITA QUINQUESPERFORATA~~	6	0.496	0	0.0
~~HCLOTURIDEA (SEA CUCUMBERS)~~				
~~LEPTOSYNAPTIA SP.~~	5	0.744	1	0.187
~~OPHIURIDEA (BRITTLE STARS)~~				
~~OPHICHRAGMUS WURDEMANI~~	2	0.165	0	0.0
~~UNIDENTIFIED SP.~~	5	0.414	1	0.187
HEMICHORDATA				
ENTEROPNEUSTA (ACORN WORMS)				
UNIDENTIFIED SP.	2	0.165	0	0.0
CEPHALOCORDATA (LANCELETS)				
BRANCHIOSTOMA FLORIDAE	5	0.744	11	2.060
VERTEBRATA				
PISCES (FISHES)				
LEPOPHIDIUM GRAELLSI	0	0.0	1	0.187
TOTALS	1209		534	
NO. SPECIES		24		60
NO. IND. PER M2		4836		2136
S-D INDEX - H*(LN)		2.8449		2.5827
EVENNESS - J		0.6421		0.6308

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
9/1/76

SPECIES	NO. OF IND. (C)		NO. OF IND. (E)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNIDARIA				
ANEMONIA (SEA ANEMONES)				
UNIDENTIFIED SP.	1	0.130	0	0.0
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	5	0.649	1	0.248
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	20	2.597	7	1.737
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	10	1.299	1	0.248
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
NATICA PLSILLA	5	0.649	0	0.0
OLIVELLA BULLULA	1	0.130	0	0.0
TEREBA DISLOCATA	2	0.260	0	0.0
TURBINILLA CORDATA	1	0.130	0	0.0
PELECYPODA (CLAMS)				
ERVILIA CONCENTRICA	30	3.896	1	0.248
LEPTON SP.	0	0.0	4	0.993
LUCINA MULTILINEATA	11	1.429	0	0.0
LYNSIA H. FLORICANA	3	0.390	0	0.0
PAPYRIDEA SOLENIIFORMIS	1	0.130	0	0.0
PITAE SIMPSONI	1	0.130	0	0.0
SEMELE PROFICUA	4	0.519	3	0.744
SYRIGILLA MIRABILIS	17	2.208	0	0.0
TELLINA IRIS	0	0.0	1	0.248
TELLINA VERSICOLOR	58	7.532	32	7.940
ANNELIDA (SEGMENTED WORMS)				
OLIGOCHAETA				
UNIDENTIFIED SP.	16	2.078	2	0.496
POLYCHAETA				
APCERICACSEIC PYGMAEA	1	0.130	0	0.0
ARICIDEA WASSI	1	0.130	0	0.0
ARMANIA MACULATA	2	0.260	0	0.0
AXICHELLA MUCOSA	2	0.260	0	0.0
BRANCHIASYCHIS AMERICANA	1	0.130	0	0.0
BRANIA BELLEFLEETENSIS	3	0.390	1	0.248
CERATOCHEILIS TERRIBILIS	16	2.078	14	3.474
CHAETILZNE GAYHEALIA	1	0.130	0	0.0
CHCNE SP.	6	0.779	3	0.744
CISTENIDES GULDII	1	0.130	0	0.0
ETECNE LACTEA	1	0.130	3	0.744
GLYCERA CIBRANCHIATA	3	0.390	1	0.248
GLYCERA SP.	2	0.260	0	0.0
GONIA LITTOREA	3	0.390	2	0.496
HAPLOSCLOPIOS FRAGILIS	4	0.519	0	0.0
LUMBERINERIS CRUZENSIS	283	36.753	207	51.365
MESOCHAETOPTERUS SAGITTARIUS	8	1.039	7	1.737
NEANTIES ACUMINATA	2	0.260	0	0.0
NEANTIES SUCCINEA	3	0.390	0	0.0
NEPHYS EUCERA	1	0.130	1	0.248
NEPHYS PICTA	17	2.208	3	0.744
CNUPIIS EREMITA OCULATA	17	2.208	3	0.744
CUNIA FUSIFORMIS	1	0.130	0	0.0
PARALNALES LYRA	1	0.130	1	0.248

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL

9/1/76
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>PARAPEIGNOSPIC PINNATA</u>	2	0.260	3	0.744
<u>PISTA PALMATA</u>	0	0.0	1	0.248
<u>POECILOCEPHEUS JOHNSONI</u>	0	0.0	2	0.496
<u>PRINCIPIDIA CRISTATA</u>	32	4.156	37	5.181
<u>PULLIDINEREIS MEXICANA</u>	21	2.727	9	2.233
<u>SCOLELEPIS TEXANA</u>	1	0.130	0	0.0
<u>SCOLLELIS ARMIGER</u>	50	6.494	17	4.218
<u>SIGAMERA TENTACULATA</u>	0	0.0	3	0.744
<u>SPIC PETTICORAE</u>	17	2.208	1	0.248
<u>SPIDICHAETOPTERUS OCULATUS</u>	3	0.390	0	0.0
<u>SPICEPLANES BOMBAYX</u>	6	0.779	2	0.496
 <u>SIPUNCULICA (PEANUT WORMS)</u>				
<u>GOLFINGIA TRICHOCEPHALA</u>	1	0.130	0	0.0
 <u>ARTHROPODA (CRUSTACEANS)</u>				
<u>AMPHIPODA</u>				
<u>ACANTHOCALANUS SP.</u>	9	1.169	0	0.0
<u>AMPELISCA ABDITA</u>	1	0.130	0	0.0
<u>AMPELISCA VERRILLI</u>	13	1.688	7	1.737
<u>COROPHUM SP.</u>	1	0.130	0	0.0
<u>MONOCLODES SP.</u>	0	0.0	4	0.993
<u>PROTOMALSTORUS SP.</u>	3	0.390	0	0.0
<u>PSEUDOCALANUS SP.</u>	4	0.519	0	0.0
<u>PSEUDOPALATIUS SP.</u>	2	0.260	3	0.744
<u>SYNGELIDUM SP.</u>	2	0.260	0	0.0
<u>ANOMURA</u>				
<u>ALBUEA PARETI</u>	1	0.130	0	0.0
<u>BRACHYLRA</u>				
<u>CALLINECTES SP.</u>	2	0.260	0	0.0
<u>PINNIXIA RETINENS</u>	1	0.130	0	0.0
<u>CALLINASSIDAE</u>				
<u>CALLINASSA JAMAICENSE</u>	2	0.260	0	0.0
<u>CARIDEA</u>				
<u>LATHEUTES PARVULUS</u>	1	0.130	0	0.0
<u>PROCESSA VICINA</u>	1	0.130	0	0.0
<u>CLMACEA</u>				
<u>CYCLAPSIS VARIANS</u>	1	0.130	2	0.496
<u>ORYZOPSIS SMITHI</u>	2	0.260	4	0.993
<u>MYSIDACEA</u>				
<u>UNIDENTIFIED SP.</u>	0	0.0	4	0.993
<u>OSTRACODA</u>				
<u>UNIDENTIFIED SP.</u>	6	0.779	4	0.993
<u>STOMATOPODA</u>				
<u>ACANTHOSQUILLA BIRMINGHAMI</u>	2	0.260	0	0.0
 <u>ECHINODERMATA</u>				
<u>ASTEROIDEA (STARFISHES)</u>				
<u>ASTROPECIEN ARTICULATUS</u>	1	0.130	0	0.0
<u>ECHINOIDEA (SAND DOLLARS, URCHINS)</u>				
<u>HELLITA QUINQUESPERFORATA</u>	3	0.390	0	0.0
<u>HOLOTHUROIDEA (SEA CUCUMBERS)</u>				
<u>LEPTOSYNAPTIA SP.</u>	1	0.130	2	0.496
<u>OPHIURIDEA (BRITTLE STARS)</u>				
<u>OPHIOBRANCHIUS BURDEMANI</u>	4	0.519	0	0.0
 <u>HEMICHORDATA</u>				
<u>ENTEROPNEUSTA (ACORN WORMS)</u>				
<u>UNIDENTIFIED SP.</u>	4	0.519	0	0.0
 <u>CEPHALOCORDATA (LANCELETS)</u>				

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
5/1/76
(CONTINUED)

SPECIES	NO. CF IND. (C)		NO. CF IND. (E)	
	TOTAL	PERCENT	TOTAL	PERCENT
BRANCHIOSIOMA FLORIDAE	4	0.519	0	0.0
TOTALS	770		403	
NO. SPECIES		74		38
NO. IND. PER 42		3080		1612
S-M INDEX - H' (LN)		2.8922		2.1365
EVENNESS - J		0.6720		0.5873

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
9/8/76

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNICARIA				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	4	0.708	0	0.0
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	1	0.177	0	0.0
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	23	4.071	6	1.786
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	4	0.708	0	0.0
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
ACTECCINA CANDEI	2	0.354	0	0.0
CYLICHNELLA BIDENTATA	1	0.177	1	0.298
NATICA PUSILLA	2	0.354	1	0.298
OLIVELLA BULLULA	1	0.177	0	0.0
PELECYPODA (CLAMS)				
ERVILIA CONCENTRICA	12	2.124	1	0.298
LEPTON SP.	1	0.177	2	0.595
LUCINA MULTIILINEATA	17	3.009	0	0.0
PAPYRIDEA SOLENIFORMIS	1	0.177	0	0.0
PERIPLOMA MARGARITACEUM	6	1.062	0	0.0
PITAR SIMPSONI	1	0.177	0	0.0
SEMELE PROFICUA	2	0.354	1	0.298
STRIGILLA MIRABILIS	3	0.531	0	0.0
TELLINA IRIS	0	0.0	1	0.298
TELLINA TEXANA	0	0.0	9	2.679
TELLINA VERSICOLOR	37	6.549	15	4.464
VENERIDAE UNIDENTIFIED SP.	3	0.531	1	0.298
ANNELIDA (SEGMENTED WORMS)				
OLIGOCHAETA				
UNIDENTIFIED SP.	10	1.770	1	0.298
POLYCHAETA				
APOPELICUSPIC PYGMAEA	1	0.177	4	1.190
ARICIDEA SP.	1	0.177	0	0.0
ARMANDIA AGILIS	0	0.0	2	0.595
ARMANDIA MACULATA	3	0.531	1	0.298
AXICHELLEA MUCCSA	1	0.177	0	0.0
FRANIA WELFLEETENSIS	1	0.177	0	0.0
CAPITELLIDES JONESI	1	0.177	0	0.0
CERATONEREIS IRRITABILIS	18	3.186	11	3.274
CERATONEREIS MIRABILIS	1	0.177	2	0.595
CHAETILINE GAYHEADIA	3	0.531	0	0.0
CHONE SP.	5	0.885	0	0.0
CISTENIDES GULDII	4	0.708	0	0.0
ETHECNE LACTEA	1	0.177	0	0.0
EULALIA SANGUINEA	1	0.177	0	0.0
GLYCERA CIERANCHIATA	5	0.885	1	0.298
GONIATA LITTOREA	15	2.655	1	0.298
EAPLESCLOPUS ROBUSTUS	1	0.177	0	0.0
HARMOTHE LUNULATA	1	0.177	0	0.0
ISOLIA PULCELLA	2	0.354	0	0.0
LUMENINERIS CRUZENSIS	150	26.549	112	32.333
LUMENINERIS TETRAURA	3	0.531	0	0.0
PAGELCNA SP.	3	0.531	0	0.0

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
9/8/76
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>MEDICMASTUS CALIFORNIENSIS</u>	3	0.531	0	0.0
<u>MESOCYCTOPTERUS SAGITTARIUS</u>	4	0.708	1	0.298
<u>NEANTHES SUCCINEA</u>	0	0.0	2	0.595
<u>NEPHYS PICTA</u>	15	2.655	1	0.298
<u>NOTOMASTUS HEMIPODUS</u>	0	0.0	1	0.298
<u>NOTOMASTUS LATERICEUS</u>	3	0.531	0	0.0
<u>ONUPHIS EREMITA OCULATA</u>	28	4.956	14	4.167
<u>ONUPHIS NEBULOSA</u>	1	0.177	0	0.0
<u>OWENIA FUSIFORMIS</u>	2	0.354	0	0.0
<u>PARANATHES SPECIOSA</u>	1	0.177	1	0.298
<u>PARACNIDES LYRA</u>	9	1.593	0	0.0
<u>PARACNIS FULGENS</u>	1	0.177	0	0.0
<u>PARAPRIONOSPIO PINNATA</u>	4	0.708	9	2.679
<u>PHYLLODOCE ARENAE</u>	10	1.770	8	2.381
<u>POECILOCHAETUS JOHNSONI</u>	0	0.0	2	0.595
<u>PRIONOSPIO CRISTATA</u>	17	3.009	26	7.738
<u>RULLIERINEREIS MEXICANA</u>	15	3.363	16	4.762
<u>SCOLOPLUS ARMIGER</u>	22	3.894	33	9.821
<u>SIGALICA ARENICOLA</u>	2	0.354	0	0.0
<u>SIGAMMEA BASSI</u>	1	0.177	0	0.0
<u>SIGAMBRA TENTACULATA</u>	2	0.354	0	0.0
<u>SPIC FETTERBONEAE</u>	1	0.177	14	4.167
<u>SPICHAETOPTERUS OCULATUS</u>	0	0.0	1	0.298
<u>SPICPHANES BOMBIX</u>	8	1.416	7	2.083
 <u>SIPUNCULICA (PEANUT WORMS)</u>				
<u>GOLFINGIA TRICHOCEPHALA</u>	2	0.354	0	0.0
 <u>ARTHROPODA (CRUSTACEANS)</u>				
<u>AMPHIPCCA</u>				
<u>AMPELISCA ABDITA</u>	1	0.177	1	0.298
<u>AMPELISCA VERRILLI</u>	2	0.354	4	1.190
<u>ARGISSA SP.</u>	0	0.0	1	0.298
<u>COROPHIUM SP.</u>	0	0.0	1	0.298
<u>MONICULODES SP.</u>	2	0.354	0	0.0
<u>PARAPHOXUS SP.</u>	2	0.354	0	0.0
<u>PSEUDOPLATYISCHNOPUS SP.</u>	1	0.177	1	0.298
<u>SYNCELICUM SP.</u>	3	0.531	0	0.0
<u>ANOMURA</u>				
<u>ALPHEA PAREII</u>	1	0.177	2	0.595
<u>ERACHYURA</u>				
<u>PINNIXIA RETINENS</u>	1	0.177	0	0.0
<u>CARIDEA</u>				
<u>PROCESSA HEMPHILLI</u>	3	0.531	0	0.0
<u>PROCESSA VICINA</u>	3	0.531	0	0.0
<u>CUMACEA</u>				
<u>CYCLAPSIS SP.</u>	6	1.062	1	0.298
<u>CYCLAPSIS VARIANS</u>	2	0.354	1	0.298
<u>CYLOSTYLIS SMITHI</u>	5	0.885	4	1.190
<u>LEPTOSTRACA</u>				
<u>NEBALIA SP.</u>	2	0.531	1	0.298
<u>MYSIDACEA</u>				
<u>UNIDENTIFIED SP.</u>	3	0.531	0	0.0
<u>OSTRACCCA</u>				
<u>UNIDENTIFIED SP.</u>	5	0.885	1	0.298
<u>PENAEIDAE</u>				
<u>PENAEUS OLICRAEUM</u>	1	0.177	0	0.0
<u>TRACHYPENAEUS CONSTRICTUS</u>	1	0.177	2	0.595
<u>TANAIDACEA</u>				
<u>UNIDENTIFIED SP.</u>	1	0.177	0	0.0

ECHINODERMATA

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
5/8/76
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
HOLOTHUROIDEA (SEA CUCUMBERS) <u>LEPTOSYNAPTE SP.</u>	10	1.770	0	0.0
HEMICHORDATA ENTEROPNEUSTA (ACORN WORMS) UNIDENTIFIED SP.	1	0.177	0	0.0
CEPHALCHORDATA (LANCELETS) <u>BRANCHIOSTOMA FLORIDAE</u>	1	0.177	7	2.083
TOTALS	565		336	
NO. SPECIES	83		47	
NO. IND. PER M2	2260		1344	
S-W INDEX - H' (LN)	3.3627		2.7387	
EVENNESS - J	0.7610		0.7113	

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
9/21/76

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNIDARIA				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	2	0.256	0	0.0
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	1	0.128	1	0.433
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	21	2.685	7	3.030
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	9	1.151	0	0.0
PHORONIDA (PHORONIDS)				
<u>PHORONIS ARCHITECTA</u>	2	0.256	0	0.0
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
<u>CYLICHNELLA BIDENTATA</u>	0	0.0	1	0.433
<u>NATICA PUSILLA</u>	3	0.384	0	0.0
<u>CLIVELLA BULLULA</u>	3	0.384	1	0.433
PELECYFODA (CLAMS)				
<u>ANADARA FLORIDANA</u>	1	0.128	0	0.0
<u>ERVILIA CONCENTRICA</u>	3	0.384	0	0.0
<u>LAEVICARDIUM PICTUM</u>	1	0.128	0	0.0
<u>LUCINA MULTILINEATA</u>	3	0.384	3	1.299
<u>SYRIGILLA MIRABILIS</u>	5	0.639	0	0.0
<u>TELLINA IRIS</u>	4	0.512	3	1.299
<u>TELLINA TEXANA</u>	6	1.023	1	0.433
<u>TELLINA VERSICOLOR</u>	31	3.964	16	6.926
ANNELIDA (SEGMENTED WORMS)				
CLIGGCHAETA				
UNIDENTIFIED SP.	15	1.918	0	0.0
POLYCHAETA				
<u>AMPHARETE ACUTIFRONS</u>	1	0.128	0	0.0
<u>APORRIONOSPION CYGMAEA</u>	0	0.0	3	1.299
<u>ARICIDEA FRAGILIS</u>	1	0.128	0	0.0
<u>ARICIDEA SP.</u>	2	0.256	0	0.0
<u>ARMANDIA AGILIS</u>	2	0.256	0	0.0
<u>ARMANDIA MACULATA</u>	3	0.384	1	0.433
<u>AXIOHELLA MUCOSA</u>	1	0.128	0	0.0
<u>BRANIA BELLEFLEETENSIS</u>	6	1.023	0	0.0
<u>CAPITELLA CAPITATA</u>	0	0.0	8	3.463
<u>CAULIERELLA SP.</u>	2	0.256	0	0.0
<u>CERATOCEREIS IRREGULARIS</u>	17	2.174	7	3.030
<u>CERATOCEREIS MIRABILIS</u>	2	0.256	0	0.0
<u>CHAETOCEREIS GAYHEADIA</u>	1	0.128	0	0.0
<u>CHONE SP.</u>	17	2.174	1	0.433
<u>CISTENIDES GULDII</u>	1	0.128	0	0.0
<u>DICPATRA CUPREA</u>	1	0.128	1	0.433
<u>EUCLEIA LACINIA</u>	3	0.384	0	0.0
<u>GLYCERA AMERICANA</u>	4	0.512	0	0.0
<u>GLYCERA DIBRANCHIATA</u>	2	0.256	1	0.433
<u>GLYCINE SOLITARIA</u>	1	0.128	0	0.0
<u>GONIATA LITTOREA</u>	3	0.384	0	0.0
<u>GRUENLEPIS MEXICANA</u>	1	0.128	0	0.0
<u>GYPSIS VITTATA</u>	1	0.128	0	0.0
<u>FABRICIOLUS FOLIOSUS</u>	2	0.256	0	0.0

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
9/21/76
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>HAPLICSCILLICLOS FRAGILIS</u>	6	0.767	0	0.0
<u>HAPLICSCILLICLOS ROBUSTUS</u>	4	0.512	0	0.0
<u>HEPTACMASTUS FILIFORMIS</u>	1	0.128	1	0.433
<u>LUMBERNERIS CRUZENSIS</u>	315	40.281	70	30.303
<u>LUMBERNERIS TETRAURA</u>	9	1.151	0	0.0
<u>MAGELCMA LONGICORNIS</u>	0	0.0	1	0.433
<u>MEDICMASTUS CALIFORNIENSIS</u>	2	0.256	0	0.0
<u>MESOCMAETICTEFUS SAGITTARIUS</u>	2	0.256	1	0.433
<u>NEANTHES SUCCINEA</u>	0	0.0	1	0.433
<u>NEPHIYS BUCFA</u>	1	0.128	0	0.0
<u>NEPHIYS FICIA</u>	7	0.895	1	0.433
<u>NOTCMASTUS HEMIPODUS</u>	1	0.128	0	0.0
<u>NOTCMASTUS LATERICEUS</u>	1	0.128	0	0.0
<u>ONUPHIS EREMITA CCULATA</u>	26	3.325	7	3.030
<u>ONUPHIS NEBULOSA</u>	1	0.128	0	0.0
<u>OWENIA FUSIFORMIS</u>	3	0.384	0	0.0
<u>PARANATHES SPECIOSA</u>	1	0.128	0	0.0
<u>PARACNIDES LYRA</u>	1	0.128	1	0.433
<u>PARACNIS FULGENS</u>	3	0.384	0	0.0
<u>PARACNIS SP.</u>	3	0.384	0	0.0
<u>PARAPRIONOSPIO PINNATA</u>	0	0.0	2	0.866
<u>PHYLLODOCE ARENAE</u>	7	0.895	10	4.329
<u>PRIONOSPIO CRISTATA</u>	25	3.197	12	5.195
<u>RULLIERINEREIS MEXICANA</u>	25	3.197	7	3.030
<u>SCOLELEPIS TEXANA</u>	1	0.128	0	0.0
<u>SCOLCPLCS ARMIGER</u>	40	5.115	7	3.030
<u>SIGAMBRA BASSI</u>	3	0.384	0	0.0
<u>SIGAMBRA TENTACULATA</u>	0	0.0	5	2.165
<u>SPIO PETTIBONEAE</u>	5	1.151	4	1.732
<u>SPIOPHANES BCMBYX</u>	5	0.639	4	1.732
<u>STHENELAIS BOA</u>	0	0.0	1	0.433
<u>STREPTOSYLLIS ARENAE</u>	1	0.128	0	0.0
SIPUNCULICA (PEANUT WORMS)				
<u>GOLFINGIA TRICHOCEPHALA</u>	2	0.256	0	0.0
ARTHROPOCA (CRUSTACEANS)				
AMPHIPCCA				
<u>ACANTHOHALSTORIUS SP.</u>	10	1.279	0	0.0
<u>AMPELISCA ABDITA</u>	4	0.512	0	0.0
<u>AMPELISCA VERRILLI</u>	19	2.430	6	2.597
<u>ARGISSA SP.</u>	0	0.0	1	0.433
<u>CARCIDIUM SP.</u>	3	0.384	0	0.0
<u>LISTEJELLA SP.</u>	1	0.128	1	0.433
<u>MONOCULODES SP.</u>	4	0.512	0	0.0
<u>PROTODHAUSTORIUS SP.</u>	1	0.128	0	0.0
<u>PSEUDODHAUSTORIUS SP.</u>	1	0.128	0	0.0
<u>PSEUDOPLATYSCHEMUS SP.</u>	7	0.895	7	3.030
<u>SYNCHELIDIUM SP.</u>	2	0.256	0	0.0
ANOMURA				
<u>ALBUNEA PARETII</u>	1	0.128	0	0.0
CARIDEA				
<u>PROCESSA FEMPHILLI</u>	2	0.256	1	0.433
CUMACEA				
<u>CYCLAESIS SP.</u>	2	0.256	0	0.0
<u>CYCLAESIS VARIANS</u>	0	0.0	2	0.866
<u>XYUETSTYLIS SMITHI</u>	3	0.384	1	0.433
LEPTOSTRACA				
<u>NERALIA SP.</u>	1	0.128	1	0.433
OSTRACCCA				
UNIDENTIFIED SP.	7	0.895	14	6.061

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
5/21/76
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
PENAEIDAE				
<u>SICYPTIA BREVIROSTRIS</u>	1	0.128	2	0.866
<u>TRACYPENAEUS CONSTRICTUS</u>	4	0.512	1	0.433
ECHINODERMATA				
ECHINOIDEA (SAND DOLLARS; URCHINS)				
<u>MELLITA QUINQUESPERFORATA</u>	4	0.512	0	0.0
HOLOTHURIDEA (SEA CUCUMBERS)				
<u>LEPTOSYNAPTE SP.</u>	5	0.639	3	1.299
OPHIURIDEA (BRITTLE STARS)				
UNIDENTIFIED SP.	1	0.128	0	0.0
HEMICHORCATA				
ENTEROPNEUSTA (ACORN WORMS)				
UNIDENTIFIED SP.	1	0.128	0	0.0
CEPHALOCORDATA (LANCELETS)				
<u>BRANCHIOSTOMA FLORIDAE</u>	5	0.639	1	0.433
TOTALS	782		231	
NO. SPECIES		89		45
NO. IND. PER M2		3128		524
S-W INDEX - H' (LN)		2.9755		2.9440
EVENNESS - J		0.6629		0.7734

TREASLE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
10/4/76

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	0	0.0	1	0.164
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	27	3.466	17	2.787
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	4	0.513	1	0.164
PHORONIDA (PHORONIDS)				
<u>PHORONIS ARCHITECTA</u>	1	0.128	0	0.0
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
<u>CYLICHNELLA BIDENTATA</u>	0	0.0	1	0.164
<u>OLIVA SAYANA</u>	1	0.128	0	0.0
<u>OLIVELLA FUSILLA</u>	1	0.128	0	0.0
<u>POLINICES DUPLICATUS</u>	1	0.128	0	0.0
PELECYFEDA (CLAMS)				
<u>ANADARA FLORIDANA</u>	3	0.385	0	0.0
<u>ERVILIA CONCENTRICA</u>	3	0.385	1	0.164
<u>LUCINA MULTILINEATA</u>	19	2.439	25	4.754
<u>PERIPLCMA MARGARITACEUM</u>	8	1.027	4	0.656
<u>PITAR SIMPSONI</u>	1	0.128	0	0.0
<u>STRIGILLA MIRABILIS</u>	1	0.128	0	0.0
<u>TELLINA AEGUISIRIATA</u>	0	0.0	1	0.164
<u>TELLINA IRIS</u>	3	0.385	2	0.328
<u>TELLINA TEXANA</u>	5	0.642	4	0.656
<u>TELLINA VESICOLOR</u>	33	4.236	19	3.115
VENERIDAE UNIDENTIFIED SP.	1	0.128	1	0.164
ANNELIDA (SEGMENTED WORMS)				
OLIGOCHAETA				
UNIDENTIFIED SP.	23	2.953	21	3.443
POLYCHAETA				
<u>AGLAPHAMUS VERRILLI</u>	1	0.128	0	0.0
<u>AMPHARETE ACUTICORNIS</u>	1	0.128	1	0.164
<u>APOPRIONOSPIO PYGMAEA</u>	1	0.128	0	0.0
<u>ARICIDEA FRAGILIS</u>	1	0.128	1	0.164
<u>ARICIDEA SP.</u>	1	0.128	0	0.0
<u>ARMANDIA AGILIS</u>	0	0.0	1	0.164
<u>ARMANDIA MACULATA</u>	0	0.0	3	0.492
<u>ASYCHIS CAROLINAE</u>	2	0.257	19	3.115
<u>BRANIA BELLEFLEETENSIS</u>	2	0.257	3	0.492
<u>CAPITELLA CAPITATA</u>	0	0.0	2	0.328
<u>CALLIERIELLA SP.</u>	0	0.0	1	0.164
<u>CERATONEREIS IRRITABILIS</u>	27	3.466	44	7.213
<u>CERATONEREIS MIRABILIS</u>	0	0.0	2	0.328
<u>CHAETONE GAYHEADIA</u>	0	0.0	2	0.328
<u>CHAETONE SETOSA</u>	1	0.128	0	0.0
<u>CHONE SP.</u>	6	0.770	3	0.492
<u>CISTENIDES GOULDII</u>	4	0.513	2	0.328
<u>DASYBRANCHIUS LUMBRICOIDES</u>	0	0.0	1	0.164
<u>DIOPATRA CIPREA</u>	1	0.128	1	0.164
<u>EUDONE LACSEA</u>	1	0.128	4	0.656
<u>GLYCERA AMERICANA</u>	1	0.128	4	0.656
<u>GLYCERA DIBRANCHIATA</u>	5	0.642	2	0.328
<u>GLYCERA SP.</u>	0	0.0	1	0.164
<u>GONIACA LITIGREA</u>	15	1.926	1	0.164

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
10/4/76
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>GYPTIS VITIATA</u>	2	0.257	2	0.328
<u>HAPLISCILLIUS FOLIOSUS</u>	2	0.257	5	0.820
<u>HAPLISCILLIUS FRAGILIS</u>	10	1.284	1	0.164
<u>HAPLISCILLIUS ROBUSTUS</u>	1	0.128	1	0.164
<u>HABRILICE LUNULATA</u>	1	0.128	0	0.0
<u>HEYTELMASTUS FILIFORMIS</u>	0	0.0	4	0.656
<u>ISCHIA PULCHELLA</u>	1	0.128	0	0.0
<u>LUMBERNEIS CRUZENSIS</u>	196	25.160	48	7.869
<u>LUMBERNEIS TETRAURA</u>	46	5.905	27	4.426
<u>MACRICLYMENE ZONALIS</u>	2	0.257	1	0.164
<u>MAGELMA SP.</u>	1	0.128	0	0.0
<u>MEDICMASTUS CALIFORNIENSIS</u>	1	0.128	3	0.492
<u>MEGALCMA BIOCUATUM</u>	0	0.0	4	0.656
<u>MESOCYPTOPTERUS SAGITTARIUS</u>	1	0.128	1	0.164
<u>MYCHESPIO PIGMENTATA</u>	1	0.128	0	0.0
<u>MYRICELE SP.</u>	0	0.0	1	0.164
<u>NEANTHES SUCCINEA</u>	1	0.128	0	0.0
<u>NEPHYS PICTA</u>	11	1.412	2	0.328
<u>NOTICMASTUS HEMIPODUS</u>	5	0.642	3	0.492
<u>NOTICMASTUS LATERICEUS</u>	1	0.128	0	0.0
<u>ONUPHIS EREMITA OCULATA</u>	36	4.621	22	3.607
<u>ONUPHIS NEBULOSA</u>	1	0.128	0	0.0
<u>OWENIA FUSIFORMIS</u>	1	0.128	7	1.148
<u>PARANICES LYRA</u>	15	1.926	7	1.148
<u>PARANICES FULGENS</u>	6	0.770	1	0.164
<u>PARANICES SP.</u>	4	0.513	1	0.164
<u>PARAFICINIS PINNATA</u>	4	0.513	20	3.279
<u>PHYLLICE ARENAE</u>	2	0.257	9	1.475
<u>POLYCHITES LUPINA</u>	1	0.128	0	0.0
<u>PRICINIS CRISTATA</u>	55	7.060	51	8.361
<u>PULLERINERIS MEXICANA</u>	29	3.723	17	2.787
<u>SCILLIUS ARMIGER</u>	47	6.033	11	1.803
<u>SCILLIUS EUERA</u>	1	0.128	2	0.328
<u>SIGALIN ARENICOLA</u>	1	0.128	0	0.0
<u>SIGAMERA TENTACULATA</u>	0	0.0	21	3.443
<u>SPIO PETTIBONEAE</u>	3	0.385	3	0.492
<u>SPIOPTANES BOMBYX</u>	13	1.669	11	1.803
<u>SIMENILAISS BDA</u>	0	0.0	1	0.164
 <u>SIPUNCULIDA (PEANUT WORMS)</u>				
<u>GOLFINGIA TRICHOCEPHALA</u>	3	0.385	6	0.984
 <u>ARTHROPODA (CRUSTACEANS)</u>				
<u>AMPHIPODA</u>				
<u>AMPELISCA AEDITA</u>	5	0.642	12	1.967
<u>AMPELISCA VERRILLI</u>	16	2.054	15	2.459
<u>CONCELIUM SP.</u>	0	0.0	1	0.164
<u>LISTEELLA SP.</u>	0	0.0	3	0.492
<u>PONCULIDES SP.</u>	2	0.257	1	0.164
<u>PARAPLOXUS SP.</u>	1	0.128	0	0.0
<u>PSEUDHAUSTORIUS SP.</u>	0	0.0	2	0.328
<u>PSEUDOPALYPSCHNOPUS SP.</u>	2	0.257	4	0.656
<u>ANOMURA</u>				
<u>ALBUEA PARETI</u>	4	0.513	12	1.967
<u>EUCERAMUS PRAELONGUS</u>	1	0.128	0	0.0
<u>BRACHYURA</u>				
<u>CALLINectes SAPIDUS</u>	1	0.128	0	0.0
<u>PERSEPHIA P. AQUILONARIS</u>	1	0.128	0	0.0
<u>PINNIXIA SAYANA</u>	0	0.0	3	0.492
<u>CALLINANASSIDAE</u>				
<u>CALLINANASSA JAMAICENSE</u>	2	0.257	1	0.164

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
10/4/76
(CONTINUED)

SPECIES	NO. OF IND. (C)		NO. OF IND. (E)	
	TOTAL	PERCENT	TOTAL	PERCENT
CARIDEA				
<u>PROCESSA HEMPHILLI</u>	2	0.257	2	0.328
CUMACEA				
<u>CYCLAPSIS SP.</u>	6	0.770	2	0.328
<u>CYCLAPSIS VARIANS</u>	0	0.0	2	0.328
<u>OXYUDOSTYLIS SMITHI</u>	3	0.385	5	0.820
MYSTACEA				
UNIDENTIFIED SP.	2	0.257	4	0.656
OSTRACODA				
UNIDENTIFIED SP.	6	0.770	14	2.295
PENAEIDAE				
<u>SICYONIA BREVIROSTRIS</u>	1	0.128	1	0.164
<u>TRACYPENAEUS CONSTRICTUS</u>	6	0.770	1	0.164
TANAIDACEA				
UNIDENTIFIED SP.	0	0.0	1	0.164
ECHINODERMATA				
ASTEROIDEA (STARFISHES)				
<u>ASTROPLECTEN ARTICULATUS</u>	2	0.257	3	0.492
HCLOTHUROIDEA (SEA CUCUMBERS)				
<u>LEPTOSYNAPTEA SP.</u>	5	0.642	26	4.262
OPHIURIDEA (BRITTLE STARS)				
<u>OPHIOPHRAGMUS MURDEMANNI</u>	1	0.128	0	0.0
CEPHALOCHOERDATA (LANCELETS)				
<u>BRANCHIOSTOMA FLORIDAE</u>	0	0.0	2	0.328
TOTALS	779		610	
NO. SPECIES		87		85
NO. IND. PER M2		3116		2440
S-B INDEX - H*(LN)		3.2650		3.7160
EVENNESS - J		0.7311		0.8364

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
10/18/76

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNIDARIA				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	1	0.102	0	0.0
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	2	0.204	0	0.0
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	15	1.534	4	1.423
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	32	3.272	3	1.068
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
OLIVA SAYANA	1	0.102	0	0.0
CLIVELLA BULLULA	0	0.0	1	0.356
TEREBA DISLOCATA	3	0.307	0	0.0
PELECYFODA (CLAMS)				
ERVILIA CONCENTRICA	6	0.613	1	0.356
LUCINA MULTILINEATA	5	0.511	11	3.915
MACRACALLISTA NIMBOSA	1	0.102	0	0.0
PERIPLOMA MARGARITACEUM	1	0.102	0	0.0
STRIGILLA MIRABILIS	17	1.738	0	0.0
TELLINA IRIS	2	0.204	0	0.0
TELLINA TEXANA	5	0.520	1	0.356
TELLINA VERSICOLOR	26	2.658	5	1.779
ANNELIDA (SEGMENTED WORMS)				
CLIGGCHAETA				
UNIDENTIFIED SP.	19	1.943	3	1.068
POLYCHAETA				
APOPRIONOSPION PYGMAEA	0	0.0	3	1.068
ARICIDEA CERRETTI	1	0.102	0	0.0
ARICIDEA SUECICA	7	0.716	0	0.0
ARMANCIA AGILIS	1	0.102	1	0.356
ARMANDIA MACULATA	10	1.022	0	0.0
BRANIA BELLIFLETENSIS	13	1.329	0	0.0
CERATONEREIS IRRITABILIS	5	0.511	11	3.915
CHAELOZOE SETIOSA	0	0.0	2	0.712
CHONE SP.	10	1.022	0	0.0
DIOPATRA CUPREA	1	0.102	0	0.0
DORVILLEA SOCIABILIS	3	0.307	0	0.0
EUCONE LACTEA	4	0.409	1	0.356
GLYCERA AMERICANA	9	0.920	9	3.203
GLYCERA DIBRANCHIATA	2	0.204	0	0.0
GONIADA LITICREA	1	0.102	0	0.0
GYPTIS VITTATA	2	0.204	0	0.0
HAPLOSCILLIPUS FOLIOSUS	4	0.409	2	0.712
HAPLOSCILLIPUS FRAGILIS	1	0.102	0	0.0
HAPLOSCILLIPUS ROBUSTUS	1	0.102	0	0.0
HARMONIE LUNULATA	0	0.0	1	0.356
LUMBERNERS CRUZENSIS	44	45.194	72	25.623
LUMBERNERS TETRAURA	2	0.204	0	0.0
LAGELINA SP.	1	0.102	0	0.0
MEDICINASTUS CALIFORNIENSIS	1	0.102	1	0.356
NEANITES ACUMINATA	1	0.102	1	0.356
NEPTIS PICTA	10	1.022	2	0.712
NOTOMASTUS HEMIPODUS	2	0.204	4	1.423

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
10/18/76
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>NOTICMASTUS LATERICEUS</u>	0	0.0	1	0.356
<u>ENUPHIS EREMITA CCULATA</u>	1	0.102	6	2.135
<u>CRIBRIA FISERI</u>	1	0.102	0	0.0
<u>LEVENIA FUSIECRMIS</u>	1	0.102	0	0.0
<u>PARACALLES SPECIOSA</u>	2	0.204	0	0.0
<u>PARACNIDES LYRA</u>	1	0.102	0	0.0
<u>PARACNIS FULGENS</u>	2	0.204	0	0.0
<u>PARACNIS SP.</u>	1	0.102	0	0.0
<u>PARAPRIONOSPIO PINNATA</u>	1	0.102	3	1.068
<u>PHYLLODOCE ARENAE</u>	0	0.0	2	0.712
<u>PRIONOSPIO CRISTATA</u>	77	7.873	27	9.609
<u>RULLIERINEREIS MEXICANA</u>	15	1.534	14	4.982
<u>SCOLELEPIS TEXANA</u>	1	0.102	0	0.0
<u>SCOLOPLOS ARMIGER</u>	37	3.783	6	2.135
<u>SCOLOPLOS RUBRA</u>	1	0.102	0	0.0
<u>SIGAMBRA BASSI</u>	3	0.307	0	0.0
<u>SIGAMBRA TENTACULATA</u>	0	0.0	3	1.068
<u>SPIO PETTIBONEAE</u>	2	0.204	2	0.712
<u>EPIOPLANES BOMBYX</u>	3	0.307	4	1.423
ARTHOPODIA (CRUSTACEANS)				
AMPHIPODA				
<u>AMPELISCA VERRILLI</u>	14	1.431	29	10.320
<u>ARGISSA SP.</u>	1	0.102	0	0.0
<u>LISIBIELLA SP.</u>	5	0.511	3	1.068
<u>MONOCLODES SP.</u>	19	1.943	6	2.135
<u>PARAPHOXUS SP.</u>	1	0.102	0	0.0
<u>PHOTIS SP.</u>	1	0.102	0	0.0
<u>PROTICHAUSTORIUS SP.</u>	7	0.716	0	0.0
<u>PSEUDICHAUSTORIUS SP.</u>	1	0.102	0	0.0
<u>PSEUDOPLATYLSCHNOPUS SP.</u>	28	2.863	20	7.117
<u>SYNCELIDIUM SP.</u>	3	0.307	1	0.356
<u>TIRON BISCOCELLATUS</u>	0	0.0	1	0.356
ANCMURA				
<u>ALBUEA FARETII</u>	3	0.307	4	1.423
<u>EUCERATUS PRAELONGUS</u>	1	0.102	2	0.712
<u>PAGUELS SP.</u>	1	0.102	1	0.356
CALLINANASSIDAE				
<u>CALLINANASSA JAMAICENSE</u>	0	0.0	1	0.356
CARTOEA				
<u>UGYRIDES LIMICOLA</u>	0	0.0	1	0.356
CUMACEA				
<u>CYCLAPSIS SP.</u>	31	3.170	1	0.356
<u>OXYUROSSTYLIS SMITHI</u>	0	0.0	1	0.356
MYSTICACEA				
<u>UNIDENTIFIED SP.</u>	1	0.102	0	0.0
OSTRACCA				
<u>UNIDENTIFIED SP.</u>	8	0.818	1	0.356
ECHINODERMATA				
ASTEROIDEA (STARFISHES)				
<u>ASTROPECTEN ARTICULATUS</u>	1	0.102	0	0.0
ECHINOIDEA (SAND DOLLARS; URCHINS)				
<u>MELLITA QUINQUESPERFORATA</u>	4	0.409	0	0.0
HOLOTHUROIDEA (SEA CUCUMBERS)				
<u>LEPTOSYNDAETA SP.</u>	1	0.102	1	0.356
OPHIURICEA (BRITTLE STARS)				
<u>OPHIOPHAGMUS BURDEMANI</u>	2	0.204	0	0.0
<u>UNIDENTIFIED SP.</u>	1	0.102	0	0.0
CEPHALOCORDATA (LANCELETS)				

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
11/1/76

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNIDARIA				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	3	0.397	0	0.0
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	2	0.265	2	0.391
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	18	2.384	6	1.174
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	8	1.060	1	0.196
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
NATICA PUSILLA	1	0.132	0	0.0
OLIVELLA BULLATA	1	0.132	3	0.587
TEREBRA DISLOCATA	1	0.132	0	0.0
PELECYFODA (CLAMS)				
CARDIOMYA COSTELLATA	0	0.0	1	0.196
ERVILIA CONCENTRICA	1	0.132	0	0.0
LUCINA MULTILINEATA	2	0.265	14	2.740
SEMELE PROFICUA	0	0.0	1	0.196
STRIGILLA MIRABILIS	10	1.325	0	0.0
TELLINA A. TAYLORIANA	0	0.0	1	0.196
TELLINA IRIS	0	0.0	3	0.587
TELLINA TEXANA	4	0.530	0	0.0
TELLINA VERSICOLOR	16	2.119	7	1.370
ANNELIDA (SEGMENTED WORMS)				
OLIGOCHAETA				
UNIDENTIFIED SP.	35	4.636	1	0.196
POLYCHAETA				
APOPEIRONOSPLO PYGMAEA	1	0.132	0	0.0
ARICIDEA SUECICA	5	0.662	0	0.0
ARMANDIA MACULATA	5	0.662	2	0.391
BRANIA CLAVATA	1	0.132	0	0.0
BRANIA WELFLEETIENSIS	12	1.589	0	0.0
CAULLERYELLA SP.	1	0.132	0	0.0
CERATONEREIS IRRITABILIS	11	1.457	18	3.523
CHAELOZOE SEIOSA	1	0.132	1	0.196
CHONE SP.	7	0.927	0	0.0
ETEOPE LACTEA	2	0.265	0	0.0
EUNICE ANTENNATA	1	0.132	0	0.0
EURYTHOE COMPLANATA	0	0.0	1	0.196
GLYCERA AMERICANA	7	0.927	2	1.566
GLYCERA DIERBRANCHIATA	2	0.265	1	0.196
GONIADA LITTORAEA	2	0.265	1	0.196
GYPTIS VITIATA	7	0.927	0	0.0
HAPLISCILLICHLIS FOLIOSUS	7	0.927	0	0.0
HAPLISCILLICHLIS FRAGILIS	5	0.662	2	0.391
HAPLISCILLOPS ROBUSTUS	0	0.0	1	0.196
LUMBRINERIS CALZENSIS	342	45.298	216	42.270
LUMBRINERIS TETRAURA	4	0.530	1	0.196
MAGELCNA LONGICORNIS	0	0.0	2	0.391
NEOICHAELIS CALIFORNIENSIS	2	0.265	2	0.391
NEANTHES SUCCINEA	0	0.0	1	0.196
NEPHYS BUCERA	2	0.265	0	0.0
NEPHYS PICTA	5	1.192	1	0.196

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
10/18/76
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>BRANCHIOSTICMA FLORIDAE</u>	21	2.147	1	0.356
TOTALS	978		281	
NO. SPECIES		77		46
NO. IND. PER M2		3912		1124
S-W INDEX - H*(LN)		2.6227		2.9372
EVENNESS - J		0.6038		0.7672

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
11/1/76
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
NOTOMASTUS HEMIPODUS	1	0.132	1	0.196
ONUPHIS EREMITA OCLATA	11	1.457	6	1.174
ONUPHIS NEBULOSA	2	0.265	2	0.391
ORBINIA RISERI	1	0.132	0	0.0
OBENIA FLSTIFORMIS	1	0.132	0	0.0
PARACNIDES LYRA	1	0.132	0	0.0
PARACNIS FULGENS	6	0.795	1	0.196
PARAFRICNSPIC PINNATA	0	0.0	8	1.566
PHYLLIDICE ARENAE	2	0.265	1	0.196
PRICNSPIC CRISTATA	15	1.987	14	2.740
RULLIERINEREIS MEXICANA	11	1.457	17	3.327
SCOLICELLS ARMIGER	30	3.974	10	1.957
SCOLICELLS RUBEA	3	0.397	0	0.0
SIGAMERA BASSI	1	0.132	0	0.0
SIGAMERA TENTACULATA	0	0.0	1	0.196
SPIC PETTIBONEAE	1	0.132	2	0.391
SPIDOPHONES BOMBYX	0	0.0	7	1.370
ARTHROPODA (CRUSTACEANS)				
AMPHIPODA				
ACANTHOFAUSTORIUS SP.	2	0.265	0	0.0
AMPELISCA VERRILLI	5	0.662	29	5.675
LISTEHLA SP.	3	0.397	3	0.587
MONOCULODES SP.	8	1.060	18	3.523
PROTOFAUSTORIUS SP.	6	0.795	0	0.0
PSEUDOFALSTORIUS SP.	18	2.384	0	0.0
PSEUDOPLATYSCHNOPUS SP.	65	8.609	66	12.916
SYNCELIDIUM SP.	0	0.0	1	0.196
ANOMURA				
ALBINEA PARETI	0	0.0	6	1.174
BRACHYURA				
OVALISES CELLATUS	3	0.397	1	0.196
PINNIXIA SAYANA	0	0.0	1	0.196
CALLINANASSIDAE				
CALLINANASSA JAMAICENSE	1	0.132	0	0.0
CARIDEA				
LEPTOCHELA SERRATORBITA	0	0.0	1	0.196
PROCESSA HEMPHILLI	2	0.265	2	0.391
CUMACEA				
CYCLAPSIS SP.	2	0.265	1	0.196
CYCLAPSIS VARIANS	2	0.265	0	0.0
MYSTICACEA				
UNIDENTIFIED SP.	1	0.132	2	0.391
OSTRACCA				
UNIDENTIFIED SP.	1	0.132	2	0.391
ECHINODERMATA				
ASTEROIDEA (STARFISHES)				
ASTROPECTEN AFICULATUS	2	0.265	2	0.391
ECHINOIDEA (SAND DOLLARS; URCHINS)				
MELLITA QUINQUESPERFORATA	1	0.132	0	0.0
HOLOTHUROIDEA (SEA CUCUMBERS)				
LEPTISYNAPTA SP.	1	0.132	2	0.391
OPHIURICEA (BRITTLE STARS)				
OPHIOPHRAGMUS MURDEMANI	0	0.0	1	0.196
CEPHALOCORDATA (LANCELETS)				
BRANCHIOSTOMA FLORIDAE	8	1.060	5	0.978

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
11/1/76
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
TOTALS	755		511	
NO. SPECIES		67		55
NO. IND. PER M2		3020		2044
S-H INDEX - H' (LN)		2.6057		2.4953
EVENNESS - J		0.6197		0.6227

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
12/1/76

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	29	3.766	15	1.695
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	26	3.377	2	0.226
PHORONIDA (PHORONIDS)				
<u>PHORONIS ARCHITECTA</u>	3	0.390	0	0.0
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
<u>NASSARIUS ACUTUS</u>	1	0.130	0	0.0
<u>POLINICES DUPLICATUS</u>	2	0.260	2	0.226
PELECYPODA (CLAMS)				
<u>ERVILIA CONCENTRICA</u>	1	0.130	0	0.0
<u>LUCINA MULTILINEATA</u>	12	1.558	6	0.678
<u>PERIELCMA MARGARITACEUM</u>	4	0.519	0	0.0
<u>SEMELE FRIGIDA</u>	2	0.260	0	0.0
<u>STRIGILLA MIRABILIS</u>	2	0.260	6	0.678
<u>TELLINA TEXANA</u>	0	0.0	5	0.565
<u>TELLINA VESICOLOR</u>	13	1.688	6	0.678
ANNELIDA (SEGMENTED WORMS)				
OLIGOCHAETA				
UNIDENTIFIED SP.	45	5.844	9	1.017
POLYCHAETA				
<u>AGLAPHAMUS VERRILLI</u>	1	0.130	0	0.0
<u>AMPHARETE ACUTIFRONS</u>	1	0.130	0	0.0
<u>APPELICATSEIC PYGMAEA</u>	3	0.390	3	0.339
<u>ARICIDEA CERUTI</u>	11	1.429	1	0.113
<u>ARICIDEA FRAGILIS</u>	1	0.130	1	0.113
<u>ARMANDIA AGILIS</u>	2	0.260	6	0.678
<u>ARMANDIA MACULATA</u>	5	0.649	9	1.017
<u>AXIOTELLA MUCOSA</u>	1	0.130	0	0.0
<u>ERANIA CLAVATA</u>	2	0.260	1	0.113
<u>ERANIA WELFLEETENSIS</u>	2	0.260	0	0.0
<u>CABIRA INCERTA</u>	0	0.0	1	0.113
<u>CERATONEREIS IRRITABILIS</u>	2	0.260	6	0.678
<u>CHAETIZONE SETOSA</u>	0	0.0	1	0.113
<u>CHONE SP.</u>	8	1.039	0	0.0
<u>CISTENICES GOULDII</u>	2	0.260	0	0.0
<u>ECKE LACTEA</u>	5	0.649	4	0.452
<u>GLYCERA AMERICANA</u>	3	0.390	8	0.904
<u>GLYCERA DIBRANCHIATA</u>	1	0.130	0	0.0
<u>GONIACA LITTOREA</u>	7	0.909	1	0.113
<u>GYPSIS VITTATA</u>	6	0.779	0	0.0
<u>HAPLOSCOLOPLOS FOLIOLIS</u>	8	1.039	11	1.243
<u>HAPLOSCOLOPLOS FRAGILIS</u>	10	1.299	3	0.339
<u>HAPLOSCOLOPLOS ROLLSIS</u>	1	0.130	0	0.0
<u>ISOCHA PULCHELLA</u>	1	0.130	0	0.0
<u>LUGDUNINERIS CRUZENSIS</u>	240	31.169	429	48.475
<u>LUGDUNINERIS TENUIS</u>	21	2.727	5	0.565
<u>MACELONA LONGICORNIS</u>	0	0.0	2	0.226
<u>MACELONA PEYRONIAE</u>	1	0.130	0	0.0
<u>MACELONA RIGIDA</u>	0	0.0	1	0.113
<u>MACELONATUS CALIFORNIENSIS</u>	2	0.260	0	0.0
<u>MACELONATUS CUBICERA</u>	1	0.130	0	0.0
<u>MACELONATUS ACUMINATA</u>	0	0.0	1	0.113
<u>MACELONATUS</u>	1	0.130	1	0.113
<u>MACELONATUS</u>	0	0.0	2	0.226

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
12/1/76
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>NOTOMASTUS HEMIPODUS</u>	1	0.130	4	0.452
<u>NOTOMASTUS LATERICEUS</u>	2	0.260	0	0.0
<u>ONUPHIS FREMITA OCULATA</u>	28	3.636	15	1.695
<u>ONUPHIS NEBULOSA</u>	4	0.519	0	0.0
<u>OWENIA FUSIFORMIS</u>	1	0.130	0	0.0
<u>PARACNIDES LYRA</u>	4	0.519	0	0.0
<u>PARACNIS FLIGENS</u>	1	0.130	3	0.339
<u>PARAFELICNSPIO PINNATA</u>	0	0.0	3	0.339
<u>PHYLLIDICE ARENAE</u>	0	0.0	4	0.452
<u>PRICNCSPIC CRISTATA</u>	41	5.325	55	6.215
<u>RULLIERINEREIS MEXICANA</u>	12	1.558	9	1.017
<u>SCOLICELIS ARMIGER</u>	34	4.416	44	4.972
<u>SCOLCPOS RUERA</u>	1	0.130	0	0.0
<u>SIGAMBRA TENTACULATA</u>	1	0.130	0	0.0
<u>SPIO PETTIBONEAE</u>	1	0.130	1	0.113
<u>SPIOPLANES BOMBYX</u>	4	0.519	5	0.565
ARTHROPODA (CRUSTACEANS)				
AMPHIPCA				
<u>ACANTHCHAUSTORIUS SP.</u>	1	0.130	3	0.339
<u>AMPELISCA VERRILLI</u>	10	1.299	23	2.599
<u>HYPERIA SP.</u>	14	1.818	0	0.0
<u>LISTRIELLA SP.</u>	4	0.519	4	0.452
<u>MONCCULDES SP.</u>	4	0.519	7	0.791
<u>PROTOKAUSTORIUS SP.</u>	0	0.0	9	1.017
<u>PSEUDOKAUSTORIUS SP.</u>	2	0.260	8	0.904
<u>PSEUDOPLATYISCHNORUS SP.</u>	79	10.260	117	13.220
<u>SYNCHERIDIUM SP.</u>	1	0.130	1	0.113
<u>TIRON SP.</u>	1	0.130	0	0.0
ANCHURA				
<u>ALBUNEA PARETI</u>	4	0.519	6	0.678
<u>PAGURUS LONGICARPUS</u>	2	0.260	0	0.0
EFACHYURA				
<u>GVALIFES CCELLATUS</u>	2	0.260	1	0.113
CALLIANASSIDAE				
<u>CALLIANASSA JAMAICENSE</u>	1	0.130	0	0.0
CARIDEA				
<u>HIPPOLYTE PLEURACANTHA</u>	1	0.130	0	0.0
<u>LEPTICELLA SERRATORBITA</u>	0	0.0	2	0.226
CUMACEA				
<u>CYCLAFSIS SP.</u>	1	0.130	1	0.113
<u>CYCLAFSIS VARIANS</u>	2	0.260	0	0.0
OSTRACCA				
<u>UNIDENTIFIED SP.</u>	3	0.390	2	0.226
PENAEIDA				
<u>TRACHYPENAEUS CONSTRICTUS</u>	1	0.130	1	0.113
ECHINODERMATA				
HOLOTHUROIDEA (SEA CUCUMBERS)				
<u>LEPTCSYNAPIA SP.</u>	3	0.390	0	0.0
CEPHALOCHERDATA (LANCELETS)				
<u>BRANCHIOSTOMA FLORIDAE</u>	10	1.299	9	1.017
TOTALS				
NO. SPECIES	770	74	885	54
NO. IND. PER M2		3080		3540
S-W INDEX - H'(LN)		2.9874		2.2595
EVENNESS - J		0.6941		0.5664

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
1/5/77

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNIDARIA				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	3	0.696	0	0.0
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	9	2.088	13	2.372
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	8	1.856	1	0.182
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
ACTECCINA CANDEI	1	0.232	0	0.0
POLINICES DUPLICATUS	0	0.0	1	0.182
PELECYPODA (CLAMS)				
LUCINA MULTILINEATA	1	0.232	3	0.547
PERIPLOMA MARGARITACEUM	0	0.0	3	0.547
STRIGILLA MIRABILIS	5	1.160	0	0.0
TELLINA TEXANA	1	0.232	0	0.0
TELLINA VERSICOLOR	18	4.176	1	0.182
ANNELIDA (SEGMENTED WORMS)				
OLIGOCHAETA				
UNIDENTIFIED SP.	57	13.225	2	0.365
POLYCHAETA				
AGLAOPHANTUS VERRILLI	1	0.232	0	0.0
APOPRIONOSPION PYGMAEA	0	0.0	1	0.182
ARICIDEA CERRUTI	2	0.464	0	0.0
ARMANDIA AGILIS	1	0.232	3	0.547
ARMANDIA MACULATA	3	0.696	0	0.0
BRANIA WELFLEETENSIS	1	0.232	0	0.0
CERATONEURIS IRRITABILIS	0	0.0	6	1.095
CHONE SP.	6	1.392	3	0.547
CISTENIODES GOULDII	0	0.0	1	0.182
DICPATRA CUPREA	1	0.232	0	0.0
GLYCERA AMERICANA	8	1.856	4	0.730
GONIADA LITIGEA	1	0.232	0	0.0
GYPTIS VITTATA	2	0.464	0	0.0
HAPLISCOCLOPUS FOLIOSUS	2	0.464	0	0.0
HAPLISCOCLOPUS FRAGILIS	0	0.0	1	0.182
LUMBERNEIS ACUTUS	4	0.928	0	0.0
LUMBERNEIS CRUZENSIS	15	3.480	343	62.591
LUMBERNEIS TENUIS	2	0.464	0	0.0
LUMBERNEIS TETRAURA	4	0.928	0	0.0
MACRICLYMENE ZONALIS	1	0.232	1	0.182
PAGELCNA LONGICORNIS	1	0.232	0	0.0
PAGELCNA SP.	2	0.464	0	0.0
NEPHTYS BUCERA	1	0.232	0	0.0
NEPHTYS PICTA	4	0.928	0	0.0
NOTOMASTUS HEMIPODUS	2	0.464	0	0.0
NOTOMASTUS LATERICEUS	0	0.0	1	0.182
CNUPTIS FREMIATA OCOLATA	0	0.0	3	0.547
PARACNICES LYRA	0	0.0	1	0.182
PARACNICES FULGENS	1	0.232	0	0.0
PARAPRIONOSPION PINNAIA	0	0.0	3	0.547
PHYLLODOCE ARENAE	0	0.0	3	0.547
PRIONOSPION CRISTATA	16	3.712	35	6.387
RULLIERINEREIS MEXICANA	13	3.016	2	0.365
SCOLELEPIS SQUAMATA	2	0.464	0	0.0
SCOLELEPIS TEXANA	3	0.696	0	0.0

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
1/5/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>SCOLOPLOS ARMIGER</u>	35	8.121	23	4.197
<u>SIGAMBRA BASSI</u>	1	0.232	0	0.0
<u>SPIO PETTIBONEAE</u>	1	0.232	0	0.0
<u>SPIOPHANES BOMBYX</u>	6	1.392	11	2.007
ARTHROPODA (CRUSTACEANS)				
AMPHIPODA				
<u>ACANTHHAUSTORIUS SP.</u>	3	0.696	0	0.0
<u>AMPELISCA VERRILLI</u>	1	0.232	10	1.825
<u>ERICHTHONIUS SP.</u>	2	0.464	0	0.0
<u>LISTRIELLA SP.</u>	2	0.464	1	0.182
<u>MONOCULODES SP.</u>	1	0.232	1	0.182
<u>PROTOKHALSTORIUS SP.</u>	15	3.480	4	0.730
<u>PSEUDOKHALSTORIUS SP.</u>	40	9.281	4	0.730
<u>PSEUDOPLATYISCHNOPUS SP.</u>	55	22.042	45	8.212
<u>SYNCHELIDIUM SP.</u>	0	0.0	3	0.547
ANOMURA				
<u>PAGURUS LONGICARPUS</u>	2	0.464	0	0.0
BRACHYURA				
<u>OVALIPES OCELLATUS</u>	2	0.464	2	0.365
<u>PINNIXIA SAYANA</u>	2	0.464	1	0.182
CARIDEA				
<u>HIPPILYTE FLEURACANTHA</u>	1	0.232	0	0.0
CUMACEA				
<u>CYCLAPSIS VARIANS</u>	1	0.232	0	0.0
ECHINODERMATA				
ECHINOIDEA (SAND DOLLARS; URCHINS)				
<u>MELLITA QUINQUESPERFORATA</u>	1	0.232	0	0.0
HOLOTHURIOIDEA (SEA CUCUMBERS)				
<u>LEPTISYNAPIA SP.</u>	2	0.464	3	0.547
HEMICHORDATA				
ENTEROPNEUSTA (ACORN WORMS)				
UNIDENTIFIED SP.	1	0.232	0	0.0
CEPHALOCORDATA (LANCELETS)				
<u>BRANCHIOSTOMA FLORIDAE</u>	15	3.480	5	0.912
TOTALS				
NO. SPECIES	431	56	548	36
NO. IND. PER M2		1724		2192
S-W INDEX - H ² (LN)		3.0102		1.7037
EVENNESS - J		0.7478		0.4754

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
2/2/77

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	16	4.222	16	2.893
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	18	4.749	0	0.0
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
ANACIS FLORIDANA	1	0.264	0	0.0
NATICA PUSILLA	4	1.055	1	0.181
OLIVA SAYANA	0	0.0	1	0.181
POLINICES DUPLICATUS	1	0.264	0	0.0
TURBOCILLA CONRADI	1	0.264	0	0.0
PELECYPODA (CLAMS)				
LUCINA MULTILINEATA	6	1.583	15	2.712
MACROCALLISTA NIMBOSA	2	0.528	0	0.0
PERIPLOMA MARGARITACEUM	1	0.264	0	0.0
PITAR SIMPSONI	1	0.264	0	0.0
STRIGILLA MIRABILIS	6	1.583	0	0.0
TELLINA TEXANA	2	0.528	1	0.181
TELLINA VESICOLOR	7	1.847	3	0.542
ANNELIDA (SEGMENTED WORMS)				
OLIGOCHAETA				
UNIDENTIFIED SP.	35	9.235	4	0.723
POLYCHAETA				
AGLAPLAMUS VERRILLI	0	0.0	1	0.181
APOLYCHOSPID PYGMAEA	1	0.264	0	0.0
ARICICEA CERRUTI	0	0.0	1	0.181
ARICICEA FRAGILIS	0	0.0	2	0.362
ARMANCIA AGILIS	1	0.264	1	0.181
ARMANDIA MACULATA	6	1.583	0	0.0
BRANIA CLAVATA	3	0.792	0	0.0
BRANIA WELFLEETENSIS	1	0.264	0	0.0
CHAETIZONE SETOSA	0	0.0	2	0.362
CHONE SP.	4	1.055	4	0.723
DIOPATRA CUPREA	0	0.0	2	0.362
EYONE LACTEA	0	0.0	1	0.181
GLYCERA AMERICANA	1	0.264	0	0.0
GLYCERA DIBRANCHIATA	0	0.0	2	0.362
GONIADA LITTOREA	1	0.264	2	0.362
GYPTIS VITIATA	1	0.264	0	0.0
HAPLCSOPOLOS FOLIOLUS	1	0.264	0	0.0
HAPLCSOPOLOS FRAGILIS	2	0.528	1	0.181
LUMBRINERIS CRUZENSIS	6	1.583	325	58.770
LUMBRINERIS TETRAURA	5	1.319	1	0.181
MAGELONA LONGICORNIS	0	0.0	4	0.723
MAGELONA SP.	1	0.264	0	0.0
MEDICASTYUS CALIFORNIENSIS	0	0.0	2	0.362
NEANTHES SP.	0	0.0	2	0.362
NEPHIYS BUCERA	1	0.264	0	0.0
NEPHIYS PICIA	3	0.792	3	0.542
NOTOMASTYS HEMIPODUS	1	0.264	0	0.0
OLYPIIS EREMITA OCULATA	2	0.528	32	5.787
PARACNIDES LYRA	1	0.264	0	0.0
PARACNIDES FULGENS	1	0.264	0	0.0
POLYDORA TETRAERANCHIA	2	0.528	1	0.181
PRICATSPIC CRISTATA	15	3.958	31	5.606
SCOLELERINERIS MEXICANA	7	1.847	7	1.266
SCOLELERIS SQUAMATA	5	1.319	2	0.362

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
2/2/77
(CONTINUED)

SPECIES	NO. CF IND. (C.)		NO. CF INC. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>SCOLELEPIS TEXANA</u>	5	2.375	4	0.723
<u>SCOLOPLOS ARMIGER</u>	28	7.388	2	0.362
<u>SIGAMBRA TENTACULATA</u>	0	0.0	1	0.181
<u>SPIO PETTIBONEAE</u>	1	0.264	0	0.0
<u>SPIOPHANES BCMBYX</u>	11	2.902	26	4.702
ARTHROPOCA (CRUSTACEANS)				
<u>AMPHIPCCA</u>				
<u>ACANTHOHAUSTORIUS SP.</u>	16	4.222	2	0.362
<u>AMPELISCA VERRILLI</u>	2	0.528	0	0.0
<u>LITORELLA SP.</u>	3	0.792	0	0.0
<u>MONACULODES SP.</u>	1	0.264	0	0.0
<u>PROCTHAUSTORIUS SP.</u>	34	8.971	2	0.362
<u>PSEUDOPHAUSTORIUS SP.</u>	0	0.0	1	0.181
<u>PSEUDOPLATYSCHEPUS SP.</u>	88	23.219	35	6.329
<u>SYNCALIDIDUM SP.</u>	2	0.528	0	0.0
<u>ANOMURA</u>				
<u>ALBUEA PARETI</u>	0	0.0	4	0.723
<u>ERACHYURA</u>				
<u>CYALIFES CELLATUS</u>	0	0.0	1	0.181
<u>CUMACEA</u>				
<u>CYCLAPSID VARIANS</u>	0	0.0	1	0.181
<u>OSTRACODA</u>				
<u>UNIDENTIFIED SP.</u>	3	0.792	1	0.181
<u>PENALIDEA</u>				
<u>SICYONIA BREVIROSTRIS</u>	0	0.0	1	0.181
ECHINODERMATA				
<u>ASTEROIDEA (STARFISHES)</u>				
<u>ASTROPECTEN ARTICULATUS</u>	1	0.264	0	0.0
<u>ECHINOIDEA (SAND DOLLARS; URCHINS)</u>				
<u>MOIRA ATEOPS</u>	1	0.264	0	0.0
<u>HOLOTHURIDEA (SEA CUCUMBERS)</u>				
<u>LEPTOSYNAPTIA SP.</u>	0	0.0	1	0.181
CEPHALOCORDATA (LANCELETS)				
<u>BRANCHIOSTOMA FLORIDAE</u>	6	1.583	1	0.181
TOTALS	379		553	
NO. SPECIES	53		44	
NO. INC. PER M2	1516		2212	
S-W INDEX - H' (LN)	3.0609		1.8770	
EVENNESS - J	0.7710		0.4960	

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
3/1/77

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	17	2.881	17	1.822
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	6	1.017	10	1.072
PHORONIDA (PHORONIDS)				
<u>PHORONIS ARCHITECTA</u>	1	0.169	0	0.0
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
<u>ACTECCINA CANALICULATA</u>	1	0.169	0	0.0
<u>ANACTIS FLORICANA</u>	1	0.169	0	0.0
<u>NATICA PUSILLA</u>	3	0.508	6	0.643
<u>CLIVA SAYANA</u>	1	0.169	0	0.0
<u>POLYCHITES DUPLICATUS</u>	4	0.678	1	0.107
<u>TURBENILLA CONRADI</u>	4	0.678	2	0.214
PELECYPOCA (CLAMS)				
<u>LUCINA MULTILINEATA</u>	8	1.356	6	0.643
<u>PERITELCA MARGARITACEUM</u>	3	0.508	1	0.107
<u>STRIGILLA MIRABILIS</u>	0	0.0	3	0.322
<u>TELLINA ACQUISTRATA</u>	1	0.169	0	0.0
<u>TELLINA VERSICOLOR</u>	10	1.695	9	0.965
ANNELIDA (SEGMENTED WORMS)				
OLIGOCHAETA				
UNIDENTIFIED SP.	55	10.000	5	0.536
POLYCHAETA				
<u>AGLAPHAMUS VERRILLI</u>	2	0.339	1	0.107
<u>AMPHARETE ACUTIGENS</u>	0	0.0	1	0.107
<u>APOEIONOSPIO PYGMAEA</u>	0	0.0	1	0.107
<u>ARICIDEA CERRETI</u>	2	0.339	5	0.536
<u>ARICIDEA FRAGILIS</u>	1	0.169	0	0.0
<u>ARICIDEA PHILBINAE</u>	4	0.678	0	0.0
<u>ARICIDEA SLEICHA</u>	1	0.169	0	0.0
<u>ARICIDEA TAYLORI</u>	1	0.169	0	0.0
<u>ARNANDIA AGILIS</u>	4	0.678	10	1.072
<u>BRANIA CLAVATA</u>	0	0.0	5	0.536
<u>BRANIA WELFLEETENSIS</u>	0	0.0	2	0.214
<u>CAPITELLA CAPITATA</u>	0	0.0	1	0.107
<u>CERATONEURIS IRRITABILIS</u>	0	0.0	2	0.214
<u>CERATONEURIS MIRABILIS</u>	0	0.0	3	0.322
<u>CHCNE SP.</u>	1	0.169	0	0.0
<u>EYELINE LACIEA</u>	1	0.169	0	0.0
<u>GLYCERA AMERICANA</u>	0	0.0	3	0.322
<u>GLYCERA DIBRANCHIATA</u>	3	0.508	1	0.107
<u>GONIATA LITTOREA</u>	2	0.339	0	0.0
<u>GYPTIS VITTATA</u>	3	0.508	2	0.214
<u>LAPLOSCHOLOPLOS FOLIOSUS</u>	2	0.339	0	0.0
<u>LAPLOSCHOLOPLOS FRAGILIS</u>	5	0.847	0	0.0
<u>LAPLOSCHOLOPLOS ROBUSTUS</u>	0	0.0	1	0.107
<u>LOIMIA VIRIDIS</u>	1	0.169	0	0.0
<u>LUMBERINERIS CRUZENSIS</u>	127	21.525	271	29.046
<u>LUMBERINERIS TETRAURA</u>	29	4.915	3	0.322
<u>MACHOCLYMENE ZONALIS</u>	1	0.169	2	0.214
<u>NEOMICASTUS CALIFORNIENSIS</u>	5	0.847	0	0.0
<u>MICHOPELTHALMUS ABERRANS</u>	0	0.0	1	0.107
<u>MINUSPIO CIRRIFERA</u>	1	0.169	0	0.0
<u>NEPHYS BUCERA</u>	0	0.0	2	0.214
<u>NEPHYS PICTA</u>	25	4.237	28	3.001

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
3/1/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>NOTICHAUSTORIUS HEMIPODUS</u>	3	0.508	2	0.214
<u>ONUPHIS EREMITA OCULATA</u>	11	1.864	26	2.787
<u>CRIBRIA RISEI</u>	0	0.0	1	0.107
<u>PARACNIDES LYRA</u>	4	0.678	1	0.107
<u>PARACNIS FULGENS</u>	1	0.169	4	0.429
<u>PHYLLODOCE ARENAE</u>	1	0.169	1	0.107
<u>POLYCORA SOCIALIS</u>	0	0.0	1	0.107
<u>PRIONOSPID CRISTATA</u>	29	4.915	146	15.648
<u>PSEUDOCURYTHOE AMBIGUA</u>	0	0.0	1	0.107
<u>RULLITERINEREIS MEXICANA</u>	12	2.034	3	0.322
<u>SCOLELEPIS SQUAMATA</u>	5	0.847	6	0.643
<u>SCOLELEPIS TEXANA</u>	0	0.0	9	0.965
<u>SCOLOPLOS ARMIGER</u>	43	7.288	48	5.145
<u>SCOLOPLOS RUBRA</u>	2	0.339	0	0.0
<u>SIGAMBRA TENTACULATA</u>	0	0.0	3	0.322
<u>SPID PETTIBONEAE</u>	0	0.0	4	0.429
<u>SPIOCHAETOPTERUS OCULATUS</u>	1	0.169	0	0.0
<u>SPIOPHANES BOMBYX</u>	57	9.661	112	12.004
<u>SIPHACULLICA (PEANUT WORMS)</u>				
<u>ASPIDOSIPHON SP.</u>	0	0.0	1	0.107
<u>ARTHROPODA (CRUSTACEANS)</u>				
<u>AMPHIPODA</u>				
<u>ACANTHOHAUSTORIUS SP.</u>	0	0.0	7	0.750
<u>AMPELISCA VERRILLI</u>	3	0.508	3	0.322
<u>LISTRIELLA SP.</u>	2	0.339	4	0.429
<u>MONOCLODUS SP.</u>	1	0.169	4	0.429
<u>PROTOMALSTORIUS SP.</u>	0	0.0	12	1.286
<u>PSEUDOMALSTORIUS SP.</u>	2	0.339	0	0.0
<u>PSEUDOPLATYSCHEIDOPUS SP.</u>	42	7.119	104	11.147
<u>SYNCELEIDIUM SP.</u>	1	0.169	0	0.0
<u>TIRCA BISCCELLATUS</u>	0	0.0	1	0.107
<u>ANCURA</u>				
<u>ALBINEA PARELLI</u>	1	0.169	6	0.643
<u>EUCERANUS PRAELONGUS</u>	1	0.169	0	0.0
<u>PAGURUS LONGICARPUS</u>	1	0.169	1	0.107
<u>BRACHYURA</u>				
<u>OVALIPES OCELLATUS</u>	2	0.339	5	0.536
<u>PINNIPEDIA SAYANA</u>	0	0.0	1	0.107
<u>CARIDEA</u>				
<u>HIPPELYTE PLEURACANTHA</u>	1	0.169	0	0.0
<u>CUMACEA</u>				
<u>CYCLAPSIS SP.</u>	1	0.169	0	0.0
<u>CYCLAPSIS VARIANS</u>	1	0.169	7	0.750
<u>OSTRACODA</u>				
<u>UNIDENTIFIED SP.</u>	8	1.356	2	0.214
<u>ECHINODERMATA</u>				
<u>ASTEROCIDEA (STARFISHES)</u>				
<u>ASTROPECTEN ARTICULATUS</u>	2	0.339	0	0.0
<u>HELOTHURIDEA (SEA CUCUMBERS)</u>				
<u>LEPTOSYNAPTIA SP.</u>	9	1.525	0	0.0
<u>OPHIURIDEA (BRITTLE STARS)</u>				
<u>OPHIOPHRAGMUS WARDENANI</u>	0	0.0	1	0.107
<u>CEPHALOCORDATA (LANCELETS)</u>				
<u>BRANCHIOSIGMA FLORIDAE</u>	3	0.508	1	0.107

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
3/1/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
TOTALS	590		933	
NO. SPECIES		64		62
NO. IND. PER M2		2360		3732
S-B INDEX - H' (LN)		3.0592		2.6117
EVENNESS - J		0.7356		0.6328

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
4/1/77

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNICARIA				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	2	0.304	0	0.0
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	11	1.672	17	2.163
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	27	4.103	0	0.0
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
NATICA PUSILLA	0	0.0	1	0.127
TEREBERA DISLOCATA	1	0.152	1	0.127
TURBOCILLA CONRADI	1	0.152	0	0.0
PELECYPODA (CLAMS)				
ANATINA ANATINA	1	0.152	0	0.0
LUCINA MULTILINEATA	7	1.064	3	0.382
PERILECMA MARGARITACEUM	3	0.456	0	0.0
SOLENYA SP.	0	0.0	5	0.636
STRIGILLA MIRABILIS	2	0.304	0	0.0
TELLINA VERTICOLOR	11	1.672	13	1.654
ANNELIDA (SEGMENTED WORMS)				
CLIGGCHAETA				
UNIDENTIFIED SP.	31	4.711	10	1.272
POLYCHAETA				
AGLAPHAMUS VERRILLI	1	0.152	0	0.0
APOBRIONOSPIO PYGMAEA	1	0.152	2	0.254
ARICIDEA FAUVELI	3	0.456	5	0.636
ARMANDIA AGILIS	5	0.760	5	0.636
ARMANDIA MACULATA	2	0.304	1	0.127
ERANIA WELFLEETENSIS	5	0.760	0	0.0
CERATOCEREIS MIRABILIS	0	0.0	2	0.254
CHONE SP.	1	0.152	2	0.254
CIRRATULIDAE UNIDENTIFIED SP.	0	0.0	1	0.127
DIOPATRA CUPREA	0	0.0	1	0.127
DISPIO UNCINATA	20	3.040	4	0.509
EUDONE LACTEA	1	0.152	2	0.254
GLYCERA AMERICANA	6	0.912	3	0.382
GLYCERA DIDRANCHIATA	1	0.152	1	0.127
GONIACA LITTOREA	1	0.152	0	0.0
GYDITE BREVIHALPA	5	0.760	0	0.0
HAPLISCLOPLOS FRAGILIS	0	0.0	1	0.127
LUMBRINERIS CRUZENSIS	32	4.863	126	23.664
LUMBRINERIS ERECTA	0	0.0	1	0.127
LUMBRINERIS TETRAURA	10	1.520	1	0.127
MAGELCNA LONGICORNIS	1	0.152	3	0.382
MAGELCNA SP.	3	0.456	1	0.127
MEDICMASTILS CALIFORNENSIS	1	0.152	3	0.382
NEPHIYS BUGERA	0	0.0	1	0.127
NEPHIYS PICTA	75	11.398	37	4.707
NOTOMASTILS HEMIPODUS	0	0.0	2	0.254
ONUPHIS EREMITA OCULATA	0	0.0	23	2.926
ONUPHIS PALLICA	0	0.0	1	0.127
PARACNIDES LYRA	1	0.152	0	0.0
PARACNIS FULGENS	8	1.216	1	0.127
PARAFICINSPIC PINNATA	3	0.456	4	0.509
PHYLLIDICE ARENAE	0	0.0	6	0.763
POLYCIUREUS EXIMIUS	0	0.0	1	0.127

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
4/1/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>POLYDORA SOCIALIS</u>	1	0.152	1	0.127
<u>POLYDORA TETRABRANCHIA</u>	3	0.456	0	0.0
<u>PRIONOSPIO CRISTATA</u>	26	3.951	51	6.489
<u>PSEUDOCERYTHOE AMBIGUA</u>	1	0.152	0	0.0
<u>RULLIERINEREIS MEXICANA</u>	6	0.912	2	0.254
<u>SCOLELEPIS TEXANA</u>	39	5.927	24	3.053
<u>SCOLOPLOS ARMIGER</u>	24	3.647	25	3.181
<u>SCOLICELUS RUERA</u>	7	1.064	0	0.0
<u>SIGAMBRA BASSI</u>	4	0.608	1	0.127
<u>SIGAMBRA TENTACULATA</u>	0	0.0	2	0.254
<u>SPIC PETTIBONEAE</u>	0	0.0	1	0.127
<u>SPICIFRANES BOMBYX</u>	102	15.502	316	40.204
 SIPUNCULIDA (PEANUT WORMS)				
<u>GOLFINGIA TRICHOCEPHALA</u>	0	0.0	1	0.127
 ARTHROPODA (CRUSTACEANS)				
AMPHIPODA				
<u>ACANTHCHAUSTORIUS SP.</u>	14	2.128	0	0.0
<u>AMPELISCA VERRILLI</u>	0	0.0	1	0.127
<u>ERICHTHINIUS SP.</u>	13	1.976	0	0.0
<u>MONACULODES SP.</u>	1	0.152	1	0.127
<u>PROTODAUSTORIUS SP.</u>	28	4.255	0	0.0
<u>PSEUDOPALATYISCHNOPUS SP.</u>	89	13.526	4	0.509
UNIDENTIFIED SP.	3	0.456	0	0.0
ANOMURA				
<u>ALBONEA PARETI</u>	2	0.304	2	0.254
CARIDEA				
<u>HIPPELYTIE PLEURACANTHA</u>	1	0.152	0	0.0
CUMACEA				
<u>CYCLAPSIUS SP.</u>	1	0.152	0	0.0
<u>CYCLAPSIUS VARIANS</u>	4	0.608	1	0.127
OSTRACODA				
UNIDENTIFIED SP.	1	0.152	1	0.127
TANAIDACEA				
UNIDENTIFIED SP.	1	0.152	0	0.0
 ECHINODERMATA				
ECHINOIDEA (SAND DOLLARS; URCHINS)				
<u>MOIRA ATROPS</u>	1	0.152	0	0.0
 CEPHALOCHOORDATA (LANCELETS)				
<u>BRANCHIOSTOMA FLORIDAE</u>	3	0.456	1	0.127
 TOTALS	658		786	
NO. SPECIES		57		52
NO. IND. PER M2		2632		3144
S-B INDEX - H'(LN)		3.0944		2.1706
EVENNESS - J		0.7654		0.5493

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
5/2/77

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	0	0.0	1	0.242
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	15	2.333	16	3.865
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	8	1.244	0	0.0
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
NATICA PUSILLA	0	0.0	1	0.242
TURBENILLA CONRADI	2	0.311	0	0.0
PELECYPODA (CLAMS)				
LEPTIS SP.	1	0.156	2	1.932
LUCINA MULTILINEATA	4	0.622	2	0.483
SOLEMYA VELUM	1	0.156	0	0.0
STRIGILLA MIRABILIS	0	0.0	1	0.242
TELLINA TEXANA	0	0.0	1	0.242
TELLINA VERSICOLOR	9	1.400	0	0.0
ANNELIDA (SEGMENTED WORMS)				
CLIGOCHAETA				
UNIDENTIFIED SP.	29	4.510	7	1.691
POLYCHAETA				
AMPHARETE ACUTIERCENS	0	0.0	1	0.242
APODEIONOSPIO PYGMAEA	0	0.0	2	0.483
ARICIDEA FAUVELI	4	0.622	5	1.208
ARICIDEA FRAGILIS	5	0.778	4	0.966
ARICIDEA PHILBINAE	0	0.0	1	0.242
ARMANDIA AGILIS	1	0.156	0	0.0
BRANIA WELFLEETENSIS	2	0.311	2	0.483
CAPITELLA CAPITATA	0	0.0	3	0.725
CERATINERIS MIRABILIS	2	0.311	5	1.208
CHCNE SP.	2	0.311	1	0.242
DISPAT UNGINATA	4	0.622	4	0.966
ETECNE LACTEA	0	0.0	2	0.483
GLYCERA AMERICANA	2	0.311	3	0.725
GYPTIS BREVIDALPA	3	0.467	0	0.0
HAPLISCLOPLCS FOLIOSUS	13	2.022	7	1.691
HAPLISCLOPLCS FRAGILIS	4	0.622	0	0.0
LOPIA MEDUSA	1	0.156	0	0.0
LUMBERINERIS CRUZENSIS	81	12.597	62	14.976
LUMBERINERIS TETRAURA	15	2.333	3	0.725
MAGELLANA LONGICORNIS	1	0.156	2	0.483
MAGELLANA SP.	5	0.778	3	0.725
MEDICINASTUS CALIFORNIENSIS	3	0.467	2	0.483
NEPHYS BUCERA	1	0.156	1	0.242
NEPHYS PICTA	90	13.997	58	14.010
NOTOMASTUS HEMIPODUS	1	0.156	2	0.483
ONUPPIS EREMITA OCULATA	0	0.0	5	1.208
OPHELIA SP.	2	0.311	0	0.0
PARACNICES LYRA	5	0.778	1	0.242
PARACNIS FULGENS	1	0.156	0	0.0
PARAPHIONOSPIO PINNATA	3	0.467	50	12.077
PHYLLODOCE ARENAE	3	1.244	9	2.174
PODARKE OBSCURA	0	0.0	1	0.242
POECILOCHAETUS JOHNSONI	0	0.0	1	0.242
PRIONOSPIO CRISTATA	14	2.177	10	2.415

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
5/2/77
(CONTINUED)

SPECIES	NO. OF INC. (C.)		NO. OF INC. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>BULLI ERINEREIS MEXICANA</u>	4	0.622	2	0.483
<u>SCOLELEPIS TEXANA</u>	2	0.311	1	0.242
<u>SCOLCELOS ARMIGER</u>	0	0.0	3	0.725
<u>SCOLOPLOS RUBRA</u>	0	0.0	14	3.382
<u>SIGAMERA BASSI</u>	1	0.156	0	0.0
<u>SIGAMERA TENTACULATA</u>	1	0.156	0	0.0
<u>SPIO PETTIBONEAE</u>	11	1.711	0	0.0
<u>SPIOPANES BOMBYX</u>	157	24.417	89	21.498
<u>STHELELAIS BOA</u>	0	0.0	1	0.242
ARTHROPODA (CRUSTACEANS)				
AMPHIPODA				
<u>AMPELISCA ABDITA</u>	0	0.0	1	0.242
<u>AMPELISCA VERRILLI</u>	9	1.400	0	0.0
<u>LISTRILLA SP.</u>	3	0.467	1	0.242
<u>MICROPOTOPUS SP.</u>	0	0.0	1	0.242
<u>PROCEAUSTORIUS SP.</u>	0	0.0	2	0.483
<u>PSAUDOAUSTORIUS SP.</u>	1	0.156	0	0.0
<u>PSAUDOPALATYISCHNOPUS SP.</u>	92	14.308	0	0.0
<u>SYNGELIDIUM SP.</u>	4	0.622	2	0.483
BRACHYURA				
<u>MEICECHAPIS CALCARATA</u>	0	0.0	1	0.242
<u>PILIDIA LUNZI</u>	1	0.156	0	0.0
<u>PILIDIA SAYANA</u>	1	0.156	0	0.0
CANIOLA				
<u>PROCESSA MEMPHILLI</u>	1	0.156	0	0.0
CLAPSEA				
<u>CYCLAPSIS SP.</u>	1	0.156	1	0.242
<u>CYCLAPSIS VARIANS</u>	4	0.622	1	0.242
OSTRACODA				
<u>CAPOCYTHERIDEA SEPTIPUNCTATA</u>	3	0.467	0	0.0
<u>UNIDENTIFIED SP.</u>	1	0.156	1	0.242
ECHINODERMATA				
ASTEROIDEA (STARFISHES)				
<u>ASTROPECTEN ARTICULATUS</u>	1	0.156	2	0.483
OPHTHIOIDEA (BRITTLE STARS)				
<u>UNIDENTIFIED SP.</u>	0	0.0	1	0.242
HEMICHORCATA				
ENTEROPNEUSTA (ACORN WORMS)				
<u>UNIDENTIFIED SP.</u>	1	0.156	0	0.0
CEPHALOCHORDATA (LANCELETS)				
<u>BRANCHIOSTOMA FLORIDAE</u>	2	0.311	3	0.725
TOTALS				
NO. SPECIES	643	55	414	54
NO. INC. PER M2		2572		1656
S-B INDEX - H'(LN)		2.7186		2.8260
EVENNESS - J		0.6784		0.7085

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
6/1/77


SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	15	3.846	26	3.194
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	4	0.810	0	0.0
PHORONIDA (PHORONIDS)				
PHORONIS ARCHITECIA	0	0.0	2	0.246
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
CYLICNELLA BIDENTATA	2	0.405	1	0.123
DIASICMA VARIUM	0	0.0	1	0.123
PELECYPOCA (CLAMS)				
ANADAEA FLORIDANA	0	0.0	1	0.123
CUMINGIA TELLINOIDES	0	0.0	4	0.491
LEPTA SP.	0	0.0	57	7.002
LUCINA MULTILINEATA	16	3.239	33	4.054
MACTEA SP.	0	0.0	1	0.123
PITAR SIMPSONI	4	0.810	0	0.0
STRIGILLA MIRABILIS	1	0.202	0	0.0
TELLINA TEXANA	0	0.0	2	0.246
TELLINA VERSICOLOR	17	3.441	59	7.248
VENERIDAE UNIDENTIFIED SP.	0	0.0	3	0.369
ANNELIDA (SEGMENTED WORMS)				
CLIGOCCHAETA				
UNIDENTIFIED SP.	23	4.656	24	2.948
PCLYCHAETA				
AMPHARETE ACUTIFRONS	0	0.0	1	0.123
APOPRIONOSPIO PYGMAEA	0	0.0	3	0.369
ARICIDEA FAUVELI	4	0.810	3	0.369
ARICIDEA FRAGILIS	1	0.202	2	0.246
ARMANDOIA MACULATA	0	0.0	3	0.369
BRANIA WELFLEETENSIS	1	0.202	0	0.0
CAPITELLA CAPITATA	0	0.0	1	0.123
CERATONEREIS IRRITABILIS	0	0.0	1	0.123
CERATONEREIS MIRABILIS	0	0.0	2	0.246
CHONE SP.	9	1.822	4	0.491
DIOPATRA CUPREA	0	0.0	11	1.351
DISPIO UNCINATA	0	0.0	4	0.491
ETEONE LACTEA	4	0.810	5	0.614
GLYCERA AMERICANA	12	2.429	14	1.720
GLYCERA DIBRANCHIATA	0	0.0	1	0.123
GLYCINE SOLITARIA	0	0.0	3	0.369
GONIADA LITTOREA	3	0.607	0	0.0
GYPTIS VITATA	0	0.0	1	0.123
HAPLOSCOLOPLOS FOLIOSUS	7	1.417	11	1.351
HAPLOSCOLOPLOS FRAGILIS	0	0.0	1	0.123
HARMOTHOE LUNULATA	1	0.202	0	0.0
LUMBERNERIS CRUZENSIS	29	5.870	51	6.265
LUMBERNERIS TETRAURA	9	1.822	6	0.737
MAGELLANA SP.	4	0.810	0	0.0
MEDICASTUS CALIFORNIENSIS	0	0.0	6	0.737
MINUSPIC CIRRIFERA	0	0.0	3	0.369
MYRISCHELE SP.	0	0.0	1	0.123
NEPHYS BUCERA	10	2.024	0	0.0
NEPHYS PICTA	99	20.040	150	18.428
NOTICASTUS HEMIPODUS	2	0.405	0	0.0
CNUPHIS EREMITA OCULATA	27	5.466	26	3.194

NATIONAL MARINE FISHERIES SERVICE PANAMA CITY BEACH F--ETC F/G 8/1
BENTHIC COMMUNITY RESPONSE TO DREDGING BORROW PITS, PANAMA CITY--ETC(U)
MAR 82 C H SALOMAN, S P NAUGHTON, J L TAYLOR DACW72-81-M-0198
CERC-MR-82-3 NL

CEBC-HR-A2-3

[illegible]

END
DATE
FILMED
8 82
RIV



TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
6/1/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>ONLPHIS PALLIDA</u>	1	0.202	0	0.0
<u>PARANAITES SPECIOSA</u>	1	0.202	0	0.0
<u>PARACNIDES LYRA</u>	6	1.215	6	0.737
<u>PARAPRIONOSPION PINNATA</u>	0	0.0	27	3.317
<u>PHYLLODOCE ARENAE</u>	2	0.405	5	0.614
<u>PRIONOSPION CRISTATA</u>	21	4.251	84	10.319
<u>RUILLIERINEREIS MEXICANA</u>	5	1.012	1	0.123
<u>SCOLELEPIS TEXANA</u>	5	1.012	7	0.860
<u>SIGAMBRA BASSI</u>	1	0.202	0	0.0
<u>SIGAMBRA TENTACULATA</u>	0	0.0	5	1.106
<u>SPIO PETTIBONEAE</u>	8	1.619	0	0.0
<u>SPIOPHANES DOMOYX</u>	18	3.644	50	6.143
ARTHROPOCCA (CRUSTACEANS)				
AMPHIPODA				
<u>ACANTHOMAUSTORIUS SP.</u>	20	4.049	0	0.0
<u>AMPELISCA ABDITA</u>	0	0.0	3	0.369
<u>AMPELISCA VAGORUM</u>	0	0.0	2	0.246
<u>AMPELISCA VERRILLI</u>	14	2.834	3	0.369
<u>ARGISSA SP.</u>	0	0.0	2	0.246
<u>LEPTACTYLUS SP.</u>	0	0.0	1	0.123
<u>LITRIELLA SP.</u>	3	0.607	0	0.0
<u>LYSIANDOPSIS SP.</u>	1	0.202	1	0.123
<u>PROTOHAUSTORIUS SP.</u>	10	2.024	1	0.123
<u>PSEUDOMAUSTORIUS SP.</u>	3	0.607	0	0.0
<u>PSEUDOPLATYSCHEILES SP.</u>	16	3.239	22	2.703
<u>SYNCELIDIUM SP.</u>	9	1.822	4	0.491
ANOMURA				
<u>ALBONEA PARETII</u>	0	0.0	1	0.123
BRACHYURA				
<u>QUALIFES OCELLATUS</u>	1	0.202	1	0.123
<u>PINNIXIA CYLINDRICA</u>	0	0.0	1	0.123
<u>PINNIXIA RETINENS</u>	2	0.405	1	0.123
<u>PINNIXIA SAYANA</u>	1	0.202	0	0.0
CARIDEA				
<u>PROCESSA HEMPHILLI</u>	2	0.405	5	0.614
CUMACEA				
<u>CYCLAPSIS SP.</u>	3	0.607	1	0.123
<u>CYCLAPSIS VARIANS</u>	9	1.822	10	1.229
<u>OXYURASTYLIS SMITHI</u>	0	0.0	14	1.720
LEPTOSTRACA				
<u>NEBALIA SP.</u>	1	0.202	1	0.123
MYSIDACEA				
<u>UNIDENTIFIED SP.</u>	1	0.202	0	0.0
OSTRACOCA				
<u>SARSIELLA CHILDI</u>	0	0.0	1	0.123
PENAIODEA				
<u>SICYCNIA BREVIROSTRIS</u>	0	0.0	1	0.123
ECHINODERMATA				
ASTEROIDEA (STARFISHES)				
<u>ASTROPECTEN ARTICULATUS</u>	1	0.202	0	0.0
ECHINOIDEA (SAND DOLLARS; URCHINS)				
<u>LYTECHINUS VARIEGATUS</u>	1	0.202	0	0.0
<u>MELLITA QUINQUESPERFORATA</u>	7	1.417	8	0.983
OPHIUROIDEA (BRITTLE STARS)				
<u>UNIDENTIFIED SP.</u>	5	1.012	12	1.474
CEPHALOCORDATA (LANCELETS)				
<u>BRANCHIOSTOMA FLORIDAE</u>	8	1.619	1	0.123

APPENDIX B (CONTINUED)

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
6/1/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
TOTALS	494		814	
NO. SPECIES	55		69	
NO. IND. PER M2	1976		3256	
S-B INDEX - H' (LN)	3.3330		3.1585	
EVENNESS - J	0.8317		0.7554	

TREASURE ISLAND HOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
7/5/77

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNIDARIA				
ACTINIDARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	0	0.0	1	0.204
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	3	0.368	0	0.0
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	25	3.064	15	3.055
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	5	0.613	0	0.0
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
ACTEOCINA CANDEI	28	3.431	0	0.0
CYLICINELLA BIDENTATA	0	0.0	1	0.204
NATICA PUSILLA	0	0.0	1	0.204
POLINICES DUPLICATUS	0	0.0	1	0.204
PELECYPODA (CLAMS)				
LAEVICARDIUM MORTONI	1	0.123	0	0.0
LEPICK SP.	10	1.225	0	0.0
LUCINA MULTILINEATA	73	8.946	25	5.092
MACRCCALLISTA MACULATA	2	0.245	0	0.0
PERIPLOMA MARGARITACEUM	3	0.368	0	0.0
TELLINA AEGUSTRATA	3	0.368	2	0.407
TELLINA TEXANA	14	1.716	14	2.851
TELLINA VERSICOLOR	81	9.926	58	11.813
VENERIDAE UNIDENTIFIED SP.	2	0.245	0	0.0
ANNELIDA (SEGMENTED WORMS)				
CLIGOCMAETA				
UNIDENTIFIED SP.	14	1.716	1	0.204
POLYCHAETA				
APOEIONOSPIO PYGMAEA	2	0.245	2	0.407
ARICIDEA CERRUTI	0	0.0	1	0.204
ARICIDEA FAUVELI	3	0.368	6	1.222
ARICIDEA FRAGILIS	4	0.490	2	0.407
ARICIDEA SUECICA	0	0.0	1	0.204
ARMANDIA AGILIS	0	0.0	2	0.407
ARMANDIA MACULATA	0	0.0	1	0.204
CERATONEREIS IRRITABILIS	0	0.0	1	0.204
CHONE SP.	14	1.716	3	0.611
CIRROPHORUS LYRIFORMIS	0	0.0	1	0.204
DISPIO UNCINATA	0	0.0	3	0.611
EYBINE LACIEA	5	0.613	1	0.204
GLYCERA AMERICANA	41	5.025	16	3.259
GLYCERA DIBRANCHIATA	0	0.0	1	0.204
GONIADA LITTOREA	8	0.980	11	2.240
GRUPELEPIS MEXICANA	1	0.123	0	0.0
GYPHIS VITIATA	1	0.123	0	0.0
HAPLISCLOPLOS FOLIOSUS	6	0.735	1	0.204
HAPLISCLOPLOS FRAGILIS	1	0.123	0	0.0
LUMBRINERIS CRUZENSIS	154	18.873	90	18.330
LUMBRINERIS TETRAURA	24	2.941	1	0.204
NEODINASTYUS CALIFORNIENSIS	3	0.368	1	0.204
MICROPHIALMUS SP.	2	0.245	0	0.0
NEPHYS BUCERA	2	0.245	0	0.0
NEPHYS PICTA	112	13.725	129	26.273

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
7/5/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>NOTOMASTUS HEMIPODUS</u>	2	0.245	1	0.204
<u>ONUPHIS EREMITA OCULATA</u>	22	2.696	10	2.037
<u>OWENIA FUSIFORMIS</u>	9	1.103	0	0.0
<u>PARANAITES SPECIOSA</u>	1	0.123	0	0.0
<u>PARACNIDES LYRA</u>	10	1.225	26	5.295
<u>PARACNIS FULGENS</u>	3	0.368	0	0.0
<u>PARAPRIONOSPIO PINNATA</u>	2	0.245	8	1.629
<u>PHYLLODOCE ARENAE</u>	2	0.245	2	0.407
<u>PRIONOSPIO CRISTATA</u>	13	1.593	5	1.018
<u>RULLIERINEREIS MEXICANA</u>	1	0.123	0	0.0
<u>SCOLELEPIS TEXANA</u>	1	0.123	0	0.0
<u>SCOLOPLOS RUBRA</u>	2	0.245	0	0.0
<u>SIGALICA AENEICOLA</u>	1	0.123	0	0.0
<u>SPICEHANES BCMBYX</u>	1	0.123	13	2.648
 SIPUNCULIDA (PEANUT WORMS)				
UNIDENTIFIED SP.	1	0.123	0	0.0
 ARTHROPODA (CRUSTACEANS)				
AMPHIPODA				
<u>ACANTHOMAUSTORIUS SP.</u>	1	0.123	0	0.0
<u>AMPELISCA VADORUM</u>	1	0.123	0	0.0
<u>AMPELISCA VERRILLI</u>	9	1.103	7	1.426
<u>ARGISSA SP.</u>	1	0.123	0	0.0
<u>LISTRIELLA SP.</u>	4	0.490	2	0.407
<u>MICROPOTOPUS SP.</u>	3	0.368	0	0.0
<u>MONOCULODES SP.</u>	1	0.123	0	0.0
<u>PSEUDOPLATYISCHNOPUS SP.</u>	22	2.696	1	0.204
<u>SYNCELIDUM SP.</u>	21	2.574	0	0.0
UNIDENTIFIED SP.	1	0.123	0	0.0
ANCHURA				
<u>ALBINEA PAREII</u>	0	0.0	1	0.204
BRACHYLARA				
<u>PINNIXIA CHAETOPTERANA</u>	1	0.123	0	0.0
<u>PINNIXIA RETINENS</u>	3	0.368	0	0.0
CARIDEA				
<u>OGYRIDES ALPHAEROSTRIS</u>	0	0.0	1	0.204
<u>SYNALPHEUS SP.</u>	0	0.0	1	0.204
CUMACEA				
<u>CYCLAESIS SP.</u>	2	0.245	0	0.0
<u>CYCLAESIS VARIANS</u>	15	1.838	4	0.815
<u>XYULISTYLIS SMITHI</u>	0	0.0	2	0.407
PERAIDEA				
<u>PENAEUS DUORARUM</u>	1	0.123	2	0.407
 ECHINODERMATA				
ASTEROIDEA (STARFISHES)				
<u>ASTROPECTEN ARTICULATUS</u>	0	0.0	1	0.204
ECHINOIDEA (SAND DOLLARS; URCHINS)				
<u>MELITA QUINQUESPERECRATA</u>	2	0.245	0	0.0
HOLOTHURIDEA (SEA CUCUMBERS)				
<u>LEPTOSYNAPIA SP.</u>	0	0.0	10	2.037
OPHIOURIDEA (BRITTLE STARS)				
UNIDENTIFIED SP.	0	0.0	1	0.204
 HEMICHORCATA				
ENTEROPNEUSTA (ACRON WORMS)				
UNIDENTIFIED SP.	2	0.245	0	0.0
 CEPHALOCORDATA (LANCELETS)				
<u>BRANCHIOSTOMA FLORIDAE</u>	98	0.980	0	0.0

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
7/5/77
(CONTINUED)

SPECIES	NO. OF IND. (C)	NO. OF IND. (E)
	TOTAL PERCENT	TOTAL PERCENT
TOTALS	816	491
NO. SPECIES	64	49
NO. IND. PER M2	3264	1964
S-B INDEX - H'(LN)	3.0767	2.6678
EVENNESS - J	0.7398	0.6855

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
8/2/77

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNICARIA				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	0	0.0	1	0.137
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	11	0.851	3	0.411
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	33	2.554	16	2.192
BRACHIOPODA (LAMP SHELLS)				
GLOTTIDIA PYRAMIDATA	3	0.232	0	0.0
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
ACTECCINA CANDEI	9	0.697	8	1.096
CYLICNELLA BIDENTATA	0	0.0	73	10.000
DIASOMA VARIUM	24	1.858	0	0.0
NATICA PUSILLA	10	0.774	2	0.274
POLINICES DUPLICATUS	1	0.077	0	0.0
TEREERA DISLOCATA	1	0.077	0	0.0
TURBINILLA CONRADI	3	0.232	1	0.137
PELECYPODA (CLAMS)				
ANADARA FLORICANA	0	0.0	3	0.411
ERYLLIA CONCENTRICA	26	2.012	0	0.0
LUCINA MULTILINEATA	31	2.399	15	2.055
NUCULANA ACUTA	0	0.0	6	0.822
TELLINA AEGUISTRIATA	6	0.464	5	0.685
TELLINA TAMPAENSIS	1	0.077	9	1.233
TELLINA TEXANA	0	0.0	6	0.822
TELLINA VERSICOLOR	180	13.932	102	13.973
TRACHYCARDIUM MURICATUM	4	0.310	1	0.137
ANNELIDA (SEGMENTED WORMS)				
OLIGOCHAETA				
UNIDENTIFIED SP.	18	1.393	0	0.0
POLYCHAETA				
APOPRIONOSPIO PYGMAEA	3	0.232	1	0.137
ARICIDEA CERRUTI	1	0.077	0	0.0
ARICIDEA FRAGILIS	3	0.232	0	0.0
ARMANDIA MACULATA	12	0.929	3	0.411
BRANCHIOASYCHIS AMERICANA	0	0.0	5	0.685
CERATIONEREIS IRRITABILIS	2	0.155	18	2.466
CHONE SP.	31	2.399	0	0.0
CISTENIDES GOULDII	0	0.0	1	0.137
DIOPATRA CLPREA	0	0.0	33	4.521
DORVILLEA SOCIABILIS	3	0.232	0	0.0
ENOPLORHYNCHUS SANGUINEUS	1	0.077	0	0.0
EUCONE LACTEA	9	0.697	0	0.0
GLYCERA AMERICANA	35	2.709	0	0.0
GLYCERA DIBRANCHIATA	7	0.542	4	0.548
GLYCINE SOLITARIA	0	0.0	4	0.548
GONIADA LITTOREA	10	0.774	1	0.137
GYPIIS BREVIPALPA	4	0.310	0	0.0
HAPLISCLOPELOS FRAGILIS	1	0.077	0	0.0
HARMOTHOE IMBRICATA	0	0.0	1	0.137
LODINIA MEDUSA	3	0.232	3	0.411
LUMBRINERIS CRUZENSIS	373	28.870	2	1.096
LUMBRINERIS TETRAURA	21	1.625	1	0.137

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL

8/2/77
(CONTINUED)

SPECIES	NO. OF IND. (C)		NO. OF IND. (E)	
	TOTAL	PERCENT	TOTAL	PERCENT
MAGELCNA LONGICORNIS	0	0.0	2	0.274
MEDIONASTUS CALIFORNIENSIS	0	0.0	10	1.370
NEOSCHAEYOPTERUS SAGITTARIUS	10	0.774	24	3.288
MINUSPID CIRRIFERA	0	0.0	3	0.411
NEANTLES SUCCINEA	0	0.0	14	1.918
NEPHIYS BUCERA	4	0.310	0	0.0
NEPHIYS PICTA	65	5.031	7	0.959
NEREIS SP.	0	0.0	1	0.137
NOTOMASTUS HEMIPODUS	4	0.310	3	0.411
ONUPPIS EREMITA OCULATA	16	1.238	4	0.548
OWENIA FUSIFORMIS	1	0.077	1	0.137
PARACNIDES LYRA	15	1.161	2	0.274
PARACNIDES FULGENS	4	0.310	0	0.0
PARABRIONOSPID PINNATA	0	0.0	11	1.507
PHYLLODOCE ARENAE	9	0.697	4	0.548
POLYDORA SOCIALIS	0	0.0	2	0.274
POLYDORA TETRABRANCHIA	1	0.077	0	0.0
PRIONOSPID CRISTATA	96	7.430	75	10.274
PSEUDOSYTHOE AMBIGUA	1	0.077	0	0.0
RULLIERINEREIS MEXICANA	2	0.155	1	0.137
SIGAMBRA BASSI	1	0.077	0	0.0
SIGAMBRA TENTACULATA	2	0.155	109	14.932
SPID PETTIBONEAE	6	0.464	0	0.0
SPIOPHANES BOMBYX	6	0.464	0	0.0
STHELELAIS BOA	0	0.0	3	0.411
STREPTOSYLLIS ARENAE	1	0.077	0	0.0
SIPUNCULICA (PEANUT WORMS)				
UNIDENTIFIED SP.	1	0.077	1	0.137
ECHIURIDA (ECHIURIDS)				
UNIDENTIFIED SP.	0	0.0	3	0.411
ARTHOPODA (CRUSTACEANS)				
AMPHIPCCA				
ACANTHOHAUSTORIUS SP.	1	0.077	0	0.0
AMPELISCA VERRILLI	46	3.560	9	1.233
ERICHTHONIUS SP.	1	0.077	2	0.274
LISIBELLA SP.	0	0.0	2	0.274
PSEUDOPLATYSCYNOPUS SP.	20	1.548	1	0.137
SYNCELIDIUM SP.	20	1.548	6	1.096
ANOMURA				
ALBONEA PARETII	2	0.155	0	0.0
BRACHYURA				
CSACHILA TUBEFOSA	2	0.155	0	0.0
PANCREUS HERBERTI	0	0.0	8	1.096
PINNIXIA RELINENS	2	0.155	0	0.0
PINNIXIA SP.	0	0.0	2	0.274
PINNIXIAES OSTREUM	1	0.077	0	0.0
PORTULUS SAYI	0	0.0	6	0.822
CALLINASSIDAE				
CALLINASSA JAMAICENSE	1	0.077	1	0.137
CARIDEA				
LATREUTES PARVULUS	0	0.0	3	0.411
PROCESSA FEMPHILLI	6	0.464	4	0.548
CUMACEA				
CYCLASSIS SP.	6	0.464	1	0.137
CYCLASSIS VARIANS	8	0.619	9	1.233
CRYDERSYLLIS SHIIMI	9	0.697	1	0.137
ISCPODA				
APANTHURA MAGNIFICA	1	0.077	0	0.0

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
8/2/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
LEPTOSTRACA				
<u>NEBALIA SP.</u>	8	0.619	2	0.274
MYSIDACEA				
<u>MYSIDOPSIS BIGELOWI</u>	3	0.232	0	0.0
OSTRACCA				
UNIDENTIFIED SP.	4	0.310	7	0.959
PENAEIDAE				
<u>ACETES AMERICANUS</u>	2	0.155	0	0.0
<u>SICYPTIA SP.</u>	1	0.077	1	0.137
<u>BRACHYDONTES CONRICTUS</u>	0	0.0	11	1.507
ECHINODERMATA				
ECHINIDEA (SAND DOLLARS; URCHINS)				
<u>LYTECHINUS VARIEGATUS</u>	2	0.155	0	0.0
<u>NOVA ATRIPS</u>	2	0.155	34	4.658
HOLOTHUROIDEA (SEA CUCUMBERS)				
<u>LEPTOSYNAPTIA SP.</u>	4	0.310	2	0.274
OPHIURIDEA (BRITTLE STARS)				
<u>HEMIPHOLIS ELCNGATA</u>	0	0.0	1	0.137
<u>MICROPHOLIS GRACILLINA</u>	2	0.155	0	0.0
<u>OPHIOPHAGUS BURDEMANI</u>	1	0.077	0	0.0
HEMICHORCATA				
ENTEROPNEUSTA (ACORN WORMS)				
UNIDENTIFIED SP.	1	0.077	1	0.137
CEPHALOCORDATA (LANCELETS)				
<u>BRANCHIOSTOMA FLORIDAE</u>	10	0.774	0	0.0
VERTEBRATA				
PISCES (FISHES)				
Gobiidae, UNIDENTIFIED SP.	1	0.077	0	0.0
TOTALS	1292		730	
NO. SPECIES	80		70	
NO. IND. PER M2	5168		2920	
S-W INDEX - H'(LN)	3.0096		3.2331	
EVENNESS - J	0.6868		0.7610	

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
9/1/77

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNICARIA				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	1	0.112	2	1.818
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	1	0.112	0	0.0
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	26	2.912	5	4.545
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	5	0.560	0	0.0
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
ACTEOPCINA CANDEI	4	0.448	0	0.0
ANACIS FLORIDANA	3	0.336	0	0.0
CYLICHNELLA BIDENTATA	3	0.336	0	0.0
DIASOMA VARIUM	116	12.990	0	0.0
NASSARIUS ACUTUS	0	0.0	9	8.182
NATICA PLSILLA	4	0.448	2	1.818
TEREBRA DISLOCATA	4	0.448	2	1.818
TURBNILLA CONRADI	1	0.112	0	0.0
PELECYPODA (CLAMS)				
ANADARA FLORIDANA	0	0.0	1	0.909
ERVILIA CONCENTRICA	28	3.135	0	0.0
LUCINA MULTILINEATA	9	1.008	0	0.0
NUCULANA ACUTA	0	0.0	1	0.909
PERIPLOMA MARGARITACEUM	15	1.680	0	0.0
TELLINA AEGUISTRIATA	4	0.448	0	0.0
TELLINA TEXANA	6	0.672	7	6.364
TELLINA VERSICOLOR	138	15.454	13	11.818
ANNELIDA (SEGMENTED WORMS)				
OLIGOCHAETA				
UNIDENTIFIED SP.	18	2.016	1	0.909
POLYCHAETA				
AGLAPHAMUS VERRILLI	1	0.112	0	0.0
APOEIONOSPION PYGMAEA	2	0.224	1	0.909
ARICIDEA FAUVELI	4	0.448	0	0.0
ARICIDEA FRAGILIS	2	0.224	0	0.0
ARICIDEA SUECICA	1	0.112	0	0.0
ARMANCIA AGILIS	1	0.112	0	0.0
BRANIA WELFLEETENSIS	3	0.336	0	0.0
CAPITELLA CAPITATA	0	0.0	1	0.909
CAULLERYELLA SP.	1	0.112	0	0.0
CERATONEREIS IRRADIABILIS	0	0.0	4	3.636
CHONE SP.	13	1.456	0	0.0
DIOPATRA CUPREA	0	0.0	3	2.727
DRIESCHIA PELLUCIDA	0	0.0	1	0.909
EIEONE LACIEA	3	0.336	0	0.0
GLYCERA AMERICANA	3	0.336	0	0.0
GLYCERA DIBRANCHIATA	10	1.120	2	1.818
GLYCINE SOLITARIA	0	0.0	1	0.909
GONIACA LITTOREA	7	0.784	0	0.0
HAPLOSCELICUS FOLIOSUS	2	0.224	0	0.0
LOPIA VIRIDIS	1	0.112	0	0.0
LUMBERNERIS CRUZENSIS	252	28.219	1	0.909
LUMBERNERIS TETRAURA	8	0.896	0	0.0

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL

9/1/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
MEDICMASTUS CALIFORNIENSIS	1	0.112	0	0.0
MESOCALOPTERUS SAGITTARIUS	1	0.112	0	0.0
NEANIMES ACUMINATA	1	0.112	0	0.0
NEANIMES SUCCINEA	0	0.0	1	0.909
NEPHIYS PICTA	14	1.568	1	0.909
NOTOMASTUS HEMIPODUS	5	0.560	1	0.909
ONUPHIS EREMITA OCULATA	22	2.464	1	0.909
OMENIA FLISIFORMIS	1	0.112	0	0.0
PARACNIDES LYRA	25	2.800	0	0.0
PARAFRICOSPID PINNATA	0	0.0	5	4.545
PHYLLOCOCE ARENAE	6	0.672	0	0.0
FRICOSPIC CRISTATA	2	0.224	0	0.0
ESEUDEURYTHOE AMBIGUA	1	0.112	0	0.0
BULLYERIDEIS MEXICANA	10	1.120	2	1.818
SCCLELEPIS TEXANA	3	0.336	0	0.0
SIGAMEA BASSI	1	0.112	0	0.0
SIGAMBRA TENTACULATA	0	0.0	1	0.909
SPIOPHANES BOMBYX	1	0.112	0	0.0
SIPLUNCULIDA (PEANUT WORMS)				
GOLFINGIA TRICHOCEPHALA	1	0.112	0	0.0
ARTHROPODA (CRUSTACEANS)				
AMPHIPODA				
ACANTHOMHAUSTORIUS SP.	3	0.336	0	0.0
AMPELISCA VERRILLI	6	0.672	0	0.0
LEPTACTYLUS SP.	2	0.224	0	0.0
LISTERIELLA SP.	1	0.112	0	0.0
PSEUDOPLATYISCHNOPUS SP.	28	3.135	0	0.0
SYNCELIDION SP.	13	1.456	0	0.0
ANCHURA				
ALBUNEA PARETI	2	0.224	0	0.0
PAGURUS LONGICARPUS	5	0.560	0	0.0
BRACHYURA				
PLANILIA SAYANA	0	0.0	1	0.909
CUMACEA				
CYCLAPSIS SP.	2	0.224	0	0.0
CYCLAPSIS VARIANS	14	1.568	0	0.0
ISCOPEA				
EDOTEA MCNIOSA	1	0.112	0	0.0
MYSIDACEA				
BOWMANIELLA SP.	0	0.0	2	1.818
MYSIDOPSIS BIGELOWI	0	0.0	2	1.818
UNIDENTIFIED SP.	1	0.112	0	0.0
OSTRACODA				
UNIDENTIFIED SP.	1	0.112	0	0.0
PENAEIDEA				
ACEIES AMERICANUS	0	0.0	2	1.818
ECHINODERMATA				
ECHINOIDEA (SAND DOLLARS; URCHINS)				
MOIRA AISCPS	1	0.112	0	0.0
PELLITA QUINQUESPERFORATA	6	0.672	0	0.0
HOLOTHURIDEA (SEA CUCUMBERS)				
LEPTOSYNAPTIA SP.	0	0.0	1	0.909
OPHIUROIDEA (BRITTLE STARS)				
HEMIPOLOIS ELONGATA	0	0.0	28	25.455
MICROPHOLIS GRACILLIMA	3	0.336	0	0.0
HEMICHORDATA				
ENTEROPNEUSTA (ACRON WORMS)				

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
9/1/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
UNIDENTIFIED SP.	1	0.112	5	4.545
CEPHALCCHCRDATA (LANCELETS) BRANCHIOSTOMA FLORIDAE	6	0.672	0	0.0
VERTEBRATA PISCES (FISHES) SYMPLEUS SP.	2	0.224	0	0.0
TOTALS	893		110	
NO. SPECIES		70		32
NO. IND. PER M2		3572		440
S-B INDEX - H ² (LN)		2.8562		2.8449
EVENNESS - J		0.6723		0.8209

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
10/3/77

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNICARIA				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	2	0.379	2	0.504
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	2	0.379	4	1.008
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	17	3.220	13	3.275
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	4	0.758	1	0.252
PHORONIDA (PHORONIDS)				
PHORONIS ARCHITECTA	0	0.0	1	0.252
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
ACTECCINA CANDEI	0	0.0	1	0.252
NASSARIUS ACUTUS	0	0.0	8	2.015
NATICA PUSILLA	1	0.189	0	0.0
TURBNILLA CONRADI	0	0.0	1	0.252
PELECYPODA (CLAMS)				
CHIONE CANCELLATA	1	0.189	0	0.0
ERVILIA CONCENTRICA	17	3.220	2	0.504
LUCINA MULTILINEATA	6	1.136	16	4.030
PERILOMA MARGARITACEUM	2	0.379	6	1.511
TELLINA AEGUSTRATA	0	0.0	1	0.252
TELLINA TEXANA	8	1.515	0	0.0
TELLINA VERSICOLOR	39	7.386	32	8.060
TRACHECARDIUM MURICATUM	1	0.189	0	0.0
ANNELIDA (SEGMENTED WORMS)				
OLIGOCHAETA				
UNIDENTIFIED SP.	24	4.545	12	3.023
POLYCHAETA				
APCPRICUSPIO PYGMAEA	1	0.189	1	0.252
ARICIDEA FRAGILIS	0	0.0	1	0.252
ARICIDEA SUECICA	3	0.568	1	0.252
ARMANDIA AGILIS	1	0.189	0	0.0
ARMANDIA MACULATA	1	0.189	1	0.252
FRANIA WELFLEETENSIS	3	0.568	0	0.0
CERATONEREIS IRRITABILIS	0	0.0	1	0.252
CHONE SP.	7	1.326	5	1.259
ETEONE LACTEA	2	0.379	9	2.267
GLYCERA AMERICANA	1	0.189	5	1.259
GLYCERA DIORANCHIATA	0	0.0	12	3.023
GONIADA LITTOREA	11	2.083	5	1.259
GRUBEULEPIS MEXICANA	1	0.189	0	0.0
HAPLCSOLOPLOS FOLIOLUS	6	1.136	3	0.756
LUMBRINERIS CRUZENSIS	207	39.205	107	26.952
LUMBRINERIS TETRAURA	6	1.136	27	6.801
MEDICMASTUS CALIFORNIENSIS	0	0.0	1	0.252
MESOCFAETOPTERUS SAGITTARIUS	1	0.189	4	1.008
NEANTLES ACUMINATA	2	0.379	0	0.0
NEANTLES SUCCINEA	1	0.189	0	0.0
NEPHYS PICTA	15	2.841	4	1.008
NOTOMASTUS HEMIPODUS	0	0.0	2	0.504
ONUPETIS FREMITA UCLATA	4	0.758	6	1.511

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
10/3/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>PARANAITES SPECIOSA</u>	1	0.189	0	0.0
<u>PARACNICES LYRA</u>	18	3.409	25	6.297
<u>PARACNIS FULGENS</u>	1	0.189	0	0.0
<u>PARAPRIONOSPION PINNATA</u>	1	0.189	0	0.0
<u>PHYLLODOCE ARENAE</u>	2	0.379	1	0.252
<u>PRIONOSPION CRISTATA</u>	6	1.136	2	0.504
<u>RULLIERINEREIS MEXICANA</u>	11	2.083	5	1.259
<u>SCOLELEPIS TEXANA</u>	0	0.0	1	0.252
<u>SCOLICLOS RUBRA</u>	0	0.0	2	0.504
<u>SIGAMBRA BASSI</u>	2	0.379	0	0.0
<u>SPIONA PETTIBONEAE</u>	4	0.758	1	0.252
<u>SPIOCAETOPTERUS OCULATUS</u>	7	1.326	1	0.252
 SIPUNCULICA (PEANUT WORMS)				
UNIDENTIFIED SP.	2	0.379	1	0.252
 ARTHROPODA (CRUSTACEANS)				
AMPHIPCA				
<u>ACANTHOGHAUSTORIUS SP.</u>	3	0.568	0	0.0
<u>AMPELISCA VERRILLI</u>	2	0.379	1	0.252
<u>GIANTOPSIS SP.</u>	1	0.189	0	0.0
<u>LEPIDACTYLUS SP.</u>	14	2.652	1	0.252
<u>LISTRILLA SP.</u>	2	0.379	2	0.504
<u>PSEUDOPLATYISCHNOPUS SP.</u>	16	3.030	19	4.786
<u>SYNCELIDILM SP.</u>	4	0.758	7	1.763
ANCURA				
<u>ALBINEA PARETII</u>	0	0.0	1	0.252
<u>PAGURUS LONGICARPUS</u>	3	0.568	3	0.756
BRACHYURA				
<u>PINNIXIA SAYANA</u>	0	0.0	1	0.252
<u>PINNOTHERES OSTREUM</u>	3	0.568	0	0.0
CARIDEA				
<u>PROCESSA TEMPHILLI</u>	1	0.189	1	0.252
CUMACEA				
<u>CYCLAESIS SP.</u>	4	0.758	7	1.763
<u>CYCLAESIS VARIANS</u>	2	0.379	2	0.504
<u>EXYUCSTYLIS SMITHI</u>	1	0.189	2	0.504
<u>SPILICUMA SALLMANI</u>	0	0.0	1	0.252
MYSIDACEA				
<u>MYSIDOPSIS BIGELOWI</u>	2	0.379	2	0.504
OSTRACODA				
UNIDENTIFIED SP.	1	0.189	0	0.0
PENAEIDAE				
<u>LUCIFER FAXONI</u>	1	0.189	1	0.252
<u>TRACHYPENAEUS CONSTRICTUS</u>	1	0.189	2	0.504
 ECHINODERMATA				
ECHINOIDEA (SAND DOLLARS; URCHINS)				
<u>MOIRA ATROPS</u>	2	0.379	0	0.0
OPHIURIDAE (BRITTLE STARS)				
<u>HENIOPHLLIS ELONGATA</u>	0	0.0	1	0.252
<u>HICHTHLLIS GRACILLIMA</u>	7	1.326	5	1.259
<u>OPHIOPHAGUS BURDEMANI</u>	1	0.189	0	0.0
UNIDENTIFIED SP.	2	0.379	1	0.252
 CEPHALOCORDATA (LANCELETS)				
<u>BRANCHIOSTOMA FLORIDAE</u>	3	0.568	3	0.756

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
10/3/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
TOTALS	528		397	
NO. SPECIES		64		61
NO. IND. PER M2		2112		1588
S-W INDEX - H'(LN)		2.8345		3.1138
EVENNESS - J		0.6815		0.7575

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
11/1/77

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNIDARIA				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	0	0.0	1	0.328
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	2	0.275	1	0.328
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	29	3.994	27	8.852
PHORONIDA (PHORONIDS)				
<u>PHORONIS ARCHITECTA</u>	1	0.138	0	0.0
MOLLUSCA (SHELLFISH)				
GASTROPOCA (SNAILS)				
<u>NASSARILS ACUTUS</u>	4	0.551	4	1.311
<u>NATICA FLUILLA</u>	1	0.138	1	0.328
<u>OLIVA SAYANA</u>	0	0.0	1	0.328
<u>OLIVELLA MUTICA</u>	4	0.551	0	0.0
<u>TEREBEA DISLOCATA</u>	4	0.551	1	0.328
<u>TURBICALLA CINEADI</u>	3	0.413	0	0.0
PELECYFIDA (CLAMS)				
<u>CHIONE CANCELLATA</u>	0	0.0	2	0.656
<u>DIPODCNTA SEMIASPERA</u>	1	0.138	0	0.0
<u>ERYLIA CONCENTRICA</u>	4	0.551	0	0.0
<u>LUCINA MULTILINEATA</u>	10	1.377	5	1.639
<u>NUCULANA ACUTA</u>	0	0.0	1	0.328
<u>PERIPLCMA MARGARITACEUM</u>	10	1.377	0	0.0
<u>TELLIDORA CRISTATA</u>	1	0.138	0	0.0
<u>TELLINA IRIS</u>	0	0.0	3	0.984
<u>TELLINA TEXANA</u>	1	0.138	0	0.0
<u>TELLINA VERSICOLOR</u>	29	3.994	6	1.967
ANNELIDA (SEGMENTED WORMS)				
CLIGGCHAETA				
UNIDENTIFIED SP.	19	2.617	4	1.311
POLYCHAETA				
ANTINCE SP.	1	0.138	1	0.328
<u>ARICIDEA FAUVELI</u>	0	0.0	1	0.328
<u>ARICIDEA FRAGILIS</u>	3	0.413	6	1.967
<u>ARICIDEA PHILGINAE</u>	1	0.138	0	0.0
<u>ARMANDIA MACULATA</u>	7	0.964	0	0.0
<u>CAULLERIELLA SP.</u>	1	0.138	0	0.0
<u>CHAELOZONE SETOSA</u>	1	0.138	0	0.0
<u>CHLOEIA VIRIDIS</u>	1	0.138	0	0.0
<u>CHONE SP.</u>	10	1.377	1	0.328
<u>CYSTENIDES GOLLOII</u>	1	0.138	0	0.0
<u>EYONE LACTEA</u>	13	1.791	1	0.328
<u>EULALIA SANGUINEA</u>	0	0.0	1	0.328
<u>GLYCERA AMERICANA</u>	8	1.102	2	0.656
<u>GLYCERA DIBRANCHIATA</u>	2	0.275	25	8.197
<u>GONIADA LITICEA</u>	4	0.551	2	0.656
<u>GRUBEULEFIS MEXICANA</u>	1	0.138	0	0.0
<u>GYPTIS BREVIPALPA</u>	1	0.138	0	0.0
<u>HAPLISCELLUS FOLIOSUS</u>	2	0.275	0	0.0
<u>HARPATICE LUNULATA</u>	0	0.0	1	0.328
<u>LUMBICINERIS CRUZENSIS</u>	235	32.369	47	15.410
<u>LUMBICINERIS TETRAURA</u>	20	2.755	3	0.984
<u>NAGELINA FIDJAI</u>	0	0.0	1	0.328

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
11/1/77
(CONTINUED)

SPECIES	NO. CF IND. (C.)		NO. CF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>MEDIONASTILS CALIFORNIENSIS</u>	1	0.138	2	0.656
<u>NEANTHES SUCCINEA</u>	2	0.275	0	0.0
<u>NEPHIYS PICTA</u>	11	1.515	7	2.295
<u>NOTOMASTILS HEMIPODUS</u>	5	0.689	6	1.967
<u>ONUPHIS EREMITA OCULATA</u>	8	1.102	7	2.295
<u>PARANATHES SPECIOSA</u>	0	0.0	1	0.328
<u>PARACNIDES LYRA</u>	37	5.096	75	24.590
<u>PARACNIS FULGENS</u>	5	0.689	0	0.0
<u>PARAPRICACSPIC PINNATA</u>	2	0.275	1	0.328
<u>PHERUS EHLERSI</u>	1	0.138	0	0.0
<u>PHYLLIDICE ARENAE</u>	1	0.138	0	0.0
<u>POLYACIDAE UNIDENTIFIED SP.</u>	1	0.138	1	0.328
<u>PRIONOSPIO CRISTATA</u>	38	5.234	2	0.656
<u>RULLIERINEREIS MEXICANA</u>	23	3.168	4	1.311
<u>SCOLOPLOS RUBRA</u>	1	0.138	0	0.0
<u>SPIO PETTIBONEAE</u>	3	0.413	1	0.328
<u>THARIX ANNULOSUS</u>	1	0.138	0	0.0
 SIPUNCULICA (PEANUT WORMS)				
<u>GOLFINGIA TRICHOCEPHALA</u>	2	0.275	0	0.0
 ARTHROPODA (CRUSTACEANS)				
AMPHIPODA				
<u>AMPELISCA VERRILLI</u>	3	0.413	1	0.328
<u>ERICHTHONIS SP.</u>	2	0.275	1	0.328
<u>PARAPHOXUS SP.</u>	3	0.413	0	0.0
<u>PSEUDOPLATYLSCHNOPUS SP.</u>	22	3.030	26	8.525
<u>SYNCHELIDUM SP.</u>	10	1.377	2	0.656
ANCMURA				
<u>ALBOLNEA PARETII</u>	3	0.413	1	0.328
<u>EUCERAMUS PRAELONGUS</u>	0	0.0	1	0.328
<u>PAGURUS LONGICARPUS</u>	7	0.964	0	0.0
BRACHYLARA				
<u>OVALIPES OCELLATUS</u>	1	0.138	1	0.328
<u>PINNULIA SAYANA</u>	0	0.0	1	0.328
CALLINASSIDAE				
<u>CALLINASSA JAMAICENSE</u>	0	0.0	1	0.328
CUMACEA				
<u>CYCLAPSID SP.</u>	10	1.377	1	0.328
<u>CYCLAPSID VARIANS</u>	3	0.413	2	0.656
<u>OXYUECSTYLIS SMITHI</u>	8	1.102	1	0.328
LEPTOSTRACA				
<u>NEBALIA SP.</u>	1	0.138	3	0.984
MYSTACEA				
<u>ROMMANIELLA SP.</u>	1	0.138	0	0.0
<u>MYSTICOPSIS BIGELOWI</u>	1	0.138	0	0.0
OSTRACODA				
UNIDENTIFIED SP.	3	0.413	1	0.328
PENAIDEA				
<u>TRACYPENAEUS CONSTRICTUS</u>	1	0.138	0	0.0
TANAIDACEA				
UNIDENTIFIED SP.	1	0.138	0	0.0
 ECHINODERMATA				
ECHINOIDEA (SAND DOLLARS; URCHINS)				
<u>MOIRA AIROPS</u>	2	0.275	0	0.0
<u>MELLITA OLINQUIE SPERFECRATA</u>	55	7.576	0	0.0
OPHTHURIDEA (BRITTLE STARS)				
<u>MICROPHOLIS GRACILLIMA</u>	3	0.413	2	0.656
<u>OPHIOPHRAGMUS WURDEMANI</u>	0	0.0	1	0.328

CEPHALOCHORDATA (LANCELETS)

TREASURE ISLAND MOTEL (STATION 1) - CONTROL AND EXPERIMENTAL
11/1/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>FRANCIOSIOMA FLORIDAE</u>	9	1.240	2	0.656
TOTALS	726		305	
NO. SPECIES		72		54
NO. IND. PER M2		2904		1220
S-W INDEX - H ² (LN)		3.0299		2.8764
EVENNESS - J		0.7085		0.7211

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL
7/11/77

SPECIES	NO. OF IND. (C ₁)		NO. OF IND. (E ₂)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNIDARIA				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	1	0.048	5	0.330
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	2	0.095	5	0.330
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	51	2.425	52	3.435
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	9	0.428	0	0.0
BRACHIOPOCA (LAMP SHELLS)				
<u>GLOTTIDIA PYRAMIDATA</u>	8	0.380	4	0.264
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
<u>ACTECCINA CANALICULATA</u>	1	0.048	0	0.0
<u>ACTECCINA CANDEI</u>	39	1.854	1	0.066
<u>BULLA STRIATA</u>	1	0.048	0	0.0
<u>CAECUM IMBRICATUM</u>	2	0.095	0	0.0
<u>CYLICENELLA BIDENTATA</u>	2	0.095	6	0.396
<u>DIASICMA VARIUM</u>	18	0.856	1	0.066
<u>NATICA PUSILLA</u>	6	0.285	0	0.0
<u>OLIVA SAYANA</u>	1	0.048	0	0.0
<u>POLINICES DUPLICATUS</u>	3	0.143	3	0.198
<u>TEREBRA DISLOCATA</u>	1	0.048	0	0.0
PELECYPODA (CLAMS)				
<u>ERVILIA CONCENTRICA</u>	52	2.473	0	0.0
<u>LEPTON SP.</u>	0	0.0	3	0.198
<u>LUCINA MULTILINEATA</u>	167	7.941	45	2.972
<u>PERIPLECHA MARGARITACEUM</u>	8	0.380	1	0.066
<u>PITAR SIMPSONI</u>	4	0.190	0	0.0
<u>SOLEN VIRIDIS</u>	0	0.0	1	0.066
<u>TELLINA AEGULISTRIATA</u>	1	0.048	1	0.066
<u>TELLINA TAMPAENSIS</u>	3	0.143	0	0.0
<u>TELLINA TEXANA</u>	20	0.951	9	0.594
<u>TELLINA VERSICOLOR</u>	182	8.654	68	4.491
ANNELIDA (SEGMENTED WORMS)				
OLIGOCHAETA				
UNIDENTIFIED SP.	35	1.664	17	1.123
POLYCHAETA				
<u>AGLACPHANUS VERRILLI</u>	1	0.048	0	0.0
<u>APROPILACSPID PYGMAEA</u>	7	0.333	12	0.793
<u>ARICIDEA CEREBUTI</u>	1	0.048	0	0.0
<u>ARICIDEA FRAGILIS</u>	10	0.476	16	1.255
<u>ARICIDEA PHILEINAE</u>	5	0.238	0	0.0
<u>ARICIDEA SUCCICA</u>	0	0.0	11	0.727
<u>ARMADIA AGILIS</u>	0	0.0	5	0.330
<u>ARMADIA MACULATA</u>	0	0.0	1	0.066
<u>CARAZZIELLA SP.</u>	6	0.285	0	0.0
<u>CAULLERIELLA SP.</u>	1	0.048	0	0.0
<u>CHONE SP.</u>	53	2.520	15	0.991
<u>CISTENIDES GOULDII</u>	0	0.0	1	0.066
<u>DISPID UNCINATA</u>	0	0.0	5	0.330
<u>ETECNE LACTEA</u>	10	0.476	0	0.0
<u>GLYCERA AMERICANA</u>	104	4.945	102	6.737

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL
7/11/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
GLYCEFA DIBRANCHIATA	0	0.0	2	0.132
GLYCEFA LITTYCEA	18	0.856	15	0.991
GRUBEULEPIS MEXICANA	2	0.095	0	0.0
GYPHIS VITTATA	4	0.190	4	0.264
HAPLISCELLIPUS FOLIOSUS	1	0.048	4	0.264
HAPLISCELLIPUS FRAGILIS	3	0.143	3	0.198
HAPLISCELLIPUS ROBUSTUS	0	0.0	2	0.132
HARMITHCE LUNULATA	1	0.048	0	0.0
LUMBFINERIS CRUZENSIS	391	18.592	249	16.446
LUMBFINERIS TETRAURA	22	1.046	6	0.396
MAGELINA LONGICORNIS	0	0.0	1	0.066
MAGELINA SP.	3	0.143	2	0.132
MEDICMASTUS CALIFORNIENSIS	5	0.238	1	0.066
MESOCFAETOPTERUS SAGITTARIUS	0	0.0	1	0.066
MICROPHthalmus SP.	1	0.048	0	0.0
NEANTHES ACUMINATA	3	0.143	2	0.132
NEANTHES SUCCINEA	0	0.0	2	0.132
NEPHYS BUCERA	4	0.190	4	0.264
NEPHYS PICTA	280	13.314	391	25.826
NEREIS LAMELLOSA	0	0.0	2	0.132
NOTICMASTUS HEMIPODUS	1	0.048	2	0.132
ONUPHIS EREMITA OCULATA	54	2.568	37	2.444
ONUPHIS NEBULOSA	5	0.238	0	0.0
OWENIA FUSIFORMIS	5	0.238	4	0.264
PARANATHES SPECIOSA	1	0.048	0	0.0
PARACNIDES LYRA	53	2.520	148	9.775
PARACNIDES FULGENS	6	0.285	0	0.0
PARAPHRONOSPID PINNATA	2	0.095	6	0.396
PHYLLODOCE ARENAE	3	0.143	2	0.132
PODARKE OBSCURA	1	0.048	0	0.0
POECILOCHAETUS JOHNSONI	1	0.048	2	0.132
PRICNISPEL CRISTATA	27	1.284	12	0.793
PSEUDEURYTHCE AMBIGUA	1	0.048	0	0.0
RULLIENINERIS MEXICANA	4	0.190	0	0.0
SABELLA MICROPHthALMA	0	0.0	1	0.066
SCOLELEPIS SCUANATA	4	0.190	0	0.0
SCOLELEPIS TEXANA	0	0.0	1	0.066
SCICELPUS ARMIGER	6	0.285	6	0.396
SCICELPUS RUERA	2	0.095	0	0.0
SIGALIN ARENICOLA	2	0.095	0	0.0
SIGAMERA BASSI	2	0.095	0	0.0
SPIO PETTIBONEAE	0	0.0	1	0.066
SPIOCAETOPTERUS OCULATUS	0	0.0	1	0.066
SPIOPLANES DOMBYX	18	0.856	27	1.783
SIPUNCULIDA (PEANUT WORMS)				
GOLFINGIA TRICHOCEPHALA	1	0.048	0	0.0
ARTHROPODA (CRUSTACEANS)				
AMPHIPODA				
ACANTHOAUSTORIUS SP.	2	0.095	0	0.0
AMPELISCA ADDITA	11	0.523	2	0.132
AMPELISCA VADORUM	5	0.238	0	0.0
AMPELISCA VERRILLI	40	1.902	37	2.444
ARGISSA SP.	1	0.048	2	0.132
CAPRELLIDAE UNIDENTIFIED SP.	3	0.143	2	0.132
ERICHTHENIUS SP.	2	0.095	0	0.0
GAMMAROPELIS SP.	1	0.048	0	0.0
LISTRIELLA SP.	12	0.571	3	0.198
LYSIANCA SP.	1	0.048	1	0.066
MONICULIDES SP.	2	0.095	1	0.066

TREASURE ISLAND MOTEL (STATION 1) - CONTROL & EXPERIMENTAL
7/11/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>PAGETIS SP.</u>	6	0.285	0	0.0
<u>PROCTAUSTORIUS SP.</u>	0	0.0	1	0.066
<u>PSEUDOPALATYISCHNOPUS SP.</u>	26	1.236	15	0.991
<u>SYNCELIUM SP.</u>	68	3.233	40	2.642
ANOMURA				
<u>ALBUEA FARETII</u>	0	0.0	1	0.066
BRACHYURA				
<u>PINNIXIA CYLINDRICA</u>	4	0.190	0	0.0
<u>PORTUNUS SP.</u>	0	0.0	12	0.793
PORTUNIDAE UNIDENTIFIED SP.	6	0.285	0	0.0
CARIDEA				
<u>OGYRIDES ALPHAEOSTRIS</u>	0	0.0	6	0.396
<u>OGYRIDES LIMICULA</u>	0	0.0	2	0.132
<u>PERCELIENES LONGICAUCATUS</u>	0	0.0	1	0.066
<u>PROCESSA HEMPHILLI</u>	5	0.238	3	0.198
CUMACEA				
<u>CYCLAPSIS SP.</u>	16	0.903	4	0.264
<u>CYCLAPSIS VARIANS</u>	82	3.899	22	1.453
<u>OXYLOSTYLIS SMITHI</u>	24	1.141	5	0.330
LEPTOSTRACA				
<u>NEBALIA SP.</u>	1	0.048	1	0.066
MYSTACEA				
UNIDENTIFIED SP.	1	0.048	1	0.066
OSTRACODA				
UNIDENTIFIED SP.	1	0.048	0	0.0
PENAEIDA				
<u>TRACHYPENAEUS CONSTRICTUS</u>	0	0.0	2	0.132
STOMATOPODA				
<u>ACANTHOSQUILLA BIRNIEI</u>	1	0.048	0	0.0
ECHINODERMATA				
HOLOTHURIDAE (SEA CUCUMBERS)				
<u>LEPTOSYNAPTIS SP.</u>	3	0.143	11	0.727
OPHIURIDAE (BRITTLE STARS)				
<u>OPHIOPHRAGMUS BURDEMANI</u>	1	0.048	0	0.0
UNIDENTIFIED SP.	6	0.285	3	0.198
HEMICHORDATA				
ENTEROPNEUSTA (ACORN WORMS)				
UNIDENTIFIED SP.	1	0.048	0	0.0
CEPHALOCHORDATA (LANCELETS)				
<u>BRANCHIOSTOMA FLORIDAE</u>	14	0.666	2	0.132
VERTEBRATA				
PISCES (FISHES)				
<u>HEMIFRENCIUS NOVACULA</u>	2	0.095	0	0.0
TOTALS	2103		1514	
NO. SPECIES	99		81	
NO. IND. PER M2	3365		2422	
S-W INDEX - M'(LN)	3.2301		2.8504	
EVENNESS - J	0.7029		0.6577	

SUN & SWIM MOTEL (STATION 2) - CONTROL & EXPERIMENTAL
7/15/77

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
Cnidaria				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	1	0.043	0	0.0
Platyhelminthes				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	12	0.512	16	0.663
Nemertinea (Ribbon Worms)				
UNIDENTIFIED SP.	57	2.432	45	1.864
Nematoda (Roundworms)				
UNIDENTIFIED SP.	16	0.683	10	0.414
Brachiozoa (Lamp Shells)				
GLOTHICIA PYRAMIDATA	10	0.427	10	0.414
Mollusca (Shellfish)				
GASTROPODCA (SNAILS)				
ACTECCINA CANALICULATA	0	0.0	1	0.041
ACTECCINA CANDEI	19	0.811	32	1.326
ANACIS FLORICANA	0	0.0	1	0.041
BULLA STRIATA	1	0.043	4	0.166
CYLLINELLA BIDENTATA	4	0.171	4	0.166
LIASICMA VARIUM	21	0.896	16	0.663
NATICA PUSILLA	10	0.427	21	0.870
OLIVELLA BULLULA	1	0.043	1	0.041
OLIVELLA MINUTA	0	0.0	3	0.124
OLIVELLA MUTICA	3	0.128	0	0.0
POLINICES DUPLICATUS	1	0.043	2	0.083
TEREBRA DISLOCATA	2	0.085	0	0.0
TURBNILLA ELEGANTULA	1	0.043	1	0.041
PELECYPODA (CLAMS)				
ANATIA ANATINA	3	0.128	2	0.083
ERVILIA CONCENTRICA	41	1.749	44	1.823
LAEVICAEDUM LAEVIGATUM	5	0.213	16	0.663
LEPICA SP.	2	0.085	0	0.0
LUCINA MULTILINEATA	191	8.148	19	0.787
MACRALLISTA NIMBOSA	0	0.0	1	0.041
MUSCULUS LATERALIS	1	0.043	0	0.0
PERIELCMA MARGARITACEUM	5	0.213	18	0.746
PYAS SIMPSONI	0	0.0	1	0.041
TELLINA AEGUISIRIATA	6	0.256	4	0.166
TELLINA TEXANA	19	0.811	5	0.373
TELLINA VERSICOLOR	262	11.177	196	8.119
TRACFYCARDIUM MURICATUM	3	0.128	9	0.373
VARICORRULA OPERCULATA	5	0.213	0	0.0
VENERIDAE UNIDENTIFIED SP.	53	2.261	45	1.864
Annelida (Segmented Worms)				
OLIGOCHAETA				
UNIDENTIFIED SP.	46	1.962	55	2.278
POLYCHAETA				
APOEIGNOSPLO PYGMAEA	2	0.085	6	0.249
ARICILEA FRAGILIS	9	0.384	2	0.083
ARICILEA PHILBINAE	7	0.299	0	0.0
ARICILEA SUECICA	0	0.0	1	0.041
ARICILEA SP.	0	0.0	2	0.083
ARMANCIA AGILIS	10	0.427	13	0.539
ARMANCIA MACULATA	13	0.555	21	0.870

SUN & SWIM MOTEL (STATION 2) - CONTROL & EXPERIMENTAL
7/15/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
BRANIA WELFLEETENSIS	1	0.043	0	0.0
CAPITELLA CAPITATA	0	0.0	3	0.124
CERATONEURIS IRRITABILIS	0	0.0	1	0.041
CHONE SP.	28	1.195	30	1.243
CYRRIPIEUS LYRIFORMIS	9	0.384	15	0.621
CISTERIDES GOULDII	2	0.085	1	0.041
DISPID UNCINATA	0	0.0	2	0.083
EYECHE LACTEA	10	0.427	6	0.249
EULALIA SANGUINEA	1	0.043	0	0.0
GLYCERA AMERICANA	87	3.712	92	3.811
GONIACA LITTOREA	18	0.768	13	0.539
GRUBELEPIS MEXICANA	3	0.128	1	0.041
GYPTIS VITTATA	7	0.299	3	0.124
HAPLISCILLICUS FOLICUS	2	0.085	2	0.083
HAPLISCILLOPS FRAGILIS	1	0.043	2	0.083
HARMITHOE LUNULATA	1	0.043	0	0.0
LUMBRINERIS CRUZENSIS	397	16.937	437	18.103
LUMBRINERIS TETRAURA	16	0.683	13	0.539
MAGELLANA SP.	2	0.085	5	0.207
MEDICASTILUS CALIFORNIENSIS	0	0.0	1	0.041
MICROCEPHALUS SCZELKOWII	1	0.043	0	0.0
MICROCEPHALUS SP.	3	0.128	0	0.0
NEANTHES ACUMINATA	1	0.043	1	0.041
NEPHYS BUCERA	5	0.213	3	0.124
NEPHYS PICTA	206	8.788	122	5.054
NEREIS LAMELLICSA	2	0.085	0	0.0
NOTICASTIUS HEMIPODUS	0	0.0	1	0.041
NOTICASTIUS LATERICEUS	1	0.043	0	0.0
OPHIS EREMITA OCULATA	34	1.451	51	2.113
OPHELIA SP.	1	0.043	0	0.0
OWENIA FUSIFORMIS	21	0.896	9	0.373
PARACNICES LYRA	83	3.541	42	1.740
PARACNICES FUIGENS	6	0.256	3	0.124
PARAPRIONOSPIO PINNATA	0	0.0	1	0.041
PHYLLODOCE ARENAE	1	0.043	11	0.456
PRIONOSPIO CRISTATA	44	1.877	46	1.906
REUTERINERIS MEXICANA	5	0.213	7	0.290
SABELLA MICROPHYTHALMA	5	0.213	0	0.0
SCOLELEPIS TEXANA	0	0.0	2	0.083
SCOLELUS ARMIGER	10	0.427	14	0.580
SCOLOPLOS RUDEA	1	0.043	1	0.041
SIGAMERA BASSI	0	0.0	1	0.041
SPHAEROSYLLIS SP.	0	0.0	1	0.041
SPIO PETTIBONEAE	1	0.043	1	0.041
SPIOPHANES BOMBYX	9	0.384	6	0.249
STREPTOSYLLIS ARENAE	0	0.0	1	0.041
WEBSTERINERIS TRIDENTATA	0	0.0	1	0.041
SIPUNCULICA (PEANUT WORMS)				
GOLFINGIA TRICHOCEPHALA	1	0.043	0	0.0
ARTHOPODA (CRUSTACEANS)				
AMPHIPODA				
ACANTHOHAUSTORIUS SP.	7	0.299	12	0.497
AMPELISCA ADDITA	2	0.085	4	0.166
AMPELISCA VACCUM	4	0.171	4	0.166
AMPELISCA VERRILLI	32	1.365	56	2.320
ARGISSA SP.	4	0.171	6	0.249
CAPRELLIDAE UNIDENTIFIED SP.	4	0.171	2	0.083
CARINOBATEA SP.	1	0.043	0	0.0
ERICHTHONIS SP.	2	0.085	1	0.041

SLN & SWIM MOTEL (STATION 2) - CONTROL & EXPERIMENTAL
7/15/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
HIPPOMEDON SP.	1	0.043	1	0.041
LYSTRIELLA SP.	9	0.384	11	0.456
LYSIANOPSIS SP.	1	0.043	6	0.249
MELITA APPENDICULATA	4	0.171	1	0.041
MICRODEUTOPUS SP.	1	0.043	2	0.083
MONOCULODES SP.	3	0.128	7	0.290
PHOTIS SP.	1	0.043	1	0.041
PSEUDCHAUSTORIUS SP.	3	0.128	9	0.373
PSEUDCHAUSTORIUS SP.	38	1.621	36	1.491
PSEUDCELATYLSCHNIPUS SP.	57	2.432	72	2.983
SYNCELIDIUM SP.	69	2.944	81	3.355
UNIDENTIFIED SP.	4	0.171	1	0.041
ANOMURA				
ALBUEA FAREII	3	0.128	2	0.083
PAGUEUS LONGICARPUS	1	0.043	1	0.041
BRACHYLFA				
PINNIXIA SAYANA	1	0.043	6	0.249
PINNITHERES OSTREUM	6	0.256	0	0.0
PORTUNUS SP.	16	0.683	17	0.704
CARIDEA				
LATREUTES PARVULUS	0	0.0	1	0.041
OGYRIDES LIMICOLA	3	0.128	7	0.290
PROCESSA HEMPHILLI	10	0.427	20	0.829
CUMACEA				
CYCLAPSIS SP.	45	1.920	83	3.438
CYCLAPSIS VARIANS	59	2.517	229	9.486
GYROSTYLIS SMITHI	39	1.664	59	2.444
UNIDENTIFIED SP.	1	0.043	1	0.041
LEPTOSTRACA				
NEBALIA SP.	10	0.427	50	2.071
MYSIDACEA				
MYSIDOPSIS BIGELOWI	1	0.043	0	0.0
UNIDENTIFIED SP.	5	0.213	8	0.331
OSTRACCA				
UNIDENTIFIED SP.	1	0.043	6	0.249
PENAEIDAE				
SICYCNIA BREVIROSTRIS	0	0.0	2	0.083
TRACHYPENAEUS CONSTRICTUS	2	0.085	1	0.041
STOMATOPODA				
ACANTHOSQUILLA BIMINIENSIS	2	0.085	0	0.0
CORONIS EXCAVATRIX	0	0.0	1	0.041
ECHINODERMATA				
ASTEROIDEA (STARFISHES)				
LUIDIA ALTERNATA	1	0.043	0	0.0
HOLOTUROIDAE (SEA CUCUMBERS)				
LERISCYNAPIA SP.	1	0.043	2	0.083
OPHIURIDEA (BRITTLE STARS)				
UNIDENTIFIED SP.	10	0.427	4	0.166
HEMICHORCATA				
ENTEROPNEUSTA (ACORN WORMS)				
UNIDENTIFIED SP.	0	0.0	1	0.041
CEPHALOCORDATA (LANCELETS)				
BRANCHISTOMA FLORIDAE	12	0.512	15	0.621
VERTEBRATA				
PISCES (FISHES)				
HEMIPTERONOTUS NOVACULA	0	0.0	1	0.041
LEPOPHIDIUM GRAELLI	0	0.0	1	0.041

SLN & SWIM MOTEL (STATION 2) - CONTROL & EXPERIMENTAL
7/15/77
(CONTINUED)

SPECIES	NO. OF IND. (C)		NO. OF IND. (E)	
	TOTAL	PERCENT	TOTAL	PERCENT

TOTALS	2344		2414	
NO. SPECIES	112		114	
NO. IND. PER M2	3750		3662	
S-M INDEX - H' (LN)	3.4273		3.5029	
EVENNESS - J	0.7264		0.7396	

MILTON HOLICAY INN (STATION 3) - CONTROL & EXPERIMENTAL
7/25/77

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNICARIA				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	0	0.0	5	0.198
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	9	0.333	37	1.467
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	49	1.812	57	2.259
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	18	0.666	22	1.110
PHORONIDA (PHORONIDS)				
<u>PHORONIS ARCHITECTA</u>	1	0.037	3	0.119
BRACHIOPODA (LAMP SHELLS)				
<u>GLOTTIDIA PYRAMIDATA</u>	0	0.0	4	0.159
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
<u>ACTECCINA CANALICULATA</u>	0	0.0	1	0.040
<u>ACTECCINA CANDEI</u>	27	0.999	10	0.396
<u>ANACIS FLORICANA</u>	1	0.037	1	0.040
<u>BULLA STELLATA</u>	2	0.074	0	0.0
<u>CAECUM IMBRICATUM</u>	1	0.037	0	0.0
<u>CAECUM PULCHELLUM</u>	3	0.111	0	0.0
<u>CYLICNELLA BIDENTATA</u>	11	0.407	0	0.0
<u>CIASIMA VARIUM</u>	83	3.070	23	0.912
<u>HELANELLA JAMAICENSIS</u>	0	0.0	1	0.040
<u>NATICA PUSILLA</u>	46	1.701	30	1.189
<u>OLIVA SAYANA</u>	1	0.037	0	0.0
<u>OLIVELLA MINUTA</u>	5	0.185	4	0.159
<u>OLIVELLA MUTICA</u>	7	0.259	7	0.277
<u>PHILINE SAGRA</u>	0	0.0	4	0.159
<u>POLINICES DUPLICATUS</u>	1	0.037	1	0.040
<u>TURBOCILLA CONRADI</u>	6	0.222	11	0.436
PELECYPODA (CLAMS)				
<u>ANATINA ANATINA</u>	4	0.148	4	0.159
<u>ERVILIA CONCENTRICA</u>	27	0.999	18	0.713
<u>LAEVICARDIUM LAEVIGATUM</u>	1	0.037	0	0.0
<u>LEPTIS SP.</u>	10	0.370	4	0.159
<u>LUCINA MULTILINEATA</u>	53	1.960	69	2.735
<u>LYONSIA H. FLORICANA</u>	0	0.0	1	0.040
<u>MACOMA CONSTRICTA</u>	2	0.074	0	0.0
<u>PANDORA TRILINEATA</u>	0	0.0	1	0.040
<u>PERIPLOMA MARGARITACEUM</u>	2	0.074	2	0.079
<u>PITAE SIMPSONI</u>	47	1.738	29	1.149
<u>STRIGILLA MIRABILIS</u>	4	0.148	8	0.317
<u>TELLINA AEGUISIRIATA</u>	2	0.074	18	0.713
<u>TELLINA TEXANA</u>	363	13.425	349	13.833
<u>TELLINA VERSICOLOR</u>	203	7.507	166	6.579
<u>TRACLYCARDIUM MURICATUM</u>	3	0.111	2	0.079
VENERIDAE UNIDENTIFIED SP.	12	0.444	37	1.467
ANNELIDA (SEGMENTED WORMS)				
OLIGOCHAETA				
UNIDENTIFIED SP.	39	1.442	8	0.317
POLYCHAETA				

HILTON HOLIDAY INN (STATION 3) - CONTROL & EXPERIMENTAL
7/25/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
AMERICCAUPEHIS MAGNA	2	0.074	0	0.0
APPOPEICUSPIC PYGMAEA	1	0.037	2	0.079
ARICIDEA EFAGILIS	9	0.333	37	1.467
ARICIDEA PHILBINAE	0	0.0	1	0.040
ARICIDEA SUECICA	2	0.074	2	0.079
ARMANDIA AGILIS	6	0.222	3	0.119
ARMANDIA MACULATA	35	1.294	23	0.912
BRANIA CLAVATA	1	0.037	1	0.040
BRANIA WELFLEETENSIS	6	0.222	6	0.238
CHONE SP.	33	1.220	19	0.753
CIRRIPHORUS LYRIFORMIS	0	0.0	15	0.555
CISTENICES GOULDII	1	0.037	1	0.040
DIOPATRA CUPREA	0	0.0	4	0.159
DISPIC UNCINATA	0	0.0	5	0.198
ETECNE LACTEA	6	0.222	12	0.476
GLYCERA AMERICANA	20	0.740	19	0.753
GONIACA LITTOREA	6	0.222	0	0.0
GRUBBELLIPS MEXICANA	2	0.074	11	0.436
GYPTIS VITTATA	1	0.037	0	0.0
LAEONEREIS CULVERI	0	0.0	5	0.198
LOXIMIA MEDUSA	3	0.111	0	0.0
LUMBERINERIS CRUZENSIS	653	24.149	500	19.818
LUMBERINERIS TETRAURA	2	0.074	0	0.0
LYSIDICE NINETTA	1	0.037	0	0.0
MAGELCNA SP.	17	0.629	20	0.793
MEDICMASTUS CALIFORNIENSIS	1	0.037	0	0.0
MESOCHEILOPTERUS SAGITTARIUS	10	0.370	13	0.515
NEANTHES ACUMINATA	1	0.037	2	0.079
NEPHIYS BUCERA	9	0.333	12	0.476
NEPHIYS PICTA	81	2.996	71	2.814
NEREIS LAMELLCSA	1	0.037	0	0.0
NOTOMASTUS HEMIPODUS	2	0.074	3	0.119
NOTOMASTUS LATERICEUS	1	0.037	0	0.0
ONUPHIS EREMITA OCULATA	8	0.296	21	0.832
PARACNIDES LYRA	46	1.701	6	0.238
PARANIS FULGENS	10	0.370	3	0.119
PARAFICUSPID PINNATA	1	0.037	0	0.0
PARAFICUSYLLIS LONGICIRRATA	0	0.0	4	0.159
PHYLLIDICE ARENAE	1	0.037	6	0.238
PTECILCHAETUS JOHNSONI	0	0.0	1	0.040
PRICUSPIC CRISTATA	56	2.071	55	2.180
RULLIEFINEFIS MEXICANA	4	0.148	1	0.040
SABELLA MICROPHALMA	1	0.037	0	0.0
SCOLELEPIS TEXANA	3	0.111	3	0.119
SCOLLEPIS ARMIGER	2	0.074	1	0.040
SCOLLEPIS RUERA	1	0.037	1	0.040
SIGAMBRA BASSI	11	0.407	13	0.515
SPIO PETTIBONEAE	2	0.074	2	0.079
SPIOCHAETOPTERUS OCULATUS	1	0.037	1	0.040
SPIOPTANES DOMBYX	1	0.037	2	0.079
STREPTOSYLLIS ARENAE	0	0.0	1	0.040
ARTHROPODA (CRUSTACEANS)				
AMPHIPODA				
ACANTHOCHAUSTORIUS SP.	1	0.037	0	0.0
AMPELISCA AEDITA	4	0.148	3	0.119
AMPELISCA VERRILLI	16	0.592	13	0.515
ARGISSA SP.	3	0.111	7	0.277
ELASMOPOUS SP.	1	0.037	0	0.0
LISTERIELLA SP.	9	0.333	7	0.277
MELITA APPENDICULATA	3	0.111	0	0.0

HILTON HOLIDAY INN (STATION 3) - CONTROL & EXPERIMENTAL
7/25/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>MICRODEUTOPUS SP.</u>	10	0.370	4	0.159
<u>MONOCULODES SP.</u>	22	0.814	30	1.189
<u>PROTOFAUSTORIUS SP.</u>	146	5.399	242	9.592
<u>PSEUDOFALSTORILS SP.</u>	9	0.333	4	0.159
<u>PSEUDOPLATYISCHNOPLS SP.</u>	118	4.364	115	4.558
<u>SYNCELIDUM SP.</u>	26	0.962	41	1.625
ANOMURA				
<u>ALBUNEA PARETII</u>	3	0.111	5	0.198
<u>PAGURUS LONGICARPUS</u>	3	0.111	1	0.040
BRACHYURA				
<u>HEPATIS EPHELITICUS</u>	0	0.0	1	0.040
<u>OVALIPES OCELLATUS</u>	0	0.0	1	0.040
<u>PINNIXIA SAYANA</u>	12	0.444	0	0.0
<u>PINNIXIA OSTRUM</u>	1	0.037	0	0.0
<u>PORTUNUS SP.</u>	9	0.333	2	0.079
CARIDEA				
<u>LABRELITES PARVULUS</u>	1	0.037	0	0.0
<u>PROCESSA MEMPHILLI</u>	7	0.259	2	0.079
CLMACEA				
<u>CYCLAPSIS SP.</u>	22	0.814	55	2.338
<u>CYCLAPSIS VARIANS</u>	55	2.034	61	2.418
<u>OXYUFOSTYLIS SMITHI</u>	6	0.222	13	0.515
ISOPODA				
<u>EDOTEA MONTOSA</u>	1	0.037	0	0.0
LEPTOSTRACA				
<u>NERALIA SP.</u>	13	0.481	11	0.436
MYSIDACEA				
UNIDENTIFIED SP.	10	0.370	4	0.159
OSTRACODA				
UNIDENTIFIED SP.	14	0.518	17	0.674
PENAEIDEA				
<u>TRACHYPENAEUS CONSTRICTUS</u>	0	0.0	1	0.040
ECHINODERMATA				
OPHIURCIDEA (BRITTLE STARS)				
UNIDENTIFIED SP.	5	0.185	8	0.317
CEPHALOCHORDATA (LANCELETS)				
<u>BRANCHIOSTOMA FLORIDAE</u>	69	2.552	19	0.753
VERTEBRATA				
PISCES (FISHES)				
<u>HEMIPYRONOTUS NOVACULA</u>	1	0.037	1	0.040
TOTALS	2704		2523	
NO. SPECIES	105		98	
NO. IND. PER M2	4326		4037	
S-W INDEX - H' (LN)	3.1958		3.2651	
EVENNESS - J	0.6867		0.7121	

SANDPIPER MOTEL (STATION 4) - CONTROL & EXPERIMENTAL
7/26/77

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNICARIA				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	2	0.079	1	0.062
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	11	0.435	1	0.062
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	36	1.422	34	2.103
NEMATODA (ROUNDWORMS)				
UNIDENTIFIED SP.	0	0.0	1	0.062
PHORONIDA (PHORONIDS)				
PHORONIS ARCHITECTA	0	0.0	1	0.062
BRACHIOPODA (LAMP SHELLS)				
GLYPTIDIA PYRAMICATA	1	0.040	8	0.495
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
ACTEOCINA CANDEI	0	0.0	1	0.062
CYLICHNELLA BIDENTATA	51	2.015	23	1.422
NATICA PUSILLA	36	1.422	10	0.618
OLIVELLA MINUTA	7	0.277	2	0.124
OLIVELLA MUTICA	7	0.277	3	0.186
PHILINE SAGRA	1	0.040	1	0.062
TURBENILLA CONRADI	5	0.198	1	0.062
TURBENILLA SP.	3	0.119	1	0.062
PELECYPODA (CLAMS)				
ERVILIA CONCENTRICA	15	0.593	1	0.062
LEPTON SP.	29	1.146	1	0.062
LUCINA MULTILINEATA	13	0.514	157	5.709
PERIPLOMA MARGARITACEUM	0	0.0	1	0.062
PITAR SIMPSONI	114	4.504	18	1.113
STRIGILLA MIRABILIS	14	0.553	5	0.309
TELLINA AEGISIRATA	0	0.0	4	0.247
TELLINA IRIS	0	0.0	11	0.680
TELLINA TEXANA	443	17.503	89	5.504
TELLINA VERSICOLOR	102	4.030	120	7.421
TRACHYCARDIUM MURICATUM	4	0.158	4	0.247
ANNELIDA (SEGMENTED WORMS)				
OLIGOCHAETA				
UNIDENTIFIED SP.	5	0.198	1	0.062
POLYCHAETA				
AONIDES MAYAQUEZENSIS	0	0.0	16	0.989
APOEIGNOSPION PYGMAEA	0	0.0	2	0.124
ARICIDEA FRAGILIS	0	0.0	1	0.062
ARMANCIA AGILIS	28	1.106	57	3.525
ARMANDIA MACULATA	19	0.751	17	1.051
FRANIA WELFLEETENSIS	13	0.514	13	0.804
CAPITELLA CAPITATA	2	0.079	33	2.041
CERATONEREIS IRRITABILIS	0	0.0	1	0.062
CHONE SP.	3	0.119	2	0.124
DISPIDUNCINATA	0	0.0	1	0.062
EYEDNE ALBA	1	0.040	0	0.0
EYEDNE LACTEA	9	0.356	1	0.062
GLYCERA AMERICANA	28	1.106	65	4.020

SANDPIPER MOTEL (STATION 4) - CONTROL & EXPERIMENTAL

7/26/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
GONIACA LITTOREA	0	0.0	2	0.124
GRUEULEPIS MEXICANA	0	0.0	4	0.247
GYPTIS VITTATA	0	0.0	10	0.618
HAPLCSCOLOPLOS FOLIOSUS	0	0.0	7	0.433
HAPLCSCOLOPLOS FRAGILIS	2	0.079	2	0.124
HAPLCSCOLOPLOS ROBUSTUS	0	0.0	4	0.247
HEMIPODUS ROSEUS	1	0.040	0	0.0
LOIMIA MEDUSA	0	0.0	4	0.247
LUMBERINERIS CRUZENSIS	521	20.585	129	7.978
MAGELCNA SP.	10	0.395	4	0.247
MESOCFAETOPTERUS SAGITTARIUS	13	0.514	30	1.855
NEANITES ACUMINATA	0	0.0	3	0.186
NEPHIYS BUCERA	26	1.027	12	0.742
NEPHIYS PICTA	2	0.079	143	8.844
NOTOMASTUS HEMIPODUS	0	0.0	1	0.062
ONUPFIS EREMITA OCVLATA	22	0.869	10	0.618
OPHELIA SP.	0	0.0	2	0.124
ORBINIA RISER	2	0.079	3	0.186
PARANAITES SPECIOSA	1	0.040	21	1.299
PARACNIS FULGENS	61	2.410	4	0.247
PARAPRIONOSPID PINNATA	0	0.0	1	0.062
PHYLLODOCE ARENAE	4	0.158	5	0.309
PHYL C ORNATUS	1	0.040	0	0.0
POLYDORA SOCIALIS	1	0.040	0	0.0
POLYDORA TETRABRANCHIA	2	0.079	0	0.0
PRIONOSPID CRISTATA	18	0.711	7	0.433
RULLIERINEREIS MEXICANA	0	0.0	8	0.495
SCOLCPLCS ARMIGER	8	0.316	10	0.618
SIGALION ARENICOLA	0	0.0	3	0.186
SIGAMBRA BASSI	0	0.0	33	2.041
SPIO PETTIBONEAE	6	0.237	21	1.299
SPIOFANES BOMBYX	6	0.237	9	0.557
STREPTOSYLLIS ARENAE	1	0.040	0	0.0
SIPUNCULICA (PEANUT WORMS)				
UNIDENTIFIED SP.	5	0.198	3	0.186
ARTROPOCOA (CRUSTACEANS)				
AMPHIPODA				
AMPELISCA ABDITA	2	0.079	6	0.371
AMPELISCA VERRILLI	8	0.316	1	0.062
ARGISSA SP.	2	0.079	0	0.0
LISTRIELLA SP.	2	0.079	1	0.062
MONOCILLIDES SP.	10	0.395	1	0.062
PROTHALSTORIUS SP.	385	15.211	38	2.350
PSEUDOPHAUSTORIUS SP.	15	0.593	25	1.546
PSEUDOPLATYISCHNOPUS SP.	141	5.571	38	2.350
SYNCELIDIUM SP.	52	2.055	5	0.309
ANCHURA				
ALBULEA PARETII	1	0.040	5	0.309
PAGURUS LONGICARPUS	3	0.119	5	0.309
BRACHYURA				
PINNIXIA CRISTATA	1	0.040	0	0.0
PINNIXIA LEPTOSYNAPIAE	0	0.0	3	0.186
PINNIXIA PEARSEI	0	0.0	1	0.062
PINNICTERES OSTREUM	0	0.0	3	0.186
PORTUNUS SP.	4	0.158	1	0.062
CALLINANASSIDAE				
CALLINANASSA JAMAICENSE	0	0.0	1	0.062
CARIDEA				
HIPPCLYTE PLEURACANTHA	1	0.040	0	0.0

SANDPIPER MOTEL (STATION 4) - CONTROL & EXPERIMENTAL
7/26/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>OGYRIDES LIMICOLA</u>	1	0.040	0	0.0
<u>PROCESSA HEMPHILLI</u>	8	0.316	4	0.247
CUMACEA				
<u>CYCLAPSIS SP.</u>	14	0.553	4	0.247
<u>CYCLAPSIS VARIANS</u>	101	3.991	17	1.051
<u>OXYUROSIS SMITHI</u>	4	0.158	2	0.124
LEPTOSTRACA				
<u>NEGALIA SP.</u>	5	0.198	9	0.557
GSTRACCA				
UNIDENTIFIED SP.	28	1.106	7	0.433
PENAEIDAE				
<u>PENAEUS DUORARUM</u>	1	0.040	0	0.0
SICMATOPODA				
<u>ACANTHOSQUILLA BIMINIENSIS</u>	0	0.0	1	0.062
ECHINODERMATA				
ASTEROIDEA (STARFISHES)				
<u>LUDIA ALTERNATA</u>	0	0.0	1	0.062
ECHINOIDEA (SAND DOLLARS; URCHINS)				
<u>PELLITA QUINQUESPERFORATA</u>	6	0.237	3	0.186
OPHTHUROIDEA (BRITTLE STARS)				
UNIDENTIFIED SP.	5	0.198	39	2.412
HEMICHORCATA				
ENTEROPNEUSTA (ACORN WORMS)				
UNIDENTIFIED SP.	0	0.0	1	0.062
CEPHALOCHORDATA (LANCELETS)				
<u>BRANCHIOSTOMA FLORICAE</u>	16	0.632	191	11.812
TOTALS	2531		1617	
NO. SPECIES		74		54
NO. IND. PER M2		4050		2587
S-W INDEX - M'(LN)		2.8718		3.4385
EVENNESS - J		0.6672		0.7568

PEPPERTREE CONDOMINIUM (STATION 5) - CONTROL & EXPERIMENTAL
7/27/77

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNICARIA				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	3	0.341	0	0.0
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	0	0.0	3	0.180
NEVERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	29	3.295	45	2.703
PHORONIDA (PHORONIDS)				
PHORONIS ARCHITECIA	0	0.0	1	0.060
BRACHIOPODA (LAMP SHELLS)				
GLOTTIDIA PYRAMICATA	0	0.0	1	0.060
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
ACTEOCINA CANDEI	1	0.114	0	0.0
ANACIS FLORICANA	0	0.0	1	0.060
CYLICHNELLA BICENTATA	3	0.341	20	1.201
NATICA FUSILLA	5	0.568	29	1.742
OLIVELLA MINUTA	0	0.0	2	0.120
OLIVELLA MUTICA	2	0.227	2	0.120
POLYICES DUPLICATUS	3	0.341	1	0.060
TEREBRA DISLOCATA	1	0.114	2	0.120
TURRITELLA SP.	2	0.227	11	0.661
PELECYPODA (CLAMS)				
CUNA DALLI	0	0.0	1	0.060
ERVILIA CONCENTRICA	2	0.227	5	0.300
LEPTA SP.	7	0.795	5	0.541
LUCINA MULTILINEATA	8	0.909	3	0.180
PITAE SIMPSONI	17	1.932	11	0.661
STRIGILLA MIRABILIS	5	0.568	6	0.360
TELLINA IRIS	0	0.0	1	0.060
TELLINA TEXANA	40	4.545	255	15.315
TELLINA VERSICOLOR	94	10.682	90	5.405
TRACYPYCARDIUM MURICATUM	0	0.0	1	0.060
ANNELIDA (SEGMENTED WORMS)				
POLYCHAETA				
ARMADILLA AGILIS	1	0.114	55	5.706
ARMADILLA MACULATA	2	0.227	3	0.180
ERANIA CLAVATA	0	0.0	1	0.060
CAPITELLA CAPITATA	0	0.0	1	0.060
CHONE SP.	0	0.0	1	0.060
DISPIDIO UNCINATA	2	0.227	1	0.060
EYDIONE LACTEA	0	0.0	1	0.060
GLYCERA AMERICANA	6	0.682	20	1.201
GRUBBULEPIS MEXICANA	0	0.0	1	0.060
GYPTIS VITTATA	0	0.0	1	0.060
HAPLOSCOLOPUS FOLIOSUS	0	0.0	1	0.060
LOXIA MEDUSA	0	0.0	1	0.060
LUMBERINERIS CRUZENSIS	62	7.045	286	17.177
MAGELONA PETTIBONEAE	1	0.114	0	0.0
MAGELONA RIOJAI	29	3.295	12	0.781
MAGELONA SP.	1	0.114	1	0.060
MESOCALOPTERUS SAGITTARIUS	13	1.477	10	0.601
NEANTHES ACUMINATA	9	1.023	3	0.180

PEPPERTREE CONDOMINIUM (STATION 5) - CONTROL & EXPERIMENTAL
7/27/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>NEPHYS BUCERA</u>	50	5.682	22	1.321
<u>NEPHYS PICTA</u>	1	0.114	5	0.300
<u>ONUPHIS EREMITA OCULATA</u>	28	3.182	15	0.901
<u>ORBINIA RISER</u>	0	0.0	9	0.541
<u>PARACNIS FULGENS</u>	11	1.250	0	0.0
<u>PHYLLODOCE ARENAE</u>	3	0.341	15	1.141
<u>PRIONOSPIO CRISTATA</u>	2	0.227	3	0.180
<u>RULLIERINEREIS MEXICANA</u>	0	0.0	1	0.060
<u>SCOLELEPIS TEXANA</u>	0	0.0	1	0.060
<u>SCOLOPLCS ARMIGER</u>	0	0.0	1	0.060
<u>SIGALON ARENICOLA</u>	0	0.0	2	0.120
<u>SIGAMBRA BASSI</u>	0	0.0	2	0.120
<u>SPIO PETTIBONEAE</u>	20	2.273	7	0.420
<u>SPIOPHANES BOMBYX</u>	6	0.682	5	0.541
<u>STREPTOSYLLIS ARENAE</u>	4	0.455	0	0.0
 SIPUNCULIDA (PEANUT WORMS)				
UNIDENTIFIED SP.	1	0.114	3	0.180
 ARTHROPODA (CRUSTACEANS)				
AMPHIPODA				
<u>ACANTHOMHAUSTORIUS SP.</u>	5	1.023	0	0.0
<u>AMPELISCA VERRILLI</u>	0	0.0	1	0.060
<u>LITRIELLA SP.</u>	0	0.0	2	0.120
<u>LYSIANOPSIS SP.</u>	0	0.0	1	0.060
<u>MONOCLODES SP.</u>	5	1.023	13	0.781
<u>PROTOMHAUSTORIUS SP.</u>	246	27.955	245	14.715
<u>PSEUDCHALSTORIUS SP.</u>	10	1.136	66	3.964
<u>PSEUDOPLATYISCHNOPUS SP.</u>	37	4.205	152	9.129
<u>SYNCHELIDIUM SP.</u>	15	1.705	15	0.901
ANCMURA				
<u>ALBINEA PAREIII</u>	3	0.341	1	0.060
BRACHYURA				
<u>PINNIXIA CRISTATA</u>	4	0.455	0	0.0
<u>PINNOTHERES SP.</u>	3	0.341	5	0.300
<u>POTUNUS SP.</u>	3	0.341	2	0.120
CALLINANASSIDAE				
<u>CALLINANASSA JAMAICENSE</u>	0	0.0	4	0.240
CARIDEA				
<u>HIPPELYTE PLEURACANTHA</u>	0	0.0	2	0.120
<u>PROCESSA HEMPHILLI</u>	8	0.909	2	0.120
CLMACEA				
<u>CYCLAPSIS SP.</u>	18	2.045	8	0.480
<u>CYCLAPSIS VARIANS</u>	14	1.591	40	2.402
<u>CYCLESTYLIS SMITHI</u>	0	0.0	4	0.240
UNIDENTIFIED SP.	0	0.0	2	0.120
ISCPCOA				
<u>ANCINA DEPRESSUS</u>	3	0.341	0	0.0
<u>CHIRIDOTEA EXCAVATA</u>	5	1.023	0	0.0
LEPTOSTRACA				
<u>NEBALIA SP.</u>	0	0.0	5	0.300
MYSIDACEA				
<u>PRAUNUS FLEXUOSUS</u>	3	0.341	0	0.0
OSTRACCA				
UNIDENTIFIED SP.	4	0.455	2	0.120
PENAEIDAE				
<u>IRACHYPENAEUS CONSTRICTUS</u>	1	0.114	3	0.180
 ECHINODERMATA				
ECHINOIDEA (SAND COLLARS; URCHINS)				
<u>MELLITA QUINQUESPERFORATA</u>	0	0.0	2	0.120

PEPPERTREE CONDOMINIUM (STATION 5) - CONTROL & EXPERIMENTAL
7/27/77
(CONTINUED)

SPECIES	NO. OF IND. (C)		NO. OF IND. (E)	
	TOTAL	PERCENT	TOTAL	PERCENT
HOLOTHUROIDEA (SEA CUCUMBERS)				
UNIDENTIFIED SP.	0	0.0	13	0.781
OPHIURCIDEA (BRITTLE STARS)				
<u>OPHIOPHRAGMUS MOOREI</u>	0	0.0	5	0.300
<u>OPHIOPHRAGMUS BURDEMANI</u>	4	0.455	0	0.0
UNIDENTIFIED SP.	1	0.114	7	0.420
HEMICHORCATA				
ENTEROPNEUSTA (ACORN WORMS)				
UNIDENTIFIED SP.	0	0.0	1	0.060
CEPHALOCHORDATA (LANCELETS)				
<u>BRANCHIOSTOMA FLORIDAE</u>	2	0.227	23	1.381
TOTALS	880		1665	
NO. SPECIES	57		80	
NO. IND. PER M2	1408		2664	
S-W INDEX - H ⁺ (LN)	2.9751		2.9427	
EVENNESS - J	0.7359		0.6715	

BLUE DOLPHIN MOTEL (STATION 6) - CONTROL & EXPERIMENTAL
7/28/77

SPECIES	NO. OF IND. (C)		NO. OF IND. (E)	
	TOTAL	PERCENT	TOTAL	PERCENT
CNICARIA				
ACTINIARIA (SEA ANEMONES)				
UNIDENTIFIED SP.	1	0.064	0	0.0
PLATYHELMINTHES				
TURBELLARIA (FLATWORMS)				
UNIDENTIFIED SP.	0	0.0	1	0.053
NEMERTINEA (RIBBON WORMS)				
UNIDENTIFIED SP.	33	2.126	57	3.006
PHORONIDA (PHORONIDS)				
<u>PHORENIS ARCHITECIA</u>	0	0.0	1	0.053
BRACHIOPODA (LAMP SHELLS)				
<u>GLOTTIDIA PYRAMIDATA</u>	0	0.0	19	1.002
MOLLUSCA (SHELLFISH)				
GASTROPODA (SNAILS)				
<u>CYLICHNELLA BIDENTATA</u>	24	1.546	31	1.635
<u>NATICA PUSILLA</u>	1	0.064	15	0.791
<u>OLIVELLA MINUTA</u>	2	0.129	2	0.105
<u>OLIVELLA MUTICA</u>	1	0.064	4	0.211
<u>POLINICES DUPLICATUS</u>	0	0.0	1	0.053
<u>TEREBRA CINCAYA</u>	1	0.064	0	0.0
<u>TURBINILLA CENRADI</u>	3	0.193	0	0.0
<u>TURBINILLA SP.</u>	11	0.709	6	0.316
PELECYPODA (CLAMS)				
<u>ERVILIA CONCENTRICA</u>	4	0.258	1	0.053
<u>LEPTON SP.</u>	13	0.838	3	0.158
<u>LUCINA MULTILINEATA</u>	18	1.160	74	3.903
<u>PERILEPTA MARGARITACEUM</u>	0	0.0	2	0.105
<u>PITAE SIMPSONI</u>	53	3.415	17	0.897
<u>STRIGILLA BIRABILLIS</u>	4	0.258	5	0.264
<u>TELLINA AEGUISIRIATA</u>	0	0.0	1	0.053
<u>TELLINA TEXANA</u>	217	13.982	137	7.226
<u>TELLINA VERSICOLOR</u>	108	6.959	58	5.169
<u>TRACHYCARDIUM MURICATUM</u>	0	0.0	1	0.053
ANNELIDA (SEGMENTED WORMS)				
OLIGCHAETA				
UNIDENTIFIED SP.	4	0.258	1	0.053
POLYCHAETA				
<u>APOPRIONOSPIO PYGMAEA</u>	2	0.129	4	0.211
<u>ARICIDEA FRAGILIS</u>	1	0.064	5	0.264
<u>ARMANDIA AGILIS</u>	36	2.320	27	4.589
<u>ARMANDIA MACULATA</u>	20	1.289	26	1.371
<u>BRANIA CLAVATA</u>	0	0.0	4	0.211
<u>BRANIA BELLEFLEETENSIS</u>	4	0.258	2	0.105
<u>CAPITELLA CAPITATA</u>	1	0.064	53	2.795
<u>CHONE SP.</u>	1	0.064	2	0.105
<u>DISPIDO UNCINATA</u>	1	0.064	0	0.0
<u>EYEDINE LACTEA</u>	1	0.064	2	0.105
<u>GLYCERA AMERICANA</u>	13	0.838	6	0.316
<u>GONIACA LITIGEA</u>	0	0.0	1	0.053
<u>GYPSIS VITTATA</u>	0	0.0	10	0.527
<u>HAPLISCELOPS ROBUSTUS</u>	0	0.0	1	0.053
<u>LOEBA MEDUSA</u>	1	0.064	1	0.053
<u>LUMBRINEIS CRUZENSIS</u>	195	12.564	208	10.970
<u>MACELINA RIOJA</u>	3	0.193	1	0.053

BLUE DOLPHIN MOTEL (STATION 6) - CONTROL & EXPERIMENTAL
7/28/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
<u>MAGELCNA SP.</u>	12	0.773	8	0.422
<u>MESOCYPTOPTERUS SAGITTARIUS</u>	9	0.580	12	0.633
<u>MINUSPIO CIRRIFERA</u>	1	0.064	0	0.0
<u>NEANTHES ACUMINATA</u>	0	0.0	28	1.477
<u>NEPHYS BUCERA</u>	35	2.255	26	1.371
<u>NEPHYS PICTA</u>	18	1.160	49	2.584
<u>ONUPHIS EREMITA Oculata</u>	8	0.515	9	0.475
<u>ORBINIA RISER</u>	2	0.129	7	0.369
<u>PARANATHES SPECIOSA</u>	0	0.0	2	0.105
<u>PARACNIS FULGENS</u>	17	1.095	8	0.422
<u>PHYLLODOCE ARENAE</u>	6	0.387	19	1.002
<u>POLYCORA SOCIALIS</u>	2	0.129	0	0.0
<u>POLYCORA TETRABRANCHIA</u>	0	0.0	1	0.053
<u>PRIONOSPION CRISTATA</u>	4	0.258	10	0.527
<u>RULLIERINEREIS MEXICANA</u>	0	0.0	2	0.105
<u>SCOLELEPIS TEXANA</u>	0	0.0	1	0.053
<u>SCOLOPLOS ARMIGER</u>	4	0.258	2	0.105
<u>SCOLOPLOS RUBRA</u>	2	0.129	0	0.0
<u>SIGAMBRA BASSI</u>	1	0.064	0	0.0
<u>SPION PETTIBONEAE</u>	7	0.451	15	0.791
<u>SPIOPHANE BOMBYX</u>	11	0.709	13	0.686
 SIPUNCULIDA (PEANUT WORMS)				
UNIDENTIFIED SP.	4	0.258	4	0.211
 ARTHROPODA (CRUSTACEANS)				
AMPHIPODA				
<u>ACANTHOHAUSTORIUS SP.</u>	1	0.064	0	0.0
<u>AMPELISCA ABDITA</u>	0	0.0	1	0.053
<u>ERICHTHONIUS SP.</u>	0	0.0	1	0.053
<u>LEMBOS SP.</u>	0	0.0	1	0.053
<u>LITTELLIA SP.</u>	5	0.322	3	0.158
<u>DELITA APPENDICULATA</u>	0	0.0	1	0.053
<u>NICEFECTOPUS SP.</u>	0	0.0	24	1.266
<u>MONOCULODES SP.</u>	9	0.580	31	1.635
<u>PROTYCHAUSTORIUS SP.</u>	307	19.781	245	12.922
<u>PSEUDOCYTHOICUS SP.</u>	20	1.289	25	1.319
<u>PSEUDOPALATYISCHNOPUS SP.</u>	114	7.345	50	2.637
<u>SYNCELIDION SP.</u>	23	1.482	4	0.211
BRACHYURA				
<u>CALLINECTES SAPIDUS</u>	0	0.0	2	0.105
<u>DISSODACTYLUS MELLITAE</u>	13	0.838	30	1.582
<u>PINAXIA SAYANA</u>	0	0.0	9	0.475
CALLINANASSIDAE				
<u>CALLINANASSA JAMAICENSE</u>	4	0.258	4	0.211
CARIDEA				
<u>HIPPOLYTE PLEURACANTHA</u>	0	0.0	5	0.264
<u>PROCESSA HEMPHILLI</u>	1	0.064	5	0.264
CUMACEA				
<u>CYCLAPES SP.</u>	25	1.611	19	1.002
<u>CYCLAPES VARIANS</u>	38	2.448	199	10.496
<u>CRYDIELLUS SMITHI</u>	3	0.193	8	0.422
LEPTOSTRACA				
<u>NEBALIA SP.</u>	4	0.258	26	1.371
OSTRACODA				
UNIDENTIFIED SP.	17	1.095	4	0.211
STOMATOPODA				
<u>ACANTHOSQUILLA BIMINIENSIS</u>	0	0.0	1	0.053

ECHINODERMATA
ECHINOIDEA (SAND DOLLARS; URCHINS)

BLUE DOLPHIN MOTEL (STATION 6) - CONTROL & EXPERIMENTAL
7/28/77
(CONTINUED)

SPECIES	NO. OF IND. (C.)		NO. OF IND. (E.)	
	TOTAL	PERCENT	TOTAL	PERCENT
~~MELLITA QUINQUESPERFORATA~~	18	1.160	35	1.846
~~OPHTHURCTIDEA (BRITTLE STARS)~~	0	0.0	3	0.158
~~OPHICHRAGMUS BURDEMANI~~	3	0.193	9	0.475
UNIDENTIFIED SP.				
HEMICHORDATA				
ENTEROPNEUSTA (ACRON WORMS)				
UNIDENTIFIED SP.	1	0.064	2	0.105
CEPHALOCHORDATA (LANCELETS)				
BRANCHIOSTOMA FLORIDAE	26	1.675	43	2.268
TOTALS	1552		1896	
NO. SPECIES		66		83
NO. IND. PER M2		2483		3034
S-W INDEX - H'(LN)		3.0020		3.3704
EVENNESS - J		0.7165		0.7627

APPENDIX D

FAUNAL SIMILARITY MATRICES


Similarity matrices for time-sequence samples at station 1, and one time sampling at stations 1 to 6 (Morisita's Index without transformations or standardizations, and with matrix values multiplied by 100)--beach restoration project, Panama City Beach, Florida (November 1974 to November 1977).

KEY

0.0 0.2 0.4 0.6 0.8 1.0



STATIONS 1-6 (7/11/77)




APPENDIX E


FAUNAL CLASSIFICATION ANALYSES

Classification analyses for time-sequence samples at station 1, and one-time sampling at stations 1 to 6 (Morisita's Index without transformations or standardizations)--beach restoration project, Panama City Beach, Florida (November 1974 to November 1977).

TREASURE ISLAND DREDGE EFFECTS (APRIL 1976- NOVEMBER 1977)



BEACH RESTORATION - ONE YEAR AFTER DREDGING AT SIX STATIONS




APPENDIX F


STABILITY ANALYSES

Stability analyses for time-sequence samples at station 1 showing sample variations and time to faunal recovery--beach restoration project, Panama City Beach, Florida (November 1974 to November 1977).

TREASURE ISLAND BEACH RESTORATION



TREASURE ISLAND BEACH RESTORATION



Saloman, Carl H.
 Benthic community response to dredging borrow pits, Panama City Beach, Florida / by Carl H. Saloman, Steven P. Naughton, and John L. Taylor.--Fort Belvoir, Va. : U.S. Army Coastal Engineering Research Center ; Springfield, Va. : available from NTIS, 1982.
 [138] p. : ill. ; 28 cm.--(Miscellaneous report ; no. 82-3)
 Prepared for Coastal Engineering Research Center by National Marine Fisheries Service, Southeast Fisheries Center; DACW72-81-M-0198.
 Report gives biological and physical oceanographic data from baseline work, and studies of dredged and undredged sediments before and after dredging (9-meter contour) at Panama City Beach, Florida. Analyses of hydrology, sediments, and benthos are included.
 1. Beach nourishment--Environmental aspects--Florida--Panama City Beach. 2. Benthos. 3. Dredging. 4. Panama City Beach (Fla.).
 I. Naughton, Steven P. II. Taylor, John L. III. Coastal Engineering Research Center (U.S.). IV. United States. National Marine Fisheries Service. V. Title. VI. Series: Miscellaneous report (Coastal Engineering Research Center (U.S.)); no. 82-3.
 TC203 .U581ar no. 82-3 627

Saloman, Carl H.
 Benthic community response to dredging borrow pits, Panama City Beach, Florida / by Carl H. Saloman, Steven P. Naughton, and John L. Taylor.--Fort Belvoir, Va. : U.S. Army Coastal Engineering Research Center ; Springfield, Va. : available from NTIS, 1982.
 [138] p. : ill. ; 28 cm.--(Miscellaneous report ; no. 82-3)
 Prepared for Coastal Engineering Research Center by National Marine Fisheries Service, Southeast Fisheries Center; DACW72-81-M-0198.
 Report gives biological and physical oceanographic data from baseline work, and studies of dredged and undredged sediments before and after dredging (9-meter contour) at Panama City Beach, Florida. Analyses of hydrology, sediments, and benthos are included.
 1. Beach nourishment--Environmental aspects--Florida--Panama City Beach. 2. Benthos. 3. Dredging. 4. Panama City Beach (Fla.).
 I. Naughton, Steven P. II. Taylor, John L. III. Coastal Engineering Research Center (U.S.). IV. United States. National Marine Fisheries Service. V. Title. VI. Series: Miscellaneous report (Coastal Engineering Research Center (U.S.)); no. 82-3.
 TC203 .U581ar no. 82-3 627

Saloman, Carl H.
 Benthic community response to dredging borrow pits, Panama City Beach, Florida / by Carl H. Saloman, Steven P. Naughton, and John L. Taylor.--Fort Belvoir, Va. : U.S. Army Coastal Engineering Research Center ; Springfield, Va. : available from NTIS, 1982.
 [138] p. : ill. ; 28 cm.--(Miscellaneous report ; no. 82-3)
 Prepared for Coastal Engineering Research Center by National Marine Fisheries Service, Southeast Fisheries Center; DACW72-81-M-0198.
 Report gives biological and physical oceanographic data from baseline work, and studies of dredged and undredged sediments before and after dredging (9-meter contour) at Panama City Beach, Florida. Analyses of hydrology, sediments, and benthos are included.
 1. Beach nourishment--Environmental aspects--Florida--Panama City Beach. 2. Benthos. 3. Dredging. 4. Panama City Beach (Fla.).
 I. Naughton, Steven P. II. Taylor, John L. III. Coastal Engineering Research Center (U.S.). IV. United States. National Marine Fisheries Service. V. Title. VI. Series: Miscellaneous report (Coastal Engineering Research Center (U.S.)); no. 82-3.
 TC203 .U581ar no. 82-3 627

Saloman, Carl H.
 Benthic community response to dredging borrow pits, Panama City Beach, Florida / by Carl H. Saloman, Steven P. Naughton, and John L. Taylor.--Fort Belvoir, Va. : U.S. Army Coastal Engineering Research Center ; Springfield, Va. : available from NTIS, 1982.
 [138] p. : ill. ; 28 cm.--(Miscellaneous report ; no. 82-3)
 Prepared for Coastal Engineering Research Center by National Marine Fisheries Service, Southeast Fisheries Center; DACW72-81-M-0198.
 Report gives biological and physical oceanographic data from baseline work, and studies of dredged and undredged sediments before and after dredging (9-meter contour) at Panama City Beach, Florida. Analyses of hydrology, sediments, and benthos are included.
 1. Beach nourishment--Environmental aspects--Florida--Panama City Beach. 2. Benthos. 3. Dredging. 4. Panama City Beach (Fla.).
 I. Naughton, Steven P. II. Taylor, John L. III. Coastal Engineering Research Center (U.S.). IV. United States. National Marine Fisheries Service. V. Title. VI. Series: Miscellaneous report (Coastal Engineering Research Center (U.S.)); no. 82-3.
 TC203 .U581ar no. 82-3 627

DATE
ILMEI
—8