


AD-A116 374

LETTERMAN ARMY INST OF RESEARCH PRESIDIO OF SAN FRANC--ETC F/6 6/5  
BODY COMPOSITION AND WORK PERFORMANCE OF CADETS AT THE UNITED S--ETC(U)  
JUN 82 R D FULTS, M S MORRIS, H L JOHNSTON

UNCLASSIFIED

NL

For  
AD-A116 374



END  
DATE  
FILMED  
7-82  
DTIC

AD A116374

142

INSTITUTE REPORT NO. 125

BODY COMPOSITION AND WORK PERFORMANCE OF CADETS  
AT THE UNITED STATES MILITARY ACADEMY,  
WEST POINT, NEW YORK

ROBERT D. FULTS, BS  
MICHAEL S. MORRIS, BA  
HERMAN L. JOHNSON, PhD  
ROBERT E. MORRIS, MD, MAJ MC  
and  
HOWERDE E. SAUBERLICH, PhD

DIVISION OF NUTRITION TECHNOLOGY

DTIC  
JUL 2 1982  
H

DTIC FILE COPY

JUNE 1982

DISTRIBUTION STATEMENT A  
Approved for public release;  
Distribution Unlimited

LETTERMAN ARMY INSTITUTE OF RESEARCH  
PRESIDIO OF SAN FRANCISCO, CALIFORNIA 94129

**Body Composition and Work Performance of Cadets at the United States Military Academy, West Point, New York--Fults et al**

Reproduction of this document in whole or in part is prohibited except with the permission of the Commander, Letterman Army Institute of Research, Presidio of San Francisco, California 94129. However, the Defense Technical Information Center is authorized to reproduce the document for United States Government purposes.

Destroy this report when it is no longer needed. Do not return it to the originator.

Citation of trade names in this report does not constitute an official endorsement or approval of the use of such items.

Human Subjects participated in these studies after giving their free and informed voluntary consent. Investigators adhered to AR 70-25 and USAMRDC Reg 50-25 on the use of volunteers in research.

This material has been reviewed by Letterman Army Institute of Research and there is no objection to its presentation and/or publication. The opinions or assertions contained herein are the private views of the author(s) and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense. (AR 360-5)

*John W. Marshall Jr.* 14 June 82  
(Signature and date)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER LAIR Institute Report No. 125	2. GOVT ACCESSION NO. AD-A116 374	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Body Composition and Work Performance of Cadets at the United States Military Academy, West Point, New York		5. TYPE OF REPORT & PERIOD COVERED Final Report April 1979 - April 1982
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Robert D. Fults, BS, Michael S. Morris, BA, Herman L. Johnston, PhD, Robert E. Morris, MD, MAJ MC, Howerde E. Sauberlich, PhD		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Nutrition Technology Division, Letterman Army Institute of Research, Presidio of San Francisco, CA 94129		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 62772A; 3M162772 A811; 00;001
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Medical Research and Development Com- mand, Fort Detrick, Frederick, MD 21701		12. REPORT DATE June 1982
		13. NUMBER OF PAGES 43
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Cadet, Army, Military Academy, nutrition, body composition, work performance, body fat, anthropometry		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A study was conducted to evaluate the cadet weight control program at the United States Military Academy, West Point, New York. During the spring academic quarter of 1979, eighty-six cadets at the Academy were en- rolled in the weight control program. This represented 2% of the male and 17% of female cadets enrolled. This report is concerned with the body composition and work performance aspects of the study. The study was conducted in two phases. Phase I, conducted during the spring quarter		

of 1979, evaluated the procedures used to estimate the body composition of the cadets. Phase II, conducted in the fall of 1979, studied the body composition and the work performance of the cadets.

In Phase I there were no significant differences in height, weight, skinfold thickness, or circumference values with respect to which entrance class the cadets were in. The recognized sex differences were observed; namely, males are generally relatively larger than females, but females are relatively fatter. However, subscapula and suprailiac skinfold values and thigh circumference measurements did not differentiate males from females. Comparisons of methods used by LAIR and USMA for the determination of the percent of body fat were highly favorable. Although the body fat estimate by the USMA method was significantly lower by 0.4 percent of body fat, this small difference in percent of body fat is of no practical significance. Hence, the procedures used at USMA to estimate the percent of body fat appear reliable.

The observations noted in phase I were found again six months later in phase II. However, a uniform weight gain that averaged 1.2 kilogram per cadet occurred during this period. In addition, subscapular, abdominal, and thigh skinfold values fell between the phases. The increase in weight could not be accounted for by an increase in percent of body fat. The male cadets exceeded the female cadets in work performance. Oxygen uptake ( $V_{O_2max}$ ) was greater for the males either on an average individual basis or per kilogram of body weight. No differences in performance were observed between the four academic classes of male or female cadets. The data were inadequate to identify an optimum body composition for the cadets.

Accession For	✓
NTIS GRAM	
DTIC TAB	
Unannounced	
Justification	
By	
Distribution/	
Availability	
Dist	
A	

DTIC  
COPY  
INSPECTED  
2

## ABSTRACT

A study was conducted to evaluate the cadet weight control program at the United States Military Academy, West Point, New York. During the spring academic quarter of 1979, eighty-six cadets at the Academy were enrolled in the weight control program. This represented 2% of the male and 17% of females cadets enrolled. This report is concerned with the body composition and work performance aspects of the study. The study was conducted in two phases. Phase I, conducted during the spring quarter of 1979, evaluated the procedures used to estimate the body composition of the cadets. Phase II, conducted in the fall of 1979, studied the body composition and the work performance of the cadets.

In phase I there were no significant differences in height, weight, skinfold thickness, or circumference values with respect to which entrance class the cadets were in. The recognized sex differences were observed; namely, males are generally relatively larger than females, but females are relatively fatter. However, subscapula and suprailiac skinfold values and thigh circumference measurements did not differentiate males from females. Comparisons of methods used by LAIR and USMA for the determination of the percent of body fat were highly favorable. Although the body fat estimate by the USMA method was significantly lower by 0.4 percent of body fat, this small difference in percent of body fat is of no practical significance. Hence, the procedures used at USMA to estimate the percent of body fat appear reliable.

The observations noted in phase I were found again six months later in phase II. However, a uniform weight gain that averaged 1.2 kilogram per cadet occurred during this period. In addition, subscapular, abdominal, and thigh skinfold values fell between the phases. The increase in weight could not be accounted for by an increase in percent of body fat. The male cadets exceeded the female cadets in work performance. Oxygen uptake ( $V_{O_2}$  max) was greater for the males either on an average individual basis or per kilogram of body weight. No differences in performance were observed between the four academic classes of male or female cadets. The data are inadequate to identify an optimum body composition for the cadets.

## TABLE OF CONTENTS

	<u>Page</u>
Abstract	i
Table of Contents	iii
BODY OF REPORT	
INTRODUCTION	1
EXPERIMENTAL DESIGN	2
METHODS	4
RESULTS AND DISCUSSION	6
CONCLUSIONS	9
RECOMMENDATION	10
REFERENCES	11
APPENDICES	
Appendix A (USCC Weight Control Program)	13
Appendix B (Tables 1-13)	14
OFFICIAL DISTRIBUTION LISTS	36

A study of body composition and work performance of cadets at the United States Military Academy, West Point, New York (USMA) was conducted at the request of the USMA Department of Physical Education (DPE)(1). Results of the study are the subject of this report.

During the spring academic quarter of 1979, eighty-six cadets (45 male, 41 female) were enrolled in the cadet weight control program (CWCP) administrated by DPE (Appendix A). These numbers represented 2% of the male and 17% of the female population in the United States Corps of Cadets (USCC) at that time. In addition, 38 other cadets (31 male, 7 female) had been enrolled and later released from the CWCP during the 1978-1979 academic year.

This situation concerned the USMA command staff because a policy was being formulated for separating obese cadets from the academy. With the cost of education near \$100,000 per cadet, a substantial financial impact would result by separating even a few cadets. Because obesity is a potentially correctable problem, the CWCP was instituted and this study as well as addenda were implemented to enhance its effectiveness.

The purpose of this study was to evaluate the CWCP and formulate recommendations for improvement. Within the context of cadet life at the USMA the number of cadets participating in the CWCP was considered to be unacceptably high and indicative of a possibly general weight control problem. Three possible reasons were indentified as explanations for the size of CWCP population. First, the cadets may indeed be overfat due to food consumption in excess of daily energy requirements. Second, the CWCP method for measuring body fat may yield incorrect results and indentify cadets as being overfat when they are not. Third, the CWCP criterion for selecting overfat cadets may be incorrect or inappropriate. Identification of any of these reasons was within the scope of this study.

Due to limitations imposed by availability of manpower, equipment, and cadet time, the study was conducted in two phases. Phase I was accomplished in the spring quarter of 1979; Phase II in the fall of 1979. Phase I was a study of body composition intended to examine the first and second considerations listed above. Phase II studied body composition and work performance in order to evaluate the third consideration as well as help indentify body composition trends that may occur during the academic year.

## EXPERIMENTAL DESIGN

It was estimated that 25 males and 25 females from each class, plus 25 males and 25 females from the CWCP, would provide adequate statistical power for the multifactor repeated measures design appropriate to this study. Thus the prospective populations were 225 subjects for Phase I (there were no females in the class of 1979) and 250 subjects for Phase II. The experimentally accessible population for Phase I was 240; however, complete data were obtained only on 221. In Phase II, complete data were obtained from 138 subjects, of which 58 had also completed Phase I of the study. As shown in Table 1, the distribution was not 25 subjects per cell as desired.

The non-uniform distribution of study subjects was due to the voluntary subject participation and the time constraints built into a field study. Collecting complete data required 5 to 7 hours from each subject. This time was divided into several sessions that the subjects were free to schedule according to their individual needs. It was important that interference with cadet academic work be avoided. Thus, a number of subjects withdrew from the study and appointments were frequently cancelled by the subjects too late for the time slot to be reassigned. The investigation team was limited to 5 weeks (Phase I) and 7 weeks (Phase II) on site and eventually time expired before all cancellations could be rescheduled or all dropouts could be replaced.

Participation was also limited by another important factor. Only 9 cadets from the CWCP volunteered. Although they were assured of the strict confidentiality of the data, cadets already under scrutiny in the CWCP were reluctant to call further attention to themselves by participation in this study. There were likewise cadets in the general population who, to avoid jeopardy under the CWCP, refused to volunteer. This was an inherent disadvantage to studying a criterion for dismissal from the academy.

Since the analysis program, BMDP2V (2), accommodates unequal group sizes, there were no problems with data analysis. However, the small number of volunteers from CWCP precluded any comparisons between cadets in CWCP and the general population. Thus, data from all cadets were pooled and the CWCP factor was removed from the design.

Data from Phase I were acquired in order to evaluate: (a) the validity and accuracy of CWCP body fat estimates; (b) measure the actual body fatness of the USMC population, and (c) identify trends in body fatness related to sex or tenure at the USMA. The following hypotheses (stated in null form) were among those tested:

- I-1. There is no difference between body fat estimates made according to the CWCP protocol and estimates made using generally accepted (and presumably more accurate) laboratory methods.
- I-2. There is no difference in relative body fatness between male and female cadets.
- I-3. There are no differences in relative body fatness among classes.
- I-4. There is no interaction between the effects of sex or class on relative body fatness.
- I-5. There is no interaction among class, sex, and methodology on body fatness measurements.

Data from Phase II were acquired for the purpose of evaluating changes in body composition over a 6-month period as well as work performance and energy expenditure of cadets. In addition, a first order attempt was made to identify objectively the optimum body composition for cadets (an admittedly optimistic undertaking). The following null hypotheses were among those tested:

- II-1. There is no difference in work performance between male and female cadets.
- II-2. There are no differences in work performance among classes.
- II-3. There is no class by sex interaction affecting work performance.

Data from both phases were used to test the following hypotheses.

- III-1. There is no difference in relative body fatness between Phase I and Phase II.
- III-2. There are no interactions among class, sex, and phase affecting relative body fatness.

Other hypotheses were tested in order to facilitate interpretation of the results. They are described at the appropriate points in the discussion.

## METHODS

During Phase I, body composition measurements were obtained before the subjects ate breakfast (between 0430 hrs and 0630 hrs) on weekdays. These measurements included body volume, weight, height, 14 skinfold thicknesses and 7 circumferences. Body volume and weight were measured on subjects dressed in nylon swimsuits. The weight and volume of these suits were determined to be within the error of reproducibility of the measurements and therefore introduced negligible error into the results.

The 14 skinfolds were measured bilaterally at seven sites: triceps (TCR, TCL); subscapula (SSR, SSL); biceps (BCR, BCL); supra-iliac (SIR, SIL); lateral abdomen (LAR, LAL); abdomen (ABR, ABL); and thigh (THR, THL). The site definitions for TCR, TCL, SSR, SSL, BCR, BCL, SIR, and SIL were taken from Durnin and Wommersley (3). The LAR and LAL measurements were devised by DPE for use in the CWCP and defined as the thickness of a vertical skinfold made at the intersection of a horizontal line across the abdomen at the level of the iliac crest and a vertical line intersecting the anterior process of the pelvis. The ABR and ABL skinfolds were defined as the thickness of horizontal skinfolds measured immediately lateral to the umbilicus. The THR and THL measurements were vertical skinfolds on the anterior thigh midway between the condyles of the femur.

The seven circumferences were measured around the neck (NK), shoulders (SH), chest (CH), abdomen (AB), buttocks (BU), and thigh (bilaterally; TR, TL). The NK measurement was taken just inferior to the larynx. The SH was measured over the deltoids. The CH was taken at the end of a passive expiration at the level of the xiphoid process. The AB was measured over the umbilicus. The BU was measured at the level of the top of the femur. Circumferences TR and TL were measured over the THR and THL skinfold sites.

Skinfolds were measured to the nearest 0.5 mm with a Lange caliper (4) and the average of three trials was recorded. Weight was recorded to the nearest 10 g from a triple beam balance (5). Height was recorded to the nearest millimeter read from a free standing anthropometer (6). Circumferences were measured to the nearest millimeter with a constant tension steel tape measure (7).

Body volume was measured in a human body volumeter using the method of Allen et al. (8) the method routinely used at the Letterman Army Institute of Research. The water level in the volumeter was recorded to the nearest 0.25 mm prior to the subject's entry. After entering the tank, each subject performed a forced expiration

to residual volume and submerged for about 10 seconds. This procedure was repeated until successive trials agreed within 1 mm on the water level scale. The smallest volume obtained was recorded.

Residual volume (RV) was measured while the subject stood submerged to the neck in the volumeter. While wearing a nose clip, the subjects performed a forced expiration to residual volume, inspired 200 ml of helium from a rubber aliquot bag fitted with a mouthpiece, then repeatedly respired into the bag to mix the helium and residual gas thoroughly. Upon the final expiration, the bag was clamped and the helium concentration ( $F_{EHe}$ ) measured with a catharometer (9). The residual volume (RV) was calculated from the helium dilution equation:  $RV = 200(1 - F_{EHe}) / F_{EHe}$  ml. This method was developed for speed and convenience of use within the confines of the volumeter. Its validity will be reported elsewhere.

In addition to the early morning body composition measurements, whole body burdens of potassium-40 ( $^{40}K$ ) were determined by gamma radiation counting. The  $^{40}K$  data were obtained using the LAIR shadow shield whole body counter (10). For this test, subjects wore duty uniforms and the half-hour scan was done sometime between 0700 and 2200 hrs.

During the procedure, the subjects were clothed and lying supine on a padded sled. The sled, under mechanical control, was slowly moved past a thallium-activated sodium-iodide crystal. The crystal, sensitive to the natural gamma radiation emitted by the body, is connected to a gamma spectrum analyzer. Data from a 30 minute scan were transferred from the analyzer to a minicomputer (MCII) (11), which quantitated the radioactivity associated with the 1.46 MeV  $^{40}K$  photopeak. From this the amount of potassium in the body was computed and from that the fat-free body mass derived (12).

Data acquired during Phase II included the anthropometry described above with the exception of circumferences. Work performance abilities were determined from a graded exercise test (GXT). Anthropometry measurements were made at the same testing session. The measurements, which required one-hour sessions, were performed between 0800 and 1700 hr on weekdays.

In the GXT protocol, the subjects walked on a motor driven treadmill (13) at a quick but comfortable pace (3.0 to 4.0 mph). At two-minute intervals, the elevation was raised 2% until the subjects voluntarily terminated the test. Throughout the test the blood pressure (BP) was monitored with an electronic sphygmomanometer (14).

The electrocardiogram (ECG) was periodically recorded and monitored on a 3-channel recorder (15) and continuously displayed on a 2-channel CRT (16). A physician was present during all GXT measurements.

Breath-by-breath oxygen consumption ( $V_{O_2}$ ) was measured with a respiratory mass spectrometer (RMS) (17). During the last minute at each grade, the breath by breath  $V_{O_2}$  values were integrated and a 10 beat average heart rate was measured from the ECG recording. The data were transferred to a MCII minicomputer where maximal oxygen uptake ( $V_{O_{2max}}$ ) as well as the relationship of  $V_{O_2}$  to heart rate from rest to maximum were computed.

For both phases of this study, data reduction, file preparation, and graphics were done on the MCII and an Eclipse C/330 minicomputer (18). Statistical analysis and archival were performed using the BMDP package (2) on a CDC 7600 computer (19).

## RESULTS AND DISCUSSION

The distribution of subjects within factors of the experimental design is shown in Table 1. With the exception of the weight control groups, the participation goal of 25 subjects per cell was essentially met in phase I. There were only 17 females representing the class of 1981, however, in light of the excess numbers in other cells the power of the design was adequate.

Participation in phase II was not up to expectation. Females in particular and the freshman class in general were reluctant to volunteer. The reason for the lack of interest on the part of the females is not clear. The freshman class had only been attending classes for two months at the start of this phase of the study. They seemed to feel that the time commitment required to participate in the study would interfere with their academic work. This would apparently be the consensus of any freshman class early in the academic year. In contrast, the freshman class showed the highest participation in phase I, which was conducted in late spring. One might conclude that 6 to 8 months after matriculation they have adjusted to the academic environment and are ready for extra-curricular activities.

The data acquired in phase I are shown in Tables 2 through 6. Height and weight data are found in Table 2. The skinfold values are presented in Table 3 and circumferences in Table 4. Body fat determination comprises Table 5, while comparison of measurements made by different investigators are shown in Table 6.

The data in Tables 2 and 4 were analyzed as a 2x4 (sex by class) factor design using multifactor analysis of variance program BMDP2V (2). The following results were obtained from the analysis:

1. There were no significant differences among classes for any of the variables measured.
2. There were no significant interactions between sex and class factors.
3. In general the effect of sex on any parameter was significant at the  $<0.05$  level.

Exceptions to the sex effect stated above are noted in the tables. The obvious sex differences were confirmed; that is, males are generally relatively larger but females are relatively fatter. It is interesting to note that the subscapula (SSL,SSR) and suprailiac (SILR,SIR) skinfold values and the thigh (TR) circumference values cannot be used to differentiate between males and females.

In order to compare methodology, the data in Table 5 were analyzed by a 2x4x4 (sex by class by method) multifactor repeated measured design using the BMDP2V program. The analysis produced the following results:

4. There were no significant differences in percent body fat among classes.
5. The effect of sex on percent body fat was significant at the  $p<0.05$  level.
6. The effect of methodology on the percent body fat measurement was significant at the  $p<0.05$  level.
7. All interactions among factors were significant at the  $p<0.05$  level.

Based upon these analysis, the following statement about the hypothesis for phase I are appropriate:

- Hypothesis I-1: (no methodology effect) may be rejected at the  $p<0.05$  level.
- Hypothesis I-2: (no sex effect) may be rejected at the  $p<0.05$  level.
- Hypothesis I-3: (no class) effect is tenable.
- Hypothesis I-4: (no class by sex interaction) is tenable.
- Hypothesis I-5: (no class by sex by methodology interaction) may be rejected at the  $p<0.05$  level.

In order to test the influence of investigator technique on the anthropometry method in use at West Point, measurements were made on a subset of the subjects by the DPE instructors who usually perform fat estimates for the CWCP. The measurements on subjects in this subset made by the DPE instructor (USMA) were pooled, and compared with the LAIR measurements. The data are shown in Table 6.

The data were analyzed as a 2x4x2 (sex by class by investigator) multifactor repeated measures of analysis of variance (ANOVA). The differences between investigators for triceps and biceps measurements were not significant. The DPE subscapular measurement (SWP) was significantly lower ( $p < .001$ ) by 0.75 mm. The DPE lateral abdominal measurements was significantly higher ( $p < .038$ ) by 0.13 mm. The resulting relative body fat estimate by DPE was significantly lower ( $p < .003$ ) by 0.4 % of body fat.

Although these differences are statistically significant (attesting to the statistical power of the experimental designs), they are not meaningful because they are quite small in the practical sense. A difference of only 0.4% body fat estimates made independently by three individuals indicates that the anthropometric measurement can be made reliably by a trained person. A regression analysis was performed (using BM DP6Q) on the investigation factor. The regression equation of  $y = 0.997x + 0.404$  and the correlation coefficient approaching unity also indicate the reliability of measurements made by different individuals. The intercept (0.4) corroborates the lower DPE estimate identified by the ANOVA. It is worthwhile to emphasize that a difference of 0.4 in percent of body fat is of no practical significance.

Data acquired during phase II of this study are shown in Tables 7 through 10. Height and weight values are given in Table 7. Skinfold thicknesses values are found in Table 8. The anthropometric fat estimates are shown in Table 9 and the results of the GXT are provided in Table 10.

These data were analyzed as a 2x4 (sex by class) design using BMDP2V. The following results were obtained:

1. There were no significant differences among classes for any variable.
2. There are no significant class by sex interactions for any variable.
3. The males and females were significantly different for all variables, except for the subscapular, lateral abdominal, and abdominal skinfolds measurements.

The obvious trends due to sex were again noted. The null hypotheses about phase II may now be evaluated.

Hypothesis II-1: (no sex difference in performance) may be rejected at the  $p = 0.002$  level.

Hypothesis II-2: (no class differences in the performance) is tenable.

Hypothesis II-3: (no class by sex interaction) is tenable.

The data from subjects completing both phase I and phase II are presented in Tables 11 through 13. The height and weight data are found in Table 11. Table 12 shows the skinfold data, and Table 13 shows the relative body fat data. When analyzed in a  $2 \times 4 \times 2$  (sex by class by phase) repeated measures design using the BMDP2V program, the following results were obtained:

1. There were no significant differences among classes for any variable.
2. The few significant differences under the sex factor showed the obvious trends.
3. There was a uniform weight gain averaging 1.2 kg from phase I to phase II.
4. A phase effect was noted by changes in the subscapular abdominal and thigh skinfold values.

The last of the null hypotheses in this study may now be evaluated:

Hypothesis III-1: (no change in percent of body fat) is tenable.

Hypothesis III-2: (no class, sex, phase interaction) is tenable.

#### CONCLUSIONS

In phase I there were no significant differences in height, weight, skinfold thickness, or circumference values with respect to which entrance class the cadets were in. The recognized sex differences were observed; namely, males are generally relatively larger than females, but females are relatively fatter. However, subscapula and suprailiac skinfold values and thigh circumference measurements could not differentiate between males and females.

Comparison of methods used by LAIR and USMA for the determination of the percent of body fat were highly favorable. Although the body fat estimate by the USMA method was significantly lower by 0.4 percent of body fat, this small difference in percent of body fat is of no practical significance. Hence, the procedures used at USMA to estimate the percent of body fat appear reliable.

The observations noted in phase I were found again six months later in phase II. However, a uniform weight gain that averaged 1.2 kilogram per cadet occurred during this period. In addition, subscapula, abdominal, and thigh skinfold values fell between the phases. The increase in weight could not be accounted for by an increase in percent of body fat.

The male cadets exceeded the female cadets in work performance. Oxygen uptake ( $V_{O_2}$  max) was greater for the males either on an average individual basis or per kilogram of body weight. No differences in performance were observed between the four academic classes of male or female cadets.

The data were too inadequate to identify an optimum body composition for the cadets.

#### RECOMMENDATION

None.

## REFERENCES

1. FULTS, R.D., R.E. MORRIS, J.H. SKALA, H.L. JOHNSON and H.E. SAUBERLICH. Evaluation of Body Composition, Energy Expenditure and Their Relationships to Work Performance of United States Military Academy Cadets, LAIR Protocol. Letterman Army Institute of Research, Presidio of San Francisco, California, 29 March 1979
2. BMDP Biomedical Computer Programs p-series, Health Sciences Computing Facility, Department of Biomathematics School of Medicine, University of California, Los Angeles, California, 1977
3. DURBIN, J.V.G.A and J. WOMMERSLEY. Body fat assessed from total body density and its estimation from skinfold thickness: measurements on 481 men and women aged from 16 to 72 years. Br J Nutr 32:77-97, 1974
4. Lange Calipers. Cambridge Scientific Industries Inc., Cambridge, Maryland
5. Homs Full Capacity Beam Scale. Douglas Homs Corporation, Belmont, California
6. WILMORE, J.H. and A.R. BEHNKE. Predictability of lean body weight through anthropometric assessment in college men. J Appl Physiol 25:349-355, 1968
7. WARD, G.M., H.J. KRZYWICKI, D.P. RAHMAN, R.L. QUAAS, R.A. NELSON and C.F. CONSOLAZIO. Relationship of anthropometric measurements to body fat as determined by densitometry, potassium-40, and body water. Am J Clin Nutr 28:162-169, 1975
8. ALLEN, T.H., H.J. KRZYWICKI, W.W. WORTH and R.M. NIMS. Human Body Volumeter Based on Water Displacement. Report No. 250. Denver, Colorado: US Army Medical Research and Nutrition Laboratory, September 1960
9. Helium Meter, Warren E. Collins, Inc., Braintree, Massachusetts 02184
10. SCAIEF, C.C., and H.J. KRZYWICKI. Development and Application of a Whole-Body Radiation System: Report No. 330. Denver, Colorado: US Army Medical Research and Nutrition Laboratory, November, 1971
11. Mod-Comp II Modular Computer Systems, Inc., Ft. Lauderdale, Florida

12. REMENCHIK, A.P., R.K. HUKKOO and C.E. MILLER. Determination of body composition by gamma spectrometry. Develop Appl Spectros 5:437-458, 1966
13. Treadmill, Model 3800. Warren E. Collins, Inc., Braintree, Massachusetts 02184
14. Infrasonde, Model 3000. Marion Scientific Corporation, Costa Mesa, California
15. Model 5330 Cardiograph. American Optical Corporation Medical Division, Bedford, Massachusetts
16. Dura Vue II. American Optical Corporation Medical Division, Bedford, Massachusetts
17. Respiratory Mass Spectrometer Model 911-5030 University of Colorado Medical School, Denver, Colorado
18. Eclipse C/330. Data General Corporation, Westboro, Massachusetts
19. CDC 7600. Control Data Corporation, Minneapolis, Minnesota

Cadets Weight Control Program

APPENDIX A

THE UNITED STATES CORPS OF CADETS

WEIGHT CONTROL PROGRAM

RESPONSIBILITY:

The Director, Department of Physical Education (DPE), is responsible for the administration of the USCC Weight Control Program.

IDENTIFICATION:

Cadets will undergo biannual height/weight surveys, typically held in conjunction with regularly scheduled DPE testing. Those cadets found to exceed, or be within 2% of exceeding, the maximum body weights in AR 600-9 will be required to report to DPE for a percent body fat measurement.

Additionally, cadets are required to report to DPE for a body fat measurement when:

- a. Volunteering for the program.
- b. Directed by an officer or another cadet in the chain of command.

PERCENT BODY FAT CALCULATION:

Percent body fat will be calculated for cadets using skinfold techniques and mathematical formulas, a relatively simple and accurate method. The normal error encountered by such measurements is taken into account in developing standards. The skinfold technique accounts for body build, muscular development, muscle tone, and bone structure. AR 600-9, Paragraph 3-4, indicates that these body parameters should be included in obesity determination.

Once the percentage of body fat is calculated, cadets will be categorized as follows:

PERCENT BODY FAT

<u>CATEGORY</u>	<u>MALE (%)</u>	<u>FEMALE (%)</u>	<u>REMARKS</u>
V	0-5	0-8	Very low, caution
IV	5.1-10	8.1-17	Excellent
III	10.1-15	17.1-22	Acceptable
II	15.1-16	22.1-23	High, bi-weekly weigh-ins required.
I	over 16.1	over 23.1	Mandatory Weight Control program.

## WEIGHT CONTROL PROGRAM

Cadets who are in Categories I or II are required to report to DPE Weight Control Clinic every 7-14 days to be weighed.

Cadets in Category I will be allowed a period of time to lose adipose tissue (fat) in order to achieve Category II or a higher category. Time allowed to lose fat will be equal to one week for every 0.5% body fat they exceed the body fat limits of Category II; however, the time period will not be less than two weeks. For example, a male cadet may weigh 200 pounds of which 19% is fat. Since his percent body fat is 3% over the 16% Category II limit, he is allowed 6 weeks to lose that 3% body fat. In this case, the 3% body fat equals 6 pounds of adipose tissue.

If a cadet has not attained the appropriate level of body fat at the end of the prescribed time period, there are two possible courses of action. At the discretion of the Director, DPE, the cadet will be recommended for dismissal, or will be granted a time extension to reach the required goal. The extension, if granted, will be no longer than one-half the originally prescribed time period. For example, the cadet described above is found to have 18% body fat at the end of the 6 weeks. He could be granted a maximum extension of 3 weeks to reach the Category II level of 16% body fat. Should the cadet still fail to reach a Category II level during the extension, the cadet may be recommended for dismissal.

Cadets who fall into Category II, or higher, and who regress into Category I two (2) or more times during any one Academic Year, may be recommended for administrative dismissal. For example, in September a female cadet may weigh 140 pounds with 24.5% body fat. She is granted 3 weeks to make the 23% Category II limit. At the end of the 3 week period she is found to have 22.5% body fat which is within the Category II limit. The cadet is then required to weigh-in at DPE bi-weekly. A weight gain is discovered in November, and a new resultant percent body fat measurement shows her to have 23.5% body fat. She is then allowed another period of two weeks to attain the Category II level. If she reaches the prescribed level in 2 weeks and a month later found to have 21.5% body fat, she would be released from the program. If the cadet is administered another percent body fat measurement in February as a result of a weight gain shown in a routine class height/weight survey, and found to be 24% body fat, she could be recommended for dismissal.

### ADMINISTRATIVE DISMISSAL:

Administrative dismissal under the Weight Control Program will be processed in accordance with Regulations for the United States Military Academy, paragraph 12.04 (proposed para 10.08, Revised Regs USMA .

LIST OF TABLES

- Table 1 - Distribution of Subjects from USCC
- Table 2 - Weight and Height Data from Phase I
- Table 3 - Skinfold Data from Phase I
- Table 4 - Circumference Data from Phase I
- Table 5 - Percent Body Fat Data from Phase I
- Table 6 - Comparison of Investigator Differences
- Table 7 - Weight and Height Data from Phase II
- Table 8 - Skinfold Data from Phase II
- Table 9 - Percent Body Fat from Phase II
- Table 10 - Oxygen Uptake Data from Phase II
- Table 11 - Weight and Height Data from Subjects Completing both Phases  
I and II
- Table 12 - Skinfold Data from Subjects Completing both Phases I and II
- Table 13 - Percent Fat Estimates from Subjects Completing both Phases  
I and II

APPENDIX B

Table 1. Distribution of Subjects from USCC

Phase	Subjects	Class (of) <sup>1</sup>					Total
		1979	1980	1981	1982	1983	
Phase I	Female	...	27	17	25	...	69
	Male	26	39	48	49	...	162
	Total	26	66	65	74	...	231
Phase II	Female	...	14	14	6	8	42
	Male	...	26	24	33	11	94
	Total	...	40	38	39	19	136
Phase I and II	Female	...	13	4	4	...	21
	Male	...	12	13	12	...	37
	Total	...	25	17	16	...	58

<sup>1</sup>There were no females in the class of 1979 and the class of 1983 did not participate in phase I.

Table 2. Weight and Height Data from Phase I

Parameter	Sex	Class <sup>1</sup>				Significant Difference (Sex Effect)
		First-79	Second-80	Third-81	Fourth-82	
Weight (kg)	F	...	61.16+ 7.90	58.95+5.76	59.30+6.13	p <.001
	M	76.32+10.32	76.96+10.49	77.55+9.78	73.93+6.24	
Height (cm)	F	...	167.52+ 6.57	165.05+7.29	166.48+7.47	p <.001
	M	180.73+ 8.44	178.86+ 6.64	178.55+6.00	177.38+6.04	

<sup>1</sup>The values represent mean + standard deviation. No significant differences among classes and no significant interactions between class and sex were identified for height or weight.

Table 3. Skinfold Data from Phase I

Parameter	Sex	Class <sup>1</sup>				Significant Difference (Sex Effect)
		First-79	Second-80	Third-81	Fourth-82	
TCR (mm)	F	...	13.44±3.37	14.24±3.45	14.22±2.96	p<.001
	M	7.78±2.91	7.96±2.65	7.88±2.91	8.47±2.71	
TCL (mm)	F	...	13.06±3.34	14.41±3.51	14.72±3.08	p<.001
	M	7.48±2.93	7.61±2.58	7.85±3.07	8.05±2.46	
SSR (mm)	F	...	10.72±3.56	10.56±1.68	10.36±2.46	NSD
	M	9.98±2.95	10.54±2.70	10.92±3.12	9.97±2.43	
SSL (mm)	F	...	10.65±3.29	10.59±1.73	10.22±2.81	NSD
	M	9.67±2.56	10.21±2.50	10.27±2.94	9.82±2.33	
BCR (mm)	F	...	5.91±2.52	5.53±1.43	6.00±2.23	p<.001
	M	3.90±1.20	3.73±1.20	3.96±1.22	3.90±1.04	
BCL (mm)	F	...	6.07±2.51	5.76±1.68	6.06±2.02	p<.001
	M	4.13±1.26	4.20±1.52	4.31±1.34	4.19±1.18	
SIR (mm)	F	...	12.33±5.13	12.35±3.18	11.32±3.91	NSD
	M	13.54±5.68	13.54±5.10	13.33±5.03	12.03±4.19	
SIL (mm)	F	...	12.57±5.78	12.76±3.61	11.72±3.50	NSD
	M	13.38±5.10	13.07±4.94	13.09±5.11	12.20±3.49	

Table 3. (continued)

Parameter	Sex	Class				Significant Difference (Sex Effect)
		First-79	Second-80	Third-81	Fourth-82	
LAR (mm)	F	...	8.17±3.88	7.76±1.95	7.88±3.06	p<.005
	M	6.73±2.87	6.70±2.68	6.95±2.87	6.56±2.06	
LAL (mm)	F	...	8.00±3.81	7.91±2.24	7.40±2.75	NSD
	M	6.79±3.30	7.17±3.11	7.23±2.97	6.55±2.34	
ABR (mm)	F	...	11.31±4.76	12.74±3.44	10.74±3.78	p<.001
	M	9.77±4.05	9.33±3.87	9.70±4.06	8.70±2.94	
ABL (mm)	F	...	11.61±5.12	13.20±3.70	11.20±3.49	p<.001
	M	9.04±4.00	8.91±4.05	9.21±3.88	8.36±3.00	
THR (mm)	F	...	21.98±5.65	22.53±4.47	23.06±3.94	p<.001
	M	10.40±3.43	11.58±5.04	11.28±3.94	11.47±4.18	
THL (mm)	F	...	22.06±5.18	22.88±4.30	23.00±3.95	p<.001
	M	10.92±3.23	11.99±5.28	11.81±4.07	12.03±4.04	

The values represent mean ± standard deviation

NSD means p>0.05. No significant differences among classes and no significant interaction between class and sex were identified for any skinfold parameter.

Table 4. Circumference Data from Phase I

Parameter	Sex	Class <sup>1</sup>				Significant Difference <sup>2</sup> (Sex Effect)
		First-79	Second-80	Third-81	Fourth-82	
NK (cm)	F	...	31.92±1.55	31.33±1.74	31.73±1.52	p<.001
	M	37.70±1.75	37.88±2.17	38.00±2.21	37.77±1.69	
SH (cm)	F	...	100.62±5.03	98.90±4.22	99.55±3.71	p<.001
	M	112.93±6.9	113.67±6.09	114.48±6.57	112.53±4.37	
SB (cm)	F	...	74.71±4.38	72.64±4.83	73.20±2.84	p<.001
	M	89.43±7.10	90.25±6.18	91.13±6.40	88.43±5.17	
AB (cm)	F	...	69.56±6.06	69.45±6.09	68.79±5.42	p<.011
	M	78.89±5.83	79.73±5.08	79.82±6.02	78.44±4.94	
BU (cm)	F	...	95.79±5.28	95.02±4.18	93.78±4.34	p<.014
	M	96.99±6.49	96.92±5.09	97.69±5.73	95.42±4.07	
TR (cm)	F	...	56.59±3.45	56.55±3.04	55.95±3.16	NSD
	M	57.44±4.14	57.58±4.34	58.01±5.35	57.00±2.92	
TL (cm)	F	...	56.21±3.58	56.44±2.63	55.53±3.26	p<.038
	M	36.91±4.14	57.43±4.40	57.90±5.30	56.60±2.88	

<sup>1</sup>The values represent mean ± standard deviation.

<sup>2</sup>NSD means p>.05. No significant differences among classes and no significant interactions between class and sex were identified for any of the parameters measured.

Table 5. Percent Body Fat Data from Phase I

Parameter	Sex	Class 1				Significant Difference <sup>2</sup> (Sex Effect)
		First-79	Second-80	Third-81	Fourth-82	
PFBV (%)	F	...	27.51±5.63	24.61±5.42	23.55±5.61	NSD
	M	18.30±5.35	18.02±5.58	17.20±5.25	18.98±6.58	
PFWD (%)	F	...	23.92±4.20	24.52±2.47	23.61±3.36	p < .001
	M	14.07±3.50	14.38±3.27	14.39±3.33	14.45±2.84	
PFWP (%)	F	...	22.44±3.98	22.90±2.35	22.33±3.26	p < .001
	M	11.60±3.20	11.94±3.16	12.18±3.18	12.31±2.72	

<sup>1</sup>The values represent mean ± standard deviation.

<sup>2</sup>NSD means p > .05. No significant differences among classes and no significant interactions between class and sex were identified for any body fat measurement.

Table 6. Comparison of Investigator Differences

Measurement	Sex	Investigator	Class <sup>1</sup>					Significant Difference <sup>2</sup>
			First-79	Second-80	Third-81	Fourth-82	Sex Investigator	
Triceps	F	LAIR	...	12.73±2.75	12.90±3.33	14.73±2.94	p < .001	
		USMA	...	12.03±3.10	13.65±2.69	14.53±2.94		
Subscapular	M	LAIR	7.64±3.26	7.40±1.98	7.70±2.75	7.94±2.16	NSD	
		USMA	7.47±3.59	7.36±2.09	8.48±3.10	7.76±1.98		
Biceps	F	LAIR	...	10.85±3.20	10.48±1.24	10.84±3.02	p < .001	
		USMA	...	10.24±3.73	8.70±1.96	10.15±3.32		
Lateral abdominal	M	LAIR	9.84±2.61	10.11±2.05	10.50±2.95	10.23±1.93	NSD	
		USMA	9.42±2.67	9.64±2.18	9.80±2.38	9.46±2.26		
Percent Fat	F	LAIR	...	6.22±2.70	5.22±1.53	6.72±2.16	p < .001	
		USMA	...	6.09±2.40	5.60±1.60	6.34±2.04		
Percent Fat	M	LAIR	4.25±1.28	3.71±0.66	4.11±1.12	4.06±1.08	p < .003	
		USMA	4.24±1.35	3.69±0.49	4.04±1.04	4.05±1.13		
Percent Fat	F	LAIR	...	7.87±3.93	7.38±1.52	8.42±3.20	p < .001	
		USMA	...	8.23±3.84	8.90±2.70	8.62±2.96		
Percent Fat	M	LAIR	6.89±3.30	6.53±1.65	6.76±2.20	7.05±2.47	p < .003	
		USMA	7.03±3.16	6.50±1.86	6.44±1.80	7.02±2.50		
Percent Fat	F	LAIR	...	22.33±4.00	21.96±2.42	23.16±3.46	p < .001	
		USMA	...	21.48±3.83	21.97±2.23	22.71±3.58		
Percent Fat	M	LAIR	11.71±3.43	11.62±2.02	12.03±2.84	12.40±2.65	p < .001	
		USMA	11.54±3.49	11.43±2.34	11.94±2.76	11.76±2.49		

<sup>1</sup>The values represent mean ± standard deviation.

<sup>2</sup>NSD means p > 0.05.

Table 7. Weight and Height Data from Phase II

Parameter Sex	Class <sup>1</sup>				Significant Difference <sup>2</sup> (Sex Effect)
	First-80	Second-81	Third-82	Fourth-83	
Weight (kg)					
F	60.88 ± 8.42	60.75 ± 7.45	59.06 ± 6.45	62.97 ± 6.46	p < .001
M	77.60 ± 10.90	74.01 ± 8.80	75.88 ± 7.92	79.70 ± 8.55	
Height (cm)					
F	165.37 ± 4.68	165.88 ± 7.11	164.85 ± 6.73	164.92 ± 4.56	p < .001
M	178.74 ± 6.07	175.19 ± 6.47	176.53 ± 7.72	179.81 ± 7.66	

<sup>1</sup>The values represent mean ± standard deviation.

<sup>2</sup>No significant differences among classes and no significant interactions between class and sex were identified for weight or height.

Table 8. Skinfold Data from Phase II

Parameter	Sex	Class <sup>1</sup>				Significant Difference <sup>2</sup> (Sex Effect)
		First-80	Second-81	Third-82	Fourth-83	
TCR (mm)	F	12.75±4.48	12.14±3.00	13.25±2.42	13.81±2.67	p <.001
	M	8.19±2.99	7.62±2.17	7.58±2.00	9.14±2.30	
TCL (mm)	F	13.07±4.23	12.64±3.48	14.33±3.34	14.56±2.50	p <.001
	M	8.13±2.84	7.79±2.44	7.56±2.00	9.14±2.55	
SSR (mm)	F	10.32±3.55	9.36±2.53	8.58±1.56	10.88±3.17	NSD
	M	10.17±2.88	10.12±2.41	10.21±2.24	9.50±2.06	
SSL (mm)	F	10.07±3.00	9.25±2.22	8.83±1.37	10.81±3.32	NSD
	M	10.21±2.61	10.10±2.08	10.14±2.27	9.09±2.31	
BCR (mm)	F	6.25±3.02	5.07±1.72	6.42±1.28	5.88±1.94	p <.001
	M	3.96±1.26	4.06±1.34	3.78±0.82	4.13±0.78	
BCL (mm)	F	6.14±3.16	5.21±1.89	6.08±1.43	6.00±1.79	p <.001
	M	4.52±1.63	4.02±1.16	3.83±0.80	4.32±0.98	
SIR (mm)	F	12.11±5.81	10.68±3.66	10.83±1.72	13.56±4.58	p <.036
	M	13.98±6.56	12.64±3.28	13.44±4.52	15.32±4.72	
SIL (mm)	F	11.89±6.06	10.93±3.69	11.00±1.26	12.62±3.96	p <.029
	M	14.31±6.20	12.48±3.47	13.32±4.41	14.68±4.67	

Table 8. (continued)

Parameter	Sex	Class				Significant Difference (Sex Effect)
		First-79	Second-80	Third-81	Fourth-82	
LAR (mm)	F	8.32±4.70	7.36±2.44	6.17±0.98	7.69±2.34	NSD
	M	7.83±3.83	7.21±2.84	6.89±2.25	7.23±1.40	
LAL (mm)	F	8.14±4.63	7.07±2.06	6.33±0.82	7.38±2.13	NSD
	M	7.58±3.80	7.02±2.70	6.62±2.14	7.18±1.54	
ABR (mm)	F	11.46±5.92	9.43±3.71	8.08±2.41	10.19±5.11	NSD
	M	10.00±4.61	9.50±3.16	8.24±3.03	8.86±1.87	
ABL (mm)	F	11.46±6.21	9.57±3.20	8.25±2.32	10.69±4.53	NSD
	M	9.58±4.61	9.08±3.48	7.97±3.02	8.32±1.74	
THR (mm)	F	21.07±6.53	18.61±4.28	21.67±3.67	21.38±5.29	p<.001
	M	11.03±4.56	9.83±3.36	10.39±2.32	10.27±2.95	
THL (mm)	F	20.93±6.59	18.21±4.78	21.42±3.20	20.38±5.37	p<.001
	M	11.44±4.51	10.42±3.04	10.73±2.40	10.32±2.86	

<sup>1</sup>The values represent mean ± standard deviation.

<sup>2</sup>No significant class effect and no significant interaction between class and sex were identified for any parameter.

Table 9. Percent Body Fat from Phase II

Parameter	Sex	Class <sup>1</sup>				Significant Difference (Sex Effect)
		First-80	Second-81	Third-82	Fourth-83	
PFWD (%)	F	23.41±4.78	22.51±3.07	23.50±1.35	24.64±3.85	p <.001
	M	14.56±3.93	14.12±2.48	14.29±2.58	15.21±2.57	
PFWP (%)	F	22.04±4.77	21.11±3.01	21.69±1.88	22.88±3.42	p <.001
	M	12.34±3.60	12.03±2.64	11.86±2.18	12.53±1.99	

<sup>1</sup>The vlaues represent mean ± standard deviation.

Table 10. Oxygen Uptake Data from Phase II

Parameter	Sex	Class <sup>1</sup>				Significant Difference (Sex Effect)
		First-80	Second-81	Third-82	Fourth-83	
V <sub>O2</sub> max (ml)	F	2473+656	2536+629	2534+ 315	2643+369	p <.001
	M	3909+739	3502+626	3873+1008	4139+762	
V <sub>O2</sub> max (ml/kg)	F	40.49+8.83	41.94+9.89	42.91+ 2.59	42.32+7.52	p <.002
	M	50.63+8.59	47.53+7.50	51.30+13.24	51.77+6.33	

<sup>1</sup>The values represent mean + standard deviation.

Table 11. Weight and Height Data from Subjects Completing both Phases I and II

Parameter	Sex	Phase	Class <sup>1</sup>			Significant Difference (Sex Effect) (Phase Effect)
			First-80	Second-81	Third-82	
Weight (kg)	F	I	58.56±6.61	60.18±5.32	55.19±3.09	p<.001
	F	II	59.94±8.08	62.41±5.62	55.34±2.88	
Weight (kg)	M	I	77.80±11.90	73.35±6.35	71.01±5.66	p<.002
	M	II	78.64±12.37	75.00±7.56	72.12±5.61	
Height (cm)	F	I	165.82±6.26	166.20±6.92	164.10±3.04	p<.001
	F	II	165.75±4.57	164.42±7.74	163.68±3.38	
Height (cm)	M	I	179.34±7.94	175.49±6.17	174.99±4.06	NSD
	M	II	179.08±7.83	175.28±6.22	175.11±4.55	

<sup>1</sup>The values represent mean ± standard deviation.

Table 12. Skinfold Data from Subjects Completing both Phases I and II

Parameter	Sex	Phase	Class <sup>1</sup>			Significant Difference <sup>2</sup> (Sex Effect)	Significant Difference <sup>3,4,5</sup> (Phase Effect)
			First-80	Second-81	Third-82		
TCR (mm)	F	I	13.62±3.04	13.87±2.87	12.38±2.40	p < .001	NSD
	F	II	13.50±4.54	13.50±3.00	11.62±2.56		
	M	I	7.86±2.56	8.27±3.65	8.62±2.58		
	M	II	7.42±3.19	7.96±3.08	7.79±1.79		
TCL (mm)	F	I	13.08±2.78	13.88±3.06	13.00±2.34	p < .001	NSD
	F	II	13.77±4.17	14.75±4.11	12.25±2.02		
	M	I	7.58±2.71	8.46±3.90	8.08±2.85		
	M	II	7.50±2.93	7.92±3.49	7.79±1.97		
SSR (mm)	F	I	10.08±3.16	12.12±2.25	8.87±1.75	NSD	p < .021
	F	II	9.85±3.07	12.00±3.16	8.00±1.63		
	M	I	10.05±2.49	10.92±3.52	10.00±2.47		
	M	II	10.08±2.46	10.15±3.29	9.71±1.75		
SSL (mm)	F	I	10.00±2.86	12.00±2.45	8.00±1.41	NSD	NSD
	F	II	9.69±2.76	11.50±2.65	8.75±1.50		
	M	I	10.45±2.04	10.19±2.83	9.79±2.07		
	M	II	10.08±2.24	9.88±2.39	9.41±1.58		

Table 12. (continued)

Parameter	Sex	Phase	Class			Significant Difference (Sex Effect)(Phase Effect)
			First-79	Second-80	Third-81	
BCR (mm)	F	I	5.69±2.49	5.50±0.91	5.25±1.19	P<.001
	F	II	6.19±3.05	6.00±1.82	5.38±1.49	
	M	I	3.83±1.23	4.00±1.11	3.92±1.00	NSD
	M	II	3.62±0.77	3.92±1.35	3.46±0.66	
BCL (mm)	F	II	6.04±2.52	5.62±1.25	5.00±0.82	p<.005
	F	II	6.15±3.06	6.38±2.56	5.12±1.31	
	M	I	4.25±1.64	4.35±1.33	4.12±1.17	NSD
	M	II	4.41±1.58	3.96±1.09	3.79±0.83	
SIR (mm)	F	I	11.50±4.06	12.12±1.31	9.12±1.75	NSD
	F	II	11.50±5.38	11.00±3.46	10.00±3.46	
	M	I	13.83±5.02	12.08±5.78	11.46±3.92	NSD
	M	II	14.25±7.05	13.11±6.00	13.42±3.99	
SIL (mm)	F	I	11.69±4.18	14.00±2.16	9.62±2.93	NSD
	F	II	11.42±5.33	12.75±2.63	10.50±2.64	
	M	I	14.74±5.56	12.04±5.70	11.79±3.42	NSD
	M	II	14.12±6.02	12.69±5.51	12.38±3.49	

Table 12. (continued)

Parameter	Sex	Phase	Class			Significant Difference (Sex Effect)	Significant Difference (Phase Effect)
			First-79	Second-80	Third-81		
LAR (mm)	F	I	7.42+3.78	8.50+1.73	6.12+0.85		NSD
	F	II	7.85+4.62	8.25+2.63	5.38+1.11		
	M	I	7.00+2.70	6.77+4.09	6.17+1.42		
	M	II	8.25+4.59	7.38+5.20	6.42+1.55		
LAL (mm)	F	I	7.23+2.72	8.88+2.59	5.75+0.50		NSD
	F	II	7.61+4.26	7.75+2.22	5.50+1.29		
	M	I	7.62+3.34	7.23+4.22	6.21+1.70		
	M	II	7.54+3.89	7.50+5.41	6.38+1.72		
ABR (mm)	F	I	10.19+3.58	15.62+1.38	7.87+0.85	P<.031	p<.017
	F	II	10.73+5.64	11.50+3.87	6.37+2.43		
	M	I	10.04+3.65	10.19+4.96	7.71+2.24		
	M	II	9.91+4.84	9.42+5.03	8.00+2.45		
ABL (mm)	F	I	10.46+3.83	15.87+2.10	8.50+1.29	p<.045	p<.005
	F	II	10.69+5.60	11.50+2.51	6.62+2.50		
	M	I	9.75+3.84	9.69+5.14	7.29+2.05		
	M	II	9.45+4.85	9.04+5.37	7.41+2.14		

Table 12. (continued)

Parameter	Sex	Phase	Class			Significant Difference (Sex Effect)(Phase Effect)
			First-79	Second-80	Third-81	
THR (mm)	F	I	21.58±5.73	20.50±2.08	20.25±3.40	p<.001
	F	II	21.31±6.73	20.25±3.68	18.00±4.69	
TAL (mm)	M	I	11.67±4.22	10.65±4.34	10.29±2.11	p<.024
	M	II	10.12±4.34	10.30±4.08	9.29±1.79	
TAL (mm)	F	I	21.54±5.59	20.62±1.89	21.12±3.42	p<.001
	F	II	20.61±6.75	19.75±3.30	18.62±3.90	
TAL (mm)	M	I	11.88±3.59	11.96±4.00	11.62±2.93	p<.001
	M	II	10.62±4.24	10.69±4.04	10.16±1.80	

<sup>1</sup>The values represent mean ± standard deviation.

<sup>2</sup>The phase by sex interaction (males increased and females decreased) was significant (p<.034).

<sup>3</sup>All interactions significant due to high values for sophomore females in phase I (phase by class: p<.022, phase by sex: p<.025, phase by class and by sex: p<.047).

<sup>4</sup>The phase by class interaction was significant (p<.016) due to high value for sophomore females in Phase I.

<sup>5</sup>NSD means p>.05.

Table 13. Percent Fat Estimates from Subjects  
Completing both Phases I and II

Estimate	Sex	Phase	Class			Significant Difference (Sex Effect)	
			First-80	Second-81	Third-82		
PFWP	F	I	21.56±3.25	23.06±2.03	20.16±1.57	p < .001	
		II	21.57±4.19	22.83±3.27	19.64±1.86		
	M	I	12.05±3.23	12.51±3.83	12.20±2.65		
		II	12.36±3.62	12.17±3.89	11.81±2.21		
PFWD	F	I	22.03±3.26	24.48±1.61	21.46±1.96		p < .001
		II	22.88±4.13	24.07±3.27	21.50±2.29		
	M	I	15.00±3.33	14.39±3.76	14.25±2.74		
		II	14.60±3.92	14.33±3.53	14.29±2.55		

The values represent mean ± standard deviation.

OFFICIAL DISTRIBUTION LIST

Commander US Army Medical Research and Development Command ATTN: SGRD-RMS/Mrs. Madigan Fort Detrick, Frederick MD 21701		Director Walter Reed Army Institute of Research Washington DC 20012
Defense Technical Information Center ATTN: DTIC-DDA Cameron Station Alexandria VA 22314	(12 copies)	Commander US Army Medical Research Institute of Infectious Diseases Fort Detrick, Frederick MD 21701
Director of Defense Research and Engineering ATTN: Assistant Director, Environmental and Life Sciences Washington DC 20301		Commander US Army Research Institute of Environmental Medicine Natick MA 01760
The Surgeon General ATTN: DASG-TLO Washington DC 20314		Commander US Army Institute of Surgical Research Brooke Army Medical Center Fort Sam Houston TX 78234
HQ DA (DASG-ZXA) WASH DC 20310		
Superintendent Academy of Health Sciences ATTN: AHS COM Fort Sam Houston TX 78234		Commander US Army Institute of Dental Research Washington DC 20012
Assistant Dean Institute and Research Support Uniformed Services University of Health Sciences 6917 Arlington Road Bethesda MD 20014		Commander US Army Medical Bioengineering Research and Development Laboratory Fort Detrick, Frederick MD 21701
Commander US Army Environmental Hygiene Agency Aberdeen Proving Ground MD 21070		Commander US Army Aeromedical Research Laboratory Fort Rucker AL 36362
US Army Research Office ATTN: Chemical and Biological Sciences Division P.O. Box 1221 Research Triangle Park NC 27709		Commander US Army Biomedical Laboratory Aberdeen Proving Ground Edgewood Arsenal MD 21010
Biological Sciences Division Office of Naval Research Arlington VA 22217		Commander Naval Medical Research Institute National Naval Medical Center Bethesda MD 20014
Director of Life Sciences USAF Office of Scientific Research (AFSC) Bolling AFB Washington DC 20332		Commander USAF School of Aerospace Medicine Aerospace Medical Division Brooks Air Force Base TX 78235

OFFICIAL DISTRIBUTION  
COOPERATING AGENCIES

Commander  
US NARADCOM  
Natick MA 01760

Commandant  
US Marine Corps  
HQ, US Marine Corps, LFS-4  
Washington DC 20389

Commander  
US NARADCOM  
ATTN: Director, Food Sci Lab  
Natick MA 01760

Commanding Officer  
Navy Food Service Systems Office  
Washington Navy Yard  
Washington DC 20374

Chairman, DOD Food Planning Board  
Director, Supply Management Policy  
OASD Manpower, Reserve Affairs/Log  
Pentagon, Room 3B730  
Washington DC 20301

Armed Forces Radiobiology Res Inst  
Building 42, NNMC  
Bethesda MD 20014

Chairman, Joint Formulation Board  
DOD Food RDT and ENG Program  
HQ, US Marine Corps LFS-4  
Washington DC 20380

Commander  
US NARADCOM  
ATTN: US Navy Rep/JTS  
Natick MA 01760

Commander  
US NARADCOM  
ATTN: US Army Rep/JTS  
Natick MA 01760

Commander  
HQDA, ODCSRDA  
ATTN: DAMA-CSS-D  
Washington DC 20310

Commander  
US NARADCOM  
ATTN: USMC REPL  
JTS, USMC DRDNA-ZF  
Natick MA 01760

Commander  
US NARADCOM  
ATTN: Chief, ORLSA  
Natick MA 01760

HQDA, ODCSLOG  
of Staff for Logistics  
ATTN: Chief, Troop Sup Div  
Washington DC 20310

Chairman, DOD Food SVC FAC  
and Equip Planning Board  
Office, Chief of Engineers  
Forrestal Building, Room 2-F-055  
Washington DC 20314

Commander  
US Army TSA  
ATTN: DALO-TAD  
Ft. Lee VA 23801

Defense Logistic Agency  
ATTN: DLA-LF, Room 3c325  
Cameron Station  
Alexandria VA 22314

Commander  
US NARADCOM  
ATTN: Technical Director  
Natick MA 01760

Commander  
US NARADCOM  
Dev Command  
ATTN: USAF Rep/JTS  
Natick MA 01760