

AD-A117 693

MATERIALS RESEARCH LABS ASCOT VALE (AUSTRALIA) F/8 6/16  
TWO NOVEL METHODS FOR THE PREPARATION OF ERYTHROCYTE STROMA FRE--ETC(U)  
NOV 81 H D CRONE, M POREYSKI, M P BLADEN  
MRL-TN-458


NL


UNCLASSIFIED

1-1  
1-1  
■



END  
DATE  
FILMED  
8-82  
DTIC





AD A 11 7 693

**DEPARTMENT OF DEFENCE  
DEFENCE SCIENCE AND TECHNOLOGY ORGANISATION  
MATERIALS RESEARCH LABORATORIES**

MELBOURNE, VICTORIA

**TECHNICAL NOTE**

**MRL-TN-458**

TWO NOVEL METHODS FOR THE PREPARATION OF ERYTHROCYTE  
STROMA FREE FROM HAEMOGLOBIN COMPARED TO THE CONVENTIONAL  
CENTRIFUGATION TECHNIQUE.

H.D. Crone, M. Poretski and M.P. Bladen

Approved for Public Release

DTIC FILE COPY



**DTIC  
ELECTE  
AUG 3 1982  
S D D**

82 48 48 050

DEPARTMENT OF DEFENCE  
MATERIALS RESEARCH LABORATORIES

TECHNICAL NOTE

MRL-TN-458

NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	



TWO NOVEL METHODS FOR THE PREPARATION OF ERYTHROCYTE  
STROMA FREE FROM HAEMOGLOBIN COMPARED TO THE CONVENTIONAL  
CENTRIFUGATION TECHNIQUE.

H.D. Crone, M. Poretski and M.P. Bladen

ABSTRACT

The removal of haemoglobin from haemolysed erythrocyte stroma by washing over a membrane filter or by passage through a column of porous gel was compared to the removal achieved by washing in the centrifuge. The column method was preferable for an initial rapid clean-up, which could be followed by a further pass through the column, or by centrifugation to concentrate the suspension. Membrane filtration was useful when total recovery of a small sample of stroma was required.

Approved for Public Release

© COMMONWEALTH OF AUSTRALIA 1981

---

POSTAL ADDRESS: Chief Superintendent, Materials Research Laboratories  
P.O. Box 60, Ascot Vale, Victoria 3032, Australia

---

## DOCUMENT CONTROL DATA SHEET

Security classification of this page:

UNCLASSIFIED

1. DOCUMENT NUMBERS:		2. SECURITY CLASSIFICATION:	
a. AR Number:	AR-002-743	a. Complete document:	UNCLASSIFIED
b. Series & Number:	MRL TECH NOTE	b. Title in isolation:	UNCLASSIFIED
c. Report Number:	MRL-TN-458	c. Abstract in isolation:	UNCLASSIFIED
3. TITLE: TWO NOVEL METHODS FOR THE PREPARATION OF ERYTHROCYTE STROMA FREE FROM HAEMOGLOBIN COMPARED TO THE CONVENTIONAL CENTRIFUGATION TECHNIQUE.			
4. PERSONAL AUTHOR(S):		5. DOCUMENT DATE:	
CRONE, H.D., PORETSKI, M. and BLADEN, M.P.		NOVEMBER, 1981	
6. TYPE OF REPORT & PERIOD COVERED:			
7. CORPORATE AUTHOR(S):		8. REFERENCE NUMBERS:	
Materials Research Laboratories		a. Task:	DST 78/025
		b. Sponsoring Agency:	
9. COST CODE: 321 250			
10. IMPRINT (Publishing establishment)		11. COMPUTER PROGRAMME(S):	
Materials Research Laboratories, P.O. Box 50, Ascot Vale, Vic 3032		(Title(s) and language(s)):	
NOVEMBER, 1981			
12. RELEASE LIMITATIONS (of the document):			
Approved for Public Release			
12-0. OVERSEAS: <input type="checkbox"/> N.O. <input type="checkbox"/> P.R. <input checked="" type="checkbox"/> 1 <input type="checkbox"/> A <input type="checkbox"/> B <input type="checkbox"/> C <input type="checkbox"/> D <input type="checkbox"/> E <input type="checkbox"/>			
13. ANNOUNCEMENT LIMITATIONS (of the information on this page):			
No Limitations on Distribution			
14. DESCRIPTORS:		CENTRIFUGATION	
ERYTHROCYTES	BLOOD		
MEMBRANES	HAEMOGLOBIN		
	HEMOLYSIS		
	MEMBRANE FILTRATION		
	GEL COLUMNS		
		15. COSATI CODES: 0601	
16. ABSTRACT (If title is classified, the announcement of this report will be similarly classified):			

The removal of haemoglobin from haemolysed erythrocyte stroma by washing over a membrane filter or by passage through a column of porous gel was compared to the removal achieved by washing in the centrifuge. The column method was preferable for an initial rapid clean-up, which could be followed by a further pass through the column, or by centrifugation to concentrate the suspension. Membrane filtration was useful when total recovery of a small sample of stroma was required.

C O N T E N T S

	<u>Page No.</u>
1. INTRODUCTION	1
2. METHODS	1
3. DISCUSSION	2
4. CONCLUSION	3
5. REFERENCES	4

TWO NOVEL METHODS FOR THE PREPARATION OF  
ERYTHROCYTE STROMA FREE FROM HAEMOGLOBIN COMPARED  
TO THE CONVENTIONAL CENTRIFUGATION TECHNIQUE

1. INTRODUCTION

The membrane of the mammalian erythrocyte is used widely as a model membrane for structural and functional investigations. The membranes are prepared by lysis of the erythrocytes in hypotonic media and then washed free of haemoglobin (Hb) by centrifugation at pH 7.4 in low ionic strength buffer solutions [1]. This method has been used in this laboratory for a number of years but since repeated washing in the centrifuge is a lengthy process, we have recently investigated other ways of removing the haemoglobin. We have found that the stroma is best prepared by rapid removal of the Hb immediately after haemolysis; prolonged contact of the membranes with the Hb solution encourages adsorption of the pigment. If the suspension is stored frozen with residual Hb present, a granular association of membranes and Hb is obtained on thawing.

This Note describes removal of the haemoglobin by a membrane filtration method and by a gel column method, and compares them with the centrifugation procedure.

2. METHODS

Freshly collected human blood was diluted into 150 mM NaCl, 10 mM Tris HCl, pH 7.4 and centrifuged to precipitate the cells. The latter were washed three times with the isotonic buffer, then haemolysed by suspension in an excess of 10 mM Tris HCl buffer at pH 7.4. The stroma was then washed free of haemoglobin by repeated centrifugation, or subjected to the two alternative procedures below. All operations were performed at 5°C.

The membrane filtration method consisted of stirring the haemolysate above a membrane of pore size sufficiently small to retain the membrane fragments but porous enough to allow rapid passage of the washing medium. Millipore type HA filters of 0.45  $\mu$ m pore size were cut to fit into an Amicon model 52 diafiltration cell. The cell was loaded with 40-50 mL of

haemolysate, stirred and subjected to light pressure ( $< 30$  KPa) by a peristaltic pump supplying air to the inlet. When the volume in the cell was reduced to 10 mL, 10 mM Tris buffer, pH 7.4, was pumped into the cell and washed through the filter at  $20 \text{ mL} \cdot \text{h}^{-1}$ , until the effluent was nearly colourless. Further concentration of the retentate was performed when necessary by pumping in air rather than buffer solution. The stroma from 6 mL of whole blood could be washed over a 40 mm diameter filter with 120 mL of buffer solution in 400 min, to reduce the Hb content to  $< 0.5\%$  of the initial value. No loss of stromal material, monitored as acetylcholinesterase (AChE) activity, was found. When desired, the washed suspension retained in the cell was solubilized by the addition of a suitable concentration of Triton X-100, and the effluent was then collected for further examination.

The column method of washing the stroma was based on the principle of passing the suspension through a bed of beaded, porous gel, so that the membrane particles passed rapidly between the beads and were eluted in the exclusion volume, but all the soluble material including the Hb entered the gel interior and was eluted later. A column of  $7 \text{ cm}^2$  cross-sectional area was packed with cross-linked agarose of bead diameter  $40\text{--}210 \mu\text{m}$  (Sephacrose CL-6B, Pharmacia Fine Chemicals) to a volume of 37 mL, using 10 mM Tris HCl pH 7.4 as the buffer solution. A sample of 10 mL of the haemolysate (equivalent to the cells from 2.5 mL of whole blood) was applied to the column, and eluted with the buffer at a flow rate of  $24 \text{ mL} \cdot \text{h}^{-1}$ . Fractions of 1.7 mL were collected and analysed for AChE [2], haemoglobin (absorbance at 412 nm) and turbidity (apparent absorbance at 700 nm). Figure 1 shows that the stroma (as represented by AChE activity and turbidity) was separated from the bulk of the Hb, and was eluted as a near square pulse not much wider than the initial 10 mL applied. From the results it can be shown that 92% of the AChE could be recovered with 11.5% of the Hb, or 60% AChE with 1% of Hb. Recovery was therefore dependent on the degree of separation desired. The stromal fraction could thus be washed in 70 min, and then concentrated by centrifugation or membrane filtration for further examination. A faster elution rate ( $50 \text{ mL} \cdot \text{h}^{-1}$ ) through the column resulted in distortion of the gel bed as the sample passed through. It was desirable to repack the column frequently, as the pressure difference occurring as the membrane fraction passed through tended to deform the gel beads and pack them more tightly.

### 3. DISCUSSION

The centrifugation method for washing the membranes has the advantages of simplicity of operation and of being concentrative. Its disadvantages are that a relatively long time is necessary (for example 4h to complete 4 washes), it is not particularly efficient, it encourages coagulation of the stroma, and the capital cost is high. Membrane filtration is very efficient, concentrative and of low capital cost, but it is relatively slow. Problems can arise due to clogging of the membrane and coagulation can also occur. The column method is rapid, of low cost, and no tendency towards coagulation of the stroma exists. On the other hand the sample is recovered slightly diluted and requires concentration by another method. Efficient separation is achieved at the expense of yield. However, the incompletely

separated Hb may be removed by a second pass of the stromal suspension through the column, resulting in a high yield of a clean suspension.

When the above factors are considered, the membrane filtration method is preferable if full recovery of a small sample is necessary (e.g. if the membranes have been labelled in some way). For general isolation of moderate quantities, the gel column procedure followed by centrifugation to concentrate the suspension is the most rapid and effective method. Our experience is that the gel column produces the best suspension, in terms of freedom from the coagulated, sticky lumps often formed by the other methods. Thus centrifugation, membrane filtration and freezing - thawing all concentrate the membrane fragments and encourage association.

Both the novel methods described can be scaled up to handle larger quantities of stroma. The critical design feature of the membrane method is to achieve efficient stirring as close as possible to the filter surface. For the column, a uniform, evenly - packed bed of gel is necessary together with a column design allowing even loading on the top and uniform drainage from the bottom.

It is recommended that researchers in membrane science should consider adapting these methods to their particular needs, as alternatives to the universally used but not always efficient centrifugation.

#### 4. CONCLUSION

Two methods are described for the removal of haemoglobin and other soluble compounds from erythrocyte membrane fragments, as alternatives to centrifugation. Continuous washing of the suspension held by a Millipore filter was efficient and allowed full recovery of the stroma, but was relatively slow (about the same duration as washing in the centrifuge). Separation on a column of beaded agarose gel (Sepharose CL-6B) was rapid and produced a homogenous suspension, but complete separation was not achieved in one passage down the column. The latter method is recommended as an alternative to centrifugation.

## 5. REFERENCES

1. Dodge, J.T., Mitchell, C. and Hanahan, D.J. (1963).  
Arch. Biochem. Biophys. 100, 119-130.
2. Ellman, G.L., Courtney, K.D., Andres, V. and Featherstone, R.M. (1961).  
Biochem, Pharmac. 7, 88-95




FIG. 1 - The separation of erythrocyte membrane particles from haemoglobin on a column of Sepharose CL-6B. Acetylcholinesterase (AChE) activity is shown on an arbitrary scale of activity on the left, Haemoglobin (Hb) concentration is expressed as the absorbance at 412 nm in a 10 mm cuvette and turbidity (T) is shown as apparent absorbance at 700 nm. For the latter parameter, the right hand scale must be reduced by a factor of 1/200.

(MRL-TN-458)

DISTRIBUTION LIST

MATERIALS RESEARCH LABORATORIES

Chief Superintendent  
Superintendent, Organic Chemistry Division  
Dr S.E. Freeman  
Library  
Dr H.C. Crone  
Mr M.P. Bladen  
Mr M. Poretski

DEPARTMENT OF DEFENCE

Chief Defence Scientist/Deputy Chief Defence Scientist/  
Controller, Projects and Analytical Studies 1 Copy  
Superintendent, Science and Technology Programme  
Controller, Service Laboratories and Trials  
Army Scientific Adviser  
Air Force Scientific Adviser  
Navy Scientific Adviser  
Librarian, (Through Officer-in-Charge), Materials Testing  
Laboratories, ALEXANDRIA, N.S.W.  
Senior Librarian, Aeronautical Research Laboratories  
Senior Librarian, Defence Research Centre Salisbury, S.A.  
Officer-in-Charge, Document Exchange Centre, (17 copies)  
Technical Reports Centre, Defence Central Library  
Central Office, Directorate of Quality Assurance -  
Air Force  
Deputy Director Scientific and Technical Intelligence,  
Joint Intelligence Organisation.  
Librarian, Bridges Library  
Defence Scientific and Technical Representative,  
Australia High Commission, London (Summary Sheets only)  
Counsellor Defence Science, (Attn: CONDS) (Summary Sheets only)

DEPARTMENT OF INDUSTRY AND COMMERCE

NASA Canberra Office, Woden, A.C.T.  
Head of Staff, British Defence Research and Supply  
Staff (Aust.)

(MRL-TN-458)

DISTRIBUTION LIST

(Continued)

OTHER FEDERAL AND STATE DEPARTMENTS AND INSTRUMENTALITIES

The Chief Librarian, Central Library, C.S.I.R.O.  
Library, Australian Atomic Energy Commission Research  
Establishment

MISCELLANEOUS - OVERSEAS

Reports Centre, Directorate of Materials Aviation, England  
Library - Exchange Desk, National of Bureau of Standards,  
U.S.A.  
UK/USA/CAN/NZ ABCA Armies Standardization Representative (4 copies)  
The Director, Defence Scientific Information and Documentation  
Centre, India  
Military, Naval and Air Adviser, High Commission of India,  
Canberra  
Director, Defence Research Centre, Kuala Lumpur, Malaysia  
Exchange Section, British Library, U.K.  
Periodicals Recording Section, Science Reference Library,  
British Library, U.K.  
Library, Chemical Abstracts Service  
INSPEC: Acquisition Section, Institution of Electrical  
Engineers, U.K.  
Engineering Societies Library, U.S.A.

ADDITIONAL DISTRIBUTION

Commander, U.S. Army Biomedical Laboratories, U.S.A.  
Directorate of Army Health Services, Canberra A.C.T.  
Director General of Air Force Health Services, Canberra A.C.T.  
Director General of Naval Health Services, Canberra A.C.T.  
Director General, Defence Research Establishment Suffield,  
Alberta, Canada.  
Director, Chemical Systems Laboratories, Maryland, U.S.A.  
Director, Microbiological Research Establishment, Wilts, England  
Prof. Sir S. Sunderland, C.M.G., Department of Experimental  
Neurology, Medical Centre, Parkville, Vic.  
The Librarian, Monash University, Clayton, Vic.  
The Librarian, La Trobe University, Bundoora, Vic.  
The Librarian, Barr Smith Library, University of Adelaide,  
Adelaide, S.A.  
Director, Chemical Defence Establishment, Wilts, England.