

AD-A119 721

OHIO STATE UNIV COLUMBUS DEPT OF CHEMISTRY
STUDIES ON THE SYNTHESSES OF BORON HYDRIDE SYSTEMS.(U)
JUL 82 S G SHORE

F/G 7/3

DAAG29-81-C-0029

ARO-15611.11-CH

NL

UNCLASSIFIED



END
DATE
FILMED
11-82
DTI

ARO 15611.11-CH

(12)

RF Project 762695/713823

Report No.

AD A119721

FINAL

REPORT

By

THE OHIO STATE UNIVERSITY
RESEARCH FOUNDATION

1314 KINNEAR RD.
COLUMBUS, OHIO 43212

To DEPARTMENT OF THE ARMY
U.S. Army Research Office
P.O. Box 12211
Research Triangle Park, North Carolina 27709
Contract No. DAAG29-81-C-0029

On STUDIES ON THE SYNTHESIS OF BORON HYDRIDE SYSTEMS

For the period July 1, 1981 - May 31, 1982

Submitted by Sheldon G. Shore
Department of Chemistry

Date July 30, 1982

DTIC
ELECTE
SEP 29 1982
S D H

CONFIDENTIAL STATEMENT
Approved for public release
Distribution Unlimited

DTIC FILE COPY

82 09 28 020

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
	AD-A119721	
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED
Studies on the Syntheses of Boron Hydride Systems		Final Report July 1, 1981-May 31, 1982
		6. PERFORMING ORG. REPORT NUMBER
		762695/713823
7. AUTHOR(s)		8. CONTRACT OR GRANT NUMBER(s)
Dr. Sheldon G. Shore		DAAG29-81-C-0029
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
The Ohio State University Research Foundation 1314 Kinnear Road, Columbus, Ohio 43214		
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE
U. S. Army Research Office Post Office Box 12211 Research Triangle Park, NC 27709		7-30-82
		13. NUMBER OF PAGES
		8
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)		
Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
NA		
18. SUPPLEMENTARY NOTES		
The view, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Boron hydride, $B_{10}H_{14}$, B_4H_{10} , 2- BrB_4H_9 , BH_3Cl^- , BrB_2H_5 , Carborane, nido-5,6- $R,R'C_2B_8H_{10}$		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
Syntheses of the new anion $[BH_3Cl]^-$, B_2H_6 , $B_{10}H_{14}$, and 2- BrB_4H_9 are discussed. The use of B_5H_9 as a starting point in the preparation of $B_9H_{13}L$, $B_{10}H_{12}L_2$ and nido-5,6- $RR'C_2B_8H_{10}$ is described.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Forward

This report covers the period July 1, 1981 - May 31, 1982. In the time frame cited above we have been concerned with several aspects of Studies on the Syntheses of Boron Hydride Systems. The work described here is an extension of the program which is discussed in an earlier final report: November 1981 No. 761119/711144 for Grant No. DAAG 29-78-G-0112.

In this report we describe the new anion $[\text{BH}_3\text{Cl}]^-$ and how it is employed to improve the yield of $\text{B}_{10}\text{H}_{14}$ in our "one pot" procedure for preparing $\text{B}_{10}\text{H}_{14}$ from B_5H_9 .

Dry syntheses of B_2H_6 and $\text{B}_2\text{H}_5\text{Br}$ are described and the preparation of $2\text{-BrB}_4\text{H}_9$ is considered in several solvents.

Our "one pot" syntheses of higher borane systems have been extended to include several thioether derivatives of the type $\text{B}_9\text{H}_{13}\text{L}$ and bis-ligand derivatives $\text{B}_{10}\text{H}_{12}(\text{P}\phi_3)_2$ and $\text{B}_{10}\text{H}_{12}(\text{SEt}_2)_2$. New preparations of carboranes of the type nido-5,6-R,R'C $_2\text{B}_8\text{H}_{10}$ are discussed.






Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

Results and Discussion

1. Preparation and Characterization of the $[\text{BH}_3\text{Cl}]^-$ Ion.

From tensimetric titrations we have demonstrated the formation of the $[\text{BH}_3\text{Cl}]^-$ ion.




Under similar reaction conditions there is no evidence for the formation of the analogous ions $[\text{BH}_3\text{Br}]^-$ and $[\text{BH}_3\text{I}]^-$.

The salts $[\text{PPN}][\text{BH}_3\text{Cl}]$, $[\text{N}(\text{n-C}_4\text{H}_9)_4][\text{BH}_3\text{Cl}]$, and $[\text{N}(\text{C}_2\text{H}_5)_4][\text{BH}_3\text{Cl}]$ readily disproportionate in CH_2Cl_2 solution at room temperature to give $[\text{B}_2\text{H}_7]^-$, $[\text{BH}_2\text{Cl}_2]^-$ and $[\text{BH}_4]^-$. However in the solid state these salts appear to be stable at room temperature for extended periods of time in the absence of air.

In collaboration with Dr. J. C. Huffman at the Indiana University Molecular Structure Center and Dr. Robert Bau of the University of Southern California, the crystal and molecular structure of $[\text{PPN}][\text{BH}_3\text{Cl}]$ has been determined. The $[\text{BH}_3\text{Cl}]^-$ ion is tetrahedral and the B-Cl distance is the longest observed (2.01Å). This of course suggests a weak bond and suggests that the anion might prove to be a ready source of BH_3 . This point is elaborated on below.



2. Addition of BH_3 to B_3H_7 and B_9H_{13} via the Anion $[\text{BH}_3\text{Cl}]^-$.

The reaction of $[\text{N}(\text{n-C}_4\text{H}_9)_4][\text{BH}_3\text{Cl}]$ with BCl_3 results in the abstraction of Cl^- to give 80-90% yields of B_2H_6 . This result suggested the possibility of using



$[\text{BH}_3\text{Cl}]^-$ as an external BH_3 source for polyhedral expansion in the absence of a solvent, rather than relying on BH_3 transfer as previously described in our earlier report¹ and publications.^{2,3} This method would eliminate the loss of boron in the polymeric residues suggested in the previously cited material.^{1,2,3} Reactions (5) and (6) below illustrate the stoichiometries used and the proposed

results for BH_3 addition to B_3H_7 and B_9H_{13} respectively.



In the case of reaction (4), the predicted yield of B_4H_{10} is not achieved, rather two independent reactions occur to generate B_4H_{10} and B_2H_6 as previously described.^{1,2,3} However, in reaction (5) the yield of $\text{B}_{10}\text{H}_{14}$ obtained is 64% based upon the stoichiometry of the reaction. This yield is nearly 71% if based solely on the boron in $[\text{B}_9\text{H}_{14}]^-$, thus indicating that reaction (5) appears to be proceeding as proposed. In our usual preparation of $\text{B}_{10}\text{H}_{14}$ from $\text{B}_9\text{H}_{14}^-$ the yield of $\text{B}_{10}\text{H}_{14}$ cannot exceed 50% of the boron in $\text{B}_9\text{H}_{14}^-$. Additionally, the quantity of B_2H_6 isolated is far less than predicted for the independent reaction of $[\text{BH}_3\text{Cl}]^-$ with BCl_3 . Further investigations into this system resulted in no further increases in yield of $\text{B}_{10}\text{H}_{14}$ when $[\text{N}(\text{n-C}_4\text{H}_9)_4][\text{BH}_4]$ was substituted for $[\text{N}(\text{n-C}_4\text{H}_9)_4][\text{BH}_3\text{Cl}]$. Thus the addition of BH_3 to B_9H_{13} is successful to a limited extent.

3. 2-BrB₄H₉ from [B₃H₇Br]⁻

The availability of the salt $[\text{N}(\text{n-C}_4\text{H}_9)_4][\text{B}_3\text{H}_7\text{Br}]^4$ suggested the possibility of preparing the 2-bromo derivative of tetraborane(10) through hydride ion abstraction from $[\text{B}_3\text{H}_7\text{Br}]^-$ followed by BH_3 transfer.

When $[\text{N}(\text{n-C}_4\text{H}_9)_4][\text{B}_3\text{H}_7\text{Br}]$ is reacted with BBr_3 in CH_2Cl_2 at -78° , 2-BrB₄H₉ and B₄H₁₀ are isolated in 15-20% yield each. Other boron containing species isolated include 2-BrB₅H₉, B₂H₆, B₃H₉, BrB₂H₃, and 1-BrB₅H₉ (1-10% yield each). The total amount of boron in the volatile products is 60-65% of boron in $[\text{B}_3\text{H}_7\text{Br}]^-$. Since both brominated and non-brominated boranes are obtained and since the solid reaction products contain both $[\text{BBr}_4]^-$ and $[\text{HBBR}_3]^-$ it appears that both hydride ion and bromide ion are susceptible to abstraction by BBr_3 . This would result in the formation of both "B₃H₇" and "B₃H₆Br" units which could undergo polyhedral expansion by transfer of either BH_3 or BH_2Br . Additionally, some decomposition must also occur to yield the B₂ and B₃ species cited above.

Various attempts were made to increase the yield of $2\text{-BrB}_4\text{H}_9$ with respect to the other boranes produced by varying the reaction conditions. Table I summarizes the yields of B_4H_{10} and $2\text{-BrB}_4\text{H}_9$ obtained and the reaction conditions employed for pertinent reactions. Of particular interest is the observation that B_4H_{10} production is directly proportional to the number of chlorine atoms in the solvent employed. This result can be rationalized from the standpoint that the solubility of the $[\text{BBr}_4]^-$ salt decreases as a more highly chlorinated solvent is employed. This method for preparing $2\text{-BrB}_4\text{H}_9$ provides a reasonable alternative to the only other reported procedure which involved the reaction of B_4H_{10} and Br_2 at -15° for 18 hours.⁵

COMPARATIVE YIELDS OF $2\text{-BrB}_4\text{H}_9$ AND B_4H_{10}

Solvent	Temperature	B_4H_{10}	$2\text{-BrB}_4\text{H}_9$
CH_2Cl_2	-78°C	15%	15%
CHCl_3	0°C	40%	Trace
CCl_4	0°C	43%	0%

Yield is calculated from boron in $\text{B}_3\text{H}_7\text{Br}^-$.

4. Convenient, Dry Syntheses of B_2H_6 and $\text{B}_2\text{H}_5\text{Br}$.

Diborane(6) was obtained in 95% yield from the reaction of a vigorously stirred mixture of NaBH_4 and BF_3 at room temperature and 1 atm. pressure.



This synthesis of B_2H_6 differs from the traditional⁵ preparation of B_2H_6 in that an ethereal solvent is not used. Furthermore, the generation of B_2H_6 probably occurs through hydride ion abstraction to give $[\text{HBF}_3]^-$ which disproportionates to $[\text{BH}_4]^-$ and $[\text{BF}_4]^-$.


Through the reaction of LiBH_4 with BBR_3 , we have developed a simple preparation of $\text{B}_2\text{H}_5\text{Br}$ in 80% yield. With vigorous stirring of reactants, this reaction occurs at room temperature. This procedure appears to be superior to the conventional⁶ method of preparing $\text{B}_2\text{H}_5\text{Br}$ which involves the reaction of

BBr_3 with B_2H_6 .


5. Preparation of $\text{B}_9\text{H}_{13}\text{L}$ and $\text{B}_{10}\text{H}_{12}\text{L}_2$ Complexes using B_5H_9 as a Starting Material.

Our final report for Grant No. DAAG 29-78-G-0112 described the use of B_5H_9 as a starting material for the preparation of $\text{B}_9\text{H}_{14}^-$ in 90% yield and its subsequent conversion to $\text{B}_9\text{H}_{13}\text{SEt}_2$ and $\text{B}_9\text{H}_{13}\text{P}\phi_3$ in effectively "one pot" synthetic procedures. We have extended these procedures to include the preparation of the complexes $\text{B}_9\text{H}_{13}\text{S}(\text{n-C}_4\text{H}_9)_2$, $\text{B}_9\text{H}_{13}\text{S}(\text{t-C}_4\text{H}_9)_2$, B_9H_{13} Tetrahydrothiophene, and B_6H_{13} Thiophene in yields of about 60%. Thus, we have established practical routes to adducts of the type $\text{B}_9\text{H}_{13}\text{L}$ using B_5H_9 as a starting material.

We have achieved a 65% conversion of $\text{B}_9\text{H}_{13}\text{P}\phi_3$ to high quality $\text{B}_{10}\text{H}_{13}(\text{P}\phi_3)_2$ through the following reaction






The analogous reaction shown below has also been observed giving 10-20% yield of $\text{B}_9\text{H}_{13}\text{SEt}_2$. Optimum conditions for conducting this reaction are currently



being studied on Grant No. DAAG-29-82-K-0112. By the same token current studies also involve conversion of $\text{B}_9\text{H}_{13}\text{S}(\text{n-C}_4\text{H}_9)_2$, $\text{B}_9\text{H}_{13}\text{S}(\text{t-C}_4\text{H}_9)_2$, B_9H_{13} Tetrahydrothiophene, and B_9H_{13} Thiophene to bis-ligand adducts: $\text{B}_{10}\text{H}_{12}\text{L}_2$. Our ultimate goal is to convert such materials to icosahedral carboranes.

6. Preparation of $\text{nido-5,6-R}_2\text{C}_2\text{B}_8\text{H}_{10}$ Carboranes using B_5H_9 as a Starting Material.

In our final report for Grant No. DAAG-29-78-G-0112 we described a "one-pot" synthesis of $\text{nido-5,6-}\phi_2\text{C}_2\text{B}_8\text{H}_{10}$ from B_5H_9 . The preparative procedure employed the formation of $\text{B}_9\text{H}_{13}\text{SEt}_2$ which was then reacted with $\phi\text{C}\equiv\text{C}\phi$ in order to form $\text{nido-5,6-}\phi_2\text{C}_2\text{B}_8\text{H}_{10}$. We have since made significant improvements in the synthetic procedure and find that it is not necessary to go through the $\text{B}_9\text{H}_{13}\text{SEt}_2$ intermediate. Three carboranes have been prepared using the procedure which is outlined below and we have obtained yields of up to 37% carborane based upon boron in the B_5H_9 starting material.



References

1. S. G. Shore, "Studies on the Syntheses of Boron Hydrides," Final Report November 1981, No. 761119/711144 for Grant No. DAAG-29-78-G-0112.
2. J. B. Leach, M. A. Toft, F. L. Himpsl, and S. G. Shore, J. Am. Chem. Soc., 1981, 103, 988.
3. M. A. Toft, J. B. Leach, F. L. Himpsl, and S. G. Shore, Inorg. Chem., 1982, 21, 1952.
4. G. E. Ryschkewitsch, and V. H. Miller, J. Am. Chem. Soc., 1975, 97, 6258.
5. J. Dobson and R. Schaeffer, Inorg. Chem., 1965, 4, 593.
6. H. I. Schlesinger, H. C. Brown, and A. B. Burg, J. Am. Chem. Soc., 1931, 53, 4321.

Publication

"New Systematic Syntheses of Boron Hydrides via Hydride Ion Abstraction Reactions:
Preparation of B_2H_6 , B_4H_{10} , B_5H_{11} , and $B_{10}H_{14}$ ", Inorg. Chem. 1982, 21, 1952.

Personnel

Dr. Sheldon G. Shore

Dr. Simon Boocock

Mark A. Toft

Steven Lawrence

Kenneth Gilbert

Michael Banks

ME
8