

AD-A120 010

DAVID W TAYLOR NAVAL SHIP RESEARCH AND DEVELOPMENT CE--ETC F/6 20/4
A VARIATIONAL PRINCIPLE ASSOCIATED WITH A LOCALIZED NUMERICAL S--ETC(U)
SEP 82 B YIM
DTNSRDC-82/077

NL


UNCLASSIFIED

Incl
AD-A
300 0



END
DATE
FILMED
DTIC

AD A120010



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER DTNSRDC-82/077	2. GOVT ACCESSION NO. AD-A120010	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A VARIATIONAL PRINCIPLE ASSOCIATED WITH A LOCALIZED NUMERICAL SOLUTION OF UNSTEADY FREE-SURFACE FLOWS		5. TYPE OF REPORT & PERIOD COVERED Final
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) B. Yim		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS David W. Taylor Naval Ship Research and Development Center Bethesda, Maryland 20084		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Program Element 61153N Task Area RR0140302 Work Unit 1542-018
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research (ONR-432) Arlington, Virginia		12. REPORT DATE September 1982
		13. NUMBER OF PAGES 22
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Variational Principle Numerical Ship Hydrodynamics Finite Element Technique Nonlinear Theory Free Surface Convolution Functional		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this report, a variational principle for unsteady body wave problems is treated both with and without a convolution integral and with both linear and nonlinear free-surface conditions. Functionals are obtained for the numerical computation of unsteady flow fields near a body that moves on or beneath the free surface. This formulation can be applied to ship hydrodynamic performance problems of water entry and body slamming, as well as to arbitrary body motion.		

DTIC
SELECTED
OCT 7 1982
H

DD FORM 1473 1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

TABLE OF CONTENTS

	Page
NOTATION.	iv
ABSTRACT.	1
ADMINISTRATIVE INFORMATION.	1
INTRODUCTION.	1
NONLINEAR PROBLEM	2
LINEAR PROBLEM.	5
USE OF CONVOLUTION.	7
SINUSOIDAL MOTION	10
ITERATIVE SCHEME.	10
CONCLUDING REMARKS.	12
REFERENCES.	13



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

NOTATION

D	Fluid domain
f_i	Function defined in Equation (23)
g	Acceleration of gravity
g_i	Function defined in Equation (24)
h	Body or free surface defined in Equation (1)
J	Functional
m	Source distribution
p	Pressure distribution
S_F	Free surface
S_J	Interface of near and far fields
S_S	Body surface
t	Time
x, y, z	Rectangular Cartesian coordinates
δ	Variation
ρ	Water density
τ	Time
ϕ	Potential
ψ	Function defined in Equation (21)
ω	Frequency defined in Equation (21)
Subscript	
i	The i th order
n	Normal derivative toward fluid
o	Projection on the $z = 0$ surface
t	The time derivative

- z The z derivative
- 1 Near field, the first order
- 2 Far field, the second order

ABSTRACT

In this report, a variational principle for unsteady body wave problems is treated both with and without a convolution integral, and with both linear and nonlinear free surface conditions. Functionals are obtained for the numerical computation of unsteady flow fields near a body that moves on or beneath the free surface. This formulation can be applied to ship hydrodynamic performance problems of water entry and body slamming, as well as to arbitrary body motion.

ADMINISTRATIVE INFORMATION

The work reported herein has been supported by the Numerical Naval Hydrodynamics Program at the David W. Taylor Naval Ship Research and Development Center. This program is jointly sponsored by the Office of Naval Research and DTNSRDC under Task Area RR0140302, Work Unit 1542-018.

INTRODUCTION

In the early 1970's the David W. Taylor Naval Ship Research and Development Center (DTNSRDC) recognized the demand for advanced numerical methods to predict the hydrodynamic performance characteristics of naval ships, particularly when classical methods proved inadequate.

Thus, in 1974 the Numerical Naval Ship Hydrodynamics Program was begun at DTNSRDC. Under this program the author previously investigated the steady ship-wave problem using a variational principle associated with a localized finite-element technique.^{1*} This method is useful particularly to analyze the flow field near the ship in detail; in the far field, the Michell approximation can be used. This report extends the problem to the unsteady case.

For both the steady and unsteady problems, the simple calculation is for a linear free surface condition with exact body boundary conditions. An iterative method is needed for a nonlinear free surface condition. However, in the unsteady problem, the variational principle requires an integration with respect to time using the initial conditions; in this instance, a convolution integral is useful. The variational principle for the unsteady body wave problem with exact body boundary conditions is treated both with and without convolution, and with both linear and

*A complete listing of references is given on page 13.

nonlinear free surface conditions. For the linear free surface problem, a functional for the variational principle is obtained with a convolution rather than a general integral. The convolution cannot be applied to a nonlinear free surface problem; the condition is required for large values of time. A nonlinear solution derived using an iteration scheme having the linear convolution form is also discussed. The time integration can be eliminated if the motion is sinusoidal.

This formulation can be applied to problems of water entry and body slamming, as well as to arbitrary body motion.

NONLINEAR PROBLEM

Since problems dealt with here can be generalized easily to three-dimensions, for simplicity we first consider a two-dimensional problem in the rectangular Cartesian (x,z) coordinate* plane. When a body whose surface is represented by

$$S_s [z=h(x,t)] \quad (1)$$

enters the water surface S_F ($z = 0, t \leq 0$; $z = h(x,t), t > 0$) at time $t = 0$, or when a semisubmerged or fully submerged body starts to move at time $t = 0$ and either exits the water or stops moving at $t = t_1$, then the boundary conditions² for a velocity potential ϕ are as follows:

$$\phi = \phi_t = 0 \text{ for } t \leq 0 \text{ everywhere}$$

$$\frac{1}{2} (\nabla\phi)^2 - \phi_t + gh = 0 \quad \text{on } S_F \quad (2)$$

$$\left. \begin{array}{l} \text{or} \\ h_t - \nabla\phi \nabla(h-z) = 0 \\ h_t - \phi_n \sqrt{h_x^2 + 1} = 0 \end{array} \right\} \text{ on } S_F \text{ and } S_s(t) \quad (3)$$

Here, $S_s(t)$ is the submerged body surface varying with time t , and n is the normal direction into the fluid. We consider potentials ϕ_1 in the domain D_1 and ϕ_2 in the

*Definition of notations are given on page iv.

domain D_2 , where D_1 is the near field including S_s , and D_2 is outside of D_1 . Then at the interface S_J of D_1 and D_2 , we need to have

$$\phi_1 = \phi_2 \quad (4)$$

$$\phi_{1n} = -\phi_{2n}$$

The outer potential ϕ_2 in D_2 is assumed to satisfy the linear free surface condition

$$\phi_{2tt} + g\phi_{2z} = 0 \quad (5)$$

For such ϕ_2 we know the time-dependent Green function.^{3,4}

Now we will construct a Lagrangian for the previously described problem, considering the Lagrangian that Luke⁵ used

$$J = \int_0^t \iint_{D_1} \left(\frac{1}{2} \nabla \phi_1 \cdot \nabla \phi_1 - \phi_{1t} \right) dz dx dt + \int_0^t \int_{S_{1F}} g \frac{h^2}{2} dx dt - \int_0^t \int_{S_J} \left(\phi_1 - \frac{1}{2} \phi_2 \right) \phi_{2n} dz dt \quad (6)$$

where ϕ_1 , ϕ_2 and h vary with time, and τ is a sufficiently large time after the body has either exited from or come to rest in the water so that we can safely assume that the variation $\delta\phi_1$, $\delta\phi_2$, and $\delta\phi_{2t}$ vanishes at $t = \tau$. It will be shown later that the use of a convolution integral necessitates only the initial condition without the condition at $t = \tau$.

Since

$$\frac{\partial}{\partial t} \int_0^{h(x,t)} \phi dz = \int_0^{h(x,t)} \phi_t ds + \phi(z=h) h_t \quad (7)$$

we have

$$\int_0^\tau \int dx \int_0^{h(x,t)} \phi_t dz dt = \int_0^\tau \int \left[\frac{\partial}{\partial t} \int_0^{h(x,t)} \phi dz - \phi(z=h) h_t \right] dx dt$$

$$= \int \left[\int_0^{h(x,t)} \phi dz \right]_0^\tau - \int_0^\tau dt \phi(z=h) h_t dx$$

or

$$\int_0^\tau \iint_{D_1(t)} \phi_t dz dx dt = - \int_0^\tau \int_{S_S US_F} h_t \phi dx dt + \iint_{D_1(t=\tau)} \phi dx dz \quad (8)$$

Now we take a variation of J in Equation (6), and use the Green theorem, and the condition at large $t = \tau$ obtaining¹

$$\delta J = \int_0^\tau \int_{S_{1F}} \left(\frac{1}{2} \nabla \phi_1 \nabla \phi_1 - \phi_{1t} + gh \right) \delta h dx dt$$

$$- \int_0^\tau \iint_{D_1} \nabla^2 \phi_1 \delta \phi_1 dz dx dt - \int_0^\tau \int_{S_{1F} US_S} \left(\phi_{1n} \sqrt{h_x^2 + 1} - h_t \right) \delta \phi_1 dx dt$$

$$- \int_0^\tau \int_{S_J} (\phi_{1n} + \phi_{2n}) \delta \phi_1 dz dt - \int_0^\tau \int_{S_J} (\phi_1 - \phi_2) \delta \phi_{2n} dz dt$$

$$+ \frac{1}{2} \int_0^\tau \int_{S_J} (\delta \phi_2 \phi_{2n} - \phi_2 \delta \phi_{2n}) dz dt = 0 \quad (9)$$

However from the identity,

$$\iint_{D_2} \{ \nabla^2 \phi_2 \delta \phi_2 - \phi_2 \nabla^2 (\delta \phi_2) \} dx dz = \int_{S_J \cup S_{2F}} \{ \phi_{2n} \delta \phi_2 - \phi_2 (\delta \phi_{2n}) \} ds = 0$$

and from the linear free surface condition on S_{2F} , Equation (5), and the condition at $t = \tau$, we have

$$\begin{aligned} \int_0^\tau \int_{S_J} (\delta \phi_2 \phi_{2n} - \phi_2 \delta \phi_{2n}) dz dt &= - \int_0^\tau \int_{S_{2F}} (\delta \phi_2 \phi_{2n} - \phi_2 \delta \phi_{2n}) dx dt \\ &= \frac{1}{g} \int_{S_{2F}} \int_0^\tau (\delta \phi_2 \phi_{2tt} - \phi_2 \delta \phi_{2tt}) dt dx = \frac{1}{g} \int_{S_{2F}} (\delta \phi_2 \phi_{2t} - \phi_2 \delta \phi_{2t})_{t=\tau} dx = 0 \end{aligned}$$

so that the last integral of Equation (9) vanishes. Since $\delta \phi_1$, $\delta \phi_2$, $\delta \phi_{2n}$, and δh are arbitrary, we obtain from Equations (9) and (10) the corresponding time-dependent, free surface boundary value problem represented by Equations (2)-(4) together with the Laplace equation for ϕ . Therefore, solving the variational problem with the functional J of Equation (6) is equivalent to solving the Laplace equation with the boundary conditions as set forth in Equations (2)-(4), provided we assume that

$$\delta \phi_1 = \delta \phi_2 = \delta \phi_{2t} = 0 \quad \text{at } t = \tau$$

LINEAR PROBLEM

If we assume a linear free surface condition in both S_{1F} and S_{2F} and keep the exact boundary condition [Equation (3)] on the body surface,

$$\phi_n = h_t / \sqrt{h_x^2 + 1} \quad (11)$$

then we may use

$$\begin{aligned}
J = & \int_0^T \iint_{D_1} \frac{1}{2} \nabla \phi_1 \nabla \phi_1 \, dz dx dt - \frac{1}{2g} \int_0^T \int_{S_{Fo}} \phi_1 \phi_{1tt} \, dx dt \\
& + \int_0^T \int_{S_s} h_t \phi_1 \, dx dt - \int_0^T \int_{S_J} \left(\phi_1 - \frac{1}{2} \phi_2 \right) \phi_{2n} \, ds dt \quad (12)
\end{aligned}$$

where S_{Fo} is the projection of S_F on $z = 0$. When we take the variation of J , we have

$$\begin{aligned}
\delta J = & - \int_0^T \iint_{D_1} \nabla^2 \phi_1 \delta \phi_1 \, dz dx dt - \int_0^T \int_{S_{1Fo}} \delta \phi_1 \left(\phi_{1n} + \frac{1}{g} \phi_{1tt} \right) \, dx dt \\
& - \int_0^T \int_{S_s} \left(\delta \phi_1 \phi_{1n} \sqrt{h_x^2 + 1} - h_t \right) \, dx dt \\
& - \int_0^T \int_{S_J} (\phi_1 - \phi_2) \delta \phi_{2n} \, dz dt - \int_0^T \int_{S_J} (\phi_{1n} + \phi_{2n}) \delta \phi_1 \, dz dt \quad (12a)
\end{aligned}$$

Here, in addition to Equation (10), the following equation holds

$$\begin{aligned}
\delta \int_0^T \int_{S_{Fo}} \phi_1 \phi_{1tt} \, dx dt &= \int_0^T \int_{S_{Fo}} \delta \phi_1 \phi_{1tt} \, dx dt + \int_0^T \int_{S_{Fo}} \phi_1 \delta \phi_{1tt} \, dx dt \\
&= 2 \int_0^T \int_{S_{Fo}} \delta \phi_1 \phi_{1tt} \, dx dt + \int_{S_{Fo}} (\phi_1 \delta \phi_{1t} - \phi_{1t} \delta \phi_1)_{t=T} \, dx = 2 \int_0^T \int_{S_{Fo}} \delta \phi_1 \phi_{1tt} \, dx dt \quad (13)
\end{aligned}$$

Thus, as in the previous section, we can easily derive the corresponding linear free surface boundary value problem using Equation (5) with the exact body boundary condition in Equation (11).

USE OF CONVOLUTION

Equations (6) and (12) are Lagrangians in a time-dependent, two-dimensional space with nonlinear and linear free surface conditions, respectively. They could be localized in space but not in time. Namely, we had to specify the conditions on S_F at $t = 0$ and $t = \tau$ with a sufficiently large τ . On the other hand, we did not require the initial condition $\phi_t = 0$ at $t = 0$. In addition, such time τ when $\phi_1 = 0$ on S_F may be too large³ for practical use. For linear free surface boundary conditions, we can treat our variational problem in the same way as those who have treated variation principles for linear initial value problems^{6,7} using convolutions defined by

$$\phi_1 * \phi_2 \equiv \int_0^\tau \phi_1(x, z, t) \phi_2(x, z; \tau - t) dt \quad (14)$$

$$\nabla \phi_1 * \nabla \phi_2 \equiv \frac{\partial \phi_1}{\partial x} * \frac{\partial \phi_2}{\partial x} + \frac{\partial \phi_1}{\partial z} * \frac{\partial \phi_2}{\partial z}$$

We change Equation (12) to

$$\begin{aligned} J = & \iint_{D_1} \frac{1}{2} \nabla \phi_1 * \nabla \phi_1 \, dz dx - \frac{1}{2g} \int_{S_{Fo}} \phi_1 * \phi_{1tt} \, dx \\ & + \int_{S_s} h_t * \phi_1 \, dx - \int_{S_J} \left(\phi_1 - \frac{1}{2} \phi_2 \right) * \phi_{2n} \, dz \end{aligned} \quad (15)$$

If we use the identity relation

$$\phi_1 * \phi_2 = \phi_2 * \phi_1$$

and the initial condition $\phi_1 = \phi_{1t} = 0$ at $t = 0$, instead of the conditions $\phi_1 = 0$ on S_F at $t = 0$ and $t = \tau$ with large τ used in the previous section, we obtain, for any time, τ

$$\begin{aligned}
 \int_{S_{Fo}} \phi_1 * \phi_{1tt} dx &= \int_0^\tau \int_{S_{Fo}} \delta\phi_1(x,t) \phi_{1tt}(x,\tau-t) dx dt + \int_0^\tau \int_{S_{Fo}} \phi_1(x,t) \delta\phi_{1tt}(x,\tau-t) dx dt \\
 &= 2 \int_0^\tau \int_{S_{Fo}} \delta\phi_1(x,t) \phi_{1tt}(x,\tau-t) dx dt \\
 &\quad + \int_{S_{Fo}} \{(\phi_1(x,t) \delta\phi_{1t}(x,t-\tau) - \phi_{1t}(x,t) \delta\phi_1(x,t-\tau))\}_{t=0}^\tau dx \\
 &= 2 \int_0^\tau \int_{S_{Fo}} \delta\phi_1(x,t) \phi_{1tt}(x,\tau-t) dx dt \\
 &= 2 \int_{S_{Fo}} \delta\phi_1 * \phi_{1tt} dx \tag{16}
 \end{aligned}$$

where τ need not be large, and x represents a point on the free surface $z = 0$. Thus, we obtain as in Equation (12a)

$$\begin{aligned}
 \delta J &= - \iint_{D_1} \nabla^2 \phi_1 * \delta\phi_1 dz dx - \int_{S_{1Fo}} \delta\phi_1 * \left(\phi_{1n} + \frac{1}{g} \phi_{1tt} \right) dx \\
 &\quad - \int_{S_s} \delta\phi_1 * \left(\phi_{1n} \sqrt{h_x^2 + 1} - h_t \right) dx \\
 &\quad - \int_{S_J} (\phi_1 - \phi_2) * \delta\phi_{2n} dz - \int_{S_J} (\phi_{1n} + \phi_{2n}) * \delta\phi_1 dz \tag{17}
 \end{aligned}$$

To obtain Equation (17), we used a convolution expression of Equation (16), where we can use the initial condition $\phi_1 = \phi_{1t} = 0$.

Since $\delta\phi_1$, $\delta\phi_2$, and $\delta\phi_{2n}$ are arbitrary, we obtain the corresponding equations for a time-dependent linear free surface boundary value problem that has a unique solution.³

If we lift out S_j so that D_1 occupies the entire fluid domain, then the last integral of Equation (15) disappears. The resulting equation appears much simpler than that obtained by Murray⁷ due to a simple difference in the treatment of the free surface condition. Equation (15) does not give the wave height as a natural boundary condition, whereas Murray's corresponding equation does. However, from $h = \phi_t/g$ the wave height can be obtained.

If we consider eigen solutions that satisfy only the Laplace equation; the linear free surface condition, Equation (5); and the radiation condition in $D_1 \cup D_2$, then ϕ , which satisfies the body boundary condition in Equation (11), can be derived from Equations (15) and (17) by using

$$J = \int_{S_s} \left(\frac{1}{2} \phi_n \sqrt{h_x^2 + 1} - h_t \right) * \phi dx \quad (18)$$

When we know a functional whose minimum value is attained by the solution, we can find the solution numerically by such methods as the finite-element technique⁸ or singularity method.⁹

For example

$$\phi = \sum_{i=1}^N m_i \phi_i \quad (19)$$

where ϕ_i is the Green function, which is available for this problem for a source distribution on the body surface. The source distribution m_i will be obtained from solution of the simultaneous equations,

$$\frac{\partial J}{\partial m_i} = 0 \quad , \quad i=1,2,\dots,N \quad (20)$$

SINUSOIDAL MOTION

If we consider a sinusoidal ship oscillation such as $h_t = f e^{i\omega t}$ for S_s of Equation (1), we substitute

$$\phi_1 = \psi_1(x, z) e^{i\omega t} \quad (21)$$

into Equation (12), integrate with respect to t , and obtain

$$\begin{aligned} e^{-i\omega t} J = J_1 = & \iint_{D_1} \frac{1}{2} \nabla \psi_1 \nabla \psi_1 \, dz dx - \frac{\omega^2}{2g} \int_{S_F} \psi_1^2 \, dx \\ & + \int_{S_s} f \psi_1 \, dx - \int_{S_J} \left(\psi_1 - \frac{1}{2} \psi_2 \right) \psi_{2n} \, dz \end{aligned} \quad (22)$$

This is exactly the same Lagrangian that Bai and Yeung⁸ used. Working from Equation (18), we can apply eigen solutions to the whole field by using

$$J = \int_{S_s} \left(\frac{1}{2} \psi_r - F \right) \psi ds$$

where

$$h_t / \sqrt{h_x^2 + 1} = F e^{i\omega t}$$

A similar functional was used by Sao et al.⁹ to solve the problem of a heaving oscillation of a dock.

ITERATIVE SCHEME

For problems with the linear free surface condition, we can completely localize the numerical scheme in D_1 with $0 \leq t < \tau$ for any τ with the help of the convolution form. However, for nonlinear problems, the finite-element technique has to rely on an iteration. We may thus use an iterative free surface condition² on $z = 0$

$$\phi_{itt} + g \phi_{iz} + f_i = 0 \quad (23)$$

$$gh_i - \phi_{it} + g_i = 0 \quad (24)$$

where, for the first order perturbation solution, $f_1 = 0$, $g_1 = 0$ and, for the n th order, f_n and g_n are known functions of ϕ_i of the $(n-1)$ th or the lower order solutions.

Then the Lagrangian for each f_i is

$$\begin{aligned} J_i = & \iint_{D_1} \frac{1}{2} \nabla \phi_{1i} * \nabla \phi_{1i} \, dz dx - \frac{1}{2g} \int_{(z=0) \cap D_1} \phi_{1i} * \phi_{1itt} \, dx \\ & - \int_{(z=0) \cap D_1} \phi_{1i} * f_i \, dx + \int h_t * \phi_{1i} \, dx \\ & - \int_{S_J} \left(\phi_{1i} - \frac{1}{2} \phi_{2i} \right) * \phi_{2in} \, dz \\ \delta J_i = & 0 \end{aligned} \quad (25)$$

where the solutions for $l = 1, 2, \dots, n-1$ should be used to determine the solution when $i = n$. Equation (24) gives the wave height h for each i .

If we specify a time-dependent, free surface pressure distribution p on the projection S_{so} of S_s to $z = 0$ instead of h_t in S_s , we may use

$$f_1 = g_1 = p/\rho \quad \text{on } S_{so}$$

in Equations (23) and (24) for the first order, where ρ is the water density.

Although we have discussed the two-dimensional time-dependent problem, Equations (19) through (24) can be extended to the three-dimensional time-dependent problem.

That is

$$\begin{aligned}
 J = & \iiint_{D_1} \frac{1}{2} \nabla \phi_1 * \nabla \phi_1 \, dx dy dz - \frac{1}{2g} \iint_{(z=0) \cap D_1} \phi_1 * \phi_{1tt} \, dx dy \\
 & - \iint_{(z=0) \cap D_1} \phi_1 * f_i \, dx dy + \iint h_t * \phi_1 \, dx dy \\
 & - \iint_{S_J} \left(\phi_1 - \frac{1}{2} \phi_2 \right) * \phi_{2n} \, ds
 \end{aligned}$$

and so on.

CONCLUDING REMARKS

We have formed functionals with both linear and nonlinear free surface boundary conditions. For the former but not the latter case, we could apply a convolution integral. However, the body boundary condition is satisfied exactly in both cases. In many cases, the flow field near an arbitrary body is of interest, and eigen solutions with linear free surface conditions are known. Even in any large unsteady motion such as ship slamming, the free surface condition for a short period in the beginning may be linear, then the convolution may be applied in the early time period. Especially in the slamming problem, the peak pressure is known to be reached early in the beginning and estimation of the early pressure distribution on the slamming body is required. With this functional, we can find the solution for an arbitrary body numerically by such methods as the finite element technique or singularity method. Thus, a wide application of such functionals can be expected.

REFERENCES

1. Yim, B., "A Variational Principle Associated with a Localized Finite-Element Technique for Steady Ship-Wave and Cavity Problems," Proceedings of the First International Conference on Numerical Ship Hydrodynamics, David Taylor Naval Ship R&D Center (1976).
2. Wehausen, J.V. and E.V. Laitone, "Surface Waves," Encyclopedia of Physics, Springer-Verlag, Berlin, Vol. IX, pp. 446-778 (1960).
3. Stoker, J.J., "Water Waves," Interscience Publishers, Inc., New York (1957), pp. 187-196.
4. Yim, B., "Investigation of Gravity and Ventilation Effects in Water Entry of Thin Foils," Proceedings of the International Union of Theoretical and Applied Mechanics Symposium held in Leningrad (1971), NAUK Publishing House, Moscow, pp. 471-475 (1973).
5. Luke, J.C., "A Variational Principle for a Fluid with a Free Surface," Journal of Fluid Mechanics, Vol. 27, Part 2, pp. 395-397 (1967).
6. Gurtin, M.E., "Variational Principles for Linear Initial-Value Problems," Quarterly Journal of Applied Mathematics, Vol. 22, pp. 252-256 (1964).
7. Murray, J.C., "A Note on Some Variational Principles for a Class of Linear Initial-Boundary Value Problems," Journal of the Institute of Mathematics and Applications, Vol. 12, pp. 119-123 (1973).
8. Bai, K.J. and R. Yeung, "Numerical Solutions of Free-Surface Flow Problems," 10th Symposium on Naval Hydrodynamics, Office of Naval Research (1974).
9. Sao, K. et al., "On the Heaving Oscillation of a Circular Dock," Journal of the Society of Naval Architects of Japan, Vol. 130, pp. 121-131 (1971).

INITIAL DISTRIBUTION

Copies		Copies	
1	CHONR/438 Lee	1	MARAD/Lib
2	NRL	4	U. of Cal/Dept Naval
	1 Code 2027		Arch, Berkeley
	1 Code 2627		1 Eng Library
4	USNA		1 Webster
	1 Tech Lib		1 Pauling
	1 Nav Sys Eng Dept		1 Wehausen
	1 Bhattacheryya	2	U. of Cal, San Diego
	1 Calisal		1 A.T. Ellis
2	NAVPGSCOL		1 Scripps Inst Lib
	1 Library	2	CIT
	1 Garrison		1 Aero Lib
1	NELC/Lib		1 T.Y. Wu
2	NOSC	1	Catholic U. of Amer/
	1 Library		Civil & Mech Eng
	1 Higdon	1	Colorado State U./Eng Res Cen
1	NADC	1	Florida Atlantic U.
1	NCEL/Code 131		1 Tech Lib
13	NAVSEA	1	U. of Hawaii/St. Denis
	1 SEA 031, R. Johnson	1	U. of Illinois/J. Robertson
	1 SEA 031, G. Kerr	2	U. of Iowa
	1 SEA 031, C. Kennel		1 Library
	1 SEA 05R, L. Benen		1 Landweber
	1 SEA 05R, Dilts	1	U. of Kansas/Civil Eng Lib
	1 SEA 05R, N. Kobitz	1	Lehigh U./Fritz Eng Lab Lib
	1 SEA 05R, J. Schuler	6	MIT
	1 SEA 312, P.A. Gale		1 Yeung
	1 SEA 312, J.W. Kehoe		1 Mandel
	1 SEA 55W, E.N. Comstock		1 Abkowitz
	1 SEA 55W, R.G. Keane, Jr.		1 Newman
	1 SEA 61433, F. Prout		1 Slavounos
	1 PMS 383, Chatterton		1 Ogilvie
12	DTIC	1	U. of Mich/NAME
1	NSF/Engineering Lib		1 Library
1	DOT/Lib, TAD-491.1		
1	NBS/Klebanoff		

Copies

1 U. of Notre Dame
1 Eng Lib

2 New York U./Courant Inst
1 A. Peters
1 J. Stoker

4 SIT
1 Breslin
1 Savitsky
1 Dalzell
1 Kim

1 U. of Texas/Arl Lib

2 Southwest Res Inst
1 Applied Mech Rev
1 Abramson

1 Stanford Res Inst/Lib

2 U. of Washington
1 Eng Lib
1 Mech Eng/Adee

3 Webb Inst
1 Library
1 Lewis
1 Ward

1 Woods Hole/Ocean Eng

1 SNAME/Tech Lib

1 Bethlehem Steel/Sparrows Point

1 Bethlehem Steel/New York/Lib

1 Exxon, NY/Design Div, Tank Dept

1 General Dynamics, EB/Boatwright

1 Gibbs & Cox/Tech Info

1 Hydronautics/Lib

1 Lockheed, Sunnyvale
1 Potash

Copies

1 Newport News Shipbuilding/Lib

1 Oceanics

1 Sperry Rand/Tech Lib

1 Sun Shipbuilding/Chief Naval Arch

2 American Bureau of Shipping
1 Lib
1 Cheng

1 Maritime Research Information
Service

CENTER DISTRIBUTION

Copies	Code	Name
1	012	D. Jewell
1	1170	R.M. Stevens
1	1170	G.R. Lamb
1	1170	S. Hawkins
1	1500	W.B. Morgan
1	1504	V.J. Monacella
1	1520	W.C. Lin
1	1521	W.G. Day
1	1521	A.M. Reed
1	1522	G.F. Dobay
1	1522	M.B. Wilson
1	1540	J.H. McCarthy
15	1540	B. Yim
1	1542	T.T. Huang
1	1542	J. Bai
1	1560	D. Cieslowski
1	1560	Division Head
1	1561	G.C. Cox
1	1561	S.L. Bales
1	1562	D.D. Moran
1	1562	E.E. Zarnick
1	1562	Y.S. Hiong

Copies	Code	Name
1	1563	W.E. Smith
1	1564	J.P. Feldman
1	1564	R.M. Curphey
10	5211.1	Reports Distribution
1	522.1	Library (C)
1	522.2	Library (A)

**LATE
LME**