

AD-A122 640

IMMUNOCHEMICAL INVESTIGATIONS OF CELL SURFACE ANTIGENS
OF NEISSERIA MENINGITIDIS AND NEISSERIA GONORRHEA(U)
CALIFORNIA UNIV IRVINE COLL OF MEDICINE F A WYLE
APR 79 DADA17-72-C-2076

1/1

UNCLASSIFIED

F/G 6/5

NL




END

DATE

FORM

25

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A122640

IMMUNOCHEMICAL INVESTIGATIONS OF CELL SURFACE ANTIGENS
OF NEISSERIA MENINGITIDIS AND NEISSERIA GONORRHEA

Final Report

April 1979

by

Frederic A. Wyle, M. D.

Supported by

US Army Medical Research and Development Command
Fort Detrick, Frederick, Maryland 21701

Contract No. DADA 17-72-C-2076

University of California School of Medicine
Irvine, California 92664

Approved for public release; distribution unlimited

The findings in this report are not to be construed as
an official Department of the Army position unless so
designated by other authorized documents.

FILE COPY

82 12 20 082


DTIC
ELECTE
DEC 21 1982
S D

D

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO. AD-A122640	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Immunochemical Investigations of Cell Surface Antigens of <u>Neisseria meningitidis</u> and <u>Neisseria gonorrhea</u>		5. TYPE OF REPORT & PERIOD COVERED Final Report 1 Dec 1971 - 28 Feb 1975
7. AUTHOR(s) Frederic A. Wyle, M. D.		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of California School of Medicine Irvine, California 92664		8. CONTRACT OR GRANT NUMBER(s) DADA 17-72-C-2076
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Medical Research and Development Command Fort Detrick, Frederick, Maryland 21701		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102A 3A161102B71Q.00.347
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE April 1979
		13. NUMBER OF PAGES 5 pages
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Neisseria gonorrhoeae Neisseria meningitidis Antigens Cell-mediated immunity		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Cell surface antigens of <u>N. gonorrhoeae</u> and <u>N. meningitidis</u> were isolated and purified. Antisera to these antigens were produced in rabbits and used to examine their species specificity. The cell mediated immune response in patients with clinical gonococcal infections were examined using these anti- gens. It was found that such patients have leukocyte transformation responses to both gonococcal and meningococcal antigens. Responses in female patients were found to be much greater than that of male patients.		

UNIVERSITY OF CALIFORNIA, IRVINE

BERKELEY • DAVIS • IRVINE • LOS ANGELES • RIVERSIDE • SAN DIEGO • SAN FRANCISCO



SANTA BARBARA • SANTA CRUZ

FREDERIC A. WYLE, M.D., VICE CHAIRMAN
DEPARTMENT OF MEDICINE
CALIFORNIA COLLEGE OF MEDICINE

Please reply to:

CHIEF, MEDICAL SERVICE
UCI-LONG BEACH MEDICAL PROGRAM
VETERANS ADMINISTRATION MEDICAL CENTER
5901 E. SEVENTH STREET
LONG BEACH, CALIFORNIA 90822

April 3, 1979

Contract No.: DADA 17-72-C-2076

Contracting Officer
HQDA (SGRD-MD)
Research and Development Command
Washington, D. C. 20314

Contractor: Regents of the University of
California
Principal Investigator:
Frederic A. Wyle, M.D.

Dear Sir:

Final Report

The work outlined in this final report falls into three categories: (1) isolation, purification and chemical analysis of cell surface antigens of Neisseria gonorrhoeae and Neisseria meningitidis; (2) determination of the serological specificity of the cell surface antigens of N. gonorrhoeae and N. meningitidis; and (3) investigations into the cellular immune response of patients with uncomplicated gonorrhea to purified cell surface antigens of N. gonorrhoeae and N. meningitidis.

Isolation of N. gonorrhoeae and N. meningitidis cell surface antigens was accomplished via a modification of the Ribl ether extraction technique. Partial purification of crude antigens was effected by ribonuclease and deoxyribonuclease treatment together with gel filtration chromatography. This procedure resulted in a two peak elution pattern which was virtually identical for the gonococcal and meningococcal antigens. Chemical analysis showed composition of the first peak to be primarily protein, while the second peak was largely nucleic acid. These purification steps were performed on numerous gonococcal and meningococcal strains stocked in the laboratory. Comparison of the antigenic structure of the outer membrane for these strains was carried out utilizing sodium dodecyl sulfate polyarylamide gel electrophoresis.

The antigens designated above were utilized to produce antisera in rabbits. The antisera were tested by immunodiffusion, countercurrent immunoelectrophoresis, and indirect hemagglutination techniques. These serological studies revealed extensive cross-reactivity between crude gonococcal and meningococcal antigenic preparations. Partial purification reduced the degree of cross-reactivity detected by the tests.

Peripheral blood lymphocytes (PBL) were isolated from blood samples obtained from patients with uncomplicated gonorrhea at either the Santa Ana Venereal Disease Clinic or the University of Southern California-Los Angeles County Medical Center Venereal Disease Clinic. PBL transformation stimulated by gonococcal and meningococcal antigens were utilized as a measure of cell-mediated immunity (CMI). Both male and female patients exhibited a broad range of blastogenic responses to gonococcal and meningococcal antigens. Control subjects demonstrated significantly lower and more uniform responses.

Contracting Officer
HQDA (SGRD-MD)

April 3, 1979

As a group, female patients displayed a greater CMI response than male patients. Cross-reactivity between N. gonorrhoeae and N. meningitidis was manifested by PBL transformation in patients with gonorrhea to non-purified meningococcal antigen. Partial purification of the meningococcal antigen by gel chromatography resulted in a reduced CMI response in both male and female patients. Female patients demonstrated marked stimulation with the purified gonococcal antigen, while male patients showed only slight stimulation. This study verified the lymphocyte transformation response in gonococcal infection. The functional role of CMI in gonococcal infection remains uncertain.

Sincerely,

FREDERIC A. WYLE, M.D.
Assistant Professor of Medicine

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	



12 Copies

Director (ATTN: SGRD-UWZ-AG)
Walter Reed Army Institute of
Research
Walter Reed Army Medical Center
Washington, D. C. 20012

4 Copies

HQDA (SGRD-SI)
Fort Detrick
Frederick, MD 21701

12 Copies

Defense Documentation Center
ATTN: DDC-DDA
Cameron Station
Alexandria, Virginia 22314

1 Copy

Dean
School of Medicine
Uniformed Services University
of the Health Sciences
4301 Jones Bridge Road
Bethesda, Maryland 20014

1 Copy

Superintendent
Academy of Health Sciences, US Army
ATTN: AHS-COM
Fort Sam Houston, Texas 78234