

AD-A124 419

A MULTIVARIATE IFR CLASS(U) PITTSBURGH UNIV PA CENTER
FOR MULTIVARIATE ANALYSIS T H SAVITS JAN 83 TR-83-02
N00014-76-C-0839

1/1


UNCLASSIFIED

F/G 12/1

NL



END
DATE
FILMED
83
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

DA 124419

(12)

A MULTIVARIATE IFR CLASS

by

Thomas H. Savits^{1,2}

January 1983

Technical Report No. 83-02

DTIC
ELECTE
FEB 15 1983
S H D

Center for Multivariate Analysis
University of Pittsburgh
Ninth Floor, Schenley Hall
Pittsburgh, PA 15260

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

DTIC FILE COPY

1. Department of Mathematics and Statistics, University of Pittsburgh, PA 15260.
2. Supported by ONR Contract N00014-76-C-0839.

83 02 014 209

A Multivariate IFR Class

by

Thomas H. Savits

ABSTRACT

nonnegative random vector T is said to have a multivariate increasing failure rate distribution (MIFR) if and only if $E[h(\underline{x}, T)]$ is log concave in $\underline{x}$ for all functions $h(\underline{x}, t)$ which are log concave in $(\underline{x}, t)$ and are non-decreasing and continuous in t for each fixed $\underline{x}$. This class of distributions is closed under deletion, conjunction, convolution and weak limits. It contains the multivariate exponential distribution of Marshall and Olkin and those distributions having a log concave density. Also, it follows that if T is MIFR and ψ is nondecreasing, nonnegative and concave then $\psi(T)$ is IFR.

A VARIATE NONDECREASING

increasing failure rate distribution

belongs to the class

AMS 1970 subject classification: Primary 60K10; Secondary 62N05.

Key words: multivariate, IFR, log concave, inequalities.

Y90) 383 217

UNCLASSIFIED

tions having a log concave density. Also, it follows that if T is MIFR and ψ is nondecreasing, nonnegative and concave then $\psi(T)$ is IFR.

UNCLASSIFIED

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 83- 02	2. SOVT ACCESSION NO. AD-A124419	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A Multivariate IFR Class		5. TYPE OF REPORT & PERIOD COVERED Technical-January 1983
		6. PERFORMING ORG. REPORT NUMBER 83- 02
7. AUTHOR(s) Thomas H. Savits		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0839
9. PERFORMING ORGANIZATION NAME AND ADDRESS Center for Multivariate Analysis University of Pittsburgh Pittsburgh, PA 15260		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, VA 22219		12. REPORT DATE January 1983
		13. NUMBER OF PAGES 14
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Multivariate, IFR, log concave, inequalities		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A nonnegative random vector T is said to have a multivariate increasing failure rate distribution (MIFR) if and only if $E[h(x, T)]$ is log concave in x for all functions $h(x, t)$ which are log concave in (x, t) and are nondecreasing and continuous in t for each fixed x . This class of distributions is closed under deletion, conjunction, convolution and weak limits. It contains the multivariate exponential distribution of Marshall and Olkin and those distribu-		

DD FORM 1 JAN 73 1473

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

References

- [1] Barlow, R.E. and Proschan, F. (1975). Statistical Theory of Reliability and Life Testing: Probability Models. Holt, Rinehart and Winston, New York.
- [2] Block, H.W. and Savits, T.H. (1980). Multivariate IFRA distributions. Ann. Prob., 8, 793-801.
- [3] Block, H.W. and Savits, T.H. (1981). Multivariate classes in reliability theory. Math of O.R., 6, 453-461.
- [4] Brascamp, H.J. and Lieb, E.H. (1975). Some inequalities for Gaussian measures and the long-range order of the one-dimensional plasma. Functional Integration and Its Application. A.M. Arthurs, ed. Clarendon Press, Oxford.
- [5] Marshall, A.W. (1975). Multivariate distributions with monotone hazard rate. Reliability and Fault Free Analysis. R.E. Barlow, J. Fussell and N.D. Singpurwalla, eds. SIAM, Philadelphia, 259-284.
- [6] Marshall, A.W. and Olkin, I. (1967). A multivariate exponential distribution. J. Amer. Statist. Assoc., 62, 30-44.
- [7] Marshall, A.W. and Shaked, M. (1982). A class of multivariate new better than used distributions. Ann. Prob., 10, 259-264.
- [8] Prekopa, A. (1971). Logarithmic concave measures with application to stochastic programming. Acta Math. Szeged, 32, 301-315.

§6. Relation to the MIFRA class.

In Block and Savits (1980), we defined a multivariate increasing failure rate average (MIFRA) class as follows. A nonnegative random vector $\underline{T}$ is said to be MIFRA if $E[h(\underline{T})] \leq E^{1/\alpha}[h^\alpha(\underline{T}/\alpha)]$ for all $0 < \alpha < 1$ and all nonnegative nondecreasing functions h ; equivalently, $P(\underline{T} \in A) \leq P^{1/\alpha}(\underline{T} \in \alpha A)$ for all $0 < \alpha < 1$ and all upper sets A . It is well known in the univariate case that if T is IFR, then T is IFRA (see Barlow and Proschan (1975)). It seems natural to expect the same conclusion in the multivariate setting. Although we believe this to be the case, we only have some partial results.

Let $\underline{T}$ be MIFR and μ its induced measure. Let A be an upper convex set. Then for $\alpha > 0$, $h(\alpha, \underline{t}) = I_A(\underline{t}/\alpha)$ is log concave in $(\alpha, \underline{t})$ and nondecreasing in $\underline{t}$ for each fixed α . We may assume without loss of generality that $P(\underline{T} > 0) = 1$; otherwise $P(T_1 = 0) > 0$ for some i . But T_1 is IFR and so $P(T_1 = 0) = 1$. We are thus reduced to the case of one less dimension. Since $\underline{T}$ is MIFR, it follows that $G(\alpha) = E[h(\alpha, \underline{T})]$ is log concave in $\alpha > 0$. Also, since $P(\underline{T} > 0) = 1$, we get that $G(0+) = 1$. Consequently, we conclude that $-\log G(\alpha)$ is star-shaped and so $\mu(\alpha A) \geq \mu^\alpha(A)$.

This result coupled with the fact that every upper set can be approximated by a finite union of upper convex sets leads us to the following conjecture.

(6.1) Conjecture. If $\underline{T}$ is MIFR, then $\underline{T}$ is MIFRA.

Now let $\epsilon \rightarrow 0$.

(5.7) Corollary. A nonnegative random vector T is MIFR if and only if $E[h(\underline{x}, T)]$ is log concave in $\underline{x}$ for all functions h which are log concave in $(\underline{x}, t)$ and are nondecreasing in t for each fixed $\underline{x}$.

fixed λ . Let $h(\lambda, \underline{y}) = \exp\{-\phi(\lambda, \underline{y})\}$. Then h is log concave in $(\lambda, \underline{y})$, and continuous and nondecreasing in $\underline{y}$ for each fixed λ . Consequently, $h_n(\lambda, \underline{y}) = h^n(\lambda, \underline{y})$ has the same properties for each n . Using the fact that T is MIFR and that $h_n(\lambda, \underline{y}) \rightarrow I_D(\lambda, \underline{y})$ as $n \rightarrow \infty$, we obtain the inequality (2.4).

(5.5) Lemma. Let μ be a probability measure and A an upper convex set. Then given $\epsilon > 0$ there exists a closed upper convex set $C \subset A$ such that $\mu(C) \geq \mu(A) - \epsilon$.

Proof. Given $\epsilon > 0$, choose a compact set $K \subset A$ such that $\mu(K) \geq \mu(A) - \epsilon$. If G is the convex hull of K , then G is a compact convex set satisfying $K \subset G \subset A$. As in Block and Savits (1980), define $C = \{\underline{y} + t: \underline{y} \in G, t \geq 0\}$. It is not hard to show that C has the desired properties; i.e., C is a closed upper convex set satisfying $K \subset G \subset C \subset A$. Consequently $\mu(C) \geq \mu(A) - \epsilon$.

(5.6) Theorem. If μ is a probability measure satisfying the inequality $\mu[\lambda A + (1-\lambda)B] \geq \mu^\lambda(A) \mu^{1-\lambda}(B)$ for all $0 < \lambda < 1$ and all closed upper convex sets A, B , then the inequality remains valid for all upper convex sets A, B .

Proof. It suffices to consider upper convex sets A and B having positive μ measure. Given $\epsilon > 0$, sufficiently small, we choose closed upper convex sets $C \subset A$, $D \subset B$ as in Lemma 5.5 such that $\mu(C) \geq \mu(A) - \epsilon$, $\mu(D) \geq \mu(B) - \epsilon$. Hence

$$\begin{aligned} \mu[\lambda A + (1-\lambda)B] &\geq \mu[\lambda C + (1-\lambda)D] \\ &\geq \mu^\lambda(C) \mu^{1-\lambda}(D) \geq [\mu(A) - \epsilon]^\lambda [\mu(B) - \epsilon]^{1-\lambda}. \end{aligned}$$

$$\begin{aligned}
h[\lambda \underline{x} + (1-\lambda)\underline{x}', \lambda \underline{w} + (1-\lambda)\underline{w}'] &= \mu\{D[\lambda \underline{x} + (1-\lambda)\underline{x}', \lambda \underline{w} + (1-\lambda)\underline{w}']\} \\
&\geq \mu\{\lambda D(\underline{x}, \underline{w}) + (1-\lambda)D(\underline{x}', \underline{w}')\} \\
&\geq \mu^\lambda[D(\underline{x}, \underline{w})] \mu^{1-\lambda}[D(\underline{x}', \underline{w}')] = h^\lambda(\underline{x}, \underline{w}) h^{1-\lambda}(\underline{x}', \underline{w}');
\end{aligned}$$

i.e., h is log concave in $(\underline{x}, \underline{w})$. Thus by Theorem 2.4, G is log concave in $\underline{x}$.

We now consider the converse. Let A and B be upper convex sets. For $0 \leq \lambda \leq 1$, let $C_\lambda = \lambda A + (1-\lambda)B$. We define $D = \{(\lambda, \underline{y}) : 0 \leq \lambda \leq 1 \text{ and } \underline{y} \in C_\lambda\}$. Then D is convex and so $h(\lambda, \underline{y}) = I_D(\lambda, \underline{y})$ is log concave. It is not hard to show that h is nondecreasing in $\underline{y}$ for each fixed λ . By hypothesis, then, $g(\lambda) = \int h(\lambda, \underline{y}) \mu(d\underline{y})$ is log concave in λ . Hence

$$\mu[\lambda A + (1-\lambda)B] = g(\lambda) \geq g^\lambda(1) g^{1-\lambda}(0) = \mu^\lambda(A) \mu^{1-\lambda}(B).$$

(5.3) Theorem. Let T be a nonnegative random vector and $\mu(d\underline{y}) = P(T \in d\underline{y})$ be its induced measure. Then T is MIFR if and only if

$$(5.4) \quad \mu[\lambda A + (1-\lambda)B] \geq \mu^\lambda(A) \mu^{1-\lambda}(B)$$

for all $0 < \lambda < 1$, all closed upper convex sets A, B .

Proof. The same proof as in Theorem 5.1 shows that if (5.4) holds then T is MIFR; if $F(\underline{x}, \underline{y})$ is also continuous in $\underline{y}$ for each fixed $\underline{x}$, then the sets $\{\underline{y} : F(\underline{x}, \underline{y}) \geq z\}$ are also closed.

Now let A, B be closed upper convex sets. As before, let $C_\lambda = \lambda A + (1-\lambda)B$ for $0 \leq \lambda \leq 1$ and set $D = \{(\lambda, \underline{y}) : 0 \leq \lambda \leq 1, \underline{y} \in C_\lambda\}$. Now D is a closed convex set. We set $\phi(\lambda, \underline{y}) = \rho(\lambda, \underline{y}, D)$ where ρ is the metric $\rho(\underline{u}, \underline{v}) = \max |u_i - v_i|$. Note that ϕ is continuous and convex. It is also a nonincreasing function in $\underline{y}$ for each

§5. An alternative condition.

In this section (which is somewhat technical) we derive an alternative condition that T be MIFR; it is directly expressable in terms of the induced measure $\mu(dy) = P(T \in dy)$. This result, coupled with some approximation ideas, allows us to remove the continuity assumption on h in Definition 4.1.

We first recall that a set $A \subset \mathbb{R}^n$ is said to be an upper set if whenever $x \in A$ and $y \geq x$, then $y \in A$.

(5.1) Theorem. Let μ be a finite measure. In order that $\int F(x,y)\mu(dy)$ be log concave in x for all log concave functions $F(x,y)$ which are nondecreasing in y for each fixed x it is necessary and sufficient that

$$(5.2) \quad \mu[\lambda A + (1-\lambda)B] \geq \mu^\lambda(A) \mu^{1-\lambda}(B)$$

for all $0 < \lambda < 1$ and all upper convex sets A, B .

Proof. We shall closely follow the argument given in Brascamp and Lieb (1975).

Suppose now that μ satisfies (5.2) and let $F(x,y)$ satisfy the conditions of the theorem. For each real $z \geq 0$, let $C(x,z) = \{y: F(x,y) \geq z\}$ and set

$g(x,z) = \mu[C(x,z)]$. Note that $C(x,z)$ is upper and convex; furthermore,

$$G(x) = \int_{-\infty}^{\infty} F(x,y)\mu(dy) = \int_0^{\infty} g(x,z)dz. \quad \text{If we make the change of variables } z = e^w,$$

then $G(x) = \int_{-\infty}^{\infty} h(x,w)e^w dw$ where $h(x,w) = g(x,e^w) = \mu[C(x,e^w)]$. We also define

$$D(x,w) = C(x,e^w). \quad \text{It is easy to show that } D[\lambda x + (1-\lambda)x', \lambda w + (1-\lambda)w'] \supset$$

$\lambda D(x,w) + (1-\lambda)D(x',w')$ for all $0 < \lambda < 1$, x, x', w, w' . Consequently, assuming (5.2),

we obtain the inequality

ponentially distributed random variables $S_1, \dots, S_n$ and subsets $J_1, \dots, J_m$ of $\{1, \dots, n\}$ such that $T_i = \min_{j \in J_i} S_j$. But $(S_1, \dots, S_n)$ is MIFR and $\psi_i(s) = \min_{j \in J_i} s_j$ ($1 \leq i \leq m$) satisfy the hypothesis of Theorem 4.3(v).

(4.6) Example 2. If T has a density which is log concave, then T is MIFR.

Apply Theorem 2.4.

$S_i = \psi_i(T)$, $1 \leq i \leq m$, with T MIFR, then $E[h(x, S)] = E[g(x, T)]$ is log concave in x .

(ii) Let $h(x, s, t)$ be log concave in (x, s, t) and continuous and nondecreasing in (s, t) for each fixed x . We first assume that h is bounded. Since T is MIFR it follows that $g(x, s) = E[h(x, s, T)]$ is log concave in (x, s) . Also, g is continuous and nondecreasing in s for each fixed x . Thus, since S is MIFR, $E[g(x, S)]$ is log concave in x . But, by Fubini, $E[g(x, S)] = E[h(x, s, T)]$. If h is not bounded, consider instead $h \wedge n$ and pass to the limit (see Theorem 2.3(i)).

(i) If $J \subset \{1, \dots, n\}$, let $\psi_j(t) = t_j$ for $j \in J$. According to Theorem 4.3(v), $\{\psi_j(T) = T_j; j \in J\}$ is MIFR.

(iii) Since S and T are independent MIFR, (S, T) is MIFR by Theorem (4.3(ii)).

Now take $\psi_j(s, t) = s_j + t_j$ and use Theorem 4.3(v).

(iv) Use Theorem 4.3(v) with $\psi_j(t) = a_j t_j$.

(v) Let $h(x, t)$ be a bounded log concave function in (x, t) which is nondecreasing and continuous in t for each fixed x . Since $T_n \rightarrow T$ in distribution, $E[h(x, T_n)] \rightarrow E[h(x, T)]$ as $n \rightarrow \infty$. But for each n , $E[h(x, T_n)]$ is log concave in x . Consequently, $E[h(x, T)]$ is log concave in x . If h is not bounded, use the argument as in the proof of part (ii).

(4.4) Corollary. If $T_1, \dots, T_n$ are independent IFR random variables and $\psi(t_1, \dots, t_n)$ is continuous, nonnegative, nondecreasing and concave, then $\psi(T_1, \dots, T_n)$ is IFR.

(4.5) Example 1. The Marshall and Olkin MVE (1967) distribution is MIFR.

This follows since if $T = (T_1, \dots, T_m)$ is MVE then there exist independent ex-

§4. A multivariate IFR class.

(4.1) Definition. Let $\underline{T}$ be a nonnegative random vector. We say that $\underline{T}$ has a multivariate increasing failure rate (MIFR) distribution if and only if $E[h(\underline{x}, \underline{T})]$ is log concave in $\underline{x}$ for all functions $h(\underline{x}, \underline{t})$ which are log concave in $(\underline{x}, \underline{t})$ and nondecreasing and continuous in $\underline{t} \geq \underline{0}$ for each fixed $\underline{x} \geq \underline{0}$.

(4.2) Remarks.

- (i) Again note that according to Theorem 2.3(iii) we need only consider functions $h(\underline{x}, \underline{t})$ with $\underline{x}$ a single variable instead of a vector.
- (ii) In Section 5 we shall show that the continuity assumption is unnecessary.

The class of MIFR distributions has many desirable closure properties.

(4.3) Theorem.

- (i) If $\underline{T}$ is MIFR, then so are all marginals.
- (ii) If $\underline{S}$ and $\underline{T}$ are independent MIFR, then $(\underline{S}, \underline{T})$ is MIFR.
- (iii) If $\underline{S}$ and $\underline{T}$ are independent MIFR of the same dimension, then $\underline{S} + \underline{T}$ is MIFR.
- (iv) If $(T_1, \dots, T_n)$ is MIFR and $a_i \geq 0$ ($i = 1, \dots, n$), then $(a_1 T_1, \dots, a_n T_n)$ is MIFR.
- (v) If $\underline{T}$ is MIFR and $\psi_1, \dots, \psi_m$ are continuous, nonnegative, nondecreasing and concave functions, then $(\psi_1(\underline{T}), \dots, \psi_m(\underline{T}))$ is MIFR.
- (vi) If $\underline{T}_1, \underline{T}_2, \dots$ are MIFR and $\underline{T}_n$ converges to $\underline{T}$ in distribution, then $\underline{T}$ is MIFR.

Proof.

(v) Let $h(\underline{x}, \underline{s})$ be log concave in $(\underline{x}, \underline{s})$ and continuous, and nondecreasing in $\underline{s}$ for each fixed $\underline{x}$. If $\psi_1, \dots, \psi_m$ are as in (v), then $g(\underline{x}, \underline{t}) = h(\underline{x}, \psi_1(\underline{t}), \dots, \psi_m(\underline{t}))$ has the same properties as h (see Theorem 2.3(ii)). Consequently, if

Proof. Let $R(t) = -\log \bar{F}(t)$ be the hazard function of T . We set $a = \sup\{t \geq 0: \bar{F}(t) = 1\}$ and $b = \inf\{t \geq 0: \bar{F}(t) = 0\}$ ($\inf \emptyset = +\infty$). If $a = b$, we simply take $\psi \equiv a$ and we are done; otherwise $0 \leq a < b \leq +\infty$. Since R is convex and finite on $(-\infty, b)$ it is continuous there; furthermore, it easily follows that R is strictly increasing on $[a, b)$. Let $A = \lim_{t \rightarrow b^-} R(t) \leq +\infty$. If ϕ denotes the restriction of R to $[a, b)$, then its inverse ϕ^{-1} is continuous, strictly increasing and concave on $[0, A)$. The function ψ is defined by $\psi(s) = \inf\{t \geq 0: R(t) > s\}$. Clearly $\psi(s) = \phi^{-1}(s)$ for $0 \leq s < A$ and $\psi(s) = b$ for $s \geq A$. It is not hard to show now that ψ has the desired properties.

(3.4) Theorem. T is IFR if and only if $E[h(\underline{x}, T)]$ is log concave in $\underline{x}$ for all functions $h(\underline{x}, t)$ which are log concave in $(\underline{x}, t)$ and are nondecreasing in t for each fixed $\underline{x} \geq 0$.

Proof. Suppose T is IFR and let h be as in the statement of the theorem. Then, according to Lemma 3.3, there exists a continuous nonnegative nondecreasing concave function ψ such that $\psi(S)$ and T have the same distribution, where S has the standard exponential distribution. Hence

$$E[h(\underline{x}, T)] = E[h(\underline{x}, \psi(S))] = \int_0^\infty h(\underline{x}, \psi(s)) e^{-s} ds$$

is log concave in $\underline{x}$. This follows from Theorems 2.3 (i), (ii) and 2.4.

To prove the converse, let $h(\underline{x}, t) = I_{(\underline{x}, \infty)}(t)$. Then h is log concave in $(\underline{x}, t)$ and nondecreasing in t for each fixed $\underline{x}$. By assumption, then, $\bar{F}(\underline{x}) = E[h(\underline{x}, T)]$ is log concave in $\underline{x}$. According to Theorem 3.1, T is IFR.

(3.5) Remark. According to Theorem 2.3(iii), we need only consider functions $h(\underline{x}, t)$ with $\underline{x}$ a single variable instead of a vector.

§3. A new univariate IFR characterization.

The univariate class of increasing failure rate (IFR) distributions has played an important role in the mathematical theory of reliability (see Barlow and Proschan (1975)). We shall first briefly review the definition. Let T be a nonnegative random variable with survival probability $\bar{F}(t) = P(T > t)$. Set $b = \inf\{t \geq 0: \bar{F}(t) = 0\}$ ($\inf \emptyset = +\infty$). We say that T has an IFR distribution if $\bar{F}(s+t)/\bar{F}(t)$ is nonincreasing in $t \in [0, b)$ for each $s \geq 0$. It is well known that the following conditions are equivalent (cf. Barlow and Proschan (1975)).

(3.1) Theorem. The following conditions are equivalent:

- (i) T is IFR.
- (ii) $\bar{F}$ is a Polya frequency function of order two (PF_2), i.e., $\bar{F} \geq 0$ and

$$\left| \begin{array}{cc} \bar{F}(x_1 - y_1) & \bar{F}(x_1 - y_2) \\ \bar{F}(x_2 - y_1) & \bar{F}(x_2 - y_2) \end{array} \right| \geq 0$$

for all $-\infty < x_1 < x_2 < \infty$, $-\infty < y_1 < y_2 < \infty$.

- (iii) $\bar{F}$ is log concave.

(3.2) Remark. If F has a density f , then T is IFR if and only if the hazard rate $r(t) = f(t)/\bar{F}(t)$ is nondecreasing on $[0, b)$.

Before we state our new characterization, we need the following simple result.

- (3.3) Lemma. If T is IFR, then there exists a continuous nonnegative non-decreasing concave function ψ on $[0, \infty)$ such that $\psi(S)$ has the same distribution as T , where S is distributed as a standard exponential.

The following facts about log concave functions are easily verifiable.

(2.3) Theorem.

- (i) If f and g are log concave, so are $f \cdot g$, $f \wedge g$, cf and f^a for all $a > 0$, $c \geq 0$.
- (ii) If f is log concave and nondecreasing and ψ is concave (and nonnegative), then the composition $f \circ \psi$ is log concave.
- (iii) f is log concave if and only if for every $\underline{x}, \underline{y} \geq \underline{0}$, the function $g(t) = f[t\underline{x} + (1-t)\underline{y}]$ is log concave on $0 \leq t \leq 1$.
- (iv) If f is log concave, then for every z , the sets $\{f > z\}$ and $\{f \geq z\}$ are convex.
- (v) If f is log concave, then f is continuous on the interior of the set $\{f > 0\}$.

We close this section with a very important result about log concave functions. This result is sometimes known as the Prékopa Theorem (1971).

An independent and simpler proof is given in Brascamp and Lieb (1975).

(2.4) Theorem. Let F be log concave on $\mathbb{R}^m \times \mathbb{R}^n$. Then $G(\underline{x}) = \int F(\underline{x}, \underline{y}) d\underline{y}$ is log concave on $\mathbb{R}^m$. (Here $d\underline{y}$ is Lebesgue measure on $\mathbb{R}^n$).

The above plays a crucial role in our development.

§2. Log concave functions.

Let A be a convex set in $\mathbb{R}^n$ and f a nonnegative function defined on A . We say that f is log concave on A if

$$(2.1) \quad f[\lambda x + (1-\lambda)y] \geq f^\lambda(x) f^{1-\lambda}(y)$$

for all $0 < \lambda < 1$, all $x, y \in A$. Sometimes we write $f(x) = e^{-Q(x)}$ where $Q(x)$ is convex, but with the understanding that Q may assume the value $+\infty$.

(2.2) Examples

- (i) If $Q(x)$ is twice continuously differentiable on an open convex set A and has the property that at each point $x \in A$, the matrix $\left[\frac{\partial^2 Q(x)}{\partial x_i \partial x_j} \right]$ is nonnegative definite, then $f(x) = e^{-Q(x)}$ is log concave on A . Hence all Gaussian densities are log concave.
- (ii) If A is any convex set in $\mathbb{R}^n$, then the indicator function $I_A(x)$ is log concave on $\mathbb{R}^n$.
- (iii) If f is a nonnegative concave function, then f is log concave.

It is convenient to make the following simple observation. Let A and B be convex sets in $\mathbb{R}^n$ and suppose that f is log concave on A . Clearly then, f is log concave on B if $B \subset A$. On the other hand suppose $A \subset B$. Then the function $\tilde{f}$, which is defined to be equal to f on A and zero on $B \setminus A$, is log concave on B . Thus without loss of generality we may assume that all log concave functions are defined on the same convex set. In the context of this paper, it is natural to work with the convex set $\mathbb{R}_+^n$. Hence, unless otherwise specified, the term log concave means log concave on $\mathbb{R}_+^n$.

§1. Introduction.

→ Various univariate classes of life distributions have been introduced in the mathematical theory of reliability, (see Barlow and Proschan (1975) for a detailed discussion). Recently there has been much interest in obtaining multivariate versions of these classes. Although there have been many different approaches, *this document mentions* ~~(see, e.g., the review paper by Block and Savits~~

~~(1981), we mention only two because of their nice closure properties:~~
the multivariate IFRA class of Block and Savits (1980) and the multivariate
NBU class of Marshall and Shaked (1982). Both of these classes are closed
under deletion, conjunction, convolution and weak limits.

~~In this paper we introduce~~ a multivariate IFR class that has similar closure properties. Its spirit also closely parallels that found in the two previously mentioned papers. Other definitions of a multivariate IFR class have been proposed by Marshall (1975), but none possess all desirable closure properties.

In Section 2 we present some preliminary facts about log concave functions. Section 3 contains a new characterization of the (univariate) IFR class. The multivariate generalization and properties thereof are given in Section 4. A useful alternative condition is delineated in Section 5. Finally, ^{is compared} ~~we compare~~ this class with the multivariate IFRA class of Block and Savits (1980) in Section 6. ←

All functions and sets in this paper are assumed to be Borel measurable with respect to $\mathbb{R}^n$. In most cases, however, we do not specify the dimension n directly; it is usually clear from the context.

[illegible]