

AD-A126 281

QUANTUM DYNAMICAL MODEL FOR LASER EXCITATION OF A
TWO-LEVEL ADATOM: SURFA. (U) ROCHESTER UNIV NY DEPT OF
CHEMISTRY J LIN ET AL. MAR 83 UROCHESTER/DC/83/TR-31
N00014-80-C-0472

1/1

UNCLASSIFIED

F/G 20/5

NL




END

FILED

11

DTM



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

DA 126201

OFFICE OF NAVAL RESEARCH

Contract N00014-80-C-0472

Task No. NR 056-749

TECHNICAL REPORT No. 31

Quantum Dynamical Model for Laser
Excitation of a Two-Level Atom:
Surface-Dressed Bloch Equations

by

Jui-teng Lin, Xi-Yi Huang
and Thomas F. George

Prepared for Publication

in

Solid State Communications

Department of Chemistry
University of Rochester
Rochester, New York 14627

March 1983

Reproduction in whole or in part is permitted for any
purpose of the United States Government.

This document has been approved for public release and
sale; its distribution is unlimited.



DTIC FILE COPY

83 03 29 067

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER UROCHESTER/DC/83/TR-31	2. GOVT ACCESSION NO. AD-4126 201	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Quantum Dynamical Model for Laser Excitation of a Two-Level Adatom: Surface-Dressed Bloch Equations		5. TYPE OF REPORT & PERIOD COVERED Interim Technical Report
7. AUTHOR(s) Jui-teng Lin, Xi-Yi Huang and Thomas F. George		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry University of Rochester Rochester, New York 14627		8. CONTRACT OR GRANT NUMBER(s) N00014-80-C-0472
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Chemistry Program Code 472 Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 056-749
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE March 1983
		13. NUMBER OF PAGES 10
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Prepared for publication in Solid State Communications.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) TWO-LEVEL ADATOM METAL SURFACE LASER REFLECTED FIELD SURFACE-DRESSED BLOCH EQUATIONS PHOTONS AND PHONONS ELECTRON-PHONON COUPLING ORIENTATION OF TRANSITION DIPOLE POPULATION INVERSION		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Surface-dressed Bloch equations are derived from a microscopic Hamiltonian which includes the inter- actions among an adatom, the laser photons and the phonon modes. The influence of the surface on the population inversion of the adatom is analyzed in terms of: (i) the reflected-field-induced damping factor which depends on the orien- tation of the transition dipole and the adatom-surface separation and (ii) the frequency shift and damping factor induced by the phonon modes.		

Quantum Dynamical Model for Laser Excitation of a Two-Level Adatom:
Surface-Dressed Bloch Equations

Jui-teng Lin
Laser Physics Branch
Optical Sciences Division
Naval Research Laboratory
Washington, DC 20375 USA

and

Xi-Yi Huang* and Thomas F. George
Department of Chemistry
University of Rochester
Rochester, NY 14627 USA

Accession For	
Final Serial	☒
Final Page	☐
Unpublished	☐
Notification	
Distribution/	
Availability Codes	
Avail and/or	Special
100	

2218
copy
inspected
2

A

Abstract

Surface-dressed Bloch equations are derived from a microscopic Hamiltonian which includes the interactions among an adatom, the laser photon and the phonon modes. The influence of the surface on the population inversion of the adatom is analyzed in terms of: (i) the reflected-field-induced damping factor which depends on the orientation of the transition dipole and the adatom-surface separation and (ii) the frequency shift and damping factor induced by the phonon modes.

*Also with the Department of Physics, University of Rochester.
Permanent address: Institute of Physics, Academia Sinica,
Beijing (Peking), The People's Republic of China.

I. Introduction

The interaction of laser radiation with matter in homogeneous systems has been extensively studied for the past several years.^{1,2} However, for heterogeneous systems (e.g., gaseous atoms near or adsorbed on a solid surface) the laser-stimulated surface phenomena (desorption, dissociation, migration and reactions) involving both multiphoton and multiphonon processes have been only recently attacked.³⁻⁵ In previous studies, we have investigated the excitation and relaxation dynamics of adspecies subjected to IR radiation in which only the vibrational degrees of freedom are concerned.⁴ In the present paper, we shall investigate the electronic excitation of a two-level atom adsorbed on a metal surface and subjected to UV or visible laser radiation. In the absence of a solid surface or when the atom is very far from the surface, the population inversion and the power spectrum of the system may be described by an ordinary optical Bloch equation (OBE). In the presence of a surface at a distance from the adatom which is comparable to the optical wavelength, the dynamical phenomena are influenced by the following factors: (i) nonradiative energy relaxation of the excited atom via electron-phonon coupling; (ii) radiative spontaneous decay and stimulated emission produced by both the applied field and the reflected field; (iii) the oscillatory behavior of the lifetime of the adatom due to the interference between the applied field and the reflected field; (iv) the reflectivity and refraction index of the surface; (v) surface-induced dephasing of the

dipole; and (vi) interaction between the adatom and plasmons; as a first step, we assume that the effects of the conduction electrons in the metal just provide a reflected field acting back to the adatom.

II. Microscopic Hamiltonian and Surface-Dressed Bloch Equations

To investigate the above surface phenomena, we start with a microscopic Hamiltonian describing a two-level adatom subjected to laser radiation,

$$H = H_A + H_B + H_F + H_{AB} + H_{AF} + H_{AR} + H_{BF}. \quad (1)$$

H_A , H_B and H_F are the unperturbed Hamiltonians for the adatom, phonon and photon (laser) modes, respectively; H_{AB} , H_{AF} and H_{AR} describe the interaction of the adatom (A) with the phonon bath (B) modes, the applied field photon (F) modes and the reflected field (R), respectively; and H_{BF} is the interaction between the B modes and the applied field. For a laser frequency in the visible to UV range, the phonon (B) modes have much weaker absorption compared to that of the adatom, and thereby the direct heating of the substrate due to H_{BF} is negligible. In this work, the effect of H_{BF} will be displayed in terms of the strength of the reflected electric field due to the reflectivity of the surface.

Employing Pauli's matrices for the two-level adatom and harmonic ladder operators for the photon and phonon modes, the total Hamiltonian may be expressed in a second-quantization

form which leads to the equations of motion, in the rotating-wave approximation, as follows:⁴

$$\dot{\sigma}_{12} = -i\omega_0\sigma_{12} + i\sum_{\nu} G_{\nu} B_{\nu} \sigma_3 + i\sum_{\lambda} C_{\lambda} a_{\lambda} \sigma_3 - \gamma_R \sigma_{12} \quad , \quad (2a)$$

$$\dot{\sigma}_3 = 2i\sum_{\lambda} C_{\lambda} (a_{\lambda}^{\dagger} \sigma_{12} + a_{\lambda} \sigma_{21}) + 2i\sum_{\nu} G_{\nu} (B_{\nu}^{\dagger} \sigma_{12} + B_{\nu} \sigma_{21}) \quad , \quad (2b)$$

$$\dot{a}_{\lambda} = -i\omega a_{\lambda} + iC_{\lambda} \sigma_{12} \quad , \quad (2c)$$

$$\dot{B}_{\nu} = -i\bar{\omega}_{\nu} B_{\nu} - i\sum_{\nu} G_{\nu} N_{\nu} \sigma_{12} \quad . \quad (2d)$$

$\sigma_{ij} = |i\rangle\langle j|$ is the atomic transition operator, $\sigma_3 = \sigma_{22} - \sigma_{11}$, and ω_0 and ω are the atomic transition and laser field frequencies, respectively; C_{λ} and G_{ν} are the coupling coefficients in the interaction Hamiltonians H_{AF} and H_{AB} , respectively; the multi-phonon processes are governed by the phonon operator, B_{ν} , frequency, $\bar{\omega}_{\nu}$, and occupation number, N_{ν} , given by⁵

$$B_{\nu} = \prod_j^{\nu} b_j, \quad \bar{\omega}_{\nu} = \sum_j^{\nu} \omega_j \quad , \quad (3a)$$

$$N_{\nu}(T) = \prod_j (\bar{n}_j + 1) - \prod_j \bar{n}_j \quad , \quad (3b)$$

$$\bar{n}_j(T) = (e^{\hbar\omega_j/kT} - 1)^{-1} \quad , \quad (3c)$$

where ω_j is the frequency of the j -th phonon mode with a thermal equilibrium occupation number at temperature T defined by Eq. (3c). The applied laser field is quantized by its ladder operator a_λ for the λ -th mode.

The effects of the quantized reflected field come in through the last term in Eq. (2a), which depends on the damping factor γ_R ,

$$\gamma_R = 2|\mu_{12}|^2 \text{Im}f/\hbar, \quad (4)$$

where μ_{12} is the electric-dipole transition matrix element between states $|i\rangle$ and $|j\rangle$. The function $f(d) \equiv \text{Re}f(d) + i\text{Im}f(d)$, depending on the orientation of the dipole with respect to the surface, is given by⁶

$$f_\perp(d) = -R\left(\frac{2}{\chi^3}\right) \left[\left(\frac{1}{(2\hat{d})^3} - \frac{i}{(2\hat{d})^2} \right) e^{2i\hat{d}} \right] \quad (5a)$$

$$f_\parallel(d) = R\left(\frac{1}{\chi^3}\right) \left[\left(\frac{1}{(2\hat{d})^3} + \frac{1}{2\hat{d}} - \frac{i}{(2\hat{d})^2} \right) e^{2i\hat{d}} \right] \quad (5b)$$

for the perpendicular and parallel cases. The reduced distance is $\hat{d} = d/\chi = 2\pi d/\lambda$, where d is the distance of the adatom from the surface and λ is the wavelength of the field. R is the reflectivity of the surface taking into account the interaction between the phonon modes and the field, H_{BF} . Since $R \approx 1$, we shall neglect the effects of H_{BF} and assume a perfectly reflective surface.

With the help of many-body techniques applied to multiphonon processes,^{5,7} we obtain the surface-dressed Bloch equations (SBE):⁸

$$\frac{d}{dt} \begin{pmatrix} u \\ v \\ w \end{pmatrix} = \begin{pmatrix} -\tilde{\gamma}_2 & -\tilde{\Delta} & 0 \\ \tilde{\Delta} & -\tilde{\gamma}_2 & \Omega \\ 0 & -\Omega & -\tilde{\gamma}_1 \end{pmatrix} \begin{pmatrix} u \\ v \\ w \end{pmatrix} + \tilde{\gamma}_1 \begin{pmatrix} 0 \\ 0 \\ w_{eq} \end{pmatrix}. \quad (6)$$

u and v are the components, in units of the transition moment, of the adatomic dipole moment "in phase" and "in quadrature" with the applied field. w is the population inversion, $w \equiv \langle \sigma_3 \rangle$, with the equilibrium value w_{eq} , where $w_{eq} = -1$ for the adatom initially in its ground state and $w_{eq} = 1$ for the excited state. $\Omega = |\mu_{12} E_0|/\hbar$ is the Rabi frequency and is proportional to the amplitude of the applied electric field, E_0 , which is assumed to have a slowly-varying envelope. $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$ are the relaxation factors for the inversion and for the dipole, respectively, and $\tilde{\Delta}$ is the effective detuning of the laser field frequency with respect to the adatomic transition frequency. These surface-dressed parameters are related to the surface-free parameters by

$$\tilde{\Delta} = \Delta + \delta\omega_F + \delta\omega_B, \quad (7a)$$

$$\tilde{\gamma}_1 = 2(\gamma_B + A/2), \quad (7b)$$

$$\tilde{\gamma}_2 = \gamma_B + A/2 + \gamma_2 + \gamma_R, \quad (7c)$$

where $\Delta = \omega_0 - \omega$ is the surface-free detuning, A is the Einstein coefficient for spontaneous emission, $\delta\omega_F$ and $\delta\omega_B$ are the frequency shifts due to the applied field and phonons, respectively, and γ_B and γ_2 are the phonon-induced energy (T_1) and phase (T_2) relaxation rates, respectively. The phonon-induced frequency shift and damping factor are given by⁵

$$\delta\omega_B = P \sum_{\nu} \frac{|G_{\nu}|^2 N_{\nu}}{\omega_0 - \bar{\omega}_{\nu}}, \quad P \equiv \text{principal part} \quad (8a)$$

$$\gamma_B = \sum_{\nu} |G_{\nu}|^2 N_{\nu} \delta(\omega_0 - \bar{\omega}_{\nu}) \quad (8b)$$

which, in contrast to that of the optically-induced parameters, are strongly temperature dependent through the multiphonon occupation number, $N_{\nu}(T)$, given by Eq. (3b). We note that these features of temperature dependence disappear when just a single-phonon process is considered in H_{AB} .

The steady-state population inversion is found analytically to be (for $w_{eq} = -1$)

$$w_{s.s.} = \left(1 + \frac{(\tilde{\gamma}_2/\tilde{\gamma}_1)\Omega^2}{\tilde{\Delta}^2 + \tilde{\gamma}_2^2} \right)^{-1}, \quad (9)$$

which is sensitive to the reduced distance and dipole orientation through the surface-dressed parameters $\tilde{\gamma}_1$, $\tilde{\gamma}_2$ and $\tilde{\Delta}$. We see from the above equation that the criterion for the weak-field limit in the SBE, given by

$$\Omega^2 \ll (\tilde{\Delta}^2 + \tilde{\gamma}_2^2) \tilde{\gamma}_1 / \tilde{\gamma}_2 ,$$

is a weaker condition than that in the surface-free OBE due to the surface-dressed parameters.

III. Discussion

The SBE displayed in Eq. (6) are different from the surface-free OBE due to the influence of the reflected field, γ_R , and the effects of the electron-phonon coupling, $\delta\omega_B$ and γ_B . We note that these surface-induced effects are characterized by various system parameters such as the temperature of the surface, the orientation and position of the induced dipole of the adatom, the transition frequencies of the adatom (ω_0) and the phonons (ω_j), and the wavelength of the applied field. For example, by the concept of the "energy gap law", $\delta\omega_B$ and γ_B increase when the temperature of the surface increases and/or the order of the multiphonon coupling, p , defined by $p \equiv (\omega_B + \delta\omega_A + \delta\omega_B) / \omega_j$ where $\delta\omega_A$ is the frequency shift due to the multimode property of the laser, decreases.^{4,5} From Eqs. (4) and (5), we see that γ_B is sensitive to the reduced distance, $\hat{d} = d/2\pi\lambda$, where for sufficiently small $\hat{d}$ the value of γ_B oscillates as a function of d , as does also the lifetime of the adatom.

In conclusion, we have shown that from a microscopic Hamiltonian the SBE can be obtained, which exhibit important surface effects on the excitation of a two-level atom. The results suggest that monochromatic laser radiation might be used not only as a characterization of surface adspecies, such as orientation, but also as a probe of the dynamics of rate processes occurring at a solid surface both for vibrational and electronic excitations.

Acknowledgements

This work was supported in part by the Air Force Office of Scientific Research (AFSC), the United States Air Force, under Grant No. AFOSR 82-0046, the Office of Naval Research and the U.S. Army Research Office. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon. Thomas F. George acknowledges the Camille and Henry Dreyfus Foundation for a Teacher-Scholar Award (1975-84).

References

1. L. Allen and J. H. Eberly, Optical Resonance and Two-Level Atoms (Wiley, New York, 1975).
2. P. A. Schultz, A. S. Sudbo, D. J. Krainovich, H. S. Kowk, Y. R. Shen and Y. T. Lee, *Annu. Rev. Phys. Chem.* 30, 379 (1979) and references therein.
3. G. S. Selwyn and M. C. Lin in Lasers as Reactants and Probes in Chemistry, ed. by W. M. Jackson and A. B. Harvey (Howard University Press, Washington, D. C., 1983), in press.
4. J. Lin, A. C. Beri, M. Hutchinson, W. C. Murphy and T. F. George, *Phys. Lett.* 79A, 233 (1980) and references therein.
5. J. Lin and T. F. George, *J. Phys. Chem.* 84, 2957 (1980).
6. R. R. Chance, A. Prock and R. Silbey, *Adv. Chem. Phys.* 38, 1 (1978).
7. W. H. Louisell, Quantum Statistical Properties of Radiation (Wiley, New York, 1973).
8. J. Lin, to be published.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 North Quincy Street Arlington, Virginia 22217	2	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Pasadena Detachment Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Dean William Tolles Naval Postgraduate School Monterey, California 93940	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
U.S. Army Research Office Attn: CRD-AA-IP P. O. Box 12211 Research Triangle Park, N.C. 27709	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Mr. Vincent Schaper DTNSRDC Code 2803 Annapolis, Maryland 21402	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1	Mr. A. M. Anzalone Administrative Librarian PLASTEC/ARRADCOM Bldg 3401 Dover, New Jersey 07801	1
Dr. David L. Nelson Chemistry Program Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1		

TECHNICAL REPORT DISTRIBUTION LIST, 056

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. G. A. Somorjai Department of Chemistry University of California Berkeley, California 94720	1	Dr. W. Kohn Department of Physics University of California (San Diego) La Jolla, California 92037	1
Dr. J. Murday Naval Research Laboratory Surface Chemistry Division (6170) 455 Overlook Avenue, S.W. Washington, D.C. 20375	1	Dr. R. L. Park Director, Center of Materials Research University of Maryland College Park, Maryland 20742	1
Dr. J. B. Hudson Materials Division Rensselaer Polytechnic Institute Troy, New York 12181	1	Dr. W. T. Peria Electrical Engineering Department University of Minnesota Minneapolis, Minnesota 55455	1
Dr. Theodore E. Madey Surface Chemistry Section Department of Commerce National Bureau of Standards Washington, D.C. 20234	1	Dr. Chia-wei Woo Department of Physics Northwestern University Evanston, Illinois 60201	1
Dr. J. M. White Department of Chemistry University of Texas Austin, Texas 78712	1	Dr. Robert M. Hexter Department of Chemistry University of Minnesota Minneapolis, Minnesota 55455	1
Dr. Keith E. Johnson Department of Metallurgy and Materials Science Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Dr. R. P. Van Duyne Chemistry Department Northwestern University Evanston, Illinois 60201	1
Dr. J. E. Demuth IBM Corporation Thomas J. Watson Research Center P. O. Box 218 Yorktown Heights, New York 10598	1	Dr. S. Sibener Department of Chemistry James Franck Institute 5640 Ellis Avenue Chicago, Illinois 60637	1
Dr. C. P. Flynn Department of Physics University of Illinois Urbana, Illinois 61801	1	Dr. M. G. Lagally Department of Metallurgical and Mining Engineering University of Wisconsin Madison, Wisconsin 53706	1

TECHNICAL REPORT DISTRIBUTION LIST, 056

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. Robert Gomar Department of Chemistry James Franck Institute 5640 Ellis Avenue Chicago, Illinois 60637	1	Dr. K. G. Spears Chemistry Department Northwestern University Evanston, Illinois 60201	1
Dr. R. G. Wallis Department of Physics University of California, Irvine Irvine, California 92664	1	Dr. R. W. Plummer University of Pennsylvania Department of Physics Philadelphia, Pennsylvania 19104	1
Dr. D. Ramaker Chemistry Department George Washington University Washington, D.C. 20052	1	Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 41106	1
Dr. P. Hansma Physics Department University of California, Santa Barbara Santa Barbara, California 93106	1	Professor D. Hercules University of Pittsburgh Chemistry Department Pittsburgh, Pennsylvania 15260	1
Dr. J. C. Hemminger Chemistry Department University of California, Irvine Irvine, California 92717	1	Professor N. Winograd The Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802	1
Dr. Martin Fleischmann Department of Chemistry Southampton University Southampton SO9 5NH Hampshire, England	1	Professor T. F. George The University of Rochester Chemistry Department Rochester, New York 14627	1
Dr. G. Rubloff IBM Thomas J. Watson Research Center P. O. Box 218 Yorktown Heights, New York 10598	1	Professor Dudley R. Herschbach Harvard College Office for Research Contracts 1350 Massachusetts Avenue Cambridge, Massachusetts 02138	1
Dr. J. A. Gardner Department of Physics Oregon State University Corvallis, Oregon 97331	1	Professor Horia Metiu University of California, Santa Barbara Chemistry Department Santa Barbara, California 93106	1
Dr. G. D. Stein Mechanical Engineering Department Northwestern University Evanston, Illinois 60201	1	Professor A. Steckl Rensselaer Polytechnic Institute Department of Electrical and Systems Engineering Integrated Circuits Laboratories Troy, New York 12181	1

TECHNICAL REPORT DISTRIBUTION LIST, 056

	<u>No. Copies</u>	<u>No. Copies</u>
Dr. John T. Yates Department of Chemistry University of Pittsburgh Pittsburgh, Pennsylvania 15260	1	
Professor G. H. Morrison Department of Chemistry Cornell University Ithaca, New York 14853	1	
Captain Lee Myers AFOSR/NC Bolling AFB Washington, D.C. 20332	1	
Dr. David Squire Army Research Office P. O. Box 12211 Research Triangle Park, NC 27709	1	
Professor Ronald Hoffman Department of Chemistry Cornell University Ithaca, New York 14853	1	

4-
DT