

AD-A128 484

ON LIMITING DISTRIBUTIONS OF ORDER STATISTICS WITH
VARIABLE RANKS FROM ST..(U) NORTH CAROLINA UNIV AT
CHAPEL HILL CENTER FOR STOCHASTIC PREC.. S CHENG

1/1

UNCLASSIFIED

JAN 83 TR-25 AFOSR-TR-83-0414

F/G 12/1

NL


END

DATE

FILED

6 83

DTIC




MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

3

AFOSR-TR- 83 - 0414

CENTER FOR STOCHASTIC PROCESSES

Department of Statistics
University of North Carolina
Chapel Hill, North Carolina



ON LIMITING DISTRIBUTIONS OF ORDER STATISTICS
WITH VARIABLE RANKS FROM STATIONARY SEQUENCES

by

Shihong Cheng

TECHNICAL REPORT #25

January 1983

DTIC
ELECTE
MAY 23 1983
S B

Approved for public release,
distribution unlimited.

DTIC FILE COPY

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR- 83-0414	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ON LIMITING DISTRIBUTIONS OF ORDER STATISTICS WITH VARIABLE RANKS FROM STATIONARY SEQUENCES		5. TYPE OF REPORT & PERIOD COVERED TECHNICAL
7. AUTHOR(s) Shihong Cheng		6. PERFORMING ORG. REPORT NUMBER TR #25
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics University of North Carolina Chapel Hill NC 27514		8. CONTRACT OR GRANT NUMBER(s) F49620-82-C-0009
11. CONTROLLING OFFICE NAME AND ADDRESS Mathematical & Information Sciences Directorate Air Force Office of Scientific Research Bolling AFB DC 20332		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS PE61102F; 2304/A5
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE JAN 83
		13. NUMBER OF PAGES 28
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		15a. DECLASSIFICATION DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Order statistics; stationary sequences; limiting distribution; variable ranks.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) SEE REVERSE		

83 05 23 046

DD FORM 1 JAN 73 1473

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ITEM #20, CONTINUED:

Let $\{X_n\}$ be a stationary sequence and $X_1^{(n)} \leq \dots \leq X_n^{(n)}$ be the order statistics of $X_1, \dots, X_n$. In this paper, the limiting distribution of $X_{k_n}^{(n)}$, where $k_n \rightarrow \infty$, $k_n/n \rightarrow \lambda$, $0 \leq \lambda \leq 1$ is discussed under distributional mixing conditions. For stationary normal sequences, the limiting distribution of $X_{k_n}^{(n)}$, where $k_n/n \rightarrow \lambda \in (0,1)$, is a normal with mean zero and variance

$$\sigma_\lambda^2 = 1 + \frac{1}{\pi\lambda(1-\lambda)} \sum_{n=0}^{\infty} \int_0^n \frac{r_n \exp\{-a_\lambda^2/(1+r)\}}{(1-r^2)^{1/2}} dr$$

if the covariance $\{r_n\}$ converges to zero as fast as $n^{-\rho}$, $\rho > 4$, a_λ being the λ -percentile of the standard normal distribution.

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	



ON LIMITING DISTRIBUTIONS OF ORDER STATISTICS
WITH VARIABLE RANKS FROM STATIONARY SEQUENCES

Shihong Cheng
Peking University and University of North Carolina

Abstract

Let $\{X_n\}$ be a stationary sequence and $X_1^{(n)} \leq \dots \leq X_n^{(n)}$ be the order statistics of $X_1, \dots, X_n$. In this paper, the limiting distribution of $X_{k_n}^{(n)}$, where $k_n \rightarrow \infty$, $k_n/n \rightarrow \lambda$, $0 \leq \lambda \leq 1$ is discussed under distributional mixing conditions. For stationary normal sequences, the limiting distribution of $X_{k_n}^{(n)}$, where $k_n/n \rightarrow \lambda \in (0,1)$, is a normal with mean zero and variance

$$\sigma_\lambda^2 = 1 + \frac{1}{\pi\lambda(1-\lambda)} \sum_{n=0}^{\infty} \int_0^n \frac{\exp\{-a_\lambda^2/(1+r)\}}{(1-r^2)^{1/2}} dr$$

if the covariance $\{r_n\}$ converges to zero as fast as $n^{-\rho}$, $\rho > 4$, a_λ being the λ -percentile of the standard normal distribution.

Keywords: Order statistics, stationary sequences, limiting distribution, variable ranks.

This research has been supported by AFOSR Contract No. F49620-82-C-0009.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DTIC
This technical report has been reviewed and is
approved for publication in the ADP/AFR 100-12.
Distribution is unlimited.
MATTHEW J. KEISER
Chief, Technical Information Division

Let $\{X_n\}$ be a sequence of random variables and $X_1^{(n)} \leq \dots \leq X_n^{(n)}$ be the order statistics of $X_1, \dots, X_n$. In this paper it is assumed that the sequence $\{X_n\}$ is stationary and that the ranks k_n of the order statistics $\{X_{k_n}^{(n)}\}$ satisfy the following condition:

$$k_n \rightarrow \infty, n - k_n \rightarrow \infty, k_n/n \rightarrow \lambda, 0 \leq \lambda \leq 1.$$

Since the case $\lambda=1$ is easily transformed to the case $\lambda=0$, we discuss only the cases:

$$(0.1) \quad k_n \rightarrow \infty, k_n/n \rightarrow \lambda, 0 \leq \lambda < 1.$$

The case $\lambda=0$ has been discussed by Watts, Rootzén and Leadbetter [7]. The case $0 < \lambda < 1$ has been discussed by the present author [2], but the mixing condition in [2] is hard to check. Here we consider the cases $\lambda=0$ and $0 < \lambda < 1$ simultaneously, under a distributional mixing condition used by Leadbetter [4].

§1. Notation, assumptions, and introduction

Let $\{X_n\}$ be a stationary sequence with finite dimensional distribution functions, $\{F_{j_1 \dots j_p}(x_1, \dots, x_p), 1 \leq j_1 < j_2 < \dots\}$, and, in particular, marginal distribution function $F_1(x) = F(x)$. Suppose that $\{u_n\}$ is a real sequence such that

$$(1.1) \quad \lim_{n \rightarrow \infty} \frac{n}{\sqrt{k_n}} [F(u_n) - \frac{k_n}{n}] = u\sqrt{1-\lambda}, \quad -\infty < u < \infty.$$

The distributional mixing coefficients of $\{X_n\}$ with $\{u_n\}$ are defined by

$$\alpha(n, \ell) = \sup \{ |F_{i_1 \dots i_p j_1 \dots j_q}(u_n) - F_{i_1 \dots i_p}(u_n) F_{j_1 \dots j_q}(u_n)| : 1 \leq i_1 < \dots < i_p < j_1 < \dots < j_q \leq n, j_1 - i_p \geq \ell \}$$

where $F_{j_1 \dots j_p}(u_n) \equiv F_{j_1 \dots j_p}(u_n, \dots, u_n)$ for any $1 \leq j_1 < j_2 < \dots < j_p \leq n$. Let $\sigma(A_1, \dots, A_k)$ denote the field generated by sets $A_1, \dots, A_k$ and

$$\beta(n, \ell) = \sup \{ |p(AB) - p(A)p(B)| : A \in \sigma(\{X_j \leq u_n\}, j=1, \dots, k), B \in \sigma(\{X_j \leq u_n\}, j=k+\ell, \dots, n), 1 \leq k < k+\ell \leq n \}$$

Lemma 1.1 For any measurable sets $A_1, \dots, A_k, B_1, \dots, B_\ell$ let

$$\alpha = \sup\{|P(A_{i_1} \dots A_{i_s} B_{j_1} \dots B_{j_t}) - P(A_{i_1} \dots A_{i_s})P(B_{j_1} \dots B_{j_t})| :$$

$$1 \leq i_1 < \dots < i_s \leq k, 1 \leq j_1 < \dots < j_t \leq \ell\} .$$

Then we have

$$(1.2) \quad |P(A_{i_1} \dots A_{i_s} B_{j_1} \dots B_{j_t}) - P(A_{i_1} \dots A_{i_s})P(B_{j_1} \dots B_{j_t})| \leq 2^{s+t} \alpha$$

for any $s, 1 \leq s \leq k$ and $t, 1 \leq t \leq \ell$, where

$$A_{i_1} \dots A_{i_s} = \left[\bigcap_{i \in \{i_1, \dots, i_s\}} \bar{A}_i \right] \cap \left[\bigcap_{i \in \{i_1, \dots, i_s\}} A_i \right]$$

$$B_{j_1} \dots B_{j_t} = \left[\bigcap_{j \in \{j_1, \dots, j_t\}} \bar{B}_j \right] \cap \left[\bigcap_{j \in \{j_1, \dots, j_t\}} B_j \right] .$$

Proof: It is easy to show that for any sets $S, S_1, \dots, S_n$,

$$P(\bar{S} \bar{S}_1 \dots \bar{S}_n) = P(S) - \sum_{p=1}^n (-1)^{p-1} \sum_{1 \leq i_1 < \dots < i_p \leq n} P(SS_{i_1} \dots S_{i_p}) .$$

Hence if $\{i_1, \dots, i_s\} = \{1, \dots, s\}$, $\{j_1, \dots, j_t\} = \{1, \dots, t\}$, (1.2) is obtained from

$$\begin{aligned} & |P(\bar{A}_1 \dots \bar{A}_s A_{s+1} \dots A_k \bar{B}_1 \dots \bar{B}_t B_{t+1} \dots B_\ell) - P(\bar{A}_1 \dots \bar{A}_s A_{s+1} \dots A_k)P(\bar{B}_1 \dots \bar{B}_t B_{t+1} \dots B_\ell)| \\ & \leq |P(A_{s+1} \dots A_k B_{t+1} \dots B_\ell) - P(A_{s+1} \dots A_k)P(B_{t+1} \dots B_\ell)| \\ & + \sum_{p=1}^s \sum_{1 \leq i_1 < \dots < i_p \leq s} |P(A_{i_1} \dots A_{i_p} A_{s+1} \dots A_k B_{t+1} \dots B_\ell) - P(A_{i_1} \dots A_{i_p} A_{s+1} \dots A_k)P(B_{t+1} \dots B_\ell)| \\ & + \sum_{q=1}^t \sum_{1 \leq j_1 < \dots < j_q \leq t} |P(A_{s+1} \dots A_k B_{j_1} \dots B_{j_q} B_{t+1} \dots B_\ell) - P(A_{s+1} \dots A_k)P(B_{j_1} \dots B_{j_q} B_{t+1} \dots B_\ell)| \\ & + \sum_{p=1}^s \sum_{1 \leq i_1 < \dots < i_p \leq s} \sum_{q=1}^t \sum_{1 \leq j_1 < \dots < j_q \leq t} |P(A_{i_1} \dots A_{i_p} A_{s+1} \dots A_k B_{j_1} \dots B_{j_q} B_{t+1} \dots B_\ell) \\ & \quad - P(A_{i_1} \dots A_{i_p} A_{s+1} \dots A_k)P(B_{j_1} \dots B_{j_q} B_{t+1} \dots B_\ell)| \\ & \leq \alpha \sum_{p=0}^s \binom{s}{p} \sum_{q=0}^t \binom{t}{q} \leq 2^{s+t} \alpha . \end{aligned}$$

Denote the indicator of the set A by I_A and write

$$I_{nj} = I_{\{X_j \leq u_n\}} \quad , \quad \tilde{I}_{nj} = I_{nj} - F(u_n) \quad , \quad \bar{I}_{nj} = \tilde{I}_{nj}/k_n^{1/2} \quad , \quad j=1, \dots, n.$$

Let ℓ_n and $\tilde{\ell}_n$ be two sequences of positive integers such that $\ell_n \leq \tilde{\ell}_n \leq n$.

Define

$$\bar{\xi}_{ni} = \sum_{j=(i-1)(\ell_n+\tilde{\ell}_n)+1}^{(i-1)(\tilde{\ell}_n+\ell_n)+\tilde{\ell}_n} \bar{I}_{nj} \quad , \quad \bar{\eta}_{ni} = \sum_{j=(i-1)(\tilde{\ell}_n+\ell_n)+\tilde{\ell}_n+1}^{i(\ell_n+\tilde{\ell}_n)} \bar{I}_{nj} \quad , \quad i=1, \dots, N_n$$

and $\bar{\zeta}_n = \sum_{j=N_n(\ell_n+\tilde{\ell}_n)+1}^n I_{nj}$, where $N_n = \lfloor \frac{n}{\ell_n+\tilde{\ell}_n} \rfloor$. To obtain our results we need

to discuss the limiting distributions of $\sum_{i=1}^{N_n} \bar{\xi}_{ni}$, $\sum_{i=1}^{N_n} \bar{\eta}_{ni}$ and $\bar{\zeta}_n$. As preliminaries, we obtain the following lemmas.

Lemma 1.2 The following inequalities hold:

$$(1.3) \quad \left| E e^{i \sum_{k=1}^{N_n} \bar{\xi}_{nk} t} - (E e^{i \bar{\xi}_{n1} t})^{N_n} \right| \leq (n/\tilde{\ell}_n) \cdot \beta(n, \ell_n)$$

$$(1.4) \quad \left| E e^{i \sum_{k=1}^{N_n} \bar{\eta}_{nk} t} - (E e^{i \bar{\eta}_{n1} t})^{N_n} \right| \leq 3^n \alpha(n, \ell_n) .$$

The above statements are still true if we use η_{nk} , $k=1, \dots, N_n$ instead of ξ_{nk} , $k=1, \dots, N_n$ in (1.3) and (1.4).

Proof: By Dvoretzky's lemma 5.3 in [3], it follows that

$$\begin{aligned} & \left| E e^{i \sum_{k=1}^{N_n} \bar{\xi}_{nk} t} - (E e^{i \bar{\xi}_{n1} t})^{N_n} \right| \\ & \leq \sum_{r=1}^{N_n} \left| E e^{i \sum_{k=1}^r \bar{\xi}_{nk} t} - (E e^{i \sum_{k=1}^{r-1} \bar{\xi}_{nk} t}) (E e^{i \bar{\xi}_{nr} t}) \right| \\ & \leq \sum_{r=1}^{N_n} \beta(n, \ell_n) \leq (n/\tilde{\ell}_n) \cdot \beta(n, \ell_n) . \end{aligned}$$

Noticing that

$$\left| E e^{i \sum_{k=1}^N \bar{\eta}_{nk} t} - (E e^{i \bar{\eta}_{n1} t})^N \right| \leq (n/\tilde{\ell}_n) \cdot B(n, \tilde{\ell}_n) \leq (n/\tilde{\ell}_n) \cdot B(n, \ell_n),$$

we see that (1.3) holds for $\bar{\eta}_{nk}$, $k=1, \dots, N_n$.

Write $A_1, A_2, \dots, A_{(r-1)\tilde{\ell}_n}$, $B_1, \dots, B_{\tilde{\ell}_n}$ for $\{X_1 \leq u_n\}, \dots, \{X_{\tilde{\ell}_n} \leq u_n\}$, $\{X_{\tilde{\ell}_n + \ell_n + 1} \leq u_n\}, \dots$

$\{X_{2\tilde{\ell}_n + \ell_n} \leq u_n\}, \dots, \{X_{(r-1)(\tilde{\ell}_n + \ell_n) + 1} \leq u_n\}, \dots, \{X_{r\tilde{\ell}_n + (r-1)\ell_n} \leq u_n\}$ respectively and

$$f_t(I_{A_1}, \dots, I_{A_{(r-1)\tilde{\ell}_n}}) = e^{i \sum_{k=1}^{r-1} \bar{\xi}_{nk} t}, \quad g_t(I_{B_1}, \dots, I_{B_{\tilde{\ell}_n}}) = e^{i \bar{\xi}_{nr} t}.$$

Let $f_t(p)$ be the value of the random variable $f_t(I_{A_1}, \dots, I_{A_{(r-1)\tilde{\ell}_n}})$ at such points that p of I_{A_i} , $i=1, \dots, (r-1)\tilde{\ell}_n$ are equal to 1 and all others are 0. Then we have

$$\begin{aligned} E f_t g_t &= f_t(0) g_t(0) P(\bar{A}_1 \dots \bar{A}_{(r-1)\tilde{\ell}_n} \bar{B}_1 \dots \bar{B}_{\tilde{\ell}_n}) \\ &+ f_t(0) \sum_{p=1}^{\tilde{\ell}_n} g_t(p) \sum_{1 \leq j_1 < \dots < j_p \leq \tilde{\ell}_n} P(\bar{A}_1 \dots \bar{A}_{(r-1)\tilde{\ell}_n} \cap B_{j_1} \dots B_{j_p}) \\ &+ g_t(0) \sum_{p=1}^{(r-1)\tilde{\ell}_n} f_t(p) \sum_{1 \leq j_1 < \dots < j_p \leq (r-1)\tilde{\ell}_n} P(A_{j_1} \dots A_{j_p} \cap \bar{B}_1 \dots \bar{B}_{\tilde{\ell}_n}) \\ &+ \sum_{p=1}^{(r-1)\tilde{\ell}_n} \sum_{1 \leq i_1 < \dots < i_p \leq (r-1)\tilde{\ell}_n} \sum_{q=1}^{\tilde{\ell}_n} \sum_{1 \leq j_1 < \dots < j_q \leq \tilde{\ell}_n} f_t(p) g_t(q) P(A_{i_1} \dots A_{i_p} \cap B_{j_1} \dots B_{j_q}) \end{aligned}$$

Since (1.2) holds (including $s=0$ or $t=0$) and $|f_t(p)| = |g_t(p)| = 1$ for any p, q , (1.4)

follows from

$$\left| E e^{i \sum_{k=1}^N \bar{\xi}_{nk} t} - (E e^{i \bar{\xi}_{n1} t})^N \right| \leq \sum_{r=1}^N |E f_t g_t - E f_t E g_t|$$

$$\leq \alpha(n, \ell_n) \sum_{r=1}^N \sum_{n=0}^{(r-1)\tilde{\ell}_n} \sum_{q=0}^{\tilde{\ell}_n} \binom{(r-1)\tilde{\ell}_n}{p} \binom{\tilde{\ell}_n}{q} 2^{p+q}$$

$$= \sum_{r=1}^N 3^{r\tilde{\ell}_n} \alpha(n, \ell_n) \leq 3^{n\alpha(n, \ell_n)} .$$

In the same way, we can show that

$$\left| E e^{i \sum_{k=1}^N \tilde{\eta}_{nk} t} - (E e^{i \tilde{\eta}_{n1} t})^N \right| \leq \sum_{r=1}^N 3^{r\tilde{\ell}_n} \alpha(n, \tilde{\ell}_n) \leq 3^{n\alpha(n, \ell_n)} ,$$

completing the proof of the lemma.

Lemma 1.3 If $\lim_n (n/k_n) \cdot F(u_n) = 1$, then

$$(1.5) \quad \left| \sum_{1 \leq i < j < k \leq \tilde{\ell}_n} E \tilde{I}_{ni} \tilde{I}_{nj} \tilde{I}_{nk} \right| \leq C_1 \tilde{\ell}_n^3 \alpha(n, \ell_n) + C_2 \frac{\tilde{\ell}_n^2 \ell_n^2}{n^2} + C_3 \tilde{\ell}_n \ell_n \left| \sum_{j=1}^{\ell_n-1} E \tilde{I}_{n1} \tilde{I}_{nj+1} \right|$$

where C_1, C_2, C_3 are constants.

Proof: Using stationarity of the process, we obtain

$$\sum_{1 \leq i < j < k \leq \tilde{\ell}_n} E \tilde{I}_{nk} \tilde{I}_{nj} \tilde{I}_{ni} = \sum_{s=1}^{\tilde{\ell}_n-2} \sum_{t=1}^{\tilde{\ell}_n-s-1} (\tilde{\ell}_n-s-t) E \tilde{I}_{n1} \tilde{I}_{ns+1} \tilde{I}_{ns+t+1} .$$

Since

$$\begin{aligned} E \tilde{I}_{n1} \tilde{I}_{ns+1} \tilde{I}_{ns+t+1} &= P(X_1 \leq u_n, X_{s+1} \leq u_n, X_{s+t+1} \leq u_n) - F(u_n) [P(X_1 \leq u_n, X_{s+1} \leq u_n) \\ &\quad + P(X_1 \leq u_n, X_{s+t+1} \leq u_n) + P(X_{s+1} \leq u_n, X_{s+t+1} \leq u_n)] + 2F^3(u_n) . \end{aligned}$$

it follows that

$$\begin{aligned} & \sum_{s=\ell_n}^{\tilde{\ell}_n-2} \sum_{t=1}^{\tilde{\ell}_n-s-1} (\tilde{\ell}_n-s-t) E \tilde{I}_{n1} \tilde{I}_{ns+1} \tilde{I}_{ns+t+1} \\ & \leq \sum_{s=\ell_n}^{\tilde{\ell}_n-2} \sum_{t=1}^{\tilde{\ell}_n-s-1} \tilde{\ell}_n \{ |F_{1,s+1,s+t+1}(u_n) - F(u_n)F_{s+1,s+t+1}(u_n)| + |F_{1,s+1}(u_n) - F^2(u_n)| \\ & \quad + |F_{1,s+t+1}(u_n) - F^2(u_n)| \} \leq 3\tilde{\ell}_n^3 \alpha(n, \ell_n) , \end{aligned}$$

and in the same way that

$$\left| \sum_{s=1}^{\ell_n-1} \sum_{t=\ell_n}^{\tilde{\ell}_n-s-1} (\tilde{\ell}_n-s-t) E \tilde{I}_{n1} \tilde{I}_{ns+1} \tilde{I}_{ns+t+1} \right| \leq 3 \tilde{\ell}_n^3 \alpha(n, \ell_n) .$$

Noticing that

$$E |\tilde{I}_{n1} \tilde{I}_{ns+1}| = [1-2F(u_n)] E \tilde{I}_{n1} \tilde{I}_{ns+1} + 4F^2(u_n) [1-F(u_n)]^2 , \text{ we have}$$

$$\begin{aligned} & \left| \sum_{s=1}^{\ell_n-1} \sum_{t=1}^{\ell_n-1} (\tilde{\ell}_n-s-t) E \tilde{I}_{n1} \tilde{I}_{ns+1} \tilde{I}_{ns+t+1} \right| \leq \tilde{\ell}_n \ell_n \sum_{s=1}^{\ell_n-1} E |\tilde{I}_{n1} \tilde{I}_{ns+1}| \\ & \leq 4 \tilde{\ell}_n \ell_n^2 F^2(u_n) [1-F(u_n)]^2 + \tilde{\ell}_n \ell_n |1-2F(u_n)| \left| \sum_{s=1}^{\ell_n-1} E \tilde{I}_{n1} \tilde{I}_{ns+1} \right| \\ & \leq C_2 \frac{\tilde{\ell}_n \ell_n^2 k^2}{n^2} + C_3 \tilde{\ell}_n \ell_n \left| \sum_{j=1}^{\ell_n-1} E \tilde{I}_{n1} \tilde{I}_{nj+1} \right| , \end{aligned}$$

so that the lemma is proved.

Lemma 1.4 If $\lim_n (n/k_n) \cdot F(u_n) = 1$, then

$$\begin{aligned} \left| \sum_{1 \leq i \leq j \leq k \leq \ell \leq \tilde{\ell}_n} E \tilde{I}_{ni} \tilde{I}_{nj} \tilde{I}_{nk} \tilde{I}_{n\ell} \right| & \leq C_1 \tilde{\ell}_n^4 \alpha(n, \ell_n) + C_2 \tilde{\ell}_n^2 \left(\sum_{s=1}^{\ell_n-1} E \tilde{I}_{n1} \tilde{I}_{ns+1} \right)^2 \\ & + C_3 \tilde{\ell}_n \ell_n^3 k^2 / n^2 + C_4 \tilde{\ell}_n \ell_n^2 \left| \sum_{s=1}^{\ell_n-1} E \tilde{I}_{n1} \tilde{I}_{ns+1} \right| \end{aligned}$$

where C_1, C_2, C_3, C_4 are constants.

Proof: Using stationarity of the process we obtain

$$\sum_{i < j < k < \ell} E \tilde{I}_{ni} \tilde{I}_{nk} \tilde{I}_{nk} \tilde{I}_{n\ell} = \sum_{s=1}^{\tilde{\ell}_n-3} \sum_{u=1}^{\tilde{\ell}_n-s-2} \sum_{t=1}^{\tilde{\ell}_n-s-u-1} (\tilde{\ell}_n-s-t-u) E \tilde{I}_{n1} \tilde{I}_{ns+1} \tilde{I}_{ns+t+1} \tilde{I}_{ns+t+u+1} .$$

Since

$$\begin{aligned} E \tilde{I}_{n1} \tilde{I}_{ns+1} \tilde{I}_{ns+t+1} \tilde{I}_{ns+t+u+1} & = F_{1,s+1,s+t+1,s+t+u+1}(u_n) - F(u_n) [F_{1,s+1,s+t+1}(u_n) \\ & + F_{1,s+1,s+t+u+1}(u_n) + F_{1,s+t+1,s+t+u+1}(u_n) + F_{1,t+1,t+u+1}(u_n)] \\ & + F^2(u_n) [F_{1,s+1}(u_n) + F_{1,s+t+1}(u_n) + F_{1,s+t+u+1}(u_n) + F_{1,t+1}(u_n) + F_{1,t+u+1}(u_n) \\ & + F_{1,u+1}(u_n)] - 3F^4(u_n) , \end{aligned}$$

it follows in the same way as in the proof of lemma 1.2 that

$$\left| \sum_{s=\ell_n}^{\ell_n-3} \sum_{u=1}^{\ell_n-s-2} \sum_{t=1}^{\ell_n-s-u-1} (\tilde{\ell}_n^{-s-t-u}) E \tilde{I}_{n1} \tilde{I}_{ns+1} \tilde{I}_{ns+t+1} \tilde{I}_{ns+t+u+1} \right| \leq 7 \tilde{\ell}_n^{4\alpha(n, \ell_n)} .$$

$$\left| \sum_{s=1}^{\ell_n-1} \sum_{u=\ell_n}^{\ell_n-s-2} \sum_{t=1}^{\ell_n-s-u-1} (\tilde{\ell}_n^{-s-t-u}) E \tilde{I}_{n1} \tilde{I}_{ns+1} \tilde{I}_{ns+t+1} \tilde{I}_{ns+t+u+1} \right| \leq 7 \tilde{\ell}_n^{4\alpha(n, \ell_n)} .$$

Writing

$$\begin{aligned} E \tilde{I}_{n1} \tilde{I}_{ns+1} \tilde{I}_{ns+t+1} \tilde{I}_{ns+t+u+1} &= [F_{1,s+1,s+t+1,s+t+u+1}(u_n) - F_{1,s+1}(u_n) F_{1,t+1}(u_n)] \\ &- F(u_n) [F_{1,s+1,s+t+1}(u_n) - F(u_n) F_{1,s+1}(u_n)] - F(u_n) [F_{1,s+t+1,s+t+u+1}(u_n) - F(u_n) F_{1,t+1}(u_n)] \\ &+ F^2(u_n) [F_{1,s+t+1}(u_n) - F^2(u_n)] + F^2(u_n) [F_{1,s+t+u+1}(u_n) - F^2(u_n)] \\ &+ F^2(u_n) [F_{1,t+1}(u_n) - F^2(u_n)] + F^2(u_n) [F_{1,t+u+1}(u_n) - F^2(u_n)] \\ &- F(u_n) [F_{1,s+1,s+t+u+1}(u_n) - F(u_n) F_{1,s+1}(u_n)] \\ &- F(u_n) [F_{1,t+1,t+u+1}(u_n) - F(u_n) F_{1,t+1}(u_n)] + [F_{1,s+1}(u_n) - F^2(u_n)] [F_{1,t+1}(u_n) - F^2(u_n)] , \end{aligned}$$

we also have

$$\left| \sum_{s=1}^{\ell_n-1} \sum_{u=1}^{\ell_n-1} \sum_{t=\ell_n}^{\ell_n-s-u-1} (\tilde{\ell}_n^{-s-t-u}) E \tilde{I}_{n1} \tilde{I}_{ns+1} \tilde{I}_{ns+t+1} \tilde{I}_{ns+t+u+1} \right| \leq 9 \tilde{\ell}_n^{4\alpha(n, \ell_n)} + \tilde{\ell}_n^2 \left(\sum_{s=1}^{\ell_n-1} E \tilde{I}_{n1} \tilde{I}_{ns+1} \right)^2$$

Finally we can show that

$$\left| \sum_{s=1}^{\ell_n-1} \sum_{t=1}^{\ell_n-1} \sum_{u=1}^{\ell_n-1} (\tilde{\ell}_n^{-s-t-u}) E \tilde{I}_{n1} \tilde{I}_{ns+1} \tilde{I}_{ns+t+1} \tilde{I}_{ns+t+u+1} \right| \leq C_4 \tilde{\ell}_n \ell_n^2 \sum_{s=1}^{\ell_n-1} E \tilde{I}_{n1} \tilde{I}_{ns+1} + C_5 \tilde{\ell}_n \ell_n^3 k_n^2 / n^2 .$$

Hence the lemma is proved.

§2. Some limit theorems

We introduce the following assumptions:

Assumption I: For some sequence $\{\ell_n\}$ of positive integers,

$$(2.1) \quad (n/k_n^{1/2}) \beta(n, \ell_n) \rightarrow 0 \quad (n \rightarrow \infty)$$

$$(2.2) \quad \lim_n (n/k_n) \cdot \sum_{j=1}^{[k_n^{1/2}] - 1} E \tilde{I}_{n1} \tilde{I}_{nj+1} = \sigma$$

$$(2.3) \quad \lim_n (n/k_n^{3/2}) \cdot \sum_{j=1}^{[k_n^{1/2}] - 1} j E \tilde{I}_{n1} \tilde{I}_{nj+1} = 0.$$

Assumption II. For some sequence $\{\ell_n\}$ of positive integers,

$$(2.4) \quad 3^n \alpha(n, \ell_n) \rightarrow 0 \quad (n \rightarrow \infty)$$

and (2.2), (2.3) hold.

It is obvious that the constant σ in (2.2) may be different if the sequence $\{u_n\}$ is changed. But we can show that σ must be the same for any $\{u_n\}$ satisfying (1.1) with some real u .

Lemma 2.1 If (2.1) or (2.4) holds for some $\ell_n = o(k_n^{1/2})$, $\ell_n \rightarrow \infty$, then (2.2) and (2.3) hold, if and only if

$$(2.2)' \quad \lim_n (n/k_n) \cdot \sum_{j=1}^{\ell_n - 1} E \tilde{I}_{n1} \tilde{I}_{nj+1} = \sigma$$

$$(2.3)' \quad \lim_n (n/k_n^{3/2}) \cdot \sum_{j=1}^{\ell_n - 1} j E \tilde{I}_{n1} \tilde{I}_{nj+1} = 0.$$

Furthermore, if (1.1) holds for some $u \in \mathbb{R}$, we can use

$$(2.2)'' \quad \lim_n (n/k_n) \cdot \sum_{j=1}^{\ell_n - 1} [P(X_1 \leq a(k_n/n), X_{j+1} \leq a(k_n/n)) - (k_n/n)^2] = \sigma$$

$$(2.3)'' \quad \lim_n (n/k_n^{3/2}) \cdot \sum_{j=1}^{\ell_n - 1} j [P(X_1 \leq a(k_n/n), X_{j+1} \leq a(k_n/n)) - (k_n/n)^2] = 0$$

instead of (2.2)' and (2.3)' respectively in the above statements. In (2.2)'' and (2.3)'',

$$a(k_n/n) = \begin{cases} a_n - 0 & \text{if } F(a_n) - k_n/n \leq k_n/n - F(a_n - 0) \\ a_n & \text{if } F(a_n) - k_n/n < k_n/n - F(a_n - 0) \end{cases}$$

where a_n is a real number such that $F(a_n - 0) \leq k_n/n \leq F(a_n)$, and the event $\{X_j \leq a_n - 0\}$ is defined as $\{X_j < a_n\}$.

Proof: The first part of the lemma follows from

$$| (n/k_n)^{\frac{1}{2}} \cdot \sum_{j=\ell_n}^{[k_n^{1/2}]-1} E \tilde{I}_{n1} \tilde{I}_{nj+1} | \leq (n/k_n)^{\frac{1}{2}} \cdot [k_n^{1/2}] \alpha(n, \ell_n) \rightarrow 0$$

$$| (n/k_n^{3/2}) \cdot \sum_{j=\ell_n}^{[k_n^{1/2}]-1} j E \tilde{I}_{n1} \tilde{I}_{nj+1} | \leq (n/k_n^{3/2}) \cdot [k_n^{1/2}]^2 \alpha(n, \ell_n) \rightarrow 0.$$

Now we show the second part. By the definition of $a(k_n/n)$, it is easy to see that

$|F(a(k_n/n)) - k_n/n| \leq |F(x) - k_n/n|$ for any x . Therefore we have

$$\begin{aligned} & | (n/k_n)^{\frac{1}{2}} \cdot \sum_{j=1}^{\ell_n-1} E \tilde{I}_{n1} \tilde{I}_{nj+1} - (n/k_n)^{\frac{1}{2}} \cdot \sum_{j=1}^{\ell_n-1} [P(X_1 \leq a(k_n/n), X_{j+1} \leq a(k_n/n)) - (k_n/n)^2] | \\ & \leq (n/k_n)^{\frac{1}{2}} \cdot \sum_{j=1}^{\ell_n-1} [|P(X_1 \leq u_n, X_{j+1} \leq u_n) - P(X_1 \leq a(k_n/n), X_{j+1} \leq a(k_n/n))| + |F^2(u_n) - (k_n/n)^2|] \\ & \leq 2(n/k_n)^{\frac{1}{2}} \cdot \sum_{j=1}^{\ell_n-1} [|F(u_n) - F(a(k_n/n))| + |F(u_n) - k_n/n|] \\ & \leq 6(\ell_n/k_n^{1/2}) \cdot (n/k_n^{1/2}) \cdot |F(u_n) - k_n/n| \rightarrow 0. \end{aligned}$$

This proves that (2.2)'' and (2.2)' are equivalent. In the same way, we can show that (2.3)'' is equivalent to (2.3)'.

Let
$$\bar{S}_n = \sum_{j=1}^n \bar{I}_{nj} = \sum_{k=1}^{N_n} \bar{\xi}_{nk} + \sum_{k=1}^{N_n} \bar{\eta}_{nk} + \bar{\zeta}_n.$$

We now start to discuss the limiting distribution of $\bar{S}_n$.

Lemma 2.2 If assumption I or II holds for some $\ell_n = o(k_n^{1/2})$ and

$$(2.5) \quad \lim_n nF(u_n)/k_n = 1,$$

then

$$(2.6) \quad P(\bar{S}_n \leq x) \xrightarrow{c} \phi_\lambda(x),$$

if and only if

$$(2.7) \quad \lim_n (E e^{i \bar{\xi}_{n1} t} N_n) = \psi(t),$$

when (2.6) or (2.7) holds, we have

$$(2.8) \quad \psi(t) = \int e^{itx} d\phi_\lambda(x).$$

Proof: Let $\tilde{\ell}_n = [k_n^{1/2}]$. If $n - N_n(\ell_n + \tilde{\ell}_n) < \ell_n$, we have

$$0 \leq E \bar{\xi}_n^2 \leq \ell_n^2/k_n \rightarrow 0.$$

If $n - N_n(\ell_n + \tilde{\ell}_n) \geq \ell_n$, we also have

$$\begin{aligned} 0 < E \bar{\xi}_n^2 &= \frac{1}{k_n} \{ [n - N_n(\ell_n + \tilde{\ell}_n)] F(u_n) [1 - F(u_n)] + 2 \sum_{j=1}^{n - N_n(\ell_n + \tilde{\ell}_n)} [n - N_n(\ell_n + \tilde{\ell}_n) - j] E \tilde{I}_{n1} \tilde{I}_{nj+1} \} \\ &\leq C/N_n + \frac{2}{N_n} \left| \frac{n}{k_n} \sum_{j=1}^{\tilde{\ell}_n - 1} E \tilde{I}_{n1} \tilde{I}_{nj+1} \right| + \frac{2}{k_n} \left| \sum_{j=1}^{\ell_n - 1} j E \tilde{I}_{n1} \tilde{I}_{nj+1} \right| + 2 \frac{(\tilde{\ell}_n + \ell_n)^2}{k_n} \alpha(n, \ell_n) \rightarrow 0. \end{aligned}$$

Hence by Chebyshev's inequality, it follows that

$$\bar{\xi} \rightarrow 0 \quad [P].$$

Since $E \bar{\eta}_{n1} = 0$, we have

$$N_n |E e^{i \bar{\eta}_{n1} t} - 1| \leq N_n E \bar{\eta}_{n1}^2 \cdot t^2/2 = (t^2/2) \cdot (N_n/k_n) \{ \ell_n F(u_n) [1 - F(u_n)] + 2 \sum_{j=1}^{\ell_n - 1} (\ell_n - j) E \tilde{I}_{n1} \tilde{I}_{nj+1} \}$$

$$t^2 \left\{ \frac{\ell_n}{\tilde{\ell}_n} \frac{n}{k_n} F(u_n) |1 - F(u_n)| + 2 \frac{\ell_n}{\tilde{\ell}_n} \left| \frac{n}{k_n} \sum_{j=1}^{\ell_n-1} E \tilde{I}_{n1} \tilde{I}_{nj+1} \right| + 2 \left| \frac{n}{k_n^{3/2}} \sum_{j=1}^{\ell_n-1} i E \tilde{I}_{n1} \tilde{I}_{nj+1} \right| \right\} \rightarrow 0,$$

i.e. $E e^{i \tilde{\eta}_{n1} t} = 1 + o\left(\frac{1}{N_n}\right)$. Therefore $\lim_n (E e^{i \tilde{\eta}_{n1} t})^{N_n} = 1$. By using lemma 1.2, this implies that

$$\lim_n E e^{i \sum_{k=1}^N \tilde{\eta}_{nk} t} = 1, \text{ i.e.}$$

$$\sum_{k=1}^N \tilde{\eta}_{nk} \rightarrow 0[P].$$

From the above argument, it follows that (2.6) is equivalent to $P\left(\sum_{k=1}^{N_n} \tilde{\xi}_{nk} \leq x\right) \xrightarrow{c} \Phi(x)$, i.e.

$$\lim_n E e^{i \sum_{k=1}^N \tilde{\xi}_{nk} t} = \psi(t),$$

where $\psi(t)$ is defined by (2.8). Using lemma 1.2 again, we see that (2.6) and (2.7) are equivalent. Hence the lemma is proved.

Lemma 2.3 If assumption I or II holds for some $\ell_n = o(\min(k_n^{1/2}, \frac{n^{1/2}}{k_n^{1/4}}))$, and (2.5) holds, then

$$\lim_n (E e^{i \tilde{\xi}_{n1} t})^{N_n} = e^{-[\sigma + \frac{1}{2}(1-\lambda)]t^2}$$

Proof: From Taylor's formula, it is easy to show that

$$|e^{it} - \sum_{k=0}^n \frac{(it)^k}{k!}| \leq \frac{|t|^{n+1}}{(n+1)!}, \quad n=0,1,2,\dots, \quad \forall t$$

Therefore, taking $n=3$, we have

$$|E e^{i \tilde{\xi}_{n1} t} - 1 - it E \tilde{\xi}_{n1} - \frac{(it)^2}{2!} E \tilde{\xi}_{n1}^2 - \frac{(it)^3}{3!} E \tilde{\xi}_{n1}^3|$$

$$\leq E |e^{i\bar{\xi}_{nl}t} - 1 - it\bar{\xi}_{nl} - \frac{(it)^2}{2!} \bar{\xi}_{nl}^2 - \frac{(it)^3}{3!} \bar{\xi}_{nl}^3| < \frac{|t|^4 E \bar{\xi}_{nl}^4}{4!}.$$

Noticing that $E\bar{\xi}_{nl} = 0$, the lemma can be proved if we show that

$$E\bar{\xi}_{nl}^2 = \frac{1}{N_n} [(1-\lambda) + 2\sigma] + o\left(\frac{1}{N_n}\right),$$

$$E\bar{\xi}_{nl}^3 = o\left(\frac{1}{N_n}\right), \quad E\bar{\xi}_{nl}^4 = o\left(\frac{1}{N_n}\right),$$

which is equivalent to

$$(2.9) \quad N_n E\bar{\xi}_{nl}^2 \rightarrow (1-\lambda) + 2\sigma,$$

$$(2.10) \quad N_n E\bar{\xi}_{nl}^3 \rightarrow 0,$$

$$(2.11) \quad N_n E\bar{\xi}_{nl}^4 \rightarrow 0.$$

Since under assumption I or II,

$$\begin{aligned} N_n E\bar{\xi}_{nl}^2 &= \frac{N_n}{k_n} \{ \tilde{\mathcal{L}}_n F(u_n) [1-F(u_n)] + 2 \sum_{j=1}^{\tilde{\mathcal{L}}_n-1} (\tilde{\mathcal{L}}_n - j) E\tilde{I}_{nl} \tilde{I}_{nj+1} \} \\ &= \frac{N_n \tilde{\mathcal{L}}_n F(u_n) [1-F(u_n)]}{k_n} + \frac{2N_n \tilde{\mathcal{L}}_n}{k_n} \sum_{j=1}^{\tilde{\mathcal{L}}_n-1} E\tilde{I}_{nl} \tilde{I}_{nj+1} - \frac{2N_n}{k_n} \sum_{j=1}^{\tilde{\mathcal{L}}_n-1} j E\tilde{I}_{nl} \tilde{I}_{nj+1} \rightarrow (1-\lambda) + 2\sigma, \end{aligned}$$

(2.9) is obvious. To prove (2.10), we expand

$$(2.12) \quad N_n E\bar{\xi}_{nl}^3 = \frac{N_n}{k_n^{3/2}} \sum_{j=1}^{\tilde{\mathcal{L}}_n} E\tilde{I}_{nj}^3 + \frac{3N_n}{k_n^{3/2}} \sum_{i \neq j} E\tilde{I}_{ni}^2 \tilde{I}_{nj} + \frac{6N_n}{k_n^{3/2}} \sum_{i < j < k} E\tilde{I}_{ni} \tilde{I}_{nj} \tilde{I}_{nk}.$$

For the first term on the right hand of (2.12), we have

$$\left| \frac{N_n}{k_n^{3/2}} \sum_{j=1}^{\tilde{\mathcal{L}}_n} E\tilde{I}_{nj}^3 \right| \leq \frac{N_n}{k_n} |E\tilde{I}_{nj}^3| \leq \frac{1}{k_n^{1/2}} |1-2F(u_n)| \cdot \frac{n}{k_n} F(u_n) [1-F(u_n)] \rightarrow 0.$$

Noticing that $E\tilde{I}_{ni}^2 \tilde{I}_{nj} = [1-2F(u_n)] E\tilde{I}_{ni} \tilde{I}_{nj}$, we have

$$\sum_{i \neq j} E \tilde{I}_{ni}^2 \tilde{I}_{nj} = 2[1-F(u_n)] \sum_{j=1}^{\tilde{\ell}_n-1} (\tilde{\ell}_n-j) E \tilde{I}_{n1} \tilde{I}_{nj+1}.$$

Therefore, for the second term on the right hand of (2.12), it follows that

$$\left| \frac{3N_n}{k_n^{3/2}} \sum_{i \neq j} E \tilde{I}_{ni}^2 \tilde{I}_{nj} \right| \leq \frac{C}{k_n^{1/2}} \left| \frac{N_n}{k_n} \sum_{j=1}^{\tilde{\ell}_n-1} (\tilde{\ell}_n-j) E \tilde{I}_{n1} \tilde{I}_{nj+1} \right| \rightarrow 0.$$

Lastly, by using lemma 1.3, we obtain

$$\left| \frac{6N_n}{k_n^{3/2}} \sum_{i < j < k} E \tilde{I}_{ni} \tilde{I}_{nj} \tilde{I}_{nk} \right| \leq C_1 \frac{\tilde{\ell}_n^2}{k_n} \cdot \frac{n}{k_n^{1/2}} \alpha(n, \ell_n) + C_2 \frac{k_n^{1/2} \ell_n^2}{n} + C_3 \frac{\ell_n}{k_n^{1/2}} \cdot \frac{n}{k_n} \sum_{j=1}^{\ell_n-1} E \tilde{I}_{n1} \tilde{I}_{nj+1} \rightarrow 0.$$

Hence (2.10) holds. To prove (2.12), expand

$$\begin{aligned} (2.13) \quad N_n E \tilde{I}_{n1}^4 &= \frac{N_n \tilde{\ell}_n}{k_n^2} E \tilde{I}_{n1}^4 + \frac{4N_n}{k_n^2} \sum_{i \neq j} E \tilde{I}_{ni}^3 \tilde{I}_{nj} + \frac{6N_n}{k_n^2} \sum_{i < j} E \tilde{I}_{ni}^2 \tilde{I}_{nj}^2 \\ &\quad + \frac{12N_n}{k_n^2} \sum_{\substack{j \neq i, k \neq i \\ j < k}} E \tilde{I}_{ni}^2 \tilde{I}_{nj} \tilde{I}_{nk} + \frac{24N_n}{k_n^2} \sum_{i < j < k < \ell} E \tilde{I}_{ni} \tilde{I}_{nj} \tilde{I}_{nk} \tilde{I}_{n\ell}. \end{aligned}$$

For the first term on the right hand of (2.13), it follows that

$$\frac{N_n \tilde{\ell}_n}{k_n^2} E \tilde{I}_{n1}^4 = \frac{N_n \tilde{\ell}_n}{k_n^2} F(u_n) [1-F(u_n)] [1-3F(u_n) + 3F^2(u_n)] \leq \frac{C}{k_n} \rightarrow 0.$$

Since $E \tilde{I}_{ni}^3 \tilde{I}_{nj} = [1-3F(u_n) + 3F^2(u_n)] E \tilde{I}_{ni} \tilde{I}_{nj}$, for the second term, we also have

$$\left| \frac{N_n}{k_n^2} \sum_{i \neq j} E \tilde{I}_{ni}^3 \tilde{I}_{nj} \right| \leq \frac{C}{k_n} \left| \frac{N_n}{k_n} \sum_{j=1}^{\tilde{\ell}_n-1} (\tilde{\ell}_n-j) E \tilde{I}_{n1} \tilde{I}_{nj+1} \right| \rightarrow 0.$$

By using $E \tilde{I}_{ni}^2 \tilde{I}_{nj}^2 = F^2(u_n) [1-F(u_n)]^2 + [1-2F(u_n)]^2 E \tilde{I}_{ni} \tilde{I}_{nj}$, it is seen that

$$\frac{N_n}{k_n^2} \sum_{i < j} E \tilde{I}_{ni}^2 \tilde{I}_{nj}^2 \leq \frac{N_n \tilde{\ell}_n^2}{k_n^2} F^2(u_n) [1-F(u_n)]^2 + \frac{N_n C}{k_n^2} \left| \sum_{j=1}^{\tilde{\ell}_n-1} (\tilde{\ell}_n-j) E \tilde{I}_{n1} \tilde{I}_{nj+1} \right| \rightarrow 0.$$

Noticing that $E\tilde{I}_{ni}^2\tilde{I}_{nj}\tilde{I}_{nk} = [1-2F(u_n)]E\tilde{I}_{ni}\tilde{I}_{nj}\tilde{I}_{nk} + 2F^2(u_n)[1-F(u_n)]E\tilde{I}_{nj}\tilde{I}_{nk}$ we obtain, for the fourth term,

$$\left| \frac{N}{k_n} \sum_{\substack{j \neq i, k \neq i \\ j < k}} E\tilde{I}_{ni}^2\tilde{I}_{nj}\tilde{I}_{nk} \right| \leq \frac{C_1}{k_n^{1/2}} \left| \frac{N}{k_n^{3/2}} \sum_{i < j < k} E\tilde{I}_{ni}\tilde{I}_{nj}\tilde{I}_{nk} \right| + \frac{C_2}{N} \cdot \frac{k_n}{n} \left| \frac{N}{k_n} \sum_{j=1}^{\ell_n-1} (\ell_n-i) \cdot \tilde{I}_{ni}\tilde{I}_{ni+1} \right| \rightarrow 0$$

Lastly, by lemma 1.4, it follows that

$$\left| \frac{N}{k_n} \sum_{i < j < k < \ell} E\tilde{I}_{ni}\tilde{I}_{nj}\tilde{I}_{nk}\tilde{I}_{n\ell} \right|$$

$$C_1 \frac{\ell_n^5}{k_n^{3/2}} \cdot \frac{n}{k_n^{1/2}} \cdot (n, \ell_n) + C_2 \frac{N}{k_n} \frac{\ell_n^2}{n} \left| \sum_{s=1}^{\ell_n-1} E\tilde{I}_{ni}\tilde{I}_{ns+1} \right|^2 + C_3 \frac{\ell_n^{1/2}}{k_n} \cdot \frac{n}{k_n^{1/2}} \cdot \frac{\ell_n}{n} + C_4 \frac{\ell_n^2}{k_n} \cdot \frac{n}{k_n} \sum_{s=1}^{\ell_n-1} E\tilde{I}_{ni}\tilde{I}_{ns+1} \rightarrow 0.$$

Hence (2.12) holds, and the lemma is proved.

From lemma 2.2 and 2.3, we obtain

Theorem 2.4 If assumption I or II holds for some $\ell_n = o(\min(k_n^{1/2}, \frac{n^{1/2}}{k_n^{1/4}}))$, and (2.5) holds, then (2.6) holds with

$$\phi_\lambda(x) = \frac{1}{(2\pi)^{1/2}\sigma_\lambda} \int_{-\infty}^x \exp(-\frac{t^2}{2\sigma_\lambda^2}) dt = \Phi(\frac{x}{\sigma_\lambda}),$$

where $\sigma_\lambda^2 = (1-\lambda)+2\sigma$, $\Phi(x)$ is the normal distribution function with mean 0 and variance 1, and when $\sigma_\lambda = 0$, $\Phi(\frac{x}{\sigma_\lambda})$ is defined to be 1 for $x \geq 0$ and 0 for $x < 0$. The above statement is still true if (2.2)', (2.3)' are used instead of (2.2), (2.3) in assumption I and II.

Furthermore, using lemma 2.1, we obtain

Theorem 2.5 If (1.1) holds, and assumption I or II holds for some

$\ell_n = o(\min(k_n^{1/2}, \frac{n^{1/2}}{k_n^{1/4}}))$, then the conclusion of Theorem 2.4 follows. The conclusion of Theorem 2.4 is still true if (2.2)", (2.3)" are used instead of (2.2), (2.3) in assumptions I and II.

It is easy to show that if for some $\ell_n = o(k_n^{1/2})$

$$(2.14) \quad \lim_{n \rightarrow \infty} \frac{n}{k_n} \sum_{j=1}^{\ell_n - 1} |E \tilde{I}_{n1} \tilde{I}_{nj+1}| = 0,$$

then (2.2) and (2.3) hold with $\sigma=0$. Therefore we obtain

Theorem 2.6 If (2.14) and one of (2.1) and (2.4) holds for some

$$\ell_n = o(\min(k_n^{1/2}, \frac{n^{1/2}}{k_n^{1/4}})), \text{ then (2.6) holds with } \Phi_\lambda(x) = \Phi\left(\frac{x}{(1-\lambda)^{1/2}}\right).$$

§5. The results for general stationary processes

An i.i.d. sequence $\{\hat{X}_n\}$ is called the associated independent sequence of a stationary sequence $\{X_n\}$ if $\hat{X}_n$ has the same marginal d.f. $F(x)$ as X_n . Smirnov [6] has shown that there are constants $a_n > 0$, b_n such that

$$(3.1) \quad P(\hat{X}_{k_n}^{(n)} \leq a_n x + b_n) \xrightarrow{c} \Psi(x)$$

if and only if

$$(3.2) \quad \frac{n}{k_n^{1/2}} [F(a_n x + b_n) - \frac{k_n}{n}] \xrightarrow{c} (1-\lambda)^{1/2} u(x)$$

where $u(x)$ is a nondecreasing, right continuous, (finite or infinite valued) real function such that $u(-\infty) = \lim_{x \rightarrow -\infty} u(x) = -\infty$, $u(\infty) = \lim_{x \rightarrow \infty} u(x) = \infty$. The relation between $\Psi(x)$ and $u(x)$ is

$$(3.3) \quad \Psi(x) = \Phi(u(x)).$$

In this paper, we will find the limiting distribution of $X_{k_n}^{(n)}$ under condition (3.2) considering only the case in which $\Phi(u(x))$ is not degenerate.

Theorem 3.1 Suppose that

1. there are $a_n > 0$, b_n such that (3.2) holds with a continuous $u(x)$,
2. for any $u_n = a_n x + b_n$, $x \in B(u(\cdot)) \equiv \{x: |u(x)| < \infty\}$, assumption I or II holds

with some $\ell_n = o(\min(k_n^{1/2}, \frac{n^{1/2}}{k_n^{1/4}}))$. Then the real σ in (2.2) is independent of x

and determined by (2.2)", and

$$(3.4) \quad P(X_{k_n}^{(n)} \leq a_n x + b_n) \rightarrow \Phi\left(\frac{(1-\lambda)^{1/2}}{\sigma_\lambda} u(x)\right), \quad \sigma_\lambda > 0.$$

Proof: According to theorem 2.5, we have

$$\begin{aligned} P(X_{k_n}^{(n)} \leq a_n x + b_n) &= P(\bar{S}_n \geq \frac{n}{k_n^{1/2}} [\frac{k_n}{n} - F(a_n x + b_n)]) \\ &\rightarrow 1 - \Phi\left(-\frac{(1-\lambda)^{1/2}}{\sigma_\lambda} u(x)\right) = \Phi\left(\frac{(1-\lambda)^{1/2}}{\sigma_\lambda} u(x)\right) \end{aligned}$$

for all $x \in B(u(\cdot))$. If $u(x) = +\infty$, then $x \geq x_0 \equiv \sup\{x: u(x) < \infty\}$. By taking $x_n \in B(u(\cdot))$, $x_n \uparrow x_0$ and using the continuity of $\Phi(\cdot)$ and $u(\cdot)$, it follows that

$$\begin{aligned} \lim_n P(X_{k_n}^{(n)} \leq a_n x + b_n) &\geq \lim_n P(X_{k_n}^{(n)} \leq a_n x_0 + b_n) \geq \lim_n P(X_{k_n}^{(n)} \leq a_n x_n + b_n) \\ &= \lim_n \Phi\left(\frac{(1-\lambda)^{1/2}}{\sigma_\lambda} u(x_n)\right) = 1, \end{aligned}$$

i.e. $\lim_n P(X_{k_n}^{(n)} \leq a_n x + b_n) = \Phi\left(\frac{(1-\lambda)^{1/2}}{\sigma_\lambda} u(x)\right)$ still holds. Similarly we can show

(3.4) also holds if $u(x) = -\infty$. This proves the theorem.

From this theorem we know that under assumption I or II, the limiting distributions of $\frac{X_{k_n}^{(n)} - b_n}{a_n}$ and $\frac{\hat{X}_{k_n}^{(n)} - b_n}{a_n}$ may be different. In fact, we have

Theorem 3.2 If there are $a_n > 0$, b_n such that (3.2) holds with a continuous $u(x)$, (2.14) and either (2.1) or (2.4) holds with some $\ell_n = o(\min(k_n^{1/2}, \frac{n^{1/2}}{k_n^{1/4}}))$ for any $u_n = a_n x + b_n$, $x \in B(u(\cdot))$, then

$$(3.5) \quad P(X_{k_n}^{(n)} \leq a_n x + b_n) \rightarrow \Phi(u(x)).$$

Proof: Notice (2.14) implies (2.2) and (2.3) with $\sigma=0$.

Theorem 3.3 If in theorem 3.1 or 3.2, $\{k_n\}$ is nondecreasing and $\lambda=0$, then (3.4) and (5.5) hold respectively.

Proof: It is proved by Wu [8] that if $\{k_n\}$ is nondecreasing then the only possible types of limiting laws of $\{\chi_{k_n}^{(n)}\}$ are $\Phi(u_i(x))$, $i=1,2,3$, where

$$u_1(x) = \begin{cases} -\alpha \log|x| & x < 0 \\ \infty & x \geq 0 \end{cases} \quad (\alpha > 0)$$

$$u_2(x) = \begin{cases} -\infty & x \leq 0 \\ \alpha \log x & x > 0 \end{cases} \quad (\alpha > 0)$$

$$u_3(x) = x.$$

The theorem follows from theorem 3.1 and 3.2, by noting that $u_i(x)$, $i=1,2,3$ are continuous.

Smirnov [6] has shown that $u(x)$ satisfying (3.2) need not be continuous if $0 < \lambda < 1$ in (0.1). Therefore theorem 3.1 and 3.2 cannot be applied to this case, and a special discussion is therefore needed.

Lemma 5.4 If $\lambda \in (0,1)$ in (0.1) and $\beta(n, \ell_n)$ (or $3^n \alpha(n, \ell_n)$) tend to zero for some $\ell_n = o(k_n^{1/2})$, $\ell_n \rightarrow \infty$, then

$$\frac{1}{k_n} \sum_{j=1}^N \tilde{I}_{nj} \rightarrow 0 \quad [P].$$

Proof: Let $\tilde{\ell}_n = \max(n\beta^{1/2}(n, \ell_n), [k_n^{1/2}])$ and

$$\tilde{\xi}_{ni} = \sum_{j=(i-1)(\tilde{\ell}_n + \ell_n) + 1}^{(i-1)(\tilde{\ell}_n + \ell_n) + \tilde{\ell}_n} \tilde{I}_{nj}, \quad \tilde{\eta}_{ni} = \sum_{j=(i-1)(\ell_n + \tilde{\ell}_n) + \tilde{\ell}_n + 1}^{i(\ell_n + \tilde{\ell}_n)} \tilde{I}_{nj},$$

$$\tilde{\zeta}_n = \sum_{j=N(\ell_n + \tilde{\ell}_n) + 1}^n \tilde{I}_{nj}.$$

We have

$$\frac{1}{k_n^2} E \tilde{\zeta}_n^2 \leq \frac{1}{k_n} (\tilde{\ell}_n + \ell_n)^2 \rightarrow 0.$$

$$\frac{1}{k_n^2} E \left(\sum_{i=1}^N \tilde{\eta}_{ni} \right)^2 \leq \frac{1}{k_n} (N \ell_n)^2 \leq \frac{n^2}{k_n^2} \cdot \left(\frac{\ell_n}{\ell_n} \right)^2 \rightarrow 0,$$

so that $\frac{1}{k_n} \tilde{\zeta}_n \rightarrow 0$ [P], $\frac{1}{k_n} \sum_{i=1}^N \tilde{\eta}_{ni} \rightarrow 0$ [P]. Noticing that under the conditions of the lemma,

$$|E e^{i \sum_{j=1}^N \frac{\tilde{\zeta}_{nj}}{k_n} t} - E e^{i \frac{\tilde{\zeta}_{n1}}{k_n} t}|^{N_n} \leq \beta^{1/2}(n, \ell_n) \rightarrow 0,$$

and that

$$\frac{N_n}{k_n^2} E |\tilde{\zeta}_{n1}|^2 \leq \frac{N_n}{k_n^2} \tilde{\ell}_n^2 \leq \frac{n}{k_n} \cdot \frac{\tilde{\ell}_n}{k_n} \rightarrow 0, \text{ we obtain}$$

$$\lim_n E e^{i \frac{1}{k_n} \sum_{j=1}^N \tilde{\zeta}_{nj} t} = \lim_n (E e^{i \frac{\tilde{\zeta}_{n1}}{k_n} t})^{N_n} = 1,$$

and hence $\frac{1}{k_n} \sum_{i=1}^N \tilde{\zeta}_{ni} \rightarrow 0$ [P]. This proves the lemma.

Lemma 3.5 Under the conditions of lemma 3.4, if for some real sequence $\{u_n\}$,

$$0 < \lim_n P(X_{k_n}^{(n)} \leq u_n) \leq \overline{\lim}_n P(X_{k_n}^{(n)} \leq u_n) < 1,$$

then (2.5) holds.

Proof: If (2.5) does not hold, from Lemma 3.4 and the fact

$$P(X_{k_n}^{(n)} \leq u_n) = P\left(\frac{1}{k_n} \sum_{j=1}^n \tilde{\eta}_{nj} \geq 1 - \frac{n}{k_n} F(u_n)\right),$$

we know that one of the two equations

$$\lim_n P(X_{k_n}^{(n)} < u_n) = 0, \quad \overline{\lim}_n P(X_{k_n}^{(n)} \leq u_n) = 1$$

must hold, contradicting the assumption of the lemma 3.5. Hence (2.5) must hold.

Theorem 3.6 Suppose that

1. $\lambda \in (0,1)$ in (0.1);
2. there are $a_n > 0$, b_n such that (3.2) holds;
3. for any $u_n = a_n x + b_n$, $x \in B_1(u(\cdot)) \equiv \{x: |u(x)| < \infty, x \text{ is a continuity point of } u(x)\}$, assumption I or II holds for some $\ell_n = o(k_n^{1/4})$, $\ell_n \rightarrow \infty$. Then the real α in (2.2) is independent of x and determined by (2.2)', and (3.4) holds. Furthermore, if conditions 1,2,(2.14) hold and either (2.1) or (2.4), with $\ell_n = o(k_n^{1/4})$, $\ell_n \rightarrow \infty$, then (3.5) holds.

Proof: It follows from theorem 2.5 that (3.4) holds for all $x \in B_1(u(\cdot))$. Thus it is sufficient to show that

$$(3.6) \quad \lim_n P(X_{k_n}^{(n)} \leq a_n x + b_n) = 1, \quad \text{if } u(x) = \infty$$

$$(3.7) \quad \lim_n P(X_{k_n}^{(n)} \leq a_n x + b_n) = 0, \quad \text{if } u(x) = -\infty.$$

If (3.6) is not true, we can choose a subsequence such that

$$\lim_{n'} P(X_{k_{n'}}^{(n')} \leq a_{n'} x_0 + b_{n'}) = \ell < 1$$

for some x_0 , $u(x_0) = \infty$. Taking $x_1 \in B_1(u(\cdot))$, we have $x_1 < x_0$, and therefore

$$0 < \lim_{n'} P(X_{k_{n'}}^{(n')} \leq a_{n'} x_1 + b_{n'}) \leq \lim_{n'} P(X_{k_{n'}}^{(n')} \leq a_{n'} x_0 + b_{n'}) = \ell < 1.$$

According to lemma 3.5, this implies (2.5) with $u_{n'} = a_{n'} x_0 + b_{n'}$. By using theorem 2.4, it follows that

$$\begin{aligned} \Phi\left(\frac{(1-\lambda)^{1/2}}{\sigma_\lambda} u(x_0)\right) &= \lim_{n'} \Phi\left(\frac{1}{\sigma_\lambda} \frac{n'}{k_{n'}} \left[F(u_{n'}) - \frac{k_{n'}}{n'}\right]\right) \\ &= \lim_{n'} P(X_{k_{n'}}^{(n')} \leq u_{n'}) = \ell \in (0,1). \end{aligned}$$

This is contrary to $\Phi(\frac{(1-\lambda)^{1/2}}{\sigma_\lambda} u(x_0)) = \Phi(\infty) = 1$. Hence (3.6) must hold, and in a similar way, we can show (3.7). Then the theorem follows.

If the rank sequence $\{k_n\}$ satisfies

$$(3.8) \quad n^{1/2}(\frac{k_n}{n} - \lambda) \rightarrow t, \quad -\infty < t < \infty, \quad 0 < \lambda < 1,$$

Smirnov [6] has shown that the only possible non-degenerate types of limiting laws of $\{\chi_{k_n}^{(n)}\}$ are $\Phi(\frac{\tilde{u}_i(x)}{\tilde{\sigma}_\lambda} - \frac{t}{\tilde{\sigma}_\lambda})$, $i=1,2,3,4$, where

$$(3.9) \quad \begin{aligned} \tilde{u}_1(x) &= \begin{cases} -\infty & x < 0 \\ Cx^\alpha & x \geq 0 \end{cases} & (C > 0, \alpha > 0) \\ \tilde{u}_2(x) &= \begin{cases} -C|x|^\gamma & x < 0 \\ \infty & x \geq 0 \end{cases} & (C > 0, \gamma > 0) \\ \tilde{u}_3(x) &= \begin{cases} -C_1|x|^\alpha & x < 0 \\ C_2x^\alpha & x \geq 0 \end{cases} & (C_1, C_2 > 0, \alpha > 0) \\ \tilde{u}_4(x) &= \begin{cases} -\infty & x < -1 \\ 0 & -1 \leq x < 1 \\ \infty & x \geq 1 \end{cases} \end{aligned}$$

and $\tilde{\sigma}_\lambda = [\lambda(1-\lambda)]^{1/2}$. For stationary processes, similar results are obtained as follows.

Theorem 3.7 If (3.8) holds, then under conditions 2 and 3 of theorem 3.6, (3.4) holds, and $u(x)$ in (3.4) is one of the four types

$$u(x) = \frac{\tilde{u}_i(x)}{\tilde{\sigma}_\lambda} - \frac{t}{\tilde{\sigma}_\lambda} \quad i=1,2,3,4,$$

and the real σ in (3.4) can be found as follows

$$(3.10) \quad \sigma = \frac{1}{\lambda} \sum_{j=1}^{\infty} [P(X_1 \leq a(\lambda), X_{j+1} \leq a(\lambda)) - \lambda^2]$$

where

$$a(\lambda) = \begin{cases} a_\lambda - 0 & \text{if } F(a_\lambda) - \lambda < \lambda - F(a_\lambda - 0) \\ a_\lambda & \text{if } F(a_\lambda) - \lambda \leq \lambda - F(a_\lambda - 0) \end{cases}$$

and a_λ is a real such that $F(a_\lambda - 0) < \lambda \leq F(a_\lambda)$, and the event $\{X_n \leq a_\lambda - 0\}$ means $\{X_n = a_\lambda\}$.

Proof: According to theorem 3.6 and Smirnov's results as above, it is sufficient to show (3.10). Writing $u_n = a_n x + b_n$, $x \in B_1(u(\cdot))$, we have

$$\begin{aligned} & \left| \sum_{j=1}^{\ell_n-1} [P(X_1 \leq u_n, X_{j+1} \leq u_n) - F^2(u_n)] - \sum_{j=1}^{\ell_n-1} [P(X_1 \leq a(\lambda), X_{j+1} \leq a(\lambda)) - \lambda^2] \right| \\ & \leq \sum_{j=1}^{\ell_n-1} [|P(X_1 \leq u_n, X_{j+1} \leq u_n) - P(X_1 \leq a(\lambda), X_{j+1} \leq a(\lambda))| + |F^2(u_n) - \lambda^2|] \\ & \leq 4\ell_n |F(u_n) - \lambda| \leq 4\ell_n (|F(u_n) - \frac{k_n}{n}| + |\frac{k_n}{n} - \lambda|) \rightarrow 0 \end{aligned}$$

so that

$$\begin{aligned} \sum_{j=1}^{\ell_n-1} [P(X_1 \leq a(\lambda), X_{j+1} \leq a(\lambda)) - \lambda^2] &= \lim_n \sum_{j=1}^{\ell_n-1} [P(X_1 \leq a(\lambda), X_{j+1} \leq a(\lambda)) - \lambda^2] \\ &= \lim_n \sum_{j=1}^{\ell_n-1} [P(X_1 \leq u_n, X_{j+1} \leq u_n) - F^2(u_n)] \\ &= \lambda \lim_n \frac{n}{k_n} \sum_{j=1}^{\ell_n-1} E \tilde{I}_{n1} \tilde{I}_{nj+1} = \lambda \sigma. \end{aligned}$$

Hence Theorem 3.7 holds.

54. Example: The Normal Case

Let $\{X_n, n=1,2,\dots\}$ be a stationary normal sequence with

$$EX_n = 0, \quad EX_n^2 = 1, \quad EX_1 X_{n+1} = r_n, \quad n=1,2,\dots$$

In this section, we give some conditions on $\{r_n\}$ such that the limiting distributions of $\{X_k^{(n)}\}$ exist for some special rank sequences $\{k_n\}$.

Lemma 4.1 If $r_n \rightarrow 0$ and

$$(4.1) \quad \lim_n \frac{n}{k_n^{1/2}} \sum_{j=\ell_n}^{n-1} j |r_j| \exp(-\frac{u_n^2}{1+|r_j|}) = 0$$

then (2.1) holds.

Proof: The method of proving this lemma is a slight extension of an argument of Leadbetter, Lindgren and Rootzén [7]. Let $\tilde{A}_{ni} = \{X_i \leq u_n\} \in R^n$, $i=1, \dots, n$. For any fixed integer k, ℓ , denote $F_k = \sigma\{\tilde{A}_{n1}, \dots, \tilde{A}_{nk}\}$, $F_{k+\ell}^* = \sigma\{\tilde{A}_{nk+\ell+1}, \dots, \tilde{A}_{nn}\}$. Then any $A \in \sigma\{\{X_i \leq u_n\}, i=1, \dots, k\}$, $B \in \sigma\{\{X_i \leq u_n\}, i=k+\ell+1, \dots, n\}$ can be represented as $A = \{X_n \in \tilde{A}\}$, $B = \{X_n \in \tilde{B}\}$ where $\tilde{A} \in F_k$, $\tilde{B} \in F_{k+\ell}^*$. Write $f_1(x_1, \dots, x_k; y_1, \dots, y_{n-k-\ell})$ for the density of $(X_1, \dots, X_k; X_{k+\ell+1}, \dots, X_n)$ and $f_0(x_1, \dots, x_k; y_1, \dots, y_{n-k-\ell}) = f_{01}(x_1, \dots, x_k) f_{02}(y_1, \dots, y_{n-k-\ell})$ where f_{01} and f_{02} are the densities of $(X_1, \dots, X_k)$ and $(X_{k+\ell+1}, \dots, X_n)$ respectively. Let R_1 and R_0 be the covariance matrices of f_1 and f_0 . It is easy to show that $R_h = hR_1 + (1-h)R_0$ is positive definite for any $h \in [0, 1]$. Writing $f_h(x_1, \dots, x_k, y_1, \dots, y_{n-k-\ell})$ for the density function of a zero-mean normal vector with covariance matrix R_h , we have

$$(4.2) \quad P(AB) - P(A)P(B) = \int_{x \in \tilde{A}} \dots \int_{y \in \tilde{B}} \left[\int_0^1 f_h' dh \right] dx dy$$

$$= \int_0^1 dh \sum_{\substack{1 \leq i \leq k \\ k+\ell+1 \leq j \leq n}} \int_{x \in \tilde{A}} \dots \int_{y \in \tilde{B}} \frac{\partial^2 f_h}{\partial x_i \partial y_{j-k-\ell}} dx dy,$$

where $x = (\begin{smallmatrix} x_1 \\ \vdots \\ x_k \end{smallmatrix})$, $y = (\begin{smallmatrix} y_1 \\ \vdots \\ y_{n-k-\ell} \end{smallmatrix})$. Split the integral $\int_{x \in \tilde{A}} \dots \int_{y \in \tilde{B}} \frac{\partial^2 f_h}{\partial x_i \partial y_{j-k-\ell}} dx dy$ into four parts: for $x \in \tilde{A} \cap \{x_i \leq u_n\}$, $y \in \tilde{B} \cap \{y_{j-k-\ell} \leq u_n\}$ and $x \in \tilde{A} \cap \{x_i > u_n\}$, $y \in \tilde{B} \cap \{y_{j-k-\ell} > u_n\}$.

and $x \in A \cap \{x_i \leq u_n\}$, $y \in B \cap \{y_{j-k-\ell} \leq u_n\}$, and $x \in \bar{A} \cap \{x_i > u_n\}$, $y \in \bar{B} \cap \{y_{j-k-\ell} > u_n\}$, where

$A = \{\tilde{A}_{n1}, \dots, \tilde{A}_{ni-1}, \tilde{A}_{ni+1}, \dots, \tilde{A}_{nk}\}$, $\bar{B} = \{\tilde{A}_{nk+\ell+1}, \dots, \tilde{A}_{n,j-1}, \tilde{A}_{n,j+1}, \dots, \tilde{A}_{nn}\}$. (This can be done since $\tilde{A}$ is a disjoint union of sets of the form $\bigcap_{j=1}^k C_{nj}$, where $C_{nj} = \tilde{A}_{nj}$ or the complement of $\tilde{A}_{nj}$, and similarly for B). Writing $x^{(i)}, y^{(i)}$ for the vectors x, y without the component x_i, y_i we have

$$\begin{aligned} & \left| \int \dots \int_{\substack{x \in \bar{A} \cap \{x_i \leq u_n\} \\ y \in \bar{B} \cap \{y_{j-k-\ell} \leq u_n\}}} \frac{\gamma^2 f_h}{|x_i y_{j-k-\ell}|} dx dy \right| \\ & \leq \int \dots \int_{R^{k-1} \times R^{n-k-\ell-1}} f_h(x_i = u_n, y_{j-k-\ell} = u_n) dx^{(i)} dy^{(i)} \\ & \leq \frac{1}{2\pi(1-r_{j-i}^2)^{1/2}} \exp\left\{-\frac{u_n^2}{1+|r_{j-i}|}\right\} \end{aligned}$$

and the same inequalities hold for other three parts of the integral. Since $r_n \rightarrow 0$ we have $\sup_{n \geq 1} |r_n| < 1$, and therefore

$$\left| \int \dots \int_{x \in \tilde{A}, y \in \tilde{B}} \frac{\gamma^2 f_h}{|x_i y_j|} dx dy \right| \leq C \exp\left(-\frac{u_n}{1+|r_{j-i}|}\right)$$

From this and (4.2) it follows that

$$|P(AB) - P(A)P(B)| \leq C \sum_{\substack{1 \leq i \leq k \\ k+\ell \leq j \leq n}} |r_{j-i}| \exp\left(-\frac{u_n}{1+|r_{j-i}|}\right) \leq C \sum_{j=\ell}^{n-1} j |r_j| \exp\left(-\frac{u_n}{1+|r_{j-1}|}\right)$$

Hence (2.1) holds if (4.1) holds.

According to theorem 3.5 and 3.7 of Cheng [1], we know that (3.2) holds for any k_n satisfying (0,1), if $F(x) = \Phi(x)$ and $a_n > 0$, b_n are defined by $\Phi(b_n) = \frac{k_n}{n}$, $a_n = \frac{k_n^{1/2}}{n} \cdot \frac{(1-\lambda)^{1/2}}{\phi(b_n)}$, where $\phi(x)$ is the density function of the standard normal distribution $\Phi(x)$. Using this fact, we discuss the limiting distribution of

order statistics from stationary normal sequences. Since the case $\lambda=0$ has been discussed in [1], we consider only the case $\lambda \in (0,1)$.

Lemma 4.2 If $\lambda \in (0,1)$ and $r_n = o(n^{-(1+\rho)})$, $\rho > 3$, then (4.1) holds for some $\ell_n = o(n^{1/4})$ and any u_n which satisfies (1.1).

Proof: To show (4.1) it is sufficient to show that

$$\lim_n n^{1/2} \sum_{j=\ell_n}^{n-1} j |r_j| \exp\left(-\frac{u_n}{1+|r_j|}\right) = 0.$$

Since $\exp\left(-\frac{u_n}{1+|r_j|}\right) \leq \exp\left(-\frac{u_n^2}{2}\right)$, this will follow if

$$\lim_n n^{1/2} \sum_{j=\ell_n}^{n-1} j |r_j| \exp\left(-\frac{u_n^2}{2}\right) = 0.$$

Since $\lambda(u_n) \rightarrow \lambda$ and the inverse function $\phi^{-1}(x)$ of $\zeta(x)$ is continuous, we have $u_n \rightarrow a_\lambda$ where a_λ is defined by $\phi(a_\lambda) = \lambda$. Hence

$$\begin{aligned} n^{1/2} \sum_{j=\ell_n}^{n-1} j |r_j| \exp\left(-\frac{u_n^2}{2}\right) &\leq \left[\exp\left(-\frac{a_\lambda^2}{2}\right) + 1\right] n^{1/2} \sum_{j=\ell_n}^{n-1} j |r_j| \\ &\leq C n^{1/2} \sum_{j=\ell_n}^{n-1} \frac{1}{j^\rho} \leq C n^{1/2} \frac{\ell_n^{-\rho+1}}{\rho-1} \end{aligned}$$

for sufficiently large n . Let $\ell_n = [n/\log n]^{1/4}$. Thus

$$n^{1/2} \sum_{j=\ell_n}^{n-1} j |r_j| \exp\left(-\frac{u_n^2}{2}\right) \leq (\log n)^{\rho-1} / n^{(\rho-3)/4} \rightarrow 0$$

completing the proof of the lemma.

Lemma 4.2 If $\lambda \in (0,1)$ in (0.1) and $\sum_{n=1}^{\infty} |r_n| < \infty$, then

$$(4.5) \quad \lim_n \sum_{j=1}^{\ell_n-1} [P(X_1 \leq b_n, X_{j+1} \leq b_n) - (\frac{k_n}{n})^2] = \frac{1}{2\pi} \sum_{n=0}^{\infty} \int_0^{r_n} \frac{\exp(-\frac{a_\lambda^2}{1+r})}{(1-r^2)^{1/2}} dr$$

$$(4.11) \quad \lim_n \frac{1}{n^{1/2}} \sum_{j=1}^{\ell_n-1} j [P(X_1 \leq b_n, X_{j+1} \leq b_n) - (\frac{k_n}{n})^2] = 0$$

for any $\ell_n = o(n^{1/4})$, $\ell_n \rightarrow \infty$, where a_λ is the solution of the equation $\phi(x) = \lambda$.

Proof: It is easy to show that

$$P(X_1 \leq b_n, X_{j+1} \leq b_n) - (\frac{k_n}{n})^2 = \frac{1}{2\pi} \int_0^{r_j} \frac{\exp(-\frac{b_n^2}{1+r})}{(1-r^2)^{1/2}} dr.$$

Notice that

$$\begin{aligned} \sum_{j=1}^{\ell_n-1} \int_0^{r_j} \frac{\exp(-\frac{b_n^2}{1+r})}{(1-r^2)^{1/2}} dr &= \sum_{j=1}^{\ell_n-1} \int_0^{r_j} \frac{\exp(-\frac{a_\lambda^2}{1+r})}{(1-r^2)^{1/2}} dr \\ &\leq \sum_{j=1}^{\ell_n-1} \int_0^{r_j} \frac{|r_j| \exp(-\frac{a_\lambda^2}{1+r})}{(1-r^2)^{1/2}} \left| \exp(-\frac{b_n^2 - a_\lambda^2}{1+r}) - 1 \right| dr \\ &\leq |b_n^2 - a_\lambda^2| \sum_{j=1}^{\ell_n-1} \int_0^{r_j} \frac{|r_j|}{(1-r^2)^{1/2} (1+r)} dr \\ &\leq C |b_n^2 - a_\lambda^2| \sum_{n=1}^{\infty} |r_n| \rightarrow 0. \end{aligned}$$

Hence to show (4.3), it is sufficient to show that the series $\sum_{n=0}^{\infty} \int_0^{r_n} \frac{\exp(-\frac{a_\lambda^2}{1+r})}{(1-r^2)^{1/2}} dr$

converges. This follows since

$$\left| \sum_{n=1}^{\infty} \int_0^1 r_n \frac{\exp(-\frac{a_{\lambda}^2}{1+r})}{(1-r^2)^{1/2}} dr \right| \leq \sum_{n=1}^{\infty} |r_n| \frac{\exp(-\frac{a_{\lambda}^2}{1+r})}{(1-r^2)^{1/2}} \leq C \sum_{n=1}^{\infty} |r_n| < \infty.$$

(4.3) is proved, and (4.4) can be shown in a similar way.

From lemma 4.2, 4.3 and theorem 3.6, we have

Theorem 4.4 If $r_n = O(n^{-(1+\rho)})$, $\rho > 3$, then

$$\lim_n P(X_{k_n}^{(n)} \leq a_n x + b_n) = \Phi\left(\frac{(1-\lambda)^{1/2}}{\sigma_{\lambda}} x\right)$$

for any k_n such that $\frac{k_n}{n} \rightarrow \lambda \in (0,1)$, where

$$\sigma_{\lambda}^2 = (1-\lambda) + \frac{1}{\pi\lambda} \sum_{n=0}^{\infty} \int_0^1 r_n \frac{\exp(-\frac{a_{\lambda}^2}{1+r})}{(1-r^2)^{1/2}} dr.$$

Acknowledgement: This paper was completed during the author's visit to the University of North Carolina at Chapel Hill. The author is grateful to Professor M.P. Leadbetter for helpful discussions.

REFERENCES

- [1] Cheng, B. (1965). The limiting distributions of order statistics. Chinese Math. 6 84-104.
- [2] Cheng, S. (1980). On the limiting distributions of middle terms of order statistics from stationary sequences. J. of Peking Univ. 1980, No. 1 14-30.
- [3] Dvoretzky, A. (1972). Asymptotic normality for sums of dependent random variables. Proc. Sixth Berkeley Symp. Math. Statist. Probability 2 513-535.
- [4] Leadbetter, M.R. (1974). On extreme values in stationary sequences. Z. Wahrsch. verw. Geb. 28 289-303.
- [5] Leadbetter, M.R., Lindgren, G. and Rootzen, H. (1979). Extremal and related properties of stationary processes. Part I: Extremes of stationary sequences. Institute of Statistics Mimeo Series #122, University of North Carolina at Chapel Hill.
- [6] Smirnov, N.V. (1949). Limit laws for terms of a variational series. Trudy Steklov Math. Inst., 25 1-60.
- [7] Watts, V., Rootzen, H. and Leadbetter, M.R. (1982). On limiting distributions of intermediate order statistics from stationary sequences. Ann. Prob. 10 653-662.
- [8] Wu, C.Y. (1966). The types of limit distributions for some terms of variational series. Scientia Sinica 15 749-762.