

AD-A131 077

BRAIN AND LIVER GLUTAMINE SYNTHETASE OF RANA
CATESBEIANA AND RANA CANCRIVORA(U) AIR FORCE ACADEMY CO
J T WEBB JUL 83 USAFA-TR-83-13


1/1

UNCLASSIFIED

F/G 6/1

NL

END
DATE
FILMED
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

12

**BRAIN AND LIVER GLUTAMINE SYNTHETASE
OF**

**RANA CATESBEIANA
AND
RANA CANCRIVORA**

MAJOR JAMES T. WEBB

**DEPARTMENT OF BIOLOGY
USAF ACADEMY, COLORADO SPRINGS, CO 80840**

**JULY 1983
FINAL REPORT**

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED



**DTIC
ELECTE
S AUG 4 1983**

D

**DEAN OF THE FACULTY
UNITED STATES AIR FORCE ACADEMY
COLORADO SPRINGS, CO 80840**

AD A 131077

DTIC FILE COPY

88 08 08 08 08

Editorial Review by Captain Hale
Department of English
USAF Academy, Colorado Springs, Colorado 80840

This research report is presented as a competent treatment of the subject, worthy of publication. The United States Air Force Academy vouches for the quality of the research, without necessarily endorsing the opinions and conclusions of the author.

This report has been cleared for open publication and/or public release by the appropriate Office of Information in accordance with AFR 190-17 and DODD 5230.9. There is no objection to unlimited distribution of this report to the public at large, or by DDC to the National Technical Information Service.

This research report has been reviewed and is approved for publication.


THOMAS E. MCCANN, Lt Col, USAF
Director of Research and
Continuing Education

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	



BRAIN AND LIVER GLUTAMINE SYNTHETASE OF
RANA CATESBEIANA
 AND
RANA CANCRIVORA

James T. Webb

DEPARTMENT OF BIOLOGY

USAF ACADEMY, COLORADO SPRINGS, COLORADO 80840

JULY 1983

DEAN OF THE FACULTY

UNITED STATES AIR FORCE ACADEMY

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER USAFA-TR 83-13	2. GOVT ACCESSION NO. 0131077	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) BRAIN AND LIVER GLUTAMINE SYNTHETASE OF <u>RANA</u> <u>CATESBEIANA</u> AND <u>RANA</u> <u>CANCRIVORA</u>		5. TYPE OF REPORT & PERIOD COVERED Final Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) James T. Webb, Major, USAF, Ph.D.		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Biology USAFA (DFB) USAF Academy, Colorado Springs, CO 80840		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 2308
11. CONTROLLING OFFICE NAME AND ADDRESS Department of Biology USAFA (DFB) USAF Academy, Colorado Springs, CO 80840		12. REPORT DATE July 1983
		13. NUMBER OF PAGES 14
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Glutamine synthetase, gamma-glutamyl transferase, glutamate, glutamine, enzyme, brain, liver, frog, urea synthesis, <u>Rana catesbeiana</u> , <u>Rana cancrivora</u> , amphibia, osmoregulation, urea synthesis.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Enzyme assays using the gamma-glutamyl transferase method provided estimates of glutamine synthetase activity in brain and liver tissues of <u>Rana catesbeiana</u> and <u>Rana cancrivora</u> . Glutamine synthetase specific activity in liver and brain tissues were not significantly different between these species. In the salt-water frog <u>R. cancrivora</u> , liver glutamine synthetase activity is insufficient to produce enough glutamine to supply the urea cycle with a nitrogen source. The pathway involving glutamine as a nitrogen source for urea production in marine chondrichthian fishes is not valid in the marine amphibian, <u>Rana cancrivora</u> .		

DD FORM 1473 1 JAN 73 EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

TABLE OF CONTENTS

<u>Title</u>	<u>Page</u>
Acknowledgements	ii
Introduction	1
Materials and Methods	2
Results and Discussion	3
Summary	5
Figure 1--Evolutionary tree of vertebrates	6
Figure 2--Proposed pathway for assimilation of ammonia into urea in marine Chondrichthyes liver	7
Table 1--Liver and brain glutamine synthetase of urea-retaining and non-urea-retaining amphibians	8
References	9

ACKNOWLEDGEMENTS

I sincerely appreciate the efforts of Captain Richard C. Postlewaite, DVM, in timely acquisition of tissue samples and for attempts to keep the live salt-water frogs in a healthy condition. Professor Angel C. Alcala was instrumental in enabling me to acquire the salt-water frogs and his assistance is greatly appreciated. Major Vicky Fogelman, DVM, was also a critical link in the transoceanic shipment of the live salt-water frogs. Many thanks to others who helped including Captain Ronald L. Merritt, and numerous people in USAF Transportation Management Offices.

INTRODUCTION

Previous work has determined that glutamine synthetase plays a role in the synthesis of urea in marine chondrichthian liver (1,15,16). This group of marine urea-retaining animals utilizes the urea as an osmolyte to counter the osmolarity of their environment (2,7,8,10,14). Figure 1 shows an evolutionary tree of vertebrates. The relationship between urea retention as an osmolyte in marine species and presence of high levels of glutamine synthetase in the liver is clear for most groups. The lungfishes (Dipnoids) do not retain urea except to avoid ammonia toxicity during estivation (11). This mechanism is not osmoregulatory and the group does not live in salt-water.

The coelacanth, Latimeria chalumnae, in the Class Osteichthyes, is a marine fish that also retains urea as an osmolyte (3,12). It too has a relatively high level of glutamine synthetase in its liver (16). The proposed connection between glutamine synthesis and urea synthesis by these marine species was discussed by Webb (15) and is depicted in Figure 2. The key enzymes of this pathway, glutamine synthetase and glutamine-dependent carbamoyl phosphate synthetase III, were shown later in a similar figure by Anderson (1).

The saltwater, crab-eating frog, Rana cancrivora, retains urea for the same purpose (9). Although Rana cates-

beiana, fresh-water bullfrogs, switch from ammonioteleism to ureotelism (4,5,6) they do not retain urea in their plasma and tissues. The question remained as to whether the salt-water frogs also use the pathway in Figure 2 as a means of scavenging ammonia for urea synthesis. This question is addressed here by analyses of the liver and brain glutamine synthetase of Rana catesbeiana and Rana cancrivora. The methods and types of comparisons between fresh-water and salt-water species used previously (15,16) were unchanged for this study.

MATERIALS AND METHODS

Rana catesbeiana, bullfrog tadpoles and adult specimens were obtained from several sources (15). Rana cancrivora, salt-water, crab-eating frogs, were obtained from Professor Angel C. Alcala, Silliman University, Philippines.

Tissues were removed from the live specimens and homogenated with water immediately prior to the assays. The Rana cancrivora specimens had been placed in full-strength artificial sea water and were highly stressed. Several of the salt-water frogs had died and the tissues from the three remaining, barely living specimens were removed, sealed in foil, and frozen. These tissues were assayed five days later.

The tissue homogenates were assayed for glutamine syn-

thetase activity according to the method of Webb (15) and Webb and Brown (16) under the following conditions: Ten minutes of incubation at 25°C; pH 6.4; 2 ml incubation mixture containing 60 mM L-glutamine, 15 mM hydroxylamine-HCl, 0.4 mM Na₂ADP; 20 mM KH₂AsO₄, 3 mM MnCl₂, and 40 mM imidazole. The gamma-glutamyl hydroxamate produced by the transferase enzyme activity was complexed with FeCl₃ (in HCl) and compared to a gamma-glutamyl hydroxamate standard (Sigma Chemical Co.) at 500nm with a B&L Spectronic 20 Spectrophotometer. A unit of glutamine synthetase activity is defined as the production of one micro-mole of gamma-glutamyl hydroxamate per minute at 25°C. Protein content was determined by the biuret method described by Zamenhof (17) using a bovine albumin standard (Calbiochem-Behring #126575).

RESULTS AND DISCUSSION

There was no significant difference between the specific activity of glutamine synthetase in the liver of urea-retaining versus non-urea-retaining species of amphibians sampled in this study (Table 1). The ratio of liver to brain specific activity also showed no difference as had been expected on the basis of an analogous study in fish (16). In that study, the urea-retaining fish had a liver to brain specific activity ratio of 1.55 ± 0.61 (19) as compared

to $0.02 \pm 0.02(48)$ for non-urea-retaining fish. This very significant difference in fish was not evident in the current study on amphibians. Activity in kidney tissue was also extremely low in both species of amphibians.

Several explanations of these unexpected data are possible. The salt-water frogs may have been over-stressed by immersion in full-strength artificial sea water and the activity of glutamine synthetase in the liver of these moribund frogs may have disappeared as a result. This however, is not consistent with an earlier report on stability of glutamine synthetase in the liver of urea-retaining fish (15). The same argument applies to the possibility of loss of enzyme activity from the time of sacrifice to the time of assay; particularly in view of the retention of activity in the brain tissue which was treated in the same manner as the liver tissue.

Another possibility involves control of the pathway of urea synthesis. Perhaps glutamine synthetase is subject to sensitive control in the liver of salt-water frogs. These specimens were shipped in a box lined with sponges soaked in fresh-water. Their exposure to the fresh-water may have been sufficient to activate a potential regulatory mechanism to stop urea synthesis since it was not needed at that time as an osmoregulatory solute. Glutamine synthetase could be the first enzyme to be repressed in the pathway of urea


synthesis (see Figure 2). Lack of reactivation of the pathway on immersion in full-strength artificial sea water could have been due to osmotic shock.

One more explanation could be the presence of a functional carbamoyl phosphate synthetase I in the salt-water frog liver which produces carbamoyl phosphate using ammonia as a substrate in place of glutamine. This possibility raises questions about the ability of the carbamoyl phosphate synthetase I to produce enough carbamoyl phosphate to feed the urea cycle at a sufficient rate to produce the high levels of osmoregulatory urea found in the salt-water frogs.

SUMMARY

Glutamine synthetase activity in liver and brain tissues from salt-water frogs (Rana cancrivora) closely resembles the activity in fresh-water frogs (Rana catesbeiana). Problems with maintaining the salt-water frogs may have lowered their liver tissue activity but probably not enough to reverse the relationship seen here. In the salt-water frog R. cancrivora, liver glutamine synthetase activity is insufficient to produce enough glutamine to supply the urea cycle with a nitrogen source. The pathway previously proposed which involved glutamine as a nitrogen source for urea production in marine chondrichthian fishes is apparently not valid in the marine amphibian, Rana cancrivora.

FIGURE 1--Evolutionary tree of vertebrates



The occurrence of high levels of glutamine synthetase in liver is indicated by presence of a circle by the group. The presence (+) or absence (-) of urea retention is shown to indicate correlations. Format after the style of Romer (13).

FIGURE 2--Proposed pathway for assimilation of ammonia into urea in marine Chondrichthyes liver (from Webb (15))




TABLE 1--Liver and brain glutamine synthetase of urea-retaining and non-urea-retaining amphibians.

Species ¹	Specific ² activity		Specific ² activity ratio
	Liver	Brain	<u>Liver</u> <u>Brain</u>
Non-urea-retaining	0.00 $\pm$ 0.00 (9)	0.62 $\pm$ 0.15 (9)	0.01 $\pm$ 0.01 (9)
Urea-retaining	0.00 $\pm$ 0.00 (2)	0.45 $\pm$ 0.13 (3)	0.01 $\pm$ 0.01 (2)

¹Non-urea-retaining amphibians: Rana catesbeiana bullfrogs (data from Webb ('79)). Urea-retaining amphibians: Rana cancrivora, salt-water, crab-eating frogs.

²Mean of activities is expressed as units of glutamine synthetase activity per mg protein $\pm$ standard deviation. Number of specimens is indicated parenthetically.

REFERENCES

1. Anderson, P.M. 1980. Glutamine- and N-acetylglutamate-dependent carbamoyl phosphate synthetase in elasmobranchs. Science, 208, 291-293.
2. Brown, G.W., Jr. 1964. Urea synthesis in elasmobranchs. In: "Taxonomic Biochemistry and Serology". (C.A. Leone, ed.) pp. 407-416. The Ronald Press Co., New York.
3. Brown, G.W., Jr. and Brown, S.G. 1967. Urea and its formation in coelacanth liver. Science, 155, 570-573.
4. Brown, G.W., Jr., Brown, W.R., and Cohen, P.P. 1959. Comparative biochemistry of urea synthesis. J. Biol. Chem. 234, 1775-1780.
5. Brown, G.W., Jr., and Cohen, P.P. 1958. Biosynthesis of urea in metamorphosing tadpoles. In: "A Symposium on The Chemical Basis of Development" (W.D. McElroy and B. Glass eds.) The Johns Hopkins Press, Baltimore. 1958. 934pp.
6. Cohen, P.P. and Brown, G.W., Jr. 1960. Ammonia metabolism and urea biosynthesis. Comp. Biochem. 2, 161-244.
7. deVlaming, V.L. and Sage, M. 1973. Osmoregulation in the euryhaline elasmobranch, Dasyatis sabina. Comp. Biochem. Physiol. 45A, 31-44.
8. Goldstein, L. and Forster, R.P. 1971. Osmoregulation and urea metabolism in the little skate Raja erinacea. Amer. J. Physiol. 220, 742-746.
9. Gordon, M.S., Schmidt-Nielsen, K., and Kelly, H.M. 1961. Osmotic regulation in the crab-eating frog (Rana cancrivora). J. Exp. Biol. 38, 659-678.
10. Holmes, W.N. and Donaldson, E.M. 1969. The body compartments and the distribution of electrolytes. In: "Fish Physiology". (W.E. Hoar and D.G. Randall, eds.) Vol. 1, pp. 1-89. Academic Press, New York.
11. Janssens, P.A. and Cohen, P.P. 1968. Nitrogen metabolism in the African lungfish. Comp. Biochem. Physiol. 24, 879-886.

12. Pickford, G.E. and Grant, F.B. 1967. Serum osmolality in the coelacanth, Latimeria chalumnae: Urea retention and ion regulation. Science, 155, 568-570.
13. Romer, A.S. 1962. "The Vertebrate Body" ed. 3. W.B. Saunders Co., Philadelphia. p. 45.
14. Smith, H.W. 1936. The retention and physiological role of urea in the elasmobranchii. Biol. Rev. Cambridge Philos. Soc. 11, 49-82.
15. Webb, J.T. 1979. "An assimilatory role of glutamine synthetase in the liver of urea-retaining marine chondrichthyes." Ph.D. Dissertation. University of Washington, Seattle. 81pp.
16. Webb, J.T. and Brown, G.W., Jr. 1980. Glutamine synthetase: Assimilatory role in liver as related to urea retention in marine chondrichthys. Science, 208, 293-295.
17. Zamenhof, S. 1957. Nucleic acids and derivatives. In: "Methods in Enzymology". (S.P. Colowick and N.O. Kaplan, eds.) Vol. 3, p. 702. Academic Press, New York.

END

DATE
FILMED

9 — 83

DTIC