

AD-A131 478

UNIFIED THEORY OF PLASMA CORRELATIONS(U) - CORNELL UNIV
ITHACA NY M A GUILLEN ET AL. 13 JUN 83 TR-R-4-83
AFOSR-TR-83-0664 AFOSR-78-3574


1/1

UNCLASSIFIED

F/G 12/1

NL

END
DATE
FILMED
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS - 1963 - A

10

AD A131478

Technical Report No. R-4-83

Project; Task No. 61102F; 2301/A3

Date: June 13, 1983

Title: Unified Theory of Plasma Correlations

Authors: Michael A. Guillen and Richard L. Liboff

Contractor: Cornell University

Principal Investigator: Richard L. Liboff

Approved for public release;
distribution unlimited.

DTIC FILE COPY

Research supported in part by the Air Force Office
of Scientific Research under grant AFOSR-78-3574.

Distribution of this report is unlimited.

83 08 08 192

DTIC
ELECTE
AUG 19 1983
S D D

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR- 83-0664	2. GOVT ACCESSION NO. DA31478	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Unified Theory of Plasma Correlations		5. TYPE OF REPORT & PERIOD COVERED Interim
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Michael A. Guillen Richard L. Liboff		8. CONTRACT OR GRANT NUMBER(s) AFOSR 78-3574
9. PERFORMING ORGANIZATION NAME AND ADDRESS Cornell University Ithaca, NY 14853		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2301/A3
11. CONTROLLING OFFICE NAME AND ADDRESS Directorate of Physics AFOSR Bolling AF Base, Washington, D.C. 20332		12. REPORT DATE June 13, 1983
		13. NUMBER OF PAGES Twenty-four
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Approved for Public Release; Distribution Unlimited.		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) One-component plasma; correlations; BBKGY; unified theory; strongly coupled; total correlation function; two-particle correlation; closed kinetic equation.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A unified approach to the theory of correlations in a plasma is presented, based on the BBKGY hierarchy. The theory is applied to a one-component plasma with the Coulomb interaction modified to include effects of the background. Closed integro-differential equations in space and time are obtained for the two- particle correlation function in both the strong and weak coupling limits. In the weak-coupling domain, $\gamma \ll 1$, the time-independent analysis returns the well- known linearized Debye-Huckle result, where γ is the plasma parameter. In the strong-coupling domain with $\gamma \gg 1$, the resulting two-particle 'total' (continued)		

DD FORM 1473 EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION (When Data Entered)

UNCLASSIFIED

20. Abstract (continued)

correlation function exhibits decaying oscillatory behavior for particle separation of the order of the effective interparticle range.

UNCLASSIFIED

Unified Theory of Plasma Correlations

Michael A. Guillen
and
Richard L. Liboff

Schools of Electrical Engineering
and
Applied Physics

Accession For	
NTIS CERA-I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFOSR)
NOTICE OF TRANSMITTAL TO DTIC
This technical report has been reviewed and is
approved for public release IAW AFS 193-12.
Distribution is unlimited.
MATTHEW J. KERPER
Chief, Technical Information Division

Cornell University
Ithaca, N.Y. 14853



Abstract

A unified approach to the theory of correlations in a plasma is presented, based on the BBKGY hierarchy. The theory is applied to a one-component plasma with the Coulomb interaction modified to include effects of the background. Closed integro-differential equations in space and time are obtained for the two-particle correlation function in both the strong and weak coupling limits. In the weak-coupling domain, $\chi \ll 1$, the time-independent analysis returns the well-known linearized Debye-Huckel result, where γ is the plasma parameter. In the strong-coupling domain with $\chi \gg 1$, the resulting two-particle 'total' correlation function exhibits decaying oscillatory behavior for particle separation of the order of the effective interparticle range.

I. Introduction

Various efforts have undertaken to describe one-component plasmas in two extreme limits. These are the weakly ($\gamma \ll 1$) and strongly ($\gamma \gg 1$) coupled domains. In the former limit many such efforts have led to the linearized Debye-Hückle result, using kinetic^{1,2,3}, statistical mechanical^{4,5,6}, and numerical^{7,8,9} formulations.

In the strong-coupling domain, studies have centered around such problems as plasma turbulence^{10,11} and electron correlations at metallic densities¹². Each approach has applied its own approximation of the correlation functions to the BBKGY hierarchy in order to obtain the dielectric response function¹³. The function thus acquired is then employed in conjunction with the fluctuation-dissipation theorem to obtain a self-consistent solution for the static form factor or, equivalently, the pair correlation function¹⁴.

A more recent study and review of calculations concerning the total internal energy of a one-component plasma in the strong-coupling limit is presented by Gould, et al.¹⁵ Comprehensive expositions on the state of the art of this subject have been given by Kalman and Carini;¹⁶ Baus and Hansen,⁸ and Ichimaru.¹⁴

The relevance of strongly-coupled plasmas to natural phenomena is illustrated in Fig. 1. We note that, in particular, the strong-coupling limit plays an important role in the description of x-ray plasmas, laser fusion devices, and in the interiors of certain super-dense stars.

In the numerical work of Brush, Sahlin and Teller⁷, the pair correlation function is evaluated over a large range of values of the plasma parameter, γ . They find that with increasing γ the

correlation function passes from a Debye-Hückle form to a decaying, oscillatory form. This numerical study further demonstrates that the effect of the background charge is to alter the effective interparticle interaction away from the bare Coulomb force.

In the present study a unified formulation for the theory of correlations in a one-component plasma is introduced which is valid for weakly and strongly-coupled plasmas. The theory is based on the BBKGY hierarchy and a sequential ordering of the correlation functions. This ordering is specified in terms of the plasma parameter γ . An ϵ -expansion imbedded within the ordering scheme permits an interactive technique of solution and renders the analysis self-consistent.

To better incorporate the role of the neutralizing background, the effective interparticle force is expanded about the bare Coulomb interaction. This inclusion is motivated by results of previous numerical studies.^{7,8} In both the weak and strongly coupled domains, closed space-time integro-differential equations are obtained for the two-particle correlation functions.

In the weak-coupling domain a time-independent analysis returns the well-known linearized Debye-Hückle correlation function.¹⁻⁹

In the strongly-coupled domain a second-order differential equation is obtained for the correlation function. The solution to this equation exhibits a decaying oscillatory structure for particle separations in excess of the order of the effective two-particle interaction range. These purely analytic results are in very good agreement with previous numerical work.⁷

The essential components of this analysis are related as shown in the flow chart given in Fig. 2.

II. Analysis

A. Basic Formulation

We consider an aggregate of charges Ze in a uniform neutralizing background. The s^{th} equation of the BBKGY hierarchy, hereafter call BY_s , is given by (see, for example, Liboff¹⁷)

$$\begin{aligned} & \left[\frac{\partial}{\partial t} + \sum_{i=1}^s \underline{p}_i \cdot \frac{\partial}{\partial \underline{x}_i} + \alpha_0 \sum_{i < j}^s \underline{G}_{ij} \cdot \left(\frac{\partial}{\partial \underline{p}_i} - \frac{\partial}{\partial \underline{p}_j} \right) \right] F_s \\ & = - \frac{\alpha_0}{4\pi\gamma_0} \sum_{i=1}^s \frac{\partial}{\partial \underline{p}_i} \cdot \left[d(s+1) \underline{G}_{i,s+1} F_{s+1} \right] \end{aligned} \quad (1)$$

Parameters are nondimensionalized as follows¹⁸, where barred variables are dimensional.

$$\begin{aligned} \bar{\underline{x}} &= r_0 \underline{x}, \quad \bar{\underline{p}} = p_0 \underline{p} \\ \bar{\underline{G}}_{ij} &= \frac{\phi_0}{r_0} \underline{G}_{ij}, \quad \phi_0 = \frac{Z^2 e^2}{r_0} \\ \bar{t} &= \frac{r_0 m}{p_0} t \end{aligned} \quad (2)$$

In addition,

$$p_0 = m C, \quad m C^2 = k_B T$$

The parameters α_0 and γ_0 in (1) which emerge as a consequence of nondimensionalization are,

$$\alpha_0 = \phi_0 / k_B T, \quad \frac{1}{\gamma_0} = 4\pi n r_0^3, \quad n = \frac{N}{V} \quad (3)$$

Here ϕ_0 and r_0 are, respectively, the strength and range of the interaction potential. For the case that $r_0 = \lambda_D$, the Debye distance, the coefficients α and γ are written without subscripts.

The nondimensional distribution is then given by¹⁹

$$F_s = (mC)^{3s} V^s \bar{f}_s \quad (4)$$

where V is the volume of the system and $\bar{f}_s$ is the s -particle joint probability distribution. The function F_s has the normalization,

$$\int F_s d\mathbf{r}_1 d\mathbf{r}_2 \dots d\mathbf{r}_s = 1 \quad (5)$$

where phase-volume elements are likewise dimensionless. The effective force on the i^{th} ion due to the j^{th} ion is $\bar{G}_{ij}$.

As noted in the introduction, it has been found in Monte Carlo calculations⁷ that the net effect of the background charge is to alter the effective interparticle force away from a simple Coulomb form. Accordingly we write

$$\bar{G}_{ij} = \hat{x}_{ij} \frac{\phi_0}{r_0} \frac{1}{x_{ij}^2} \left[1 - \sum_{n=1}^{\infty} \frac{u^{n_b} b_n^2}{x_{ij}^{n-1}} \right] \quad (6)$$

$$\bar{x}_{ij} = \bar{x}_i - \bar{x}_j \quad (6a)$$

The coefficient u is a parameter of smallness. The coefficients b_n will be addressed later in the paper. Note that the leading correction term in the series (6) has the effect of altering the magnitude but not the form of the Coulomb interaction.

The correlation functions are defined through the Mayer expansions²⁰

$$\begin{aligned} F_2(1,2) &= F_1(1) F_1(2) + C_2(1,2) \\ F_3(1,2,3) &= F_1(1) F_1(2) F_1(3) \\ &\quad + F_1(1) C_2(2,3) + F_1(2) C_2(3,1) + F_1(3) C_2(1,2) \\ &\quad + C_3(1,2,3) \\ &\quad \vdots \end{aligned} \quad (7)$$

One also writes

$$C_2(1,2) = F_1(1) F_1(2) h(1,2) \quad (8)$$

where $h(1,2)$ is the so-called 'total correlation function',²¹.

In the weak-coupling (or, 'correlationless') domain, one assumes

$$F_1 F_1 \gg C_2 \gg C_3 \dots \quad (9a)$$

By extension, in the strong-coupling domain we write

$$F_1 F_1 \ll C_2 \ll C_3 \dots \quad (9b)$$

These sequential inequalities may both be incorporated into the single set of expansions,

$$\begin{aligned} F_1 &= \gamma^0 [F_1^{(0)} + \epsilon F_1^{(1)} + \dots] \\ C_2 &= \gamma [C_2^{(0)} + \epsilon C_2^{(1)} + \dots] \\ C_3 &= \gamma^2 [C_3^{(0)} + \epsilon C_3^{(1)} + \dots] \end{aligned} \quad (10)$$

The ϵ -factors follow standard perturbation procedure and permit an iterative technique of solution. In (10) we have reintroduced the plasma parameter,

$$\gamma = \frac{1}{4\pi n \lambda_d^3} = \left(\frac{r_0}{\lambda_d} \right)^3 \gamma_0 \quad (11)$$

which in this instance serves as a bookkeeping index. With $\gamma \ll 1$, (10) is seen to return the weak-coupling sequence (9a), whereas with $\gamma \gg 1$, (10) yields the strong-coupling sequence (9b). Thus the expansion (10) serves as a unified description of plasma conditions. The Debye distance λ_d and plasma frequency ω_p which enter (11) are given by

$$\lambda_d^2 = \frac{1}{4\pi (eZ)^2} \frac{k_B T}{n}, \quad \omega_p^2 = \frac{4\pi n e^2 Z^2}{m}$$

$$\lambda_d^2 \omega_p^2 = c^2$$

With (11) we see that in the strong-coupling domain $n^{-1/3} \gg \lambda_d$, so that in this limit the Debye distance ceases to represent the range of the two-particle interaction.

It is convenient to introduce the parameter

$$\eta \equiv \left(\frac{\gamma}{\gamma_0} \right)^{1/3} = \frac{r_0}{\lambda_d} = \left(\frac{\alpha_0}{\gamma_0} \right)^{1/2} \quad (12)$$

Consequently the parameter α_0/γ_0 that multiplies the interaction integral in (1) may be rewritten as η^2 .

Another important plasma parameter found in the literature is

$$\Gamma \equiv (Ze)^2/k_B T a$$

$$\frac{4\pi}{3} n a^3 = 1$$

It is related to γ by

$$\gamma^2 = 3\Gamma^3$$

Using (12) we rewrite (1) in the more concise form

$$(\hat{K}_s + \alpha_0 \hat{B}_s) F_s = - \frac{\eta^2}{4\pi} \hat{I}_s F_{s+1} \quad (13)$$

Definition of the operators $\hat{K}_s$, $\hat{B}_s$ and $\hat{I}_s$ follows by comparison with (1). Furthermore we set

$$\begin{aligned} \hat{B}_s &= \sum_{n=0}^{\infty} u^n \hat{B}_s^{(n)} \\ \hat{I}_s &= \sum_{n=0}^{\infty} u^n \hat{I}_s^{(n)} \end{aligned} \quad (14)$$

in keeping with expansion (6).

In the context of this formalism it is possible to summarize the various possible limiting domains in the form of an α_0 - η diagram¹⁷ as shown in Fig. 3.

B. The Strongly Coupled Domain

In this limit we take $\alpha_0 \sim 1$, and the relation $\gamma = \alpha_0 \eta$ gives $\eta = \gamma$. Thus BY_1 and BY_2 become, in accord with expansion (7),

$$\hat{K}_1 F_1 = -\frac{\gamma^2}{4\pi} \hat{I}_1 [F_1(1)F_1(2) + C_2] \quad (15a)$$

$$\begin{aligned} (\hat{K}_2 + \alpha_0 \hat{B}_2) [F_1(1)F_1(2) + C_2] = \\ -\frac{\gamma^2}{4\pi} \hat{I}_2 [F_1(1)F_1(2)F_1(3) + \sum_{P(ijk)} F_1(i)C_2(j,k) \\ + C_3(1,2,3)] \end{aligned} \quad (15b)$$

Here the symbol $P(i,j,k)$ denotes summation over permutations of ijk .

Our chief aim at this point is to obtain a closed kinetic equation for $C_2^{(0)}(1,2)$. To this end a study was made of the ϵ -dependence of the parameters γ and α , with $\alpha_0 \sim 1$. It was found that $\gamma^2 \sim \epsilon^{-1/2}$ and $\alpha\gamma^2 \sim 1$ are the simplest forms which give closure. (See Appendix A.) Applying the expansions (10) and (14) to (15) with these ϵ -dependencies of α_0 , γ and α gives the following leading perturbative equations relevant to determining $C_2^{(0)}$.

$$-\frac{\gamma^3}{4\pi} \hat{I}_1^{(0)} C_2^{(0)} = 0 \quad (16a)$$

$$-\frac{\gamma^2}{4\pi} \hat{I}_1^{(0)} F_1(1)F_1(2) = 0 \quad (16b)$$

$$(\hat{K}_2 + \alpha_0 \hat{B}_2^{(0)}) C_2^{(0)} = -\frac{\gamma^2}{4\pi} \hat{I}_2^{(1)} \sum_{P(ijk)} F_1(i)C_2^{(0)}(j,k) \quad (17)$$

Here we have written F_1 for $F_1^{(0)}$. We note in passing that (17) is a closed space-time kinetic equation for $C_2^{(0)}(1,2)$. Constraints (16a,b) are consistent with an assumption of spatial homogeneity, so that we may write

$$C_2^{(0)}(i,j) = F_1(p_i)F_1(p_j)h^{(0)}(x_{ij}) \quad (18)$$

The right-hand side of (17) is simplified by virtue of (16a) and the vanishing of any space integral over an isotropic vector field. Consequently two terms survive given by

$$\text{RHS}(17) = - \frac{\gamma^2 b_1^2}{4\pi} \{ F_1(2) \frac{\partial}{\partial p_1} F_1(1) \cdot J_1^{(1,0)} + F_1(1) \frac{\partial}{\partial p_2} F_1(2) \cdot J_2^{(1,0)} \};$$

[Go to p. 8]

where

$$b_n^2 J_1^{(n,m)} \equiv \int d^3 G_{13}^{(n)} h^{(m)}(x_{23}) \quad (19a)$$

$$b_n^2 J_2^{(n,m)} \equiv \int d^3 G_{23}^{(n)} h^{(m)}(x_{13}) \quad (19b)$$

With reference to the results of Appendix B, we may write (dropping the superscript on h)

$$J_1^{(1,0)} = -4\pi \hat{x} x^{-2} \int_0^x d\rho \rho^2 h(\rho) \quad (20a)$$

$$J_2^{(1,0)} = -J_1^{(1,0)} \quad (20b)$$

Here and henceforth we set $x \equiv x_{12}$.

With (17) and (19) in (16) we obtain

$$\begin{aligned} & (\hat{K}_2 + \alpha_0 \hat{B}_2^{(0)}) F_1(p_1) F_1(p_2) h(x) \\ & = b_1^2 \gamma^2 \left[\frac{\partial}{\partial p_1} - \frac{\partial}{\partial p_2} \right] F_1(p_1) F_1(p_2) \cdot \hat{x} x^{-2} \int_0^x d\rho \rho^2 h(\rho) \end{aligned} \quad (21)$$

In the space-dependent part of $\hat{K}_2$ we use the transformation

$$x_1 + x_2 \equiv X \quad x_1 - x_2 \equiv x$$

in which case

$$\hat{K}_2 h(x) = \hat{x} \cdot (p_1 - p_2) \frac{\partial}{\partial x} h(x) \quad (22)$$

where we have neglected the time derivative in $\hat{K}_2$. With the repulsive bare Coulomb interaction, it follows from (6) and (14) that

$$\hat{B}_2^{(0)} = \hat{x} \cdot x^{-2} \left[\frac{\partial}{\partial p_1} - \frac{\partial}{\partial p_2} \right] \quad (23)$$

Employing the preceding two expressions in (21) gives

$$\begin{aligned}
\hat{x} \cdot (p_1 - p_2) F_1 F_1 \frac{\partial}{\partial x} h(x) + \alpha_0 \hat{x} \cdot x^{-2} \left[\frac{\partial}{\partial p_1} - \frac{\partial}{\partial p_2} \right] F_1(p_1) F_1(p_2) h \\
= b_1^2 \gamma^2 \left[\frac{\partial}{\partial p_1} - \frac{\partial}{\partial p_2} \right] F_1(p_1) F_1(p_2) \cdot \hat{x} x^{-2} \int_0^x d\rho \rho^2 h(\rho)
\end{aligned} \quad (24)$$

Assuming finally that F_1 is isotropic in momentum space, we may write

$$\frac{\partial}{\partial p} F_1(p) = \hat{p} \frac{\partial}{\partial p} F_1(p) \quad (25)$$

and (24) becomes

$$\begin{aligned}
\hat{x} \cdot p_1 \left[F_1 F_1 \frac{\partial}{\partial x} h(x) + \alpha_0 x^{-2} \frac{F_1(p_2)}{p_1} h(x) \frac{\partial}{\partial p_1} F_1(p_1) \right. \\
\left. - b_1^2 \gamma^2 \frac{F_1(p_2)}{p_1} \frac{\partial}{\partial p_1} F_1(p_1) x^{-2} \int_0^x d\rho \rho^2 h(\rho) \right] \\
= \hat{x} \cdot p_2 \left[F_1 F_1 \frac{\partial}{\partial x} h(x) + \alpha_0 x^{-2} \frac{F_1(p_1)}{p_2} h(x) \frac{\partial}{\partial p_2} F_1(p_2) \right. \\
\left. - b_1^2 \gamma^2 \frac{F_1(p_1)}{p_2} \frac{\partial}{\partial p_2} F_1(p_2) x^{-2} \int_0^x d\rho \rho^2 h(\rho) \right]
\end{aligned} \quad (26)$$

Operating on this equation with $\int dp_2$ eliminates the RHS, and since $\hat{x} \cdot p_1$ is generally nonvanishing, there remains

$$\begin{aligned}
F_1(p_1) \frac{\partial}{\partial x} h(x) + \frac{\alpha_0}{x^2} h(x) \frac{1}{p_1} \frac{\partial}{\partial p_1} F_1(p_1) \\
- b_1^2 \gamma^2 \frac{1}{p_1} \frac{\partial}{\partial p_1} F_1(p_1) x^{-2} \int_0^x d\rho \rho^2 h(\rho) = 0
\end{aligned} \quad (27)$$

Integration over p_1 gives

$$\frac{\partial}{\partial x} h(x) - \alpha_0 \gamma^2 x^{-2} h(x) + b_1^2 \gamma^2 x^{-2} \int_0^x d\rho \rho^2 h(\rho) = 0 \quad (28)$$

where

$$y^2 \equiv - \int dp_1 \frac{1}{p_1} \frac{3}{3p_1} F_1(p_1) \quad (29)$$

and may be assumed positive.

Differentiating (28) with respect to x gives

$$\begin{aligned} h''(x) + \frac{2\alpha_0 y^2}{x^3} h(x) - \alpha_0 y^2 x^{-2} h'(x) \\ + b_1^2 \gamma^2 y^2 \left[h(x) - \frac{2}{x^3} \int_0^x d\rho \rho^2 h(\rho) \right] = 0 \end{aligned} \quad (30)$$

Using (28) to solve for the integral term in (30) yields the differential equation

$$h'' + [2y^{-1} - \kappa y^{-2}] h' + h = 0 \quad (31)$$

where

$$y \equiv kx \quad k \equiv b_1 \gamma \quad \kappa \equiv \alpha_0 y^2 k \quad (31a)$$

Employing the transformation ²³

$$\begin{aligned} h(y) &= \frac{1}{y} e^{-\sigma/y} u(y) \\ \sigma &\equiv \kappa/2 \end{aligned} \quad (32)$$

in (31) gives

$$u'' + (1 - \frac{\sigma^2}{y^4}) u = 0 \quad (33)$$

This equation is Schroedinger-like in form and we may conclude²⁴ that for $y^2 > \sigma$, $h(y)$ is oscillatory, whereas for $y^2 < \sigma$, $h(y)$ is exponential. The explicit form of the solutions in these domains may be obtained in two extremes:

$$\underline{y^2 \gg \sigma}$$

We are left with

$$u'' + u = 0$$

which has the solution

$$u = A \cos y + B \sin y$$

With (32) we obtain

$$h^>(y) = \frac{e^{-\kappa/2y}}{y} [A \sin y + B \cos y] \quad (35)$$

$$y^2 \ll \sigma$$

In this case (33) reduces to

$$u'' - \frac{\sigma^2}{y^4} u = 0 \quad (36)$$

which has the solution²⁴

$$u(y) = y(\bar{A} e^{\sigma/y} + \bar{B} e^{-\sigma/y})$$

With (32), this gives

$$h^<(y) = \bar{B} e^{-\kappa/y} \quad (37)$$

where we have set $\bar{A} = 0$ because (31) does not allow a constant solution. As previously noted, the oscillatory behavior (35) of $h(y)$ in the domain $y^2 \gg \sigma$, is in very good agreement with previous numerical studies^{7,8}.

We now estimate the value of the leading coefficient b_1 in (6). This is accomplished through comparison of the wave number K (defined below) of $h^>(y)$ with the numerical value k_{MC} obtained in Monte Carlo studies^{7,8}. These results imply that in the vicinity of $\gamma \approx 50$ (i.e., $\Gamma \approx 10$)

$$k_{MC} \approx \frac{2\pi}{3a} = \epsilon/\lambda_d \gamma^{1/3} ; \quad a^3 = 3/4\pi n$$

$$3 = 1.5 , \quad \epsilon^2 = 8.4$$

The wavenumber K is given by

$$y = K\bar{x} \quad (38)$$

$$K = \frac{k}{r_0} = \frac{b_1 \gamma \gamma}{r_0}$$

From (11), $\lambda_d = r_0/\gamma$. Furthermore, assuming a Maxwellian form for

$F_1(p_1)$ gives $\gamma = 1$. With these values

$$K = b_1/\lambda_d$$

and comparison with k_{MC} yields

$$b_1 = \xi\gamma^{-1/3} \approx \kappa_0 3^{-0.17} \Gamma^{-1/2}$$

which with the values cited above gives $b_1 = 0.8$. With this result at hand, we may estimate the value of interparticle separation at which oscillation of $h(\bar{x})$ ensues. Such oscillation was found from (33) to occur for

$$y^2 \geq \kappa/2$$

or, equivalently

$$\left(\frac{\bar{x}}{r_0}\right)^2 \geq \frac{1}{2b_1\gamma}$$

Inserting our previous finding

$$b_1\gamma = \xi\gamma^{2/3} \approx \gamma^{2/3},$$

gives

$$\bar{x} \geq r_0/\gamma^{1/3}$$

Thus oscillation of $h(\bar{x})$ may be expected for interparticle displacement greater than or equal to the range of the two-particle interaction. The initiation of such oscillation may be interpreted as the onset of phase transition in the plasma^{7,32}. Furthermore the wavenumber K of these oscillations (38), is seen to grow with γ . This behavior may be associated with the fluid-solid phase transition³³ at large γ , found in numerical studies.^{7,34}

C. The Weak-Coupling Case

This situation has been studied by many individuals¹⁻⁹ and as we have noted previously it gives the linearized Debye-Hückle result. In the kinetic domain this limit yields the Vlasov equation, or more generally, the Balescu-Lenard Equation.²⁶⁻³³

An essential element of these studies is that the correlation functions are assumed to be perturbatively small. Accordingly we set $\gamma \sim \epsilon$ in (10). Insofar as $r_0 = \lambda_D$ in this limit, it follows that $n^2 = \alpha/\gamma = 1$, which fixes $\alpha \sim \epsilon$. We note that this limit is equivalent to the classical Rosenbluth-Rostoker ansatz.³¹ With these constraints and $u \sim \epsilon$, (6) gives the following leading equations for $C_2^{(0)}$ as derived from (13)

$$\hat{K}_1 F_1^{(0)} = - \frac{n^2}{4\pi} \hat{I}_1^{(0)} F_1^{(0)} F_1^{(0)} \quad (39a)$$

$$(39b)$$

$$\hat{K}_1 F_1^{(1)}(1) = - \frac{n^2}{4\pi} \hat{I}_1^{(0)} [C_2^{(0)}(1,2) + F_1^{(0)}(1) F_1^{(1)}(2) + F_1^{(1)}(1) F_1^{(0)}(2)]$$

$$\hat{K}_2 [C_2^{(0)}(1,2) + F_1^{(0)}(1) F_1^{(1)}(2) + F_1^{(1)}(1) F_1^{(0)}(2)] + \alpha \hat{B}_2^{(0)} F_1^{(0)}(1) F_1^{(0)}(2) =$$

$$(39c)$$

$$- \frac{n^2}{4\pi} \hat{I}_2^{(0)} \left[\sum_{P(ijk)} F_1^{(0)}(i) C_2^{(0)}(jk) + \sum_{P(ijk)} F_1^{(0)}(i) F_1^{(0)}(j) F_1^{(1)}(k) \right]$$

We must solve (39) and (40) simultaneously for $C_2^{(0)}(1,2)$. First note that (39a) may be rewritten in two equivalent ways:

$$F_1^{(1)}(2) \hat{K}_1(1) F_1^{(0)}(1) = - \frac{n^2}{4\pi} F_1^{(1)}(2) \hat{I}_1^{(0)}(1) F_1^{(0)}(1) F_1^{(0)}(3)$$

$$F_1^{(1)}(1) \hat{K}_1(2) F_1^{(0)}(2) = - \frac{n^2}{4\pi} F_1^{(1)}(1) \hat{I}_1^{(0)}(2) F_1^{(0)}(2) F_1^{(0)}(3)$$

Similarly (39b) gives

$$F_1^{(0)}(2) \hat{K}_1(1) F_1^{(1)}(1) = -\frac{\gamma^2}{4\pi} F_1^{(0)}(2) \hat{I}_1^{(0)}(1) [C_2^{(0)}(1,3) + F_1^{(0)}(1) F_1^{(1)}(3) + F_1^{(1)}(1) F_1^{(0)}(3)]$$

$$F_1^{(0)}(1) \hat{K}_1(2) F_1^{(1)}(2) = -\frac{\gamma^2}{4\pi} F_1^{(0)}(1) \hat{I}_1^{(0)}(2) [C_2^{(0)}(2,3) + F_1^{(0)}(2) F_1^{(1)}(3) + F_1^{(1)}(2) F_1^{(0)}(3)]$$

Adding these four equations and subtracting the result from (39c) yields

$$\hat{K}_2 C_2^{(0)}(1,2) + \gamma \hat{B}_2^{(0)} F_1^{(0)} F_1^{(0)} = -\frac{\gamma^2}{4\pi} \{ \hat{I}_2^{(0)} \frac{1}{P(ijk)} F_1^{(0)}(i) C_2^{(0)}(jk) - F_1^{(0)}(2) \hat{I}_1^{(0)}(1) C_2^{(0)}(1,3) - F_1^{(0)}(1) \hat{I}_1^{(0)}(2) C_2^{(0)}(2,3) \}$$

Four terms remain on the RHS of this equation. Of these, two vanish by integration of the isotropic vector field G_{ij} , leaving

$$\hat{K}_2 C_2^{(0)}(1,2) + \gamma \hat{B}_2^{(0)} F_1^{(0)} F_1^{(0)} = -\frac{\gamma^2}{4\pi} \{ \hat{I}_1^{(0)}(1) F_1^{(0)}(1) C_2^{(0)}(2,3) + \hat{I}_1^{(0)}(2) F_1^{(0)}(2) C_2^{(0)}(1,3) \} \quad (40)$$

Here we have obtained (as in the strongly-coupled analysis) a closed space-time-dependent equation of motion for $C_2^{(0)}$. The RHS of this equation simplifies to

$$\text{RHS}(40) = -\frac{\gamma^2}{4\pi} \{ F_1^{(0)}(2) \frac{1}{P_1} F_1^{(0)}(1) \cdot J_1^{(0,0)} + F_1^{(0)}(1) \frac{1}{P_2} F_1^{(0)}(2) \cdot J_2^{(0,0)} \} \quad (41)$$

where the J integrals are defined in (19), and again we find that

$$J_1^{(0,0)} = 4\pi \hat{x} x^{-2} \int_0^x d\rho \rho^2 h(\rho) \quad (42a)$$

$$J_1^{(0,0)} = - J_2^{(0,0)} \quad (42b)$$

[Go to p. 14]

Substituting (18) and (42) into (40) we obtain

$$\begin{aligned} \hat{K}_2 F_1 F_1 h(x) + \alpha \hat{B}_2^{(0)} F_1 F_1 = \\ - \eta^2 \left[\frac{\partial}{\partial p_1} - \frac{\partial}{\partial p_2} \right] F_1 F_1 \cdot \hat{x} x^{-2} \int_0^x d\rho \rho^2 h(\rho) \end{aligned} \quad (43)$$

Making use of (22) and (23) in the LHS of (43) gives

$$\begin{aligned} \hat{x} \cdot (p_1 - p_2) F_1 F_1 h + \alpha \hat{x} \cdot x^{-2} \left(\frac{\partial}{\partial p_1} - \frac{\partial}{\partial p_2} \right) F_1 F_1 = \\ - \eta^2 \left[\frac{\partial}{\partial p_1} - \frac{\partial}{\partial p_2} \right] F_1 F_1 \cdot \hat{x} x^{-2} \int_0^x d\rho \rho^2 h(\rho) \end{aligned} \quad (44)$$

Furthermore, with (25) the preceding equation becomes

$$\begin{aligned} \hat{x} \cdot p_1 \left[F_1 F_1 \frac{\partial}{\partial x} h + \frac{\alpha}{x^2} \frac{F_1(p_2)}{p_1} \frac{\partial}{\partial p_1} F_1(p_1) \right. \\ \left. + \eta^2 \frac{F_1(p_2)}{p_1} \frac{\partial}{\partial p_1} F_1(p_1) x^{-2} \int_0^x d\rho \rho^2 h(\rho) \right] \\ = \hat{x} \cdot p_2 \left[F_1 F_1 \frac{\partial}{\partial x} h + \frac{\alpha}{x^2} \frac{F_1(p_1)}{p_2} \frac{\partial}{\partial p_2} F_1(p_2) \right. \\ \left. + \eta^2 \frac{F_1(p_1)}{p_2} \frac{\partial}{\partial p_2} F_1(p_2) x^{-2} \int_0^x d\rho \rho^2 h(\rho) \right] \end{aligned} \quad (45)$$

Repeating the procedure leading to (27), we find that (45) yields

$$F_1(p_1) \frac{\partial}{\partial x} h + \frac{\alpha}{x^2} \frac{1}{p_1} \frac{\partial}{\partial p_1} F_1(p_1) + \eta^2 \frac{1}{p_1} \frac{\partial}{\partial p_1} F_1(p_1) x^{-2} \int_0^x d\rho \rho^2 h(\rho) = 0 \quad (46)$$

Integration over p_1 gives

$$\frac{\partial}{\partial x} h - \frac{\alpha}{x^2} y^2 - \eta^2 y^2 x^{-2} \int_0^x d\rho \rho^2 h(\rho) = 0 \quad (47)$$

Differentiating (47) leads to

$$h'' + \frac{2\alpha}{x^3} y^2 + 2n^2 Y^2 x^{-3} \int_0^x d\rho \rho^2 h(\rho) - n^2 Y^2 h = 0 \quad (48)$$

Combining the last two equations gives

$$h'' + \frac{2h'}{y} - h = 0 \quad (49)$$

where

$$y = nYx \quad (49a)$$

The general solution to (48) may be written

$$h(y) = D_1 \frac{e^{-y}}{y} + D_2 \frac{e^y}{y} \quad (50)$$

We need to affirm that (49) is a solution to the parent integral equation (47), which in terms of y is

$$\frac{d}{dy} h - \frac{1}{y^2} \left[\alpha Y^3 n + \int_0^y dy y^2 h(y) \right] = 0 \quad (51)$$

Inserting (49) into (50) leads to the constraint

$$D_1 + D_2 = -\alpha Y^3 n \quad (52)$$

Setting $D_2 = 0$ for physical reasons leaves the solution

$$h(y) = -\frac{\alpha Y^3 n}{y} e^{-y} \quad (53)$$

For a Maxwellian, $Y = 1$. Furthermore, in the present limit we recall that $n = 1$, so that $y = x$, and (52) may be written

$$h(x) = -x \frac{e^{-x}}{x} \quad (54)$$

This is the well-known linearized Debye-Hückle result³⁵, which follows to first-order in α from an expansion of the more precise nonlinear Debye-Hückle form

$$h(x) = e^{-\phi_d(x)} - 1 \quad (55)$$

where

$$\phi_d(x) = \alpha \frac{e^{-x}}{x} \quad (55a)$$

Whereas the Debye-Hückle form (54) agrees with the linearized result (53) for $x \gg 1$, we see that at the origin $h(x)$ diverges, but h_d has the correct value $h_d(0) = -1$.

III. Conclusions

We have described a unified formulation of correlations in a one-component plasma, based on the plasma parameter γ , which is valid in both the weakly and strongly-coupled domains. The theory is based on the BBKGY hierarchy together with expansions of the two-particle interaction and correlation functions.

In the weak-coupling domain, the analysis returns the familiar linearized Debye-Hückle correlation function. In the strong-coupling domain a second-order differential equation for the total correlation function is found which yields an exponentially damped solution near the origin. For particle separation on the order of the range of the two-particle interaction, the equation gives a Bessel-like oscillatory solution. These features of the correlation function are in excellent agreement with previous numerical studies.

In both domains, closed space-time dependent integro-differential equations were obtained for the two-particle correlation functions. We anticipate that future study will find important applications of these results both in plasma and condensed matter physics.

Appendix A

Here we examine the decoupling condition on γ employed in Section B. With expansions (6) and (10) in BY_2 (15b), we write

$$\begin{aligned} & \left[\hat{K}_2 + \alpha_0 \sum_{\tau=0}^{\infty} \mu^{\tau} \hat{B}_2^{(\tau)} \right] \left[F_1 F_1 + \gamma \sum_{n=0}^{\infty} \varepsilon^n C_2^{(n)} \right] \\ & = - \frac{\gamma^2}{4\pi} \left[\sum_{r=0}^{\infty} \mu^r \hat{I}_2^{(r)} \right] \left[F_1 F_1 F_1 + \gamma \sum_{n=0}^{\infty} \varepsilon^n C_2^{(n)} + \gamma^2 \sum_{m=0}^{\infty} \varepsilon^m C_3^{(m)} \right] \end{aligned} \quad (A1)$$

The F_1 functions are expanded as in (10). We seek a kinetic equation for $C_2^{(0)}$ consistent with $\alpha_0 = 0(1)$. In general, such an equation includes $\hat{K}$, $\hat{B}$ and $\hat{I}$ terms. The mere requirement of a $\hat{K}$ term indicates that the equation sought must be $O(\gamma)$, which leads to the condition

$$\gamma = \mu^t \gamma = \mu^r \gamma^3 \quad (A2)$$

The leftmost equality gives $t=0$, with $\mu \neq 1$. This specifies inclusion of the $\hat{B}_2^{(0)}$ term in our equation. The rightmost equality gives

$$\mu^r \gamma^2 = 1$$

The simplest choice which includes effects of the background is $r=1$ which specifies inclusion of the $\hat{I}_2^{(1)}$ term and

$$\gamma^2 \mu = 1 \quad (A3)$$

Finally, decoupling the $I_2^{(1)} C_2^{(0)}$ term from all other $C^{(n)}$ terms [in the RHS of (A1)] requires that

$$\mu \gamma \neq \mu^q \varepsilon^n \quad q > 0, \quad n > 1 \quad (A4)$$

$$\mu \gamma \neq \gamma^2 \mu^s \varepsilon^m \quad s > 0, \quad m > 0 \quad (A5)$$

These relations give, respectively,

$$\epsilon^{q-1} \neq 1 \quad q \geq 0, \quad n \geq 1 \quad (\text{A6})$$

$$\epsilon^{s-1} \neq 1 \quad s \geq 0, \quad m \geq 0 \quad (\text{A7})$$

Using (A3) in (A6) yields

$$\gamma \neq \epsilon^{n/2(q-1)} \quad q \geq 0, \quad n \geq 1 \quad (\text{A8})$$

The strong-coupling condition ($\gamma \geq 1$) requires

$$\epsilon^{n/2(q-1)} \geq 1, \quad q \geq 0, \quad n \geq 1 \quad (\text{A9})$$

This allows as the only possibility, $q=0$. Therefore, (A8) requires

$$\gamma \neq \epsilon^{-n/2} \quad n \geq 1 \quad (\text{A10})$$

Similarly, with (A3) in (A7) it follows that

$$\epsilon^{m/2s-3} \geq 1, \quad s \geq 0, \quad m \geq 0 \quad (\text{A11})$$

This leads to the cases $s=0,1$ and thus

$$\gamma \neq \epsilon^{-m/3}, \quad \epsilon^{-m} \quad m \geq 0 \quad (\text{A12})$$

Combining results (A10) and (A12), the final conditions on γ are summarized by

$$\gamma = \epsilon^{-a/b} \quad a \geq 1, \quad b \geq 4, \quad \frac{a}{b} \neq m, \quad \frac{m}{2}, \quad \frac{m}{3} \quad (\text{A13})$$

where $m \geq 0$.

Any one of these allowable orders of γ leads to the sought-after decoupled equation for $C_2^{(0)}$ in the strong-coupling limit. In this analysis we select the simplest case: $a=1$, $b=4$, corresponding to

$$\gamma^2 = \varepsilon^{-1/2} \quad (A14)$$

[Go to p. 20]

Appendix B

Consider the integral (19a). We set

$$\rho \equiv x_{23} \equiv x_2 - x_3 \quad (B1)$$

$$x_{13} = \rho + x$$

in which case for $n = 1$, $m = 0$, we find

$$J_1^{(1,0)} = \frac{\partial}{\partial x} \int d\rho \frac{h(\rho)}{|\rho + x|} \quad (B2)$$

In spherical coordinates with

$$d\rho \equiv \rho^2 \sin \theta \, d\theta \, d\phi \, d\phi$$

$$\rho \cdot x = x\rho \cos \theta \quad (B3)$$

$$|\rho + x| = (\rho^2 + 2x\rho \cos \theta + x^2)^{\frac{1}{2}}$$

and x held fixed along the z -axis throughout the integration we obtain

$$J_1^{(1,0)} = 2\pi \frac{\partial}{\partial x} \int_0^\infty d\rho \, \rho^2 h(\rho) \int_0^\pi d\theta \sin \theta (\rho^2 + 2x\rho \cos \theta + x^2)^{-\frac{1}{2}}$$

Performing the θ -integration gives

$$J_1^{(1,0)} = -2\pi \frac{\partial}{\partial x} \int_0^\infty d\rho \, \rho^2 h(\rho) \left\{ \frac{1}{x\rho} [\rho^2 + 2x\rho \cos \theta + x^2]^{\frac{1}{2}} \right\} \Big|_0^\pi$$

or, equivalently,

$$J_1^{(1,0)} = -2\pi \frac{\partial}{\partial x} \int_0^\infty d\rho \, \frac{\rho}{x} h(\rho) [(\rho^2 - 2x\rho + x^2)^{\frac{1}{2}} - (\rho^2 + 2x\rho + x^2)^{\frac{1}{2}}]$$

$$= -2\pi \frac{\partial}{\partial x} \int_0^\infty d\rho \, \frac{\rho}{x} h(\rho) [|\rho - x| - |\rho + x|]$$

$$= 4\pi \frac{\partial}{\partial x} \left[\int_0^x d\rho \, \frac{\rho^2}{x} h(\rho) + \int_x^\infty d\rho \, \rho h(\rho) \right]$$

Differentiating, one finds

$$J_1^{(1,0)} = -4\pi \hat{x} x^{-2} \int_0^x d\rho \rho^2 h(\rho) \quad (B4)$$

in agreement with (20a).

For the integral (19b) we set

$$\rho \equiv x_{13} \equiv x_1 - x_3 \quad (B5)$$

$$x_{23} = \rho - x$$

which leads to

$$J_2^{(1,0)} = -\frac{\partial}{\partial x} \int d\rho \frac{h(\rho)}{|\rho - x|} \quad (B6)$$

Repeating the preceding calculations gives

$$J_2^{(1,0)} = -J_1^{(1,0)} \quad (B7)$$

We turn next to the integrals in (41), in which case $n = 0$, $m = 0$.

With (B1) we find

$$J_1^{(0,0)} = -\frac{\partial}{\partial x} \int d\rho \frac{h(\rho)}{|\rho + x|} \quad (B8)$$

and with (B5) we obtain

$$J_2^{(0,0)} = \frac{\partial}{\partial x} \int d\rho \frac{h(\rho)}{|\rho - x|} \quad (B9)$$

Following the analysis above gives

$$J_1^{(0,0)} = 4\pi \hat{x} x^{-2} \int_0^x d\rho \rho^2 h(\rho)$$

$$J_2^{(0,0)} = -J_1^{(0,0)}$$

Acknowledgements

We are indebted to our colleagues, K. C. Liu, Kenneth Gardner, and Stephen Ralph for their careful reading of the final manuscript, and to Z. Zinamon for his knowledgeable comments on this topic. This research was supported in part by Contract No. AFOSR 78-3574 between Cornell University and the United States Air Force.

Figure Captions

Figure 1

The plasma parameter γ for n and T (with $Z = 1$) showing weakly and strongly coupled (shaded) domains.

Figure 2

Flow chart explaining the analysis.


Figure 3

α_0 - γ plot showing weakly coupled (WCP) and strongly coupled (SCP) plasma domains.

References

1. E.A. Freeman and D.L. Book, Phys. Fluids 6, 1700 (1963).
2. L.C. Levitt, J.M. Richardson and E.R. Cohen, Phys. Fluids 10, 406 (1967).
3. R. Cauble and J.J. Duderstadt, Phys. Rev. A 23, 3182 (1981).
4. D.D. Carley, Phys. Rev. 131, 1406 (1963).
5. D.L. Bowers and E.E. Salpeter, Phys. Rev. 119, 1180 (1960).
6. M.S. Cooper, Phys. Rev. A 7, 1 (1973).
7. S.G. Brush, H. L. Sahlin and E. Teller, J. Chem. Phys. 45, 2102 (1966).
8. J.P. Hansen, Phys. Rev. A 8, 3096 (1973); _____ and M. Baus, Phys. Repts., 59, 1 (1980).
9. C.W. Hirt, Phys. Fluids 10, 565 (1967).
10. S. Ichimaru and T. Nakono, Phys. Lett. 25A, 168 (1967).
11. S. Ichimaru and T. Nakono, Phys. Rev. 165, 231 (1968).
12. K.I. Golden, G. Kalman and M.B. Silevitch, Phys. Rev. Lett. 33, 1954 (1974).
13. G. Kalman, T. Datta and K.I. Golden, Phys. Rev. A 12, 1125 (1975).
14. S. Ichimaru, Phys. Rev. A 2, 494 (1970), Revs. Mod. Phys. 54, 1017 (1982).
15. H. Gould, R.G. Palmer and G.A. Esteve, J. Stat. Phys. 21, 55 (1979).
16. G. Kalman and P. Carini, Strongly Coupled Plasmas (Plenum Press, New York, 1978).
17. R.L. Liboff, Introduction to the Theory of Kinetic Equations (Krieger, Melbourne, FL, 1979).
18. G. Sandri, Ann. Phys. (N.Y.) 24, 332 (1963).
19. N.N. Bogoliubov, Problemi Dynamicheskij Teorii v Statisticheskij Fizike (OGIS, Moscow, 1946).
20. J.E. Mayer and M.G. Mayer, Statistical Mechanics (Wiley, New York, 1940).
21. D.L. Goodstein, States of Matter (Prentice-Hall, Englewood Cliffs, N.J., 1975).

22. In preparation.
23. E. T. Whittaker and G.N. Watson, A Course of Modern Analysis (Cambridge University, Cambridge, 1952), 4th Ed.
24. R.L. Liboff, Introductory Quantum Mechanics (Holden Day, San Francisco, 1981).
25. E. Kamke, Differentialgleichungen (Akademische Verlagsgesellschaft, Leipzig, 1943), Vol. I, 2nd Ed.
26. R. Balescu, Phys. Fluids 3, 52 (1960).
27. A. Lenard, Ann. Phys. (N.Y.) 3, 390 (1960).
28. R. L. Liboff and A. H. Merchant, J. Math. Phys. 14, 119 (1973).
29. R. L. Liboff and G. E. Perona, J. Math. Phys. 8, 2001 (1967).
30. T. H. Dupree, Phys. Fluids 4, 696 (1961).
31. N. Rostoker and M. N. Rosenbluth, Phys. Fluids 3, 1 (1960).
32. R. L. Liboff and L. J. Caroff, J. Plasma Phys. 4, 83 (1970).
33. R. L. Liboff, Phys. Rev. 131, 2318 (1963).
34. E. L. Pollock and J. P. Hansen, Phys. Rev 8A, 3110 (1973).
35. P. Debye and E. Hückel, Phys. Z. 24, 185 (1923).



Values of the plasma parameter γ as a function of electron number density and temperature.

FIGURE 1




FIG. 2




FIG. 3

END

DATE
FILMED

9 - 83

DTIC