

AD-A131 906

ASYMPTOTIC MAXIMAL DEVIATION OF M-SMOOTHERS(U) NORTH
CAROLINA UNIV AT CHAPEL HILL DEPT OF STATISTICS
W HAERDLE APR 83 MMS-1523 AFOSR-TR-83-0688

1/1


UNCLASSIFIED

F49620-82-C-0009

F/G 12/1

NL

									END DATE FILMED 9 83 DTIC				



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A131906

AFOSR-TR- 83 - 0688

(2)



ASYMPTOTIC MAXIMAL DEVIATION OF H-SMOOTHERS

by

Wolfgang Härdle

Universität Heidelberg
Sonderforschungsbereich 123
Im Neuenheimer Feld 293
D-6900 Heidelberg 1

University of North Carolina
Department of Statistics
321 Phillips Hall 039A
Chapel Hill, NC 27514

Institute of Statistics Mimeo Series #1523

April 1983

DEPARTMENT OF STATISTICS
Chapel Hill, North Carolina

Approved for public release;
distribution unlimited.

83 08 19 07 7

DTIC FILE COPY

DTIC
ELECTE
S AUG 29 1983 D
B

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS. BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR- 83-0688	2. GOVT ACCESSION NO. AD-A131906	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) "ASYMPTOTIC MAXIMAL DEVIATION OF M-SMOOTHERS"		5. TYPE OF REPORT & PERIOD COVERED TECHNICAL
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Wolfgang Hardle		8. CONTRACT OR GRANT NUMBER(s) F49620-82-C-0009
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of North Carolina Department of Statistics Chapel Hill, NC 27514		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS PE61102F: 2304/A5
11. CONTROLLING OFFICE NAME AND ADDRESS AFOSR/NM Building 410 Bolling AFB DC 20332		12. REPORT DATE April 1983
		13. NUMBER OF PAGES 19
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Robust nonparametric regression, Kernel estimators, Uniform consistency, Robust smoothing		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		

DD FORM 1 JAN 73 1473

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

Let $(X_1, Y_1), \dots, (X_n, Y_n)$ be iid rv's with pdf $f(x, y)$ and let $m(x) = E(Y|X = x) = \int y f(x, y) dy / f_X(x)$ be the regression function of Y on X . The function $m(x)$ is estimated by $m_n(x)$ a solution of $(nh)^{-1} \sum_{i=1}^n K((x - X_i)/h) \psi(Y_i - m_n(x)) = 0$ for some odd and bounded ψ -function making $m_n(x)$ a robust estimate of $m(x)$. Probabilities of maximal deviation of $|m_n(x) - m(x)|$ are computed in a similar way as in Bickel and Rosenblatt (1973) for density estimation and in Johnston (1982) for nonparametric regression function estimation.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

ASYMPTOTIC MAXIMAL DEVIATION OF M-SMOOTHERS*

Wolfgang Härdle

Universität Heidelberg
Sonderforschungsbereich 123
Jm Neuenheimer Feld 293
D-6900 Heidelberg 1

University of North Carolina
Department of Statistics
321 Phillips Hall 039A
Chapel Hill, NC 27514

KEY WORDS: Robust nonparametric regression, Kernel estimators,
Uniform consistency, Robust smoothing

1970 AMS Subject Classification: 62G05, 62J05, 62F35

DTIC
ELECTE
S **D**
AUG 29 1983
B

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH
NOTICE OF TECHNICAL TO DTIC
This technical report has been reviewed and is
approved for publication in the AFOSR 190-12.
Distribution is unlimited.
MATTHEW J. KERPER
Chief, Technical Information Division

* Research partially supported by the "Deutsche Forschungsgemeinschaft"
SFB123, "Stochastische Mathematische Modelle".

Research also partially supported by the Air Force of Scientific
Research Contract ~~AFOSR~~-F49620 82 C 0009.

ABSTRACT

Let $(X_1, Y_1), \dots, (X_n, Y_n)$ be iid rv's with pdf $f(x, y)$ and let $m(x) = E(Y|X = x) = \int y f(x, y) dy / f_X(x)$ be the regression function of Y on X . The function $m(x)$ is estimated by $m_n(x)$ a solution of $(nh)^{-1} \sum_{i=1}^n K((x - X_i)/h) \Psi(Y_i - \cdot) = 0$ for some odd and bounded Ψ -function making $m_n(x)$ a robust estimate of $m(x)$. Probabilities of maximal deviation of $|m_n(x) - m(x)|$ are computed in a similar way as in Bickel and Rosenblatt (1973) for density estimation and in Johnston (1982) for nonparametric regression function estimation.



Accession		✓
DTIC		
DTIC		
Unannounced		
Justification		
By		
Distribution/		
Availability Codes		
Dist	Avail and/or Special	
A		

1. BACKGROUND AND INTRODUCTION

Nadaraya (1964) and Watson (1964) independently proposed the following kernel estimator

$$(1.1) \quad m_n^*(x) = (nh_n)^{-1} \sum_{i=1}^n K((x-X_i)/h_n) Y_i / [(nh_n)^{-1} \sum_{j=1}^n K((x-X_j)/h_n)]$$

of the regression function $m(x) = \int y f(x,y) dy / f_X(x)$ where $f_X(x)$ denotes the marginal density of X , $K(\cdot)$ is a kernel and $\{h_n\}$ is a sequence of positive constants ("bandwidth"). Basically this estimator averages the Y 's around $X = x$ motivated from the integral formula for $m(x)$ above. The numerator is a weighted local average of the Y 's while the denominator is a density estimate of $f_X(x)$.

It is clear that occasional outliers generated by heavy tailed conditional densities $f(y|x)$ introduce smooth peaks and troughs in the estimated curve $m_n^*(x)$. Such outliers occur quite often in practice. (Ruppert et al., 1982 Figure 7 or Bussian et al., 1982). To avoid this misleading property of $m_n^*(x)$ due to spiky Y -observations we introduce a robust estimate, the M -smoother, $m_n(x)$ as the solution of

$$(1.2) \quad (nh_n)^{-1} \sum_{i=1}^n K((x-X_i)/h_n) \Psi(Y_i - \cdot) = 0,$$

where Ψ denotes a bounded, odd and continuous function. Note that if $\Psi(u) = u$, then m_n is the Nadaraya-Watson estimator m_n^* . Bias and variance rates for $m_n(x)$ with K as the uniform window were obtained by Stuetzle and Mittal (1979), robustness properties, consistency and asymptotic normality of $m_n(x)$ were considered by Härdle (1982). For the case of nonrandom design, i.e. X_i attains fixed values, we may refer to Härdle and Gasser (1982). In this paper we show that

$$(1.3) \quad P\{(2\delta \log n)^{\frac{1}{2}} \left[\sup_{0 \leq t \leq 1} |(m_n(t) - m(t)) \cdot r(t)| / \lambda(K)^{\frac{1}{2}} - d_n \right] < x\} \\ \cdot \exp(-2 \exp(-x)) \quad ,$$

where δ , $r(t)$, $\lambda(K)$, d_n are suitable scaling parameters.

The result (1.3) improves upon that of Johnston (1982) in a number of ways. First, Johnston obtains results like (1.3), but for estimates different from the Nadaraya-Watson estimator (1.1); our result (1.3) of course applies to the Nadaraya-Watson estimator as a special case. Secondly, (1.3) holds for a much broader class of estimators. Finally, we obtain (1.3) under assumptions weaker than those needed by Johnston.

2. ASSUMPTIONS AND RESULTS

We write h for the bandwidth h_n from here on unless there is no need to do so. We make use of the following assumptions.

(A1) the kernel $K(\cdot)$ is positive has compact support $[-A, A]$ and is continuously differentiable.

(A2) $(nh)^{-\frac{1}{2}}(\log n)^{3/2} \rightarrow 0$ $(n \log n)^{\frac{1}{2}} h^{5/2} \rightarrow 0$
 $(nh^3)^{-1}(\log n)^2 \leq M$, M a constant .

(A3) $h^{-3}(\log n) \int_{|y| > a_n} f_y(y) dy = o(1)$, $f_y(y)$ the marginal density of Y , $\{a_n\}_{n=1}^{\infty}$ a sequence of constants tending to infinity as $n \rightarrow \infty$.

(A4) $\inf_{0 \leq t \leq 1} |q(t)| \geq q_0 > 0$, where $q(t) = E(\Psi'(Y-m(t)) | X=t) - f_X(t)$

(A5) the regression function $m(x)$ is twice continuously differentiable, the conditional densities $f(y|x)$ are symmetric for all x , Ψ is piecewise twice continuously differentiable.

We need some more definitions before we discuss the assumptions.

Define

$$\begin{aligned}\sigma^2(t) &= E(\Psi^2(Y-m(t))|X=t) \\ H_n(t) &= (nh)^{-1} \sum_{i=1}^n K((x-X_i)/h) \Psi(Y_i-m(t)) \\ D_n(t) &= (nh)^{-1} \sum_{i=1}^n K((x-X_i)/h) \Psi'(Y_i-m(t)).\end{aligned}$$

We further assume that $\sigma^2(t)$ and $f_X(t)$ are differentiable.

Assumption (A1) on the compact support of the kernel could possibly be relaxed introducing a cutoff technique as Csörgö and Hall (1982) for density estimators. Assumption (A2) has purely technical reasons: to keep the bias down and to ensure the vanishing of the nonlinear remainder terms. Assumption (A3) appears in a somewhat modified form also in Johnston's paper (1982). When we want to apply the following theorem to the Nadaraya-Watson estimator $m_n^*(x)$ we have actually to restate (A2) as $h^{-3} (\log n) \int_{|y|>a_n} y^2 f_y(y) dy$ (which is assumption A1 in Johnston (1982)). Assumption (A5) stating the symmetry of the conditional densities is common in robustness considerations (Huber, 1981). It guarantees that the only solution of $\int \Psi(y-\cdot) f(y|x) dy = 0$ is $m(x) = E(Y|X=x)$. If we had skew distributions then we would no longer estimate the conditional mean but rather a conditional quantile such as the median.

Theorem

$$\begin{aligned}\text{Let } h &= n^{-\delta}, \quad 1/5 < \delta < 1/3 \text{ and } \lambda(K) = \int_{-A}^A K^2(u) du \text{ and} \\ d_n &= (2\delta \log n)^{\frac{1}{2}} + (2\delta \log n)^{-\frac{1}{2}} \{ \log(c_1(K)/\pi^{\frac{1}{2}}) + \frac{1}{2} [\log \delta + \log \log n] \}, \\ &\quad \text{if } c_1(K) = K^2(A) + K^2(-A)/[2\lambda(K)] > 0 \\ d_n &= (2\delta \log n)^{\frac{1}{2}} + (2\delta \log n)^{-\frac{1}{2}} \{ \log(c_2(K)/2\pi) \} \\ &\quad \text{otherwise with } c_2(K) = \int_{-A}^A [K'(u)]^2 du / [2\lambda(K)].\end{aligned}$$

Then (1.3) holds with

$$r(t) = (nh)^{\frac{1}{2}} q(t) [\sigma^2(t) f_X(t)]^{-\frac{1}{2}}.$$

This theorem can be used to construct uniform confidence intervals for the regression function as stated in the following corollary.

Corollary: Assuming the theorem above holds, an approximate $(1-\alpha) \times 100\%$ confidence band over $[0,1]$ is

$$m_n(t) \pm (nh)^{-\frac{1}{2}} [\sigma^2(t) f_X(t) \lambda(K)]^{\frac{1}{2}} q^{-1}(t) [d_n + c(\alpha)(2s \log n)^{-\frac{1}{2}}] \cdot [\lambda(K)]^{\frac{1}{2}}$$

where $c(\alpha) = \log 2 - \log |\log(1-\alpha)|$.

The proof is essentially based on a linearization argument due to Taylor series expansion. The leading linear term will then be approximated in a similar way as in Johnston (1982), Bickel and Rosenblatt (1973). The main idea behind the proof is a strong approximation of the empirical process of $\{(X_i, Y_i)\}_{i=1}^n$ by a sequence of Brownian bridges (with two dimensional time) as provided Tusnady (1977).

It follows by Taylor expansions applied to the defining equation (1.2) that

$$(2.1) \quad m_n(t) - m(t) = (H_n(t) - EH_n(t))/q(t) + R_n(t)$$

where $[H_n(t) - EH_n(t)]/q(t)$ is the leading linear term and

$$(2.2) \quad R_n(t) = H_n(t)[q(t) - D_n(t)]/[D_n(t) \cdot q(t)] + EH_n(t)/q(t) \\ + \frac{1}{2}(m_n(t) - m(t))^2 \cdot [D_n(t)]^{-1} \cdot (nh)^{-1} \sum_{i=1}^n K((x - X_i)/h) \Psi''(Y_i - m(t) + r_n^{(i)}(t)), \\ |r_n^{(i)}(t)| < |m_n(t) - m(t)|.$$

is the remainder term. In the third section it is shown (Lemma 3.1) that

$$\|R_n\| = \sup_{0 \leq t \leq 1} |R_n(t)| = o_p((nh \log n)^{-\frac{1}{2}}).$$

Furthermore the rescaled linear part

$$Y_n(t) = (nh)^{\frac{1}{2}} [\sigma^2(t) f_X(t)]^{-\frac{1}{2}} (H_n(t) - EH_n(t))$$

is approximated by a sequence of Gaussian processes, leading finally to the following process

$$Y_{5,n}(t) = h^{-1/2} \int K((t-x)/h) dW(x),$$

as in Bickel and Rosenblatt (1973).

We also need the Rosenblatt transformation (Rosenblatt, 1952).

$$T(x,y) = (F_{X|Y}(x|y), F_Y(y))$$

which transforms (X_i, Y_i) into $T(X_i, Y_i) = (X'_i, Y'_i)$ mutually independent uniform rv's. With the aid of this transformation Theorem 1 of Tusnàdy (1977) may be applied to obtain the following lemma.

Lemma 2.1: On a suitable probability space there exists a sequence of Brownian bridges B_n such that

$$\sup_{x,y} |Z_n(x,y) - B_n(T(x,y))| = O(n^{-1/2}(\log n)^2) \text{ a.s.},$$

where $Z_n(x,y) = n^{1/2}[F_n(x,y) - F(x,y)]$ denotes the empirical process of $\{(X_i, Y_i)\}_{i=1}^n$.

Before we define the different approximating processes let us first rewrite $Y_n(t)$ as a stochastic integral with respect to the empirical process $Z_n(x,y)$.

$$Y_n(t) = h^{-1/2} g'(t)^{-1/2} \iint K((t-x)/h) \Psi(y-m(t)) dZ_n(x,y), \quad g'(t) = \sigma^2(t) f_X(t).$$

The approximating processes are now

$$Y_{0,n}(t) = (hg(t))^{-1/2} \iint_{\Gamma_n} K((t-x)/h) \Psi(y-m(t)) dZ_n(x,y),$$

$$\text{where } \Gamma_n = \{|y| \leq a_n\}, \quad g(t) = E(\Psi^2(y-m(t)) \cdot I(|y| \leq a_n) | X=t) \cdot f_X(t)$$

$$Y_{1,n}(t) = (hg(t))^{-1/2} \iint_{\Gamma_n} K((t-x)/h) \Psi(y-m(t)) dB_n(T(x,y)),$$

$\{B_n\}$ being the sequence of Brownian bridges from Lemma 2.1.

$$Y_{2,n}(t) = (hg(t))^{-1/2} \iint_{\Gamma_n} K((t-x)/h) \Psi(y-m(t)) dW_n(T(x,y))$$

$\{W_n\}$ being the sequence of Wiener processes satisfying

$$B_n(x', y') = W_n(x', y') - x'y'W_n(1,1)$$

$$Y_{3,n}(t) = (hg(t))^{-\frac{1}{2}} \int_n K((t-x)/h) \psi(y-m(x)) dW_n(T(x,y))$$

$$Y_{4,n}(t) = (hg(t))^{-\frac{1}{2}} \int g(x)^{\frac{1}{2}} K((t-x)/h) dW(x)$$

$$Y_{5,n}(t) = h^{-\frac{1}{2}} \int K((t-x)/h) dW(x),$$

(W(·)) being the Wiener process on $(-\infty, \infty)$.

Lemmata 3.2 to 3.7 ensure that all these processes have the same limit distributions. The results then follows from the following lemma

Lemma 2.2 (Bickel and Rosenblatt (1973)). Let d_n , $\lambda(K)$, δ as in the theorem.

Let

$$Y_{5,n}(t) = h^{-\frac{1}{2}} \int K((t-x)/h) dW(x).$$

Then

$$P((2 \log n)^{\frac{1}{2}} \sup_{0 \leq t \leq 1} |Y_{5,n}(t)| / [\lambda(K)]^{\frac{1}{2}} - d_n < x) \rightarrow e^{-2e^{-x}}.$$

3. PROOFS

We show first that $\|R_n\| = \sup_{0 \leq t \leq 1} |R_n(t)|$ vanishes asymptotically with the

desired rate $(nh \log n)^{-\frac{1}{2}}$.

Lemma 3.1: For the remainder term $R_n(t)$ defined in (2.2) we have

$$(3.1) \quad \|R_n\| = o_p((nh \log n)^{-\frac{1}{2}}).$$

Proof: First we have by the positivity of the kernel K and $|\psi''| \leq C_1$

$$\begin{aligned} \|R_n\| &\leq \left[\inf_{0 \leq t \leq 1} |D_n(t)| \cdot q(t) \right]^{-1} \{ \|H_n\| \cdot \|q - D_n\| + \|D_n\| \cdot \|EH_n\| \} \\ &\quad + C_1 \cdot \|m_n - m\|^2 \cdot \left[\inf_{0 \leq t \leq 1} |D_n(t)| \right]^{-1} \cdot \|f_n\|, \end{aligned}$$

where $f_n = (nh)^{-1} \sum_{i=1}^n K((x-X_i)/h)$.

The desired result (3.1) will then follow if we prove the following:

$$(3.2) \quad \|H_n\| = o_p(n^{-\frac{1}{4}} h^{-\frac{1}{4}} \cdot (\log n)^{-\frac{1}{2}}) \quad (3.2)$$

$$(3.3) \quad \|Q - D_{n,n}\| = o_p(n^{-1/4}h^{-1/2}(\log n)^{-1/2})$$

$$(3.4) \quad \|EH_n\| = O(h^2)$$

$$(3.5) \quad \|m_n - m\|^2 = o_p((nh)^{-1}(\log n)^{-1/2}).$$

Define $U_n(t) = n^{1/4}h^{1/2}(\log n)^{1/2}[H_n(t) - EH_n(t)]$.

We first show that $U_n(t) \xrightarrow{P} 0$ for all t . This follows from Markov's inequality since

$$U_n(t) = \sum_{i=1}^n U_{i,n}(t),$$

where $U_{i,n}(t) = n^{-3/4}h^{-3/4}(\log n)^{1/2}[K((t-X_i)/h)\Psi(Y_i-m(t)) - EK((t-X)/h) \cdot \Psi(Y-m(t))]$,

are iid rv's and thus

$$P(|U_n(t)| > \epsilon) \leq \epsilon^{-2} n^{-1/2} h^{-1/2} (\log n) \cdot h^{-1} EK^2((t-X)/h) \Psi^2(Y-m(t)).$$

The RHS of this inequality tends to zero since

$$\begin{aligned} h^{-1} EK^2((t-X)/h) \Psi^2(Y-m(t)) &= h^{-1} \int K^2((t-u)/h) E(\Psi^2(Y-m(t)) | X=u) f_X(u) du \\ &\sim \sigma^2(t) \cdot f_X(t) \cdot \int K^2(u) du \end{aligned}$$

by continuity of $\sigma^2(t)$ and $f_X(t)$.

Next we show the tightness of $U_n(t)$ using the following moment condition (Billingsley, 1968, Th. 15.6)

$$E\{|U_n(t) - U_n(t_1)| \cdot |U_n(t_2) - U_n(t)|\} \leq C_2 \cdot (t_2 - t_1)^2$$

where C_2 is a constant.

By the Schwarz inequality,

$$\begin{aligned} E\{|U_n(t) - U_n(t_1)| \cdot |U_n(t_2) - U_n(t)|\} \\ \leq \{E[U_n(t) - U_n(t_1)]^2 \cdot E[U_n(t_2) - U_n(t)]^2\}^{1/2}. \end{aligned}$$

It suffices to consider only the term $E\{U_n(t) - U_n(t_1)\}^2$.

Using the Lipschitz continuity of K, Ψ, m and assumption (A2) we have

$$\{E[U_n(t) - U_n(t_1)]^2\}^{1/2}$$

$$= (\log n)(nh)^{-3/2} \cdot E[A+B]^2 \cdot h^{-1/2}$$

$$= C_A(nh)^{-1/2}(\log n)^{1/2} \cdot t-t_1 + C_B(n^{-1/2}h^{-3/4}(\log n)^{1/2} \cdot t-t_1) + C_3 \cdot t-t_1$$

$$\text{where } A = \sum_{i=1}^n K((t-X_i)/h) [\psi(Y_i-m(t)) - \psi(Y_i-m(t_1))]$$

$$B = \sum_{i=1}^n \psi(Y_i-m(t_1)) [K((t_1-X_i)/h) - K((t-X_i)/h)],$$

and C_A, C_B are Lipschitz bounds for ψ, m, K .

Since (3.4) follows from the well-known bias calculation

$$EH_n(t) = h^{-1} \int K((t-u)/h) E(\psi(y-m(t)) | X=u) f_X(u) du = O(h^2),$$

where $O(h^2)$ is independent of t (Parzen, 1962) we have from assumption (A2)

$$\text{that } \|EH_n\| = o((nh)^{-1/2}(\log n)^{-1/2}).$$

Statement (3.2) thus follows using tightness of $U_n(t)$ and the inequality

$$\|H_n\| \leq \|H_n - EH_n\| + \|EH_n\|.$$

Statement (3.3) follows in the same way as (3.2) using assumption (A2)

and the continuity properties of K, ψ, m .

Finally from Härdle and Luckhaus (1982), where uniform continuity of $m_n(t) - m(t)$ is shown, we have

$$\|m_n - m\| = O_p((nh)^{-1/2}(\log n)^{1/2}),$$

which implies (3.5).

Now the assertion of the lemma follows since by tightness of $D_n(t)$,

$$\inf_{0 \leq t \leq 1} \|D_n(t)\| \xrightarrow{p} q_0 \text{ and thus}$$

$$\|R_n\| = o_p((nh)^{-1/2}(\log n)^{-1/2})(1 + \|f_n\|).$$

Finally by Theorem 3.1 of Bickel and Rosenblatt (1973) $\|f_n\| = O_p(1)$,

thus the desired result $\|R_n\| = o_p((nh)^{-1/2}(\log n)^{-1/2})$ follows. In the nonrobust case, i.e. $\psi(u) = u$, the remainder term R_n reads

$$(3.6) \quad R_n = [m_n^* - m][f_X - f_n]f_X^{-1} + E(\hat{m}_n - mf_n)/f_X,$$

$$\text{where } \hat{m}_n(x) = (nh)^{-1} \sum_{i=1}^n K((x-X_i)/h) Y_i.$$

Johnston (1982) proved that $(\hat{m}_n - E \hat{m}_n)/f$ has the desired asymptotic distribution as stated in our Theorem.

So if we apply the recent result of Mack and Silverman (1982) or Härdle and Luckhaus (1982) to $\|m_n^* - m\|$ and the well known result from Bickel and Rosenblatt (1973) to $\|f_X - f_n\|$ we may conclude that the first term on the RHS of (3.6) is $o_p((nh)^{-1/2}(\log n)^{-1/2})$. The second term in (3.6) is

$$[h^{-1} \int K((t-u)/h) \cdot m(u) f(u) du - m(t) h^{-1} \int K((t-u)/h) f(u) du] / f_X(t)$$

which is by the same calculations as mentioned above (Parzen, 1962) of the order $O(h^2)$. This shows that our result generalizes Johnston's paper. Our theorem says also that the confidence bounds are smaller. Johnston had $s^2(t) = E(Y^2|X=t)$ as a factor for the asymptotic confidence bound, we have $\sigma^2(t) = \text{var}(Y|X=t)$ which is in general smaller than $s^2(t)$. We now begin with the subsequent approximations of the processes $Y_{0,n}$ to $Y_{5,n}$.

Lemma 3.2:

$$\|Y_{0,n} - Y_{1,n}\| = O((nh)^{-1/2}(\log n)^2) \quad \text{a.s.}$$

Proof: Let t be fixed and put $L(y) = \Psi(y - m(t))$ still depending on t .

Use integration by parts and obtain:

$$\begin{aligned} & \iint_n L(y) K((t-x)/h) dZ_n(x, y) = \\ &= \int_{u=-A}^A \int_{y=-a_n}^{a_n} L(y) K(u) dZ_n(t-h \cdot u, y) = \\ &= \int_{-A}^A \int_{-a_n}^{a_n} Z_n(t-h \cdot u, y) d[L(y) K(u)] + L(a_n) \int_{-A}^A Z_n(t-h \cdot u, a_n) dK(u) \\ &\quad - L(-a_n) \int_{-A}^A Z_n(t-h \cdot u, -a_n) dK(u) + K(A) \left[\int_{-a_n}^{a_n} Z_n(t-h \cdot u, y) dL(y) \right. \\ &\quad \left. + L(a_n) Z_n(t-h \cdot A, a_n) - L(-a_n) Z_n(t-h \cdot A, -a_n) \right] \\ &\quad - K(-A) \left[\int_{-a_n}^{a_n} Z_n(t+h \cdot A, y) dL(y) + L(a_n) Z_n(t+h \cdot A, a_n) \right. \\ &\quad \left. - L(-a_n) Z_n(t+h \cdot A, -a_n) \right]. \end{aligned}$$

If we apply the same operations to $Y_{1,n}$ with $B_n(T(x,y))$ instead of $Z_n(x,y)$ and use Lemma 2.1 we finally obtain

$$\sup_{0 \leq t \leq 1} h^{\frac{1}{2}} q(t)^{\frac{1}{2}} |Y_{0,n}^{(t)} - Y_{1,n}(t)| = O((nh)^{-\frac{1}{2}} (\log n)^2) \quad \text{a.s.}$$

using the differentiability and boundedness of ψ .

Lemma 3.3:

$$\|Y_{1,n} - Y_{2,n}\| = O_p(h^{\frac{1}{2}})$$

Proof: Note that the Jacobi of $T(x,y)$ is $f(x,y)$ hence

$$|Y_{1,n}(t) - Y_{2,n}(t)| = |(q(t)h)^{-\frac{1}{2}} \int_n \int \psi(y-m(t)) K((t-x)/h) f(x,y) dx dy| \cdot |W_n(1,1)|$$

It follows that

$$h^{-\frac{1}{2}} \|Y_{1,n} - Y_{2,n}\| \cdot |W_n(1,1)| \leq \|g^{-\frac{1}{2}}\| \cdot \sup_{0 \leq t \leq 1} h^{-\frac{1}{2}} \int_n \int \psi(y-m(t)) K((t-x)/h) |f(x,y)| dx dy$$

Since $\|g^{-\frac{1}{2}}\|$ is bounded by assumption and ψ is bounded we have

$$h^{-\frac{1}{2}} \|Y_{1,n} - Y_{2,n}\| \leq |W_n(1,1)| \cdot C_4 \cdot h^{-1} \int (K((t-x)/h)) dx = O_p(1).$$

Lemma 3.4:

$$\|Y_{2,n} - Y_{3,n}\| = O_p(h^{\frac{1}{2}})$$

Proof: The difference $|Y_{2,n}(t) - Y_{3,n}(t)|$ may be written as

$$|(q(t)h)^{-\frac{1}{2}} \int_n \int [\psi(y-m(t)) - \psi(y-m(x))] K((t-x)/h) dW_n(T(x,y))|$$

If we use the fact that ψ, m are uniformly continuous this is smaller than

$$h^{-\frac{1}{2}} |q(t)|^{-\frac{1}{2}} \cdot O_p(h)$$

and the lemma thus follows.

Lemma 3.5:

$$\|Y_{4,n} - Y_{5,n}\| = O_p(h^{\frac{1}{2}})$$

Proof:

$$|Y_{4,n}(t) - Y_{5,n}(t)| = h^{-\frac{1}{2}} \left| \int \left\{ \left[\frac{q(x)}{q(t)} \right]^{\frac{1}{2}} - 1 \right\} K((t-x)/h) dW(x) \right| \leq$$

$$\begin{aligned}
 & h^{-\frac{1}{2}} \left| \int_{-A}^A W(t-hu) \left\{ \left[\frac{g(t-hu)}{g(t)} \right]^{\frac{1}{2}} - 1 \right\} K(u) du \right| \\
 & + h^{-\frac{1}{2}} \left| K(A) W(t-hA) \left\{ \left[\frac{g(t-hA)}{g(t)} \right]^{\frac{1}{2}} - 1 \right\} \right| \\
 & + h^{-\frac{1}{2}} \left| K(-A) W(t+hA) \left\{ \left[\frac{g(t+hA)}{g(t)} \right]^{\frac{1}{2}} - 1 \right\} \right| \\
 & = S_{1,n}(t) + S_{2,n}(t) + S_{3,n}(t), \text{ say.}
 \end{aligned}$$

The second term can be estimated by

$$h^{-\frac{1}{2}} \|S_{2,n}\| \leq K(A) \cdot \sup_{0 \leq t \leq 1} |W(t-hA)| \cdot \sup_{0 \leq t \leq 1} h^{-1} \left| \left[\frac{g(t-hA)}{g(t)} \right]^{\frac{1}{2}} - 1 \right|$$

by the mean value theorem it follows that

$$h^{-\frac{1}{2}} \|S_{2,n}\| = O_p(1).$$

The first term $S_{1,n}$ is estimated as follows.

$$\begin{aligned}
 h^{-1} S_{1,n}(t) &= h^{-1} \int_{-A}^A W(t-hu) \{K'(u) \left[\frac{g(t-hu)}{g(t)} \right]^{\frac{1}{2}} - 1\} du \\
 &= h^{-\frac{1}{2}} \int_{-A}^A W(t-hu) K(u) \left[\frac{g(t-hu)}{g(t)} \right]^{-\frac{1}{2}} \left[\frac{g'(t-hu)}{g(t)} \right] du \\
 &= |T_{1,n}(t) - T_{2,n}(t)|, \text{ say.}
 \end{aligned}$$

$$\|T_{2,n}\| \leq C_5 \cdot \int_{-A}^A |W(t-hu)| du = O_p(1) \text{ by assumption on } g(t) = o^2(t) \cdot f_X(t).$$

To estimate $T_{1,n}$ we again use the mean value theorem to conclude that

$$\sup_{0 \leq t \leq 1} h^{-1} \left| \left[\frac{g(t-hu)}{g(t)} \right]^{\frac{1}{2}} - 1 \right| < C_6 \cdot |u|$$

hence

$$\|T_{1,n}\| \leq C_6 \cdot \sup_{0 \leq t \leq 1} \int_{-A}^A |W(t-hu) K'(u) u| du = O_p(1).$$

Since $S_{3,n}(t)$ is estimated as $S_{2,n}(t)$ we finally obtain the desired result.

The next lemma shows that the truncation introduced through $\{a_n\}$ does not affect the limiting distribution.

Lemma 3.6:

$$\|Y_n - Y_{0,n}\| = O_p((\log n)^{-\frac{1}{2}}).$$

Proof: We shall only show that $g'(t)^{-\frac{1}{2}} h^{-\frac{1}{2}} \iint_{\mathbb{R} - \Gamma_n} \psi(y-m(t)) K((t-x)/h) dZ_n(x,y)$ fulfills the lemma.

The replacement of $q'(t)$ by $g(t)$ may be proved as in Johnston (1982). The quantity above is less than $h^{-\frac{1}{2}} \|g^{-\frac{1}{2}}\| \cdot \left\| \iint_{\{|y|>a_n\}} \psi(y-m(\cdot)) K((\cdot-x)/h) dZ(x,y) \right\|$.

It remains to show that the last factor tends to zero at a rate $O_p((\log n)^{-\frac{1}{2}})$. We show first that

$$V_n(t) = (\log n)^{\frac{1}{2}} h^{-\frac{1}{2}} \iint_{\{|y|>a_n\}} \psi(y-m(t)) K((t-x)/h) dZ_n(x,y)$$

$$\xrightarrow{P} 0 \quad \text{for all } t$$

and then we show tightness of $V_n(t)$, the result then follows.

$$\begin{aligned} V_n(t) &= (\log n)^{\frac{1}{2}} (nh)^{-\frac{1}{2}} \sum_{i=1}^n \{ \psi(Y_i - m(t)) I_{\{|y|>a_n\}}(Y_i) K((t-X_i)/h) \\ &\quad - E\psi(Y_i - m(t)) \cdot I_{\{|y|>a_n\}}(Y_i) K((t-X_i)/h) \} \\ &= \sum_{i=1}^n X_{n,i}(t) \end{aligned}$$

where $\{X_{n,i}(t)\}_{i=1}^n$ are iid for each n with $EX_{n,i}(t) = 0$ for all $t \in [0,1]$.

We have then

$$\begin{aligned} EX_{n,i}^2(t) &\leq (\log n)(nh)^{-1} E\psi^2(Y_i - m(t)) I_{\{|y|>a_n\}}(Y_i) K^2((t-X_i)/h) \\ &\leq \sup_{-A \leq u \leq A} K^2(u) \cdot (\log n)(nh)^{-1} E\psi^2(Y_i - m(t)) I_{\{|y|>a_n\}}(Y_i) \end{aligned}$$

hence

$$\begin{aligned} \text{var}\{V_n(t)\} &= E\left(\sum_{i=1}^n X_{n,i}(t)\right)^2 = n \cdot EX_{n,i}^2(t) \\ &\leq \sup_{-A \leq u \leq A} K^2(u) h^{-1} (\log n) \int_{\{|y|>a_n\}} f_y(y) dy \cdot M_\psi \end{aligned}$$

where M_ψ denotes an upper bound for ψ^2 .

This term tends to zero by assumption (A3). Thus by Markov's inequality we conclude that

$$V_n(t) \xrightarrow{P} 0 \quad \text{for all } t \in [0,1].$$

To prove tightness of $\{V_n(t)\}$ we refer again to the following moment condition as stated in Lemma 3.1.

$$E\{|V_n(t) - V_n(t_1)| \cdot |V_n(t_2) - V_n(t)|\} \leq C' \cdot (t_2 - t_1)^2$$

C' denoting a constant, $t \in [t_1, t_2]$.

We again estimate the left hand side by Schwarz's inequality and estimate each factor separately.

$$\begin{aligned} E[V_n(t) - V_n(t_1)]^2 &= (\log n)(nh)^{-1} E\left\{ \sum_{i=1}^n \psi_n(t, t_1, X_i, Y_i) \cdot I_{\{|Y_i| > a_n\}}(Y_i) - \right. \\ &\quad \left. - E(\psi_n(t, t_1, X_i, Y_i) \cdot I_{\{|Y_i| > a_n\}}(Y_i)) \right\}^2, \end{aligned}$$

where $\psi_n(t, t_1, X_i, Y_i) = \psi(Y_i - m(t))K((t - X_i)/h) - \psi(Y_i - m(t_1))K((t_1 - X_i)/h)$

Since ψ, m, K are Lipschitz continuous it follows

$$\begin{aligned} &\{E[V_n(t) - V_n(t_1)]^2\}^{1/2} \\ &\leq C_7 \cdot (\log n)^{1/2} h^{-3/2} |t - t_1| \cdot \left\{ \int_{\{|Y| > a_n\}} f_Y(y) dy \right\}^{1/2} \end{aligned}$$

If we apply the same estimations to $V_n(t_2) - V_n(t_1)$ we finally have

$$\begin{aligned} E\{|V_n(t) - V_n(t_1)| \cdot |V_n(t_2) - V_n(t_1)|\} &\leq C_7^2 (\log n) h^{-3} |t - t_1| |t_2 - t_1| \\ &\quad \cdot \int_{\{|Y| > a_n\}} f_Y(y) dy \\ &\leq C' \cdot |t_2 - t_1|^2 \text{ since } t \in [t_1, t_2]. \end{aligned}$$

by assumption (A3).

Lemma 3.7: Let $\lambda(K) = \int K^2(u) du$ and let $\{d_n\}$ as in the theorem. Then

$$(2\delta \log n)^{1/2} [\|Y_{3,n}\| / [\lambda(K)]^{1/2} - d_n]$$

has the same asymptotic distribution as

$$(2\delta \log n)^{1/2} [\|Y_{4,n}\| / [\lambda(K)]^{1/2} - d_n]$$

Proof: $Y_{3,n}(t)$ is a Gaussian process with

$$EY_{3,n}(t) = 0$$

and covariance function

$$\begin{aligned} r_3(t_1, t_2) &= EY_{3,n}(t_1)Y_{3,n}(t_2) \\ &= [g(t_1)g(t_2)]^{-\frac{1}{2}} h^{-1} \int_{\Gamma_n} \psi^2(y-m(x)) K((t_1-x)/h) K((t_2-x)/h) f(x,y) dx dy. \\ &= h^{-1} [g(t_1)g(t_2)]^{-\frac{1}{2}} \int_{\Gamma_n} \psi^2(y-m(x)) f(y|x) dy K((t_1-x)/h) K((t_2-x)/h) f_X(x) dx \\ &= h^{-1} [g(t_1)g(t_2)]^{-\frac{1}{2}} \int g(x) K((t_1-x)/h) K((t_2-x)/h) dx \\ &= r_4(t_1, t_2) \text{ the covariance function of the Gaussian process } Y_{4,n}(t), \text{ which} \\ &\text{proves the lemma.} \end{aligned}$$

ACKNOWLEDGEMENT

The author wishes to thank R.J. Carroll for helpful remarks and suggestions on a first draft of this paper.

REFERENCES

- [1] Bickel, P. and Rosenblatt, M. (1973). On some global measures of the deviation of density function estimators. Ann. Stat. 1, 1071-1095.
- [2] Billingsley, P. (1968). Convergence of Probability Measures. John Wiley & Sons.
- [3] Bussian, B.M. and Härdle, W. (1982). Robust smoothing applied to white noise and single outlier contaminated Raman Spectra. Tentatively accepted by Applied Spectroscopy.
- [4] Csörgö, S. and Hall, P. (1982). Upper and Lower Classes for Triangular Arrays. Z. F. Wahrscheinlichkeitstheorie und verw. Gebiete, 61, 207-222.
- [5] Härdle, W. (1982). Robust regression function estimation. (To appear in J. Mult. Analysis).
- [6] Härdle, W. and Gasser, T. (1982). Robust nonparametric function fitting. Submitted to JRSS(B).
- [7] Härdle, W. and Luckhaus, S. (1982). Uniform consistency of a class of regression function estimates. Preprint SFB 123 University of Heidelberg #143.
- [8] Huber, P. (1981). Robust statistics. J Wiley & Sons, New York.
- [9] Johnston, G. (1982). Probabilities of maximal deviation of nonparametric regression function estimation. J. Mult. Anal. 12, 402-414.
- [10] Mack, Y.P. and Silverman, B.W. (1982). Weak and strong uniform consistency of kernel regression estimates. Z. F. Wahrscheinlichkeitstheorie und verw. Gebiete, 61, 405-415.
- [11] Nadaraya, E.A. (1964). On estimating regression. Theor. Prob. Appl. 9, 141-142.
- [12] Parzen, M. (1962). On estimation of a probability density function and mode. Ann. Math. Stat. 32, 1065-1076.
- [13] Rosenblatt, M. (1952). Remarks on a multivariate transformation. Ann. Math. Stat. 23, 470-472.
- [14] Rosenblatt, M. (1971). Curve estimates. Ann. Math. Stat. 42, 1815-1842.
- [15] Ruppert, D., Reish, R.L., Deriso, R.B., Carroll, R.J. (1982). Monte Carlo optimization by stochastic approximation (with application to harvesting of Atlantic menhaden). Inst. of Statistics Memo Series #1500, University of North Carolina at Chapel Hill.
- [16] Stuetzle, W. and Mittal, Y. (1979). Some comments on the asymptotic behaviour of robust smoothers in "Smoothing Techniques for Curve Estimation" ed. T. Gasser and M. Rosenblatt. Springer Lecture Notes 757, Heidelberg.

- [16] Tusnady, G. (1977). A remark on the approximation of the sample distribution function in the multidimensional case. Period. Math. Hung., 8, 53-55.
- [17] Watson, G.S. (1964). Smooth regression analysis. Sankhya, Series A, Vol. 26, 359-372.