

AD-A132 181


ANALYTIC MODELS OF MAGNETIC FIELD EVOLUTION IN
LASER-PRODUCED PLASMA EXPANSIONS(U) NAVAL RESEARCH LAB
WASHINGTON DC M J KESKINEN 31 AUG 83 NRL-MR-5163


1/1

UNCLASSIFIED

F/G 4/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

ADA 132181

83 09 07 080

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NRL Memorandum Report 5163	2. GOVT ACCESSION NO. AD A132 181	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ANALYTIC MODELS OF MAGNETIC FIELD EVOLUTION IN LASER-PRODUCED PLASMA EXPANSIONS		5. TYPE OF REPORT & PERIOD COVERED Interim report on a continuing NRL problem.
7. AUTHOR(s) M.J. Keskinen		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Research Laboratory Washington, DC 20375		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS Defense Nuclear Agency Washington, DC 20305		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 62715H; 47-0889-0-3
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE August 31, 1983
		13. NUMBER OF PAGES 29
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES This research was sponsored by the Defense Nuclear Agency under Subtask S99QMXBC, work unit 00067 and work unit title "Plasma Structure Evolution."		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Analytic models Lasers Magnetic fields Plasma expansion		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Analytic models of the magnetic field evolution in laser-produced plasma expansions have been studied and applied to the NRL laser HANE (high altitude nuclear explosion) simulation experiment. Both one- and two-dimensional models have been investigated for laser plasma expansions with and without initial background magnetic fields. For the case with no initial background magnetic field, thermal source mechanisms have (Continues)		

20. ABSTRACT (Continued)

been used to discuss the two-dimensional evolution and morphology of spontaneous self-generated magnetic fields. With an initial background magnetic field, one-dimensional models with discontinuous debris density profiles give unrealistically large magnetic field compressions, while predictions from two-dimensional models with smooth profiles are in reasonable agreement with initial NRL experimental observations.

CONTENTS

I.	INTRODUCTION.....	1
II.	BASIC EQUATIONS.....	2
III.	LASER PLASMA EXPANSION WITHOUT A BACKGROUND MAGNETIC FIELD.....	4
IV.	LASER PLASMA EXPANSION WITH A BACKGROUND MAGNETIC FIELD.....	7
	A. One-dimensional models.....	7
	B. Two-dimensional models.....	9
V.	SUMMARY.....	11
	Acknowledgments.....	12
	References	17

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	



ANALYTIC MODELS OF MAGNETIC FIELD EVOLUTION IN LASER-PRODUCED PLASMA EXPANSIONS

I. INTRODUCTION

A high altitude nuclear explosion (HANE) can significantly perturb the natural ionosphere and magnetosphere by generating large-scale (tens to hundreds of kilometers), long-lived (several hours) ionization irregularities (striations). These irregularities can degrade radar and communication systems, e.g., through scintillation effects. As such, it is crucial to have a detailed understanding of the evolution of a HANE in order to achieve a reliable communications system capability in a nuclear environment. Several experiments have been conducted¹, both in space (barium cloud releases) and in the laboratory (laser-pellet explosions), in order to simulate various aspects of a HANE. Recently, renewed interest in laboratory laser-pellet HANE simulation experiments has been generated owing to detailed scaling studies.¹⁻³

An important aspect of the early time (few seconds) evolution of a HANE is the manner by which the exploding debris plasma couples into the background air plasma. The nature of this early time coupling could seed or influence the evolution and structure of late time ionization irregularities. This coupling can either be collisional (particle-particle interactions) or collisionless (wave-particle interactions) depending on ambient densities and temperatures. Collisionless coupling proceeds via plasma microturbulence which in turn is driven by various plasma instabilities⁴. Recently, a set of "turn-on" conditions for collisionless coupling, in the context of the NRL laser plasma experiment, has been derived⁵. A key ingredient in determining whether or not the aforementioned plasma instabilities will be excited is the structure and magnitude of the local magnetic field in the debris-air coupling region. If the magnetic field compressions are too small several of the coupling instabilities will be inoperative. A detailed description of the early time magnetic field evolution and morphology is also important for discussing related topics such as electron heat transport, magnetic field driven interchange instabilities, and ion leakage mechanisms. Analytic models of the evolution of the magnetic fields can be used to validate HANE numerical simulations and also be compared with experimental results.

In this report we study analytic models of the magnetic field evolution and compression in laser-produced plasma expansions. We treat both one- and two-dimensional models both with and without an initial background magnetic field. Many aspects of this problem have already been investigated⁶⁻¹². However these studies have not been discussed or applied to recent NRL laser pellet experimental observations. The outline of this report is as follows. In Section II we present and discuss the general model equation for the evolution of the magnetic field in laser-produced plasma expansions. In Section III we study laser plasma expansions without an initial ambient background magnetic field and the expected magnitudes of spontaneous self-generated magnetic fields. The calculated self-generated magnetic fields are found to be in agreement with preliminary NRL experimental values. In Section IV we investigate, using both one- and two-dimensional models, laser plasma expansions into an ambient background magnetic field. The predicted field compressions are not inconsistent with those obtained using NRL experimental observations. We find that one-dimensional models using sharp debris density profiles give unrealistically large magnetic field compressions. We show that two dimensional models with diffuse profiles can explain several experimental observations. Finally in Section V we summarize our results.

II. BASIC EQUATIONS

For high β ($B^2/8\pi \ll Nk_B T$) plasmas, the equation for the evolution of the magnetic field $\underline{B}$ can be written¹²⁻¹⁴

$$\frac{\partial \underline{B}}{\partial t} = \nabla \times \underline{V} \times \underline{B} - \frac{c^2}{4\pi} \nabla \times [\underline{\eta} \cdot (\nabla \times \underline{B})] + \underline{S} \quad (1)$$

where

$$\underline{S} = - \frac{ck_B}{eN_e} \nabla N_e \times \nabla T_e \quad (2)$$

with $\underline{\eta}$ the resistivity tensor, $\underline{V}$ the fluid velocity, c the speed of light, k_B is Boltzmann's constant, and N_e and T_e are the electron temperature and density, respectively. Radiation pressure effects have been neglected in Eq. (2) since they can be shown to be small for the laser intensities used in the current NRL experiment. Equation (1) is simply Faraday's law with the electric field determined from the force balance equation for the electrons.

The generation of a magnetic field requires that the last term in Eq. (1), the source term $\underline{S}$, be nonzero. This requires that ∇N_e and ∇T_e be nonparallel.

By defining the dimensionless quantities $\tilde{t} = V_0 t / L$, $\tilde{V} = V / V_0$, $\tilde{x} = x / L$, Eq. (1) can be written

$$\frac{\partial \underline{B}}{\partial \tilde{t}} = \tilde{V} \times \tilde{V} \times \underline{B} - R_m^{-1} \tilde{V}^2 \underline{B} \quad (3)$$

where, for the moment, we have neglected $\underline{S}$ and defined V_0 and L as a representative fluid velocity and a magnetic gradient scale length, respectively. Here R_m defines a magnetic Reynolds number $R_m = 4\pi\sigma LV_0/c^2$ with $\eta \equiv I \sigma^{-1}$ and I the unit tensor. For $R_m > 1$, magnetic field convection ($\tilde{V} \times \tilde{V} \times \underline{B}$ term) dominates over diffusion ($\tilde{V}^2 \underline{B}$ term) whereas for $R_m < 1$ the opposite is true. An effective electron collision frequency ν^{eff} can be defined by $\sigma = N_e e^2 / m_e \nu^{eff}$ where N_e and m_e are the electron density and mass, respectively. For $\nu^{eff} = \nu_{ei} = 3 \times 10^{-6} Z \ln \Lambda N_e / T_e^{3/2} \text{ sec}^{-1}$, the classical Coulomb collision frequency, with Z the charge number and $\ln \Lambda$ the Coulomb logarithm, we find $R_m \approx 10^4 L(\text{cm})$ where we have taken $T_e \approx 100 \text{ eV}$ and $V_0 \approx 4 \times 10^7 \text{ cm/sec}$ as representative NRL laser experimental parameters (B. Ripin, private communication). For $L \sim 1 \text{ cm}$, $R_m \approx 10^4$ dissipative effects are negligible. However, if the effective collision frequency is increased by other processes, e.g., plasma microturbulence, R_m will decrease and resistive effects will become more important. For example, for plasma turbulence resulting from the magnetized ion-ion instability⁴ an effective collision frequency can be written

$$\nu_{ij}^{eff} = 0.15 \omega_{Hi} \frac{\rho_i}{\rho} (\alpha_{ji}^{2/3} + 2^{-1/3} 3^{1/2} (\alpha_{ji}^{1/3} - \alpha_{ji}^{2/3})) \quad (4)$$

where i and j refer to ion species i and j , $\omega_{Hi} = \omega_{pi} (1 + \omega_{pe}^2 / \Omega_e^2)^{-1/2}$, ω_{pi} , ω_{pe} and electron plasma frequencies, ρ_i is the mass density of species i , ρ the total mass density and $\alpha_{ij} = N_j Z_j^2 m_i / N_i Z_i^2 m_j$. For aluminum (i) streaming through nitrogen (j), i.e., $N_i \approx 10^{16} \text{ cm}^{-3}$, $N_j \approx 10^{14} \text{ cm}^{-3}$, $Z_i = 10$, $Z_j = 3$ we find $R_m \approx 10 L(\text{cm}) \sim 10$ for $L \approx 1 \text{ cm}$ making resistive effects more important.

III. LASER PLASMA EXPANSION WITHOUT A BACKGROUND MAGNETIC FIELD

As shown in the previous section, spontaneous magnetic fields can be generated by nonparallel density and temperature gradients. These self-generated magnetic fields can be quite large¹⁴ near the focal spot region. Since these spontaneous fields will be carried along with the expanding plasma they could influence greatly the electron and ion dynamics in the coupling shell. In addition, large-self fields also imply asymmetric departures from completely spherical expansion and reduced coupling of the laser energy into the target.

According to Eq. (2), the magnitude and direction of the self generated magnetic field is determined by the geometrical configuration of the laser-plasma. A laser beam, which is cylindrically symmetric, will produce a plasma which expands in the direction of the normal to the target and is symmetric about its expansion direction. From symmetry considerations there can be no azimuthal density or temperature gradient. During the laser heating of the target, it is reasonable to assume that the largest contribution to the source term in Eq. (2) comes from a temperature gradient in the radial direction and a density gradient in the direction of the target normal due to expansion of the target plasma. Due to the finite radial extent of the laser beam a radial temperature gradient will exist near the edge of the focal spot. This combination of ∇T_e and ∇N_e will generate a magnetic field in the azimuthal direction in the form of a torus¹². The self-generated field will be convected radially by the expanding plasma. This scenario has been confirmed by many previous laser-pellet experiments¹⁴.

In order to approximate the self-generated magnetic fields in the NRL laser HANE experiment we assume a purely radial temperature gradient $\partial T_e / \partial r$ and a density gradient $\partial N_e / \partial z$ with r denoting distance perpendicular to the normal to the target plane and z representing distance in the axial direction perpendicular to the target plane. As a result Eq. (2) gives

$$S = (\partial B / \partial t)_{\text{self}} \approx (ck_B / eN_e) (\partial N_e / \partial z) (\partial T_e / \partial r) \quad (5)$$

The radial temperature gradient is of the order $\partial T_e / \partial r \approx T_e / r_0$ where r_0 is the radial extension of the laser heated plasma near the focal spot. We take the

density gradient in the z-direction to be given by the debris ion expansion velocity V_o , i.e., $\partial \ln N_i / \partial z \approx \partial \ln N_e / \partial z \approx [V_o \tau_L + (\Delta V_o / V_o) R]^{-1}$ where τ_L is the duration of the laser pulse, ΔV_o is the thermal debris velocity spread, and R is the approximate position of the debris density maximum. Taking $(\partial B / \partial t)_{\text{self}} \approx B_{\text{self}} / \tau_L$ we have from (5)

$$B_{\text{self}} \approx 9 \times 10^7 \frac{T_e}{V_o r_o} [1 + (\Delta V_o / V_o) (R / V_o \tau_L)]^{-1} \text{ G} \quad (6)$$

where T_e , V_o , r_o , τ_L are expressed in eV, cm/sec, cm, and sec, respectively. For $T_e \approx 100$ eV, $V_o \approx 4 \times 10^7$ cm/sec, $r_o \approx 1$ cm, $\Delta V_o / V_o = 0.2$, $R = 0.5$ cm, $\tau_L = 4 \times 10^{-9}$ sec (B. Ripin private communication), Eq. (6) gives $B_{\text{self}} \approx 100$ G which is in agreement with experimentally measured values (S. Kacendar, private communication).

To find the approximate time dependence of the self-generated magnetic field, we consider¹² the fluid variables N_e , T_e , V in Eq. (1) as consisting of zeroth order contribution plus a first order part, i.e., $N_e = N_{e0} + \Delta N_e$, $T_e = T_{e0} + \Delta T_e$, and $V = V_o + \Delta V$. The zeroth order parts N_{e0} , T_{e0} , and V_o describe a spherically symmetric expansion with the perturbations ΔN_e , ΔT_e , and ΔV representing a small departure from spherical symmetry giving rise to a source term $\underline{S}$ vanishes in the spherically symmetric case. In other words, we can linearize Eq. (1) and solve for $\partial \Delta B / \partial t$ using the zeroth order motion for $\underline{V}$ and $\underline{n}$ in the first two terms on the right hand side of Eq. (1). The perturbations ΔN_e and ΔT_e are retained in the source term $\underline{S}$.

The radius of the expanding laser plasma, r_s , is found from¹² the following

$$r_s(t) = R_o + \int_0^t V_s(t') dt' \quad (7)$$

with R_o the initial radius. The velocity of the expanding plasma, in cylindrical coordinates, follows from the continuity equation

$$V_r = \frac{r}{r_s} V_s(t) \quad (8a)$$

$$V_z = \frac{z}{r_s} V_s(t) \quad (8b)$$

The exploding plasma is assumed, in zeroth order, to have a homogeneous density $N(t)$ in the course of its assumed adiabatic expansion so that

$$\frac{T}{T_0} = \left(\frac{N}{N_0}\right)^{\gamma-1} = \left(\frac{R_0}{r_s}\right)^{3(\gamma-1)} \quad (9)$$

Furthermore, we assume that the expansion decreases the plasma temperature so that the conductivity scales as

$$\sigma = \sigma_0 \left(\frac{T}{T_0}\right)^{3/2} = \sigma_0 \left(\frac{R_0}{r_s}\right)^{9(\gamma-1)/2} \quad (10)$$

where σ_0 is the initial value and a Coulomb collisional conductivity¹⁵ has been assumed. As a result, Eq. (1)-(2) for the azimuthal component ΔB_θ of the self-generated field, assuming cylindrical symmetry ($\frac{\partial}{\partial \theta} = 0$) can be written

$$\begin{aligned} \frac{\partial B_\theta}{\partial t} = & -\frac{\partial}{\partial z} (V_z B_\theta) - \frac{\partial}{\partial r} (V_r B_\theta) + D \left(\frac{1}{r} \frac{\partial}{\partial r} r \frac{\partial B_\theta}{\partial r} - \frac{B_\theta}{r^2} + \frac{\partial^2 B_\theta}{\partial z^2} \right) \\ & + S(r, z, t) \end{aligned} \quad (11)$$

$$\text{with } S = \frac{ck_B}{eN_e} \left(\frac{\partial N_e}{\partial r} \frac{\partial T_e}{\partial z} - \frac{\partial N_e}{\partial z} \frac{\partial T_e}{\partial r} \right)$$

where, for clarity, we have dropped the Δ 's from B_θ , V , N_e , and T_e and $D(t) = D_0(r_s/R_0)^{9(\gamma-1)/2}$ with $D_0 = c^2/4\pi\sigma_0$. We wish to solve Eq. (11) as an initial value problem with $B_\theta(t=0) = 0$ and V_r , V_z given by Eq. (8).

For the expanding plasma cloud, we consider¹² the following debris density and temperature profiles which are smooth functions of position.

$$N_e = N_{e0} - \Delta N_e \exp(-r^2/r_s^2(t)) \quad (12a)$$

$$T_e = T_{e0} - (\Delta T_e/r_s) \int_0^z dz' \exp(-z'^2/r_s^2) \quad (12b)$$

From (12a) and (12b) we find

$$S(r, z, t) = (-2 c k_B/e) (\Delta N_e/N_e) (r/r_s) (\Delta T_e/r_s^2)$$

$$x \exp [-(r^2 + z^2)/r_s^2] \quad (13)$$

Tidman (1975) has solved Eq. (11) using the profiles from (12) and finds, in scaled time units,

$$B_\theta(\tilde{t}) = -8\pi (2e^2)^{-1/2} B_0 \left(\frac{\Delta N_e}{N_e}\right) \left(\frac{\Delta T_e}{T_e}\right) \left(\frac{V_d}{V_0}\right) \tilde{t} (1 + \tilde{t})^{-3} \quad (14)$$

where $B_0 = k_B T_0 \sigma_0 / ec$, $V_d = D_0 / R_0 = c^2 / 4\pi \sigma_0 R_0$, $r_s = R_0 + V_0 t$ (constant expansion velocity V_0), $\tilde{t} = V_0 t / R_0$ and $e = 2.71828$. Using parameters typical of the NRL laser experiment ($T_e \approx 100$ eV, $N_e \approx 10^{16}$ cm $^{-3}$, $V_0 \approx 4 \times 10^7$ cm/sec), $B_\theta(\tilde{t})$ is given by Fig. 1 for several values of $\Delta N_e / N_e$ and $\Delta T_e / T_e$. Both the magnitudes (several hundred and time dependence of B_θ are consistent with preliminary measurements from the current NRL laser HANE experiment.

IV. LASER PLASMA EXPANSION WITH A BACKGROUND MAGNETIC FIELD

For realistic simulations of a HANE, a background magnetic field must be introduced into laser-pellet experiments. With a background magnetic field, magnetic field compression can now take place in addition to spontaneous magnetic field creation as discussed in Section III. Magnetic field compression may be the first stage in the process leading to "pickup" of the background air ions. It is important to compute the spatial and temporal history of the magnetic field compression in order to determine where and when the peak compression is achieved.

A. One-dimensional models

We consider a one-dimensional model of laser-induced plasma expansions into a background magnetic field. The model consists of a debris plasma streaming with velocity $V_d \hat{x}$ through a stationary background (air) plasma. Choosing to work in the debris frame of reference, the ion component of the expanding debris plasma is stationary with density n_D while the background plasma is assumed uniform with density n_B and having flow velocity $-V_d \hat{x}$. The basic configuration is shown in Fig. 2. In the interaction region $-x_0 < x < 0$, continuity and quasi neutrality are imposed where $x_0 = V_D \tau_L$ with τ_L the duration of the laser pulse. These conditions determine the density $n_e = n_B + n_D$ and flow velocity $-V_e \hat{x} = -(n_B/n_B + n_D) V_d \hat{x}$ of the debris electrons. Initially, a constant background magnetic field $B_0 \hat{z}$ is

taken to be normal to the flow but excluded from the interaction region as shown in Fig. 2. We wish to determine the field compression $B_z(x,t) / B_0$.

For the case where collisions are absent, the evolution of $B_z(x,t)$ was determined by Longmire⁶ using magnetic flux conservation arguments and is illustrated in Fig. 3. Here B_z jumps discontinuously from B_0 at $X = \epsilon$ to $(V_d / V_e) B_0 > B_0$ at $X = -\epsilon$. The leading edge of the compression in the interaction region is convected with velocity V_e .

Including collisional effects Eq. (1) gives for $B_z(x,t)$

$$\frac{\partial B_z}{\partial t} = \frac{-\partial}{\partial x} (V_x B_z) + \frac{\partial}{\partial x} V_x(x) L_c(x) \frac{\partial}{\partial x} B_z \quad (15)$$

where $L_c(x) = v(x) c^2 / \omega_{pe}^2(x) V_x(x)$ with the effective collision frequency $v(x)$ being defined from $\sigma^{-1}(x) = v(x) m_e / N(x) c^2$. It should be noted that in the derivation of Eq. (1) and (2) the effects of electron inertia have been neglected. Eq. (15) has been solved for several special cases.^{7,8,10}

As a simplified model we consider the case where the effective collision frequency is non zero only in the interaction region, $-x_0 < x < 0$. This effective collision frequency is defined as the sum of the classical Coulomb collision frequency plus an anomalous part due to plasma turbulence. Let

$$L_c(x) = \begin{cases} 0 & x > 0 \\ L_{co} & -x_0 < x < 0 \end{cases}$$

where L_{co} is constant. As a result the equation governing $B_z(x,t)$ for $-x_0 < x < 0$ can be written

$$\frac{\partial B_z}{\partial t} - V_1 \frac{\partial B_z}{\partial x} - L_{co} V_1 \frac{\partial^2 B_z}{\partial x^2} = 0 \quad (16)$$

In the following dimensionless variables $\tilde{x} = x / L_{co}$, $\tilde{t} = |V_1| t / L_{co}$, and $\tilde{B} = B_z / B_0$ Eq. (16) can be written ($-x_0 < x < 0$)

$$\frac{\partial \tilde{B}}{\partial \tilde{t}} - \frac{\partial \tilde{B}}{\partial \tilde{x}} - \frac{\partial^2 \tilde{B}}{\partial \tilde{x}^2} = 0 \quad (17)$$

where, for clarity, we have dropped the tildes. The initial and boundary conditions appropriate to Eq. (17) are: $\tilde{B}(x > 0, t = 0) = 1$, $\tilde{B}(-x_0 <$

$x < 0, t = 0) = 0, \quad \tilde{B}(x = \infty, t) = 1, \quad B(x=0-\epsilon, t) + \partial \tilde{B} / \partial x (x=0-\epsilon, t) = 1.$

Eq. (17) together with these initial and boundary conditions can be solved to yield

$$B_z (-x_0 < x < 0, t) = \frac{v_d}{v_e} \left(1 + (2\pi)^{-1} \exp(-x/2) \zeta(x, t) \right) \quad (18)$$

where

$$\zeta(x, t) = (1 - 2 \partial / \partial x) \alpha(x, t)$$

and

$$\begin{aligned} \alpha(x, t) &= \int_{-\infty}^{\infty} du u (u^2 + \frac{1}{4})^{-2} \exp\left(- (u^2 + \frac{1}{4})t\right) \sin u x \\ &= \pi (x - t) \exp(-|x|/2) + (\pi^{1/2}/2) \\ &\quad \times \int_0^t dm \left(t m^{-3/2} - m^{-1/2} \right) \exp\left(\frac{-x^2}{4m} - \frac{m}{4}\right) \\ &= \pi (x-t) \exp(-|x|/2) + \pi \operatorname{erfc}\left(\frac{1}{2} x t^{-1/2}\right) \sum_{n=0}^{\infty} \frac{x^{2n} (4n+t)}{2^{3n} (2n-1)!! n!} \\ &\quad + \pi^{1/2} \exp(-x^2/4t) \sum_{n=1}^{\infty} \frac{x^{2n} (4n+t)}{2^{3n} (2n-1)!! n!} \sum_{k=0}^{n-1} (-2)^{k+1} (2k-1)!! \left(\frac{t}{2x}\right)^{k+1/2} \end{aligned}$$

Fig. 4 shows the evolution of $B_z(x, t)$ as given by Eq. (18). For times $\tilde{t} > 1$ collisions will smooth the discontinuous jump in B_z given the Longmire analysis⁶.

B. Two-dimensional models

In the previous section, a one-dimensional model of an expanding laser plasma was assumed to have sharp discontinuous debris density profile in order to simplify analytical calculations. Unrealistically large magnetic field compressions were obtained⁶ by assuming an infinitely steep density profile. In this section we show the effects of using a diffuse, smooth debris density profile by making a two-dimensional analysis¹¹ of a laser plasma expanding into a background magnetic field.

We consider a laser plasma expanding into a paraxial magnetic field $\underline{B} = B_0 \hat{z} = B_0 \cos \theta \hat{e}_r - B_0 \sin \theta \hat{e}_\theta$ which is imbedded in a stationary background plasma and assume a spherical coordinate system (r, θ, ϕ) with the polar angle θ measured from the z -axis. Assuming spherical symmetry ($\partial/\partial \theta = \partial/\partial \phi = 0$) expressions for B_r and B_θ from Eq. (1) can be written

$$\frac{\partial B_r}{\partial t} = (r \tan \theta)^{-1} V_r B_\theta + D \left(r^{-2} \frac{\partial}{\partial r} r^2 \frac{\partial B_r}{\partial r} - \frac{2B_r}{r^2} - \frac{2B_\theta \cot \theta}{r^2} \right) \quad (19)$$

$$\frac{\partial B_\theta}{\partial t} = -r^{-1} \frac{\partial}{\partial r} r V_r B_\theta + D \left(r^{-2} \frac{\partial}{\partial r} r^2 \frac{\partial B_\theta}{\partial r} - \frac{B_\theta}{r^2 \sin^2 \theta} \right) \quad (20)$$

where we have neglected the self-generated magnetic fields in comparison with B_0 and noted the B_ϕ is negligible for very early times. On time scales short compared to the resistive time scale, the diffusive terms in (19) and (20) can be neglected and we are left with

$$\frac{\partial B_r}{\partial t} = (r \tan \theta)^{-1} V_r B_\theta \quad (21)$$

$$\frac{\partial B_\theta}{\partial t} = -r^{-1} \frac{\partial}{\partial r} r V_r B_\theta \quad (22)$$

We assume uniform expansion $V_r = V_0 r / r_0$, quasi-neutrality for the electrons, and a gaussian debris density profile $N_D(r, t=0) = N_0 \exp(-r^2/r_0^2)$ where $r_0 = V_0 \tau_L$ with τ_L the duration of the laser pulse. Under these conditions, Eqs (21) and (22) can be solved¹¹ analytically for B_θ , B_r for short times δt such that the field is only slightly altered:

$$\delta B_\theta \approx \frac{2 V_0 B_\theta \delta t (N_0/N_B) \exp(-r^2/r_0^2)}{r_0 (1 + (N_0/N_B) \exp(-r^2/r_0^2))^2} \left[\frac{r^2}{r_0^2} - 1 - \frac{N_0}{N_B} \right] \exp(-r^2/r_0^2) \quad (23)$$

$$\delta B_r = (r \tan \theta)^{-1} \left(\frac{N_0 V_0}{N_B + N_0} \frac{r}{r_0} \right) B_\theta \delta t \quad (24)$$

where N_b is the background plasma density. The peak magnetic compression is achieved at $r=r_m$ where $\partial/\partial r (\delta B_\theta) = 0$ giving

$$(\delta B_0 / B_0)_{\max} \approx (V_0 \delta t / 2 r_0) \left((r_m^2 / r_0^2) - 2 \right) \quad (25)$$

with $r_m = (f(N_0, N_B))^{1/2} r_0$, $f(x,y) = \ln(x/y) - \ln g(x,y)$, $g(x,y) = (\ln(x/y) + 2) / (\ln(x/y) - 2)$. For example in $N_B \approx 10^{14} \text{ cm}^{-3}$ and $N_0 \approx 10^{16} \text{ cm}^{-3}$, we find $r_m \approx 1.9 r_0$ indicating that peak magnetic field compression occurs ahead of the expanding laser-produced plasma cloud. Furthermore, the laser plasma cloud density at r_m is given by $N_0(r = r_m) \approx 0.28 N_B$. A characteristic shell thickness, at early times, can be found using $\delta B_\theta \approx \frac{1}{2} (\delta B_\theta)_{\max}$, giving $t \approx 0.42 r_0$. These analytically obtained features that (1) the maximum field compression occurs in front of the advancing shell and (2) the maximum field intensity in the compression region is proportional to the radial displacement of the shell are consistent with preliminary NRL laser experimental results (S. Kacenjar, private communication).

Finally, by consideration of the conservation of magnetic flux in two-dimensions, the field compression scaling as given by Eq. (23) and (24) can be shown to be reasonable. Consider an annulus of compressed magnetic field with inner and outer radii of r_1 and r_2 , respectively. Magnetic flux conservation implies $B_0 r_2^2 = B_c(r_2^2 - r_1^2)$ where B_c/B_0 is the magnitude of the field compression. This gives $B_c/B_0 \approx r_2 / 2 \delta$, $\delta = r_2 - r_1$. For the NRL laser experiment $R_2 \approx 3 \text{ cm}$, $\delta \approx 1.3 \text{ cm}$, giving $B_c/B_0 \approx 1-2$ in agreement with observations (S. Kacenjar, private communication). In addition, the direct proportionality between B_c/B_0 and r_2 is not inconsistent with recent NRL experimental findings (S. Kacenjar, private communication).

V. SUMMARY

We have attempted to provide, in this preliminary report, simple analytic models for the magnetic field evolution in laser-induced plasma expansion in the context of the NRL laser plasma HANE simulation experiment. Both one- and two-dimensional models have been used for laser plasma expansions with and without initial background magnetic fields. For the case with no initial background magnetic field, we find reasonable agreement between analytic two-dimensional models and preliminary results from the NRL laser plasma experiment. It is shown that self-generated spontaneous magnetic fields are small in comparison to proposed ambient background fields. For the case of

laser plasma expansion into a background magnetic field, one dimensional models with sharp, discontinuous profiles give unrealistically large magnetic field compressions while two-dimensional models incorporating smooth debris profiles give results are not inconsistent with current NRL experimental observations. Predictions of magnetic field compressions are supported by simple conservation arguments.

Future work will include comparison of these analytic models with future experimental data and the use of these analytic models for validation of large numerical codes.

Acknowledgments

We thank B. Ripin and S. Kacenjar for useful discussions and J. Huba for a critical reading of the manuscript. This work was supported by the Defense Nuclear Agency.




Fig. 1 Self-generated magnetic field B_θ produced by an asymmetric laser-plasma expansion vs. scaled time $t = V_0 \tilde{t}/R_0$. Curve a corresponds to $\Delta N/N = \Delta T/T = 0.1$ while curve b corresponds to $\Delta N/N = \Delta T/T = 0.3$.





Fig. 2 Plot of (a) density of debris ions N_D , background ions N_B , and electrons $N_e = N_D + N_B$ as a function of x and (b) initial ($t = 0$) profile of background magnetic field. Here $x_0 \approx V_D \tau_L$ with V_D the average debris velocity and τ_L the duration of the laser pulse.




Fig. 3 Plot of $B_z(x,t)$ neglected resistance and inertia (Longmire solution).




Fig. 4 Plot of $B_z(x, t)$ with constant resistivity for $-x_0 < x < 0$. The quantities x , t , B_z are scaled by L_{co} , L_{co}/V_e , and $(V_D/V_e)B_0$, respectively.

References

1. T.F. Vesecky, J.W. Chamberlain, J.M. Cornwall, D.A. Hammer, and F.W. Perkins, SRI International, Technical Report JSR-81-31, December, 1981.
2. C. Longmire, M. Alme, R. Kilb, and L. Wright, MRC Report AMRC-R-338, 1981.
3. W. Tsai, L.L. DeRaad, Jr., and R. Lelevier, R & D Associates, 1982.
4. M. Lampe, W. Manheimer, and K. Papadopoulos, NRL Memo Report 3076, 1975.
5. R.A. Smith and J.D. Huba, NRL Memo Report (in press), 1983.
6. C.L. Longmire, The Rand Corporation Report RM-3386-PR, 1963.
7. M.L. Sloan, Bull. Am. Phys. Soc. 15, 1435, 1970.
8. R.W. Clark et al., Phys. Fluids, 16, 1097, 1973.
9. T.P. Wright, Phys. Fluids, 14, 1905, 1971.
10. M. Lampe and W. Hernandez, Bull. Am. Phys. Soc. 15, 1465, 1970.
11. D. Book and R. Clark, Phys. Fluids, 16, 720, 1973.
12. D.A. Tidman, Phys. Fluids, 18, 1454, 1975.
13. J.A. Stamper and D.A. Tidman, Phys. Fluids, 16, 2024, 1973.
14. J.A. Stamper, K. Papadopoulos, R.N. Sudan, S.O. Dean, E.A. McLean, and J.M. Dawson, Phys. Rev. Lett., 26, 1012, 1971.
15. S.I. Braginskii, Reviews of Plasma Physics, ed. M. Leontovich (Consultants Bureau, New York, 1965).

DISTRIBUTION LIST

DEPARTMENT OF DEFENSE

ASSISTANT SECRETARY OF DEFENSE
COMM, CMD, CONT 7 INTELL
WASHINGTON, D.C. 20301

DIRECTOR
COMMAND CONTROL TECHNICAL CENTER
PENTAGON RM BE 685
WASHINGTON, D.C. 20301
01CY ATTN C-650
01CY ATTN C-312 R. MASON

DIRECTOR
DEFENSE ADVANCED RSCH PROJ AGENCY
ARCHITECT BUILDING
1400 WILSON BLVD.
ARLINGTON, VA. 22209
01CY ATTN NUCLEAR MONITORING RESEARCH
01CY ATTN STRATEGIC TECH OFFICE

DEFENSE COMMUNICATION ENGINEER CENTER
1860 WIEHLE AVENUE
RESTON, VA. 22090
01CY ATTN CODE R410
01CY ATTN CODE R812

DEFENSE TECHNICAL INFORMATION CENTER
CAMERON STATION
ALEXANDRIA, VA. 22314
02CY

DIRECTOR
DEFENSE NUCLEAR AGENCY
WASHINGTON, D.C. 20305
01CY ATTN STVL
04CY ATTN TITL
01CY ATTN DDST
03CY ATTN RAAE

COMMANDER
FIELD COMMAND
DEFENSE NUCLEAR AGENCY
KIRTLAND, AFB, NM 87115
01CY ATTN FCPR

DIRECTOR
INTERSERVICE NUCLEAR WEAPONS SCHOOL
KIRTLAND AFB, NM 87115
01CY ATTN DOCUMENT CONTROL

JOINT CHIEFS OF STAFF
WASHINGTON, D.C. 20301
01CY ATTN J-3 WWMCCS EVALUATION OFFICE

DIRECTOR
JOINT STRAT TGT PLANNING STAFF
OFFUTT AFB
OMAHA, NB 68113
01CY ATTN JLTW-2
01CY ATTN JPST G. GOETZ

CHIEF
LIVERMORE DIVISION FLD COMMAND DNA
DEPARTMENT OF DEFENSE
LAWRENCE LIVERMORE LABORATORY
P.O. BOX 808
LIVERMORE, CA 94550
01CY ATTN FCPRL

COMMANDANT
NATO SCHOOL (SHAPE)
APO NEW YORK 09172
01CY ATTN U.S. DOCUMENTS OFFICER

UNDER SECY OF DEF FOR RSCH & ENGRG
DEPARTMENT OF DEFENSE
WASHINGTON, D.C. 20301
01CY ATTN STRATEGIC & SPACE SYSTEMS (OS)

WWMCCS SYSTEM ENGINEERING ORG
WASHINGTON, D.C. 20305
01CY ATTN R. CRAWFORD

COMMANDER/DIRECTOR
ATMOSPHERIC SCIENCES LABORATORY
U.S. ARMY ELECTRONICS COMMAND
WHITE SANDS MISSILE RANGE, NM 88002
01CY ATTN DELAS-EO F. NILES

DIRECTOR
BMD ADVANCED TECH CTR
HUNTSVILLE OFFICE
P.O. BOX 1500
HUNTSVILLE, AL 35807
01CY ATTN ATC-T MELVIN T. CAPPS
01CY ATTN ATC-O W. DAVIES
01CY ATTN ATC-R DON RUSS

PROGRAM MANAGER
BMD PROGRAM OFFICE
5001 EISENHOWER AVENUE
ALEXANDRIA, VA 22333
01CY ATTN DACS-BMT J. SHEA

CHIEF C-E- SERVICES DIVISION
U.S. ARMY COMMUNICATIONS CMD
PENTAGON RM 1B269
WASHINGTON, D.C. 20310
01CY ATTN C- E-SERVICES DIVISION

COMMANDER
FRADCOM TECHNICAL SUPPORT ACTIVITY
DEPARTMENT OF THE ARMY
FORT MONMOUTH, N.J. 07703
01CY ATTN DRSEL-NL-RD H. BENNET
01CY ATTN DRSEL-PL-ENV H. BOMKE
01CY ATTN J.E. QUIGLEY

COMMANDER
U.S. ARMY COMM-ELEC ENGRG INSTAL AGY
FT. HUACHUCA, AZ 85613
01CY ATTN CCC-EMEO GEORGE LANE

COMMANDER
U.S. ARMY FOREIGN SCIENCE & TECH CTR
220 7TH STREET, NE
CHARLOTTESVILLE, VA 22901
01CY ATTN DRXST-SD

COMMANDER
U.S. ARMY MATERIAL DEV & READINESS CMD
5001 EISENHOWER AVENUE
ALEXANDRIA, VA 22333
01CY ATTN DRCLDC J.A. BENDER

COMMANDER
U.S. ARMY NUCLEAR AND CHEMICAL AGENCY
7500 BACKLICK ROAD
BLDG 2073
SPRINGFIELD, VA 22150
01CY ATTN LIBRARY

DIRECTOR
U.S. ARMY BALLISTIC RESEARCH LABORATORY
ABERDEEN PROVING GROUND, MD 21005
01CY ATTN TECH LIBRARY EDWARD BAICY

COMMANDER
U.S. ARMY SATCOM AGENCY
FT. MONMOUTH, NJ 07703
01CY ATTN DOCUMENT CONTROL

COMMANDER
U.S. ARMY MISSILE INTELLIGENCE AGENCY
REDSTONE ARSENAL, AL 35809
01CY ATTN JIM GAMBLE

DIRECTOR
U.S. ARMY TRADOC SYSTEMS ANALYSIS ACTIVITY
WHITE SANDS MISSILE RANGE, NM 88002
01CY ATTN ATAA-SA
01CY ATTN TCC/F. PAYAN JR.
01CY ATTN ATTA-TAC LTC J. HESSE

COMMANDER
NAVAL ELECTRONIC SYSTEMS COMMAND
WASHINGTON, D.C. 20360
01CY ATTN NAVALEX 034 T. HUGHES
01CY ATTN PME 117
01CY ATTN PME 117-T
01CY ATTN CODE 5011

COMMANDING OFFICER
NAVAL INTELLIGENCE SUPPORT CTR
4301 SUITLAND ROAD, BLDG. 5
WASHINGTON, D.C. 20390
01CY ATTN MR. DUBBIN STIC 12
01CY ATTN NISC-50
01CY ATTN CODE 5404 J. GALET

COMMANDER
NAVAL OCCEAN SYSTEMS CENTER
SAN DIEGO, CA 92152
01CY ATTN J. FERGUSON

NAVAL RESEARCH LABORATORY

WASHINGTON, D.C. 20375

01CY ATTN CODE 4700 S. L. Ossakow
26 CYS IF UNCLASS. 1 CY IF CLASS)

01CY ATTN CODE 4701 I Vitkovitsky

01CY ATTN CODE 4780 J. Huba (100
CYS IF UNCLASS, 1 CY IF CLASS)

01CY ATTN CODE 7500

01CY ATTN CODE 7550

01CY ATTN CODE 7580

01CY ATTN CODE 7551

01CY ATTN CODE 7555

01CY ATTN CODE 4730 E. MCLEAN

01CY ATTN CODE 4108

01CY ATTN CODE 4730 B. RIPIN

20CY ATTN CODE 2628

COMMANDER

NAVAL SEA SYSTEMS COMMAND

WASHINGTON, D.C. 20362

01CY ATTN CAPT R. PITKIN

COMMANDER

NAVAL SPACE SURVEILLANCE SYSTEM

DAHLGREN, VA 22448

01CY ATTN CAPT J.H. BURTON

OFFICER-IN-CHARGE

NAVAL SURFACE WEAPONS CENTER

WHITE OAK, SILVER SPRING, MD 20910

01CY ATTN CODE F31

DIRECTOR

STRATEGIC SYSTEMS PROJECT OFFICE

DEPARTMENT OF THE NAVY

WASHINGTON, D.C. 20376

01CY ATTN NSP-2141

01CY ATTN NSSP-2722 FRED WIMBERLY

COMMANDER

NAVAL SURFACE WEAPONS CENTER

DAHLGREN LABORATORY

DAHLGREN, VA 22448

01CY ATTN CODE DF-14 R. BUTLER

OFFICER OF NAVAL RESEARCH

ARLINGTON, VA 22217

01CY ATTN CODE 465

01CY ATTN CODE 461

01CY ATTN CODE 402

01CY ATTN CODE 420

01CY ATTN CODE 421

COMMANDER

AEROSPACE DEFENSE COMMAND/DC

DEPARTMENT OF THE AIR FORCE

ENT AFB, CO 80912

01CY ATTN DC MR. LONG

COMMANDER

AEROSPACE DEFENSE COMMAND/XPD

DEPARTMENT OF THE AIR FORCE

ENT AFB, CO 80912

01CY ATTN XPDQQ

01CY ATTN XP

AIR FORCE GEOPHYSICS LABORATORY

HANSCOM AFB, MA 01731

01CY ATTN OPR HAROLD GARDNER

01CY ATTN LKB KENNETH S.W. CHAMPION

01CY ATTN OPR ALVA T. STAIR

01CY ATTN PHD JURGEN BUCHAU

01CY ATTN PHD JOHN P. MULLEN

AF WEAPONS LABORATORY

KIRTLAND AFT, NM 87117

01CY ATTN SUL

01CY ATTN CA ARTHUR H. GUENTHER

01CY ATTN NTYCE 1LT. G. KRAJEI

AFTAC

PATRICK AFB, FL 32925

01CY ATTN TF/MAJ WILEY

01CY ATTN TN

AIR FORCE AVIONICS LABORATORY

WRIGHT-PATTERSON AFB, OH 45433

01CY ATTN AAD WADE HUNT

01CY ATTN AAD ALLEN JOHNSON

DEPUTY CHIEF OF STAFF

RESEARCH, DEVELOPMENT, & ACQ

DEPARTMENT OF THE AIR FORCE

WASHINGTON, D.C. 20330

01CY ATTN AFRDQ

HEADQUARTERS

ELECTRONIC SYSTEMS DIVISION

DEPARTMENT OF THE AIR FORCE

HANSCOM AFB, MA 01731

01CY ATTN J. DEAS

HEADQUARTERS

ELECTRONIC SYSTEMS DIVISION/YSEA

DEPARTMENT OF THE AIR FORCE

HANSCOM AFB, MA 01732

01CY ATTN YSEA

HEADQUARTERS
ELECTRONIC SYSTEMS DIVISION/DC
DEPARTMENT OF THE AIR FORCE
HANSCOM AFB, MA 01731
O1CY ATTN DCKC MAJ J.C. CLARK

COMMANDER
FOREIGN TECHNOLOGY DIVISION, AFSC
WRIGHT-PATTERSON AFB, OH 45433
O1CY ATTN NICD LIBRARY
O1CY ATTN ETD P B. BALLARD

COMMANDER
ROME AIR DEVELOPMENT CENTER, AFSC
GRIFFISS AFB, NY 13441
O1CY ATTN DOC LIBRARY/TSLD
O1CY ATTN OCSE V. COYNE

SAMSO/SZ
POST OFFICE BOX 92960
WORLDWAY POSTAL CENTER
LOS ANGELES, CA 90009
(SPACE DEFENSE SYSTEMS)
O1CY ATTN SZJ

STRATEGIC AIR COMMAND/XPFS
OFFUTT AFB, NB 68113
O1CY ATTN ADWATE MAJ BRUCE BAUER
O1CY ATTN NRT
O1CY ATTN DOK CHIEF SCIENTIST

SAMSO/SK
P.O. BOX 92960
WORLDWAY POSTAL CENTER
LOS ANGELES, CA 90009
O1CY ATTN SKA (SPACE COMM SYSTEMS)
M. CLAVIN

SAMSO/MN
NORTON AFB, CA 92409
(MINUTEMAN)
O1CY ATTN MNNL

COMMANDER
ROME AIR DEVELOPMENT CENTER, AFSC
HANSCOM AFB, MA 01731
O1CY ATTN EEP A. LORENTZEN

DEPARTMENT OF ENERGY
LIBRARY ROOM G-042
WASHINGTON, D.C. 20545
O1CY ATTN DOC CON FOR A. LABOWITZ

DEPARTMENT OF ENERGY
ALBUQUERQUE OPERATIONS OFFICE
P.O. BOX 5400
ALBUQUERQUE, NM 87115
O1CY ATTN DOC CON FOR D. SHERWOOD

EG&G, INC.
LOS ALAMOS DIVISION
P.O. BOX 809
LOS ALAMOS, NM 85544
O1CY ATTN DOC CON FOR J. BREEDLOVE

UNIVERSITY OF CALIFORNIA
LAWRENCE LIVERMORE LABORATORY
P.O. BOX 808
LIVERMORE, CA 94550
O1CY ATTN DOC CON FOR TECH INFO DEPT
O1CY ATTN DOC CON FOR L-389 R. OTT
O1CY ATTN DOC CON FOR L-31 R. HAGER
O1CY ATTN DOC CON FOR L-46 F. SEWARD

LOS ALAMOS NATIONAL LABORATORY
P.O. BOX 1663
LOS ALAMOS, NM 87545
O1CY ATTN DOC CON FOR J. WOLCOTT
O1CY ATTN DOC CON FOR R.F. TASCHEK
O1CY ATTN DOC CON FOR E. JONES
O1CY ATTN DOC CON FOR J. MALIK
O1CY ATTN DOC CON FOR R. JEFFRIES
O1CY ATTN DOC CON FOR J. ZINN
O1CY ATTN DOC CON FOR P. KEATON
O1CY ATTN DOC CON FOR D. WESTERVELT
O1CY ATTN D. SAPPENFIELD

SANDIA LABORATORIES
P.O. BOX 5800
ALBUQUERQUE, NM 87115
O1CY ATTN DOC CON FOR W. BROWN
O1CY ATTN DOC CON FOR A. THORNBROUGH
O1CY ATTN DOC CON FOR T. WRIGHT
O1CY ATTN DOC CON FOR D. DAHLGREN
O1CY ATTN DOC CON FOR 3141
O1CY ATTN DOC CON FOR SPACE PROJECT DIV

SANDIA LABORATORIES
LIVERMORE LABORATORY
P.O. BOX 969
LIVERMORE, CA 94550
O1CY ATTN DOC CON FOR B. MURPHEY
O1CY ATTN DOC CON FOR T. COOK

OFFICE OF MILITARY APPLICATION
DEPARTMENT OF ENERGY
WASHINGTON, D.C. 20545
O1CY ATTN DOC CON DR. YO SONG

OTHER GOVERNMENT

DEPARTMENT OF COMMERCE
NATIONAL BUREAU OF STANDARDS
WASHINGTON, D.C. 20234
O1CY (ALL CORRES: ATTN SEC OFFICER FOR)

INSTITUTE FOR TELECOM SCIENCES
NATIONAL TELECOMMUNICATIONS & INFO ADMIN
BOULDER, CO 80303
O1CY ATTN A. JEAN (UNCLASS ONLY)
O1CY ATTN W. UTLAUT
O1CY ATTN D. CROMBIE
O1CY ATTN L. BERRY

NATIONAL OCEANIC & ATMOSPHERIC ADMIN
ENVIRONMENTAL RESEARCH LABORATORIES
DEPARTMENT OF COMMERCE
BOULDER, CO 80302
O1CY ATTN R. GRUBB
O1CY ATTN AERONOMY LAB G. REID

DEPARTMENT OF DEFENSE CONTRACTORS

AEROSPACE CORPORATION
P.O. BOX 92957
LOS ANGELES, CA 90009
O1CY ATTN I. GARFUNKEL
O1CY ATTN T. SALMI
O1CY ATTN V. JOSEPHSON
O1CY ATTN S. BOWER
O1CY ATTN D. OLSEN

ANALYTICAL SYSTEMS ENGINEERING CORP
5 OLD CONCORD ROAD
BURLINGTON, MA 01803
O1CY ATTN RADIO SCIENCES

AUSTIN RESEARCH ASSOC., INC.
1901 RUTLAND DRIVE
AUSTIN, TX 78758
O1CY ATTN L. SLOAN
O1CY ATTN R. THOMPSON

BERKELEY RESEARCH ASSOCIATES, INC.
P.O. BOX 983
BERKELEY, CA 94701
O1CY ATTN J. WORKMAN
O1CY ATTN C. PRETTIE
O1CY ATTN S. BRECHT

BOEING COMPANY, THE
P.O. BOX 3707
SEATTLE, WA 98124
O1CY ATTN G. KEISTER
O1CY ATTN D. MURRAY
O1CY ATTN G. HALL
O1CY ATTN J. KENNEY

CHARLES STARK DRAPER LABORATORY, INC.
555 TECHNOLOGY SQUARE
CAMBRIDGE, MA 02139
O1CY ATTN D.B. COX
O1CY ATTN J.P. GILMORE

COMSAT LABORATORIES
LINTHICUM ROAD
CLARKSBURG, MD 20734
O1CY ATTN G. HYDE

CORNELL UNIVERSITY
DEPARTMENT OF ELECTRICAL ENGINEERING
ITHACA, NY 14850
O1CY ATTN D.T. FARLEY, JR.

ELECTROSPACE SYSTEMS, INC.
BOX 1359
RICHARDSON, TX 75080
O1CY ATTN H. LOGSTON
O1CY ATTN SECURITY (PAUL PHILLIPS)

EOS TECHNOLOGIES, INC.
606 Wilshire Blvd.
Santa Monica, Calif 90401
O1CY ATTN C.B. GABBARD

ESL, INC.
495 JAVA DRIVE
SUNNYVALE, CA 94086
O1CY ATTN J. ROBERTS
O1CY ATTN JAMES MARSHALL

GENERAL ELECTRIC COMPANY
SPACE DIVISION
VALLEY FORGE SPACE CENTER
GODDARD BLVD KING OF PRUSSIA
P.O. BOX 8555
PHILADELPHIA, PA 19101
O1CY ATTN M.H. BORTNER SPACE SCI LAB

GENERAL ELECTRIC COMPANY
P.O. BOX 1122
SYRACUSE, NY 13201
O1CY ATTN F. REIBERT

GENERAL ELECTRIC TECH SERVICES CO., INC.
HMES
COURT STREET
SYRACUSE, NY 13201
O1CY ATTN G. MILLMAN

GEOPHYSICAL INSTITUTE
UNIVERSITY OF ALASKA
FAIRBANKS, AK 99701
(ALL CLASS ATTN: SECURITY OFFICER)
O1CY ATTN T.N. DAVIS (UNCLASS ONLY)
O1CY ATTN TECHNICAL LIBRARY
O1CY ATTN NEAL BROWN (UNCLASS ONLY)

GTE SYLVANIA, INC.
ELECTRONICS SYSTEMS GRP-EASTERN DIV
77 A STREET
NEEDHAM, MA 02194
O1CY ATTN DICK STEINHOF

HSS, INC.
2 ALFRED CIRCLE
BEDFORD, MA 01730
O1CY ATTN DONALD HANSEN

ILLINOIS, UNIVERSITY OF
107 COBLE HALL
150 DAVENPORT HOUSE
CHAMPAIGN, IL 61820
(ALL CORRES ATTN DAN MCCLELLAND)
O1CY ATTN K. YEH

INSTITUTE FOR DEFENSE ANALYSES
1801 NO. BEAUREGARD STREET
ALEXANDRIA, VA 22311
O1CY ATTN J.M. AEIN
O1CY ATTN ERNEST BAUER
O1CY ATTN HANS WOLFARD
O1CY ATTN JOEL BENGSTON

INTL TEL & TELEGRAPH CORPORATION
500 WASHINGTON AVENUE
NUTLEY, NJ 07110
O1CY ATTN TECHNICAL LIBRARY

JAYCOR
11011 TORREYANA ROAD
P.O. BOX 85154
SAN DIEGO, CA 92138
O1CY ATTN J.L. SPERLING

JOHNS HOPKINS UNIVERSITY
APPLIED PHYSICS LABORATORY
JOHNS HOPKINS ROAD
LAUREL, MD 20810
O1CY ATTN DOCUMENT LIBRARIAN
O1CY ATTN THOMAS POTEMRA
O1CY ATTN JOHN DASSOULAS

KAMAN SCIENCES CORP
P.O. BOX 7463
COLORADO SPRINGS, CO 80933
O1CY ATTN T. MEAGHER

KAMAN TEMPO-CENTER FOR ADVANCED STUDIES
816 STATE STREET (P.O. DRAWER QQ)
SANTA BARBARA, CA 93102
O1CY ATTN DASIAC
O1CY ATTN WARREN S. KNAPP
O1CY ATTN WILLIAM MCNAMARA
O1CY ATTN B. GAMBILL

LINKABIT CORP
10453 ROSELLE
SAN DIEGO, CA 92121
O1CY ATTN IRWIN JACOBS

LOCKHEED MISSILES & SPACE CO., INC
P.O. BOX 504
SUNNYVALE, CA 94088
O1CY ATTN DEPT 60-12
O1CY ATTN D.R. CHURCHILL

LOCKHEED MISSILES & SPACE CO., INC.
3251 HANOVER STREET
PALO ALTO, CA 94304
O1CY ATTN MARTIN WALT DEPT 52-12
O1CY ATTN W.L. IMHOF DEPT 52-12
O1CY ATTN RICHARD G. JOHNSON DEPT 52-12
O1CY ATTN J.B. CLADIS DEPT 52-12

MARTIN MARIETTA CORP
ORLANDO DIVISION
P.O. BOX 5837
ORLANDO, FL 32805
O1CY ATTN R. HEFFNER

M.I.T. LINCOLN LABORATORY
P.O. BOX 73
LEXINGTON, MA 02173
O1CY ATTN DAVID M. TOWLE
O1CY ATTN L. LOUGHLIN
O1CY ATTN D. CLARK

MCDONNELL DOUGLAS CORPORATION
5301 BOLSA AVENUE
HUNTINGTON BEACH, CA 92647
O1CY ATTN N. HARRIS
O1CY ATTN J. MOULE
O1CY ATTN GEORGE MROZ
O1CY ATTN W. OLSON
O1CY ATTN R.W. HALPRIN
O1CY ATTN TECHNICAL LIBRARY SERVICES

MISSION RESEARCH CORPORATION
735 STATE STREET
SANTA BARBARA, CA 93101
O1CY ATTN P. FISCHER
O1CY ATTN W.F. CREVIER
O1CY ATTN STEVEN L. GUTSCHE
O1CY ATTN R. BOGUSCH
O1CY ATTN R. HENDRICK
O1CY ATTN RALPH KILB
O1CY ATTN DAVE SOWLE
O1CY ATTN F. FAJEN
O1CY ATTN M. SCHEIBE
O1CY ATTN CONRAD L. LONGMIRE
O1CY ATTN B. WHITE

MISSION RESEARCH CORP.
1720 RANDOLPH ROAD, S.E.
ALBUQUERQUE, NEW MEXICO 87106
O1CY R. STELLINGWERF
O1CY M. ALME
O1CY L. WRIGHT

MITRE CORPORATION, THE
P.O. BOX 208
BEDFORD, MA 01730
O1CY ATTN JOHN MORGANSTERN
O1CY ATTN G. HARDING
O1CY ATTN C.E. CALLAHAN

MITRE CORP
WESTGATE RESEARCH PARK
1820 DOLLY MADISON BLVD
MCLEAN, VA 22101
O1CY ATTN W. HALL
O1CY ATTN W. FOSTER

PACIFIC-SIERRA RESEARCH CORP
12340 SANTA MONICA BLVD.
LOS ANGELES, CA 90025
O1CY ATTN E.C. FIELD, JR.

PENNSYLVANIA STATE UNIVERSITY
IONOSPHERE RESEARCH LAB
318 ELECTRICAL ENGINEERING EAST
UNIVERSITY PARK, PA 16802
(NO CLASS TO THIS ADDRESS)
O1CY ATTN IONOSPHERIC RESEARCH LAB

PHOTOMETRICS, INC.
4 ARROW DRIVE
WOBBURN, MA 01801
O1CY ATTN IRVING L. KOFISKY

PHYSICAL DYNAMICS, INC.
P.O. BOX 3027
BELLEVUE, WA 98009
O1CY ATTN E.J. FREMOUW

PHYSICAL DYNAMICS, INC.
P.O. BOX 10367
OAKLAND, CA 94610
ATTN A. THOMSON

R & D ASSOCIATES
P.O. BOX 9695
MARINA DEL REY, CA 90291
O1CY ATTN FORREST GILMORE
O1CY ATTN WILLIAM B. WRIGHT, JR.
O1CY ATTN ROBERT F. LELEVIER
O1CY ATTN WILLIAM J. KARZAS
O1CY ATTN H. ORY
O1CY ATTN C. MACDONALD
O1CY ATTN R. TURCO
O1CY ATTN L. DeRAND
O1CY ATTN W. TSAI

RAND CORPORATION, THE
1700 MAIN STREET
SANTA MONICA, CA 90406
O1CY ATTN CULLEN CRAIN
O1CY ATTN ED BEDROZIAN

RAYTHEON CO.
528 BOSTON POST ROAD
SUDBURY, MA 01776
O1CY ATTN BARBARA ADAMS

RIVERSIDE RESEARCH INSTITUTE
80 WEST END AVENUE
NEW YORK, NY 10023
O1CY ATTN VINCE TRAPANI

SCIENCE APPLICATIONS, INC.
P.O. BOX 2351
LA JOLLA, CA 92038

01CY ATTN LEWIS M. LINSON
01CY ATTN DANIEL A. HAMLIN
01CY ATTN E. FRIEMAN
01CY ATTN E.A. STRAKER
01CY ATTN CURTIS A. SMITH
01CY ATTN JACK MCDOUGALL

SCIENCE APPLICATIONS, INC
1710 GOODRIDGE DR.
MCLEAN, VA 22102
ATTN: J. COCKAYNE

SRI INTERNATIONAL
333 RAVENSWOOD AVENUE
MENLO PARK, CA 94025

01CY ATTN DONALD NEILSON
01CY ATTN ALAN BURNS
01CY ATTN G. SMITH
01CY ATTN R. TSUNODA
01CY ATTN DAVID A. JOHNSON
01CY ATTN WALTER G. CHESNUT
01CY ATTN CHARLES L. RINO
01CY ATTN WALTER JAYE
01CY ATTN J. VICKREY
01CY ATTN RAY L. LEADABRAND
01CY ATTN G. CARPENTER
01CY ATTN G. PRICE
01CY ATTN R. LIVINGSTON
01CY ATTN V. GONZALES
01CY ATTN D. MCDANIEL

TECHNOLOGY INTERNATIONAL CORP
75 WIGGINS AVENUE
BEDFORD, MA 01730
01CY ATTN W.P. BOQUIST

TOYON RESEARCH CO.
P.O. Box 6890
SANTA BARBARA, CA 93111
01CY ATTN JOHN ISE, JR.
01CY ATTN JOEL GARBARINO

TRW DEFENSE & SPACE SYS GROUP
ONE SPACE PARK
REDONDO BEACH, CA 90278

01CY ATTN R. K. PLEBUCH
01CY ATTN S. ALTSCHULER
01CY ATTN D. DEE
01CY ATTN D/ STOCKWELL
SNTF/1575

VISIDYNE
SOUTH BEDFORD STREET
BURLINGTON, MASS 01803
01CY ATTN W. REIDY
01CY ATTN J. CARPENTER
01CY ATTN C. HUMPHREY

END

FILMED

9-83

DTIC