

AD-A135 314

CONVERGENCE OF QUADRATIC FORMS IN P-STABLE RANDOM
VARIABLES AND THETA SUB. (U) NORTH CAROLINA UNIV AT
CHAPEL HILL DEPT OF STATISTICS S CAMBANIS ET AL.

1/1

UNCLASSIFIED

OCT 83 TR-41 AFOSR-TR-83-0975

F/G 12/1


NL

END

FORMED

1 84


3107



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

CENTER FOR STOCHASTIC PROCESSES

Department of Statistics
University of North Carolina
Chapel Hill, North Carolina



CONVERGENCE OF QUADRATIC FORMS IN p -STABLE RANDOM VARIABLES
AND θ_p -RADONIFYING OPERATORS

by
Stamatis Cambanis
and
Jan Rosinski
and
W. Woyczynski

TECHNICAL REPORT #41

October 1983

DTIC
ELECTE
DEC 1 1983

S D
H

DTIC FILE COPY

Approved for public release;
distribution unlimited.

88 11 29 232

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR. 33-0975	2. GOVT ACCESSION NO. AD-A135314	3. RECIPIENT'S CAT/LOG NUMBER
4. TITLE (and Subtitle) CONVERGENCE OF QUADRATIC FORMS IN p-STABLE RANDOM VARIABLES AND θ_p-RADONIFYING OPERATORS		5. TYPE OF REPORT & PERIOD COVERED TECHNICAL
7. AUTHOR(s) Stamatis Cambanis and Jan Rosinski and W. Woyczynski		6. PERFORMING ORG. REPORT NUMBER TR #41
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics University of North Carolina Chapel Hill NC 27514		8. CONTRACT OR GRANT NUMBER(s) F49620-82-C-0009
11. CONTROLLING OFFICE NAME AND ADDRESS Mathematical & Information Sciences Directorate Air Force Office of Scientific Research /NRM Bolling AFB DC 20332		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS PE61102F; 2304/A5
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE OCT 83
		13. NUMBER OF PAGES 20
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		18a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		Accession For NTIS GRA&I ☒ DTIC TAB ☐ Unannounced ☐ Justification
18. SUPPLEMENTARY NOTES		By Distribution/ Availability Codes
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Quadratic forms, p-stable random variables, double Wiener integrals, θ_p-Radonifying operators.		Avail and/or Special
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Necessary and sufficient conditions are given for the almost sure convergence of the quadratic form $\sum_{i,j} f_{ij} M_i M_j$ where (M_i) is a sequence of i.i.d. p-stable random variables. A connection is established between the convergence of the quadratic form and a radonifying property of the infinite matrix operator (f_{kj}).		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

-a-

CONVERGENCE OF QUADRATIC FORMS IN p -STABLE RANDOM VARIABLES
AND θ_p -RADONIFYING OPERATORS

Stamatis Cambanis¹
Department of Statistics
University of North Carolina
Chapel Hill, NC 27514

Jan Rosinski²
and
Wojbor A. Woyczynski¹
Department of Mathematics and Statistics
Case Western Reserve University
Cleveland, OH 44106

(M sub j)

Summary

Necessary and sufficient conditions are given for the almost sure convergence of the quadratic form $\sum \sum f_{jk} M_j M_k$ where $\{M_j\}$ is a sequence of i.i.d. p -stable random variables. A connection is established between the convergence of the quadratic form and a radonifying property of the infinite matrix operator (f_{jk}) .

(f sub k j).

¹Supported in part by the AFOSR Contract No. F49620 82 C 0009. The last named author would like to express his gratitude to the Center for Stochastic Processes of the University of North Carolina for its hospitality while this research was being carried out.

²On leave from Wroclaw University, Poland.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFOSR)
NOTICE OF TRANSMITTAL TO DTIC
This technical report has been reviewed and is
approved for public release in accordance with AFR 190-12.
Distribution is unlimited.
MATTHEW J. KEMPER
Chief, Technical Information Division

I. INTRODUCTION

The aim of this paper is to study the convergence of the random quadratic forms of the form

$$(1.1) \quad \sum_{k,j} f_{jk} M_j M_k$$

where (f_{jk}) , $j,k=1,2,3,\dots$, is a real infinite matrix and (M_j) , $j=1,2,\dots$, is a sequence of i.i.d. p -stable random variables with characteristic function $\exp(-|t|^p)$, $0 < p < 2$. Our results have obvious implications in the theory of double Wiener-type integrals of the form

$$\iint f(x,y) M(dx) M(dy)$$

where $M(x)$ is a p -stable motion (cf. Szulga and Woyczynski (1983)). We shall study them elsewhere.

We begin with a characterization of non-anticipating sequences (V_k) such that the "martingale" transform

$$T = \sum V_k M_k$$

converges almost surely. The necessary and sufficient condition here turns out to be $\sum |V_k|^p < \infty$ a.s., and moreover T converges a.s. exactly on the set $(\sum |V_k|^p < \infty)$ (Theorem 2.1). This applied to the sequence

$$V_k = \sum_{j=1}^{k-1} f_{jk} M_j$$

shows that the convergence of the off-diagonal part of the iterated sum

$$\sum_k \left(\sum_{j < k} f_{jk} M_j \right) M_k$$

is equivalent to the matrix operator $(f_{jk})_{j < k}^T$ being θ_p -radonifying from ℓ^q into ℓ^p (Theorems 2.2 and 3.1).

Finally we characterize the class of θ_p -radonifying operators $(f_{jk})^T$ from ℓ^q into ℓ^p as satisfying the condition

$$\sum_k \{ (\sum_j |f_{jk}|^p) (1 + |\log \sum_j |f_{jk}|^p|) \} < \infty$$

(Corollary 2.2). The proof of necessity of this condition was shown to us by Gilles Pisier and is included here with his permission.

The above results permit the full characterization of all matrices (f_{jk}) for which the quadratic form (1.1) converges a.s. (Theorem 4.2). We would also like to mention here that the infinite quadratic forms satisfy a fairly general 0-1 law (cf. de Acosta (1976)).

Our results should be compared with the case where the M_j 's are i.i.d. Gaussian, i.e. when $p=2$. In this case Varberg (1966) has shown that the convergence of the series $\sum_k f_{kk}$ and $\sum_{jk} f_{jk}^2$ is necessary and sufficient for the quadratic mean convergence of (1.1), and the above conditions imply a.s. convergence of (1.1) as well. Moreover Lemma 3.7 of Rosinski and Szulga (1983) in conjunction with Kahana's (1968) inequality shows that the conditions are also necessary for a.s. convergence of the Gaussian quadratic form.

In Section 5 we include some auxiliary results and a conjecture of independent interest.

2. MARTINGALE TRANSFORMS AND CONVERGENCE OF STABLE TRIANGULAR QUADRATIC FORMS

THEOREM 2.1. Let $V_k = V_k(M_1, \dots, M_{k-1})$, $k=2,3,\dots$, be a non-anticipating sequence of random variables. Then almost surely

$$\{\sum V_k M_k \text{ converges}\} = \{\sum |V_k|^p < \infty\}.$$

Proof: Let $F_k = \sigma(M_1, \dots, M_k)$, and

$$G(x) = P\{|M_1| > x\} \sim cx^{-p},$$

as $x \rightarrow \infty$, $c > 0$. Also let us set

$$A = \{\sum V_k M_k \text{ converges}\},$$

$$B = \{\sum |V_k|^p < \infty\}$$

and

$$C_n = \{|V_n M_n| > 1\}, n=2,3,\dots$$

Then we have that

$$A \subset \{C_n \text{ i.o.}\}^c = \{\sum P(C_n | F_{n-1}) < \infty\} \text{ a.s.}$$

where the equality is implied by a conditional form of a Borel-Cantelli lemma

(cf. Breiman (1968), p. 96). Since

$$P(C_n | F_{n-1}) = G(1/|V_n|),$$

we get that $G(|V_n|^{-1}) \rightarrow 0$ a.s. on A as $n \rightarrow \infty$. Therefore, for sufficiently large $N = N(\omega)$,

$$\sum_{n=N}^{\infty} |V_n|^p \leq \frac{2}{c} \sum_{n=N}^{\infty} G(|V_n|^{-1}) < \infty$$

a.s. on A . This shows that $P(A \setminus B) = 0$.

Now we shall prove that $P(B \setminus A) = 0$. Let $\pi_n = \pi_n(\cdot, \omega)$ be a regular conditional distribution of $V_n M_n$ given F_{n-1} . We have that

$$\pi_n(E, \omega) = P(V_n M_n \in E | M_1, \dots, M_{n-1}) = \mu_p(V_n^{-1}(\omega)E)$$

where μ_p is the distribution of M_1 . By a theorem of Hill (1982), $\sum V_n(\omega) M_n(\omega)$ converges for almost all ω 's for which the series $\sum X_n^{(\omega)}$ converges in probability

P' , where $\{X_n^{(\omega)}(\omega')\}$ is a sequence of independent r.v.'s (defined on another probability space (Ω', P')) with distributions $\{\pi_n(\cdot, \omega)\}$. In our case the $X_n^{(\omega)}$'s have characteristic functions $\exp(-|V_n(\omega)|^p t^p)$. Thus $P(B \setminus A) = 0$ which completes the proof of the theorem. $\square$

Theorem 2.1. enables us to translate the problem of a.s. convergence of quadratic forms into a more tractable problem of the a.s. convergence of series of independent random vectors with values in ℓ^p , whose standard basis is denoted by (e_k) , $k=1, 2, \dots$.

THEOREM 2.2. Let $(f_{jk} : 1 \leq j \leq k-1, k \geq 2)$ be a triangular matrix of real numbers, and $x_j \stackrel{\text{def}}{=} (0, \dots, 0, f_{j,j+1}, f_{j,j+2}, \dots) = \sum_{k=j+1}^{\infty} f_{jk} e_k$, $j=1, 2, \dots$. Then

$$\sum_{k=2}^{\infty} \left(\sum_{j=1}^{k-1} f_{jk} M_j \right) M_k$$

converges a.s. if and only if for every $j=1, 2, \dots$, $x_j \in \ell^p$ and the vector random series $\sum x_j M_j$ converges a.s. in ℓ^p .

Proof: Setting $V_k = \sum_{j=1}^{k-1} f_{jk} M_j$, $k=2, 3, \dots$, $V_1 = 0$, by Theorem 2.1 we have that the series $\sum V_k M_k$ converges a.s. if and only if $\sum |V_k|^p < \infty$ a.s., which is equivalent to the a.s. convergence in ℓ^p of the series $\sum e_k V_k$.

Now assume that $S = \sum e_k V_k$ converges a.s. in ℓ^p . Since

$$\sum_{k=1}^{n+1} e_k V_k = \sum_{j=1}^n \left(\sum_{k=j+1}^{n+1} f_{jk} e_k \right) M_j,$$

Proposition 5.1 applied to $x_{jn} = \sum_{k=j+1}^{n+1} f_{jk} e_k$, and $Y_j = M_j$, gives that

$x_{jn} \rightarrow x_j = \sum_{k=j+1}^{\infty} f_{jk} e_k \in \ell^p$ as $n \rightarrow \infty$, and that the series $\sum x_j M_j$ converges a.s. in ℓ^p .

Conversely, assume that $x_j = \sum_{k=j+1}^{\infty} f_{jk} e_k \in \ell^p$ and $\sum x_j M_j$ converges a.s. in ℓ^p .

The operator

$$\ell^p \ni x \rightarrow R_n(x) = \sum_{k=n+2}^{\infty} \langle x, e_k \rangle e_k \in \ell^p, \quad p > 0,$$

is continuous and linear and $R_n(x) \rightarrow 0$ in ℓ^p for every x as $n \rightarrow \infty$. Thus a.s.

$$0 + R_n \left(\sum_{j=1}^{\infty} x_j M_j \right) = \sum_{j=1}^{\infty} R_n(x_j) M_j = \sum_{j=1}^n \left(\sum_{k=n+2}^{\infty} f_{jk} e_k \right) M_j + \sum_{j=n+1}^{\infty} x_j M_j .$$

Hence,

$$\sum_{j=1}^n \left(\sum_{k=n+2}^{\infty} f_{jk} e_k \right) M_j \rightarrow 0$$

a.s. in ℓ^p as $n \rightarrow \infty$, and thus

$$\sum_{k=1}^{n+1} v_k e_k = \sum_{j=1}^n x_j M_j - \sum_{j=1}^n \left(\sum_{k=n+2}^{\infty} f_{jk} e_k \right) M_j$$

converges a.s. in ℓ^p .

□

3. CONVERGENCE OF p -STABLE RANDOM SERIES IN ℓ^p AND θ_p -RADONIFYING OPERATORS

Definition 3.1. Let $0 < p < 2$ and F be a continuous linear operator $F: s \rightarrow \ell^p$, where s is the space of sequences with finite support for $0 < p \leq 1$ and $s = \ell^q$, $1/q + 1/p = 1$, for $1 < p < 2$. We shall write $F = (f_{jk})$ where $f_{jk} = \langle Fe_j, e_k \rangle$, $j, k = 1, 2, \dots$, so that $Fe_j = \sum_k f_{jk} e_k$. We say that F is in the class $\ell^p \log \ell(\ell^p)$ if for each $j = 1, 2, \dots$, $Fe_j \in \ell^p$ and the ℓ^p -valued sequence (Fe_j) is in $\ell^p \log \ell$. In other words, $F \in \ell^p \log \ell(\ell^p)$ if and only if

$$\begin{aligned} N(F) &= \sum_j \|Fe_j\|_p^p (1 + |\log \|Fe_j\|_p|) \\ &= \sum_j \left\{ \left(\sum_k |f_{jk}|^p \right) (1 + |\log \sum_k |f_{jk}|^p|) \right\} < \infty. \end{aligned}$$

One can introduce on $\ell^p \log \ell(\ell^p)$ an Orlicz type norm in the natural fashion.

THEOREM 3.1. Let $0 < p < 2$ and $x_j = \sum_{k=1}^{\infty} f_{jk} e_k \in \ell^p$, $j = 1, 2, \dots$. Then $\sum_j x_j M_j$ converges a.s. in ℓ^p if and only if $F^T = (f_{jk})^T \in \ell^p \log \ell(\ell^p)$.

Proof: Sufficiency. Assume $F^T \in \ell^p \log \ell(\ell^p)$. We shall prove that the assumptions of Proposition 5.2 are fulfilled for the series with general term $W_k = |\sum_j f_{jk} M_j|^p$, which has the same distribution as $(\sum_j |f_{jk}|^p) |M_1|^p$. Indeed,

$$\sum_k P(W_k > 1) = \sum_k P(|M_1|^p > (\sum_j |f_{jk}|^p)^{-1/p}) \leq \text{Const} \sum_{jk} |f_{jk}|^p < \infty$$

since, by assumption, $\sum_k \sum_j |f_{jk}|^p < \infty$ and $\sum_j |f_{jk}|^p \rightarrow 0$ as $k \rightarrow \infty$. Therefore (5.1) of Proposition 5.2 is satisfied.

Now,

$$B = \sum_k E(W_k I(W_k \leq 1)) = \sum_k \left(\sum_j |f_{jk}|^p \right) E(|M_1|^p I(|M_1|^p \leq (\sum_j |f_{jk}|^p)^{-1/p}))$$

Denoting the distribution function of $|M_1|$ by ϕ , we get that for $\alpha > 1$

$$\int_0^\alpha |u|^p d\phi(u) \leq 1 + \alpha^p \phi(\alpha) - \phi(1) - \int_1^\alpha p u^{p-1} \phi(u) du$$

$$\leq 1 + \alpha^P \Phi(\alpha) - \Phi(1) - \int_1^\alpha pu^{P-1} (1 - (C/u^P)) du$$

since $\Phi(u) = 1 - P(|M_1| > u) \geq 1 - Cu^{-P}$ for $u \geq 1$ and certain constant C . Thus because $\alpha^P(\Phi(\alpha) - 1)$ remains bounded

$$\int_0^\alpha |u|^P d\Phi(u) \leq \text{Const} (1 + \log \alpha).$$

It follows that

$$B \leq \text{Const} \sum_k \sum_j |f_{jk}|^P (1 + |\log \sum_j |f_{jk}|^P|) < \infty$$

so that Condition (5.2) of Proposition 5.2 is also satisfied and as a result

$\sum_k |\sum_j f_{jk} M_j|^P < \infty$ a.s. It follows that $\sum_k (\sum_j f_{jk} M_j) e_k$ converges a.s. in ℓ^P and in view of $x_j = \sum_k f_{jk} e_k \in \ell^P$, we have that $\sum_j x_j M_j$ converges a.s. in ℓ^P .

Necessity. (The basic idea here is due to Pisier). Assume that $\sum x_j M_j = \sum (Fe_j) M_j$ converges in ℓ_p a.s. By the representation for stable processes obtained in Proposition 1.5 of Marcus and Pisier (1983) and used here in the discrete parameter case (in the case of $1 \leq p < 2$, Corollary 3 of LePage, Woodroffe and Zinn (1981) would suffice) $\sum x_j M_j$ converges in ℓ^P a.s. if and only if $\sum \varepsilon_j Y_j \Gamma_j^{-1/p}$ converges a.s. in ℓ^P , where (ε_j) are independent Bernoulli r.v.'s, (Y_j) are i.i.d. r. elements in ℓ^P with law

$$L(Y_j) = \text{Const} \sum_k \|x_k\|^P \delta_{x_k / \|x_k\|}, \quad j=1,2,\dots,$$

and $\Gamma_j = X_1 + \dots + X_k$ where $X_1, X_2, \dots$ are i.i.d. with $P(X_1 > u) = e^{-u}$. Moreover the sequences $(\varepsilon_j), (Y_j), (\Gamma_j)$ are independent of each other. By the law of large numbers, $\Gamma_j/j \rightarrow 1$ a.s. Applying twice Fubini's theorem and the comparison principle in quasi-normed spaces obtained in Theorem 4.4 of Marcus and Woyczynski (1979), we get that the series $\sum x_j M_j$ converges in ℓ^P a.s. if and only if $\sum \varepsilon_j Y_j j^{-1/p}$ does. What is useful about the latter series is that it possesses all moments in contrast to the former.

Therefore for $0 < p < 2$, and $x_j = Fe_j$, $j=1,2,\dots$,

$$\begin{aligned}
 \infty &> E \left\| \sum_j j^{-1/p} \epsilon_j Y_j \right\|_p^p = \sum_k E \left\| \sum_j j^{-1/p} \epsilon_j Y_{jk} \right\|_p^p \\
 (2.2) \quad &\geq \text{Const} \sum_k E \left(\sum_j j^{-2/p} Y_{jk}^2 \right)^{p/2} \\
 &\geq \text{Const} \sum_k E \left(\sup_j j^{-1/p} |Y_{jk}| \right)^p
 \end{aligned}$$

by Khintchine's inequality ($Y_j = (Y_{j1}, Y_{j2}, \dots)$).

If C denotes the class of all stopping times, then for all k ,

$$E \left| \sup_j j^{-1/p} Y_{jk} \right|^p \geq \sup_{\tau \in C} E \tau^{-1} |Y_{\tau k}|^p \geq \text{Const} E(|Y_{1k}|^p \log |Y_{1k}|)$$

by Chow, Robbins & Siegmund (1971), p. 97, where the last constant can be chosen greater than 0 (uniformly in k), since in our case

$$L(Y_{jk}) = \text{Const} \sum_i \|Fe_i\|_p^p \delta_{\langle Fe_i, e_k \rangle} \|Fe_i\|_p^p.$$

Therefore

$$E(|Y_{1k}|^p \log |Y_{1k}|) = \sum_i \frac{|f_{ik}|^p}{\|Fe_i\|_p^p} \log \frac{|f_{ik}|}{\|Fe_i\|_p} \cdot \|Fe_i\|_p^p$$

so that finally we get from (2.2) that

$$\sum_k \sum_j |f_{jk}|^p \log \frac{|f_{jk}|}{\sum_k |f_{jk}|^p} < \infty$$

which certainly implies that $F^T \in \ell^p \log \ell(\ell^p)$. □

COROLLARY 3.1. Let $\sum_k |f_{jk}|^p < \infty$. Then $\sum_j |\sum_k f_{jk} M_k|^p < \infty$ a.s. if and only if $\sum_j \{ (\sum_k |f_{jk}|^p) (1 + |\log \sum_k |f_{jk}|^p|) \} < \infty$.

Now let $1 < p \leq 2$ and let θ_p be a cylindrical measure on ℓ^q , $1/p + 1/q = 1$, generated by the sequence $M = (M_i)$, $i=1,2,\dots$. The characteristic functional of θ_p is given by the formula

$$\int_{\ell^q} e^{i\langle x, y \rangle} \theta_p(dy) = e^{-\|x\|_p^p}, \quad x \in \ell^p.$$

Definition 3.2. Let E be a Banach space. A linear operator $F: \ell^q \rightarrow E$ is called θ_p -radonifying if $\exp(-\|F^*y^*\|_p^p)$, $y^* \in E^*$, is the characteristic functional of a Radon measure on E . We denote by $R_p(\ell^q, E)$ the class of all such operators.

The class $R_2(\ell^2, E)$ has been extensively studied. The main result is that $\Pi_2(\ell^2, E) = R_2(\ell^2, E)$ if and only if E is of cotype 2 (cf. Linde and Pietsch (1974)), where Π_p denotes the ideal of all p -absolutely summing operators. For $1 < p < 2$ it has been proved that if $1 < r < p$ then $\Pi_r(\ell^q, E) \subset R_p(\ell^q, E)$, and that $\Pi_p(\ell^q, E) \subset R_p(\ell^q, E)$ if and only if E is isomorphic to subspace of L^p and is of stable type p (cf. Linde, Mandrekar, Weron (1980), and Thang and Tien (1982)).

From the now classical results due to Itô and Nisio (1968) it is easy to deduce the following.

THEOREM 3.2. Let $1 < p \leq 2$. The following are equivalent:

- (a) $F \in R_p(\ell^q, E)$;
- (b) $F^*: E^* \rightarrow \ell^p$ is decomposable in the sense that there exists an E -valued, strongly measurable random vector (FM) such that

$$\langle F^*y^*, M \rangle = \langle y^*, FM \rangle \quad \text{a.s., } y^* \in E^*;$$

- (c) The series $\sum F(e_j)M_j$ converges a.s. in E , where (e_j) is the standard basis in ℓ^q .

The above equivalences are heuristically better understood if one keeps in mind the following "identities":

$$E e^{i \langle T^*y^*, M \rangle} = E e^{i \sum_k M_k \langle T^*y^*, e_k \rangle} = e^{-\sum_k |\langle T^*y^*, e_k \rangle|^p} = e^{-\|T^*y^*\|_p^p}.$$

In view of the above remarks every r -absolutely summing F , $1 < r < p$, is θ_p -radonifying, but there are p -absolutely summing $F: \ell^q \rightarrow \ell^p$ which are not θ_p -radonifying since ℓ^p is not of stable type p . The corollary below, obtained immediately from Theorems 3.1 and 3.2, reflects this fact and gives an analytic description of θ_p -

radonifying operators F from $\ell^q \rightarrow \ell^p$.

COROLLARY 3.2. $[R_p(\ell^q, \ell^p)]^T = \ell^p \log \ell(\ell^p)$.

4. CONVERGENCE OF GENERAL QUADRATIC FORMS IN p -STABLE RANDOM VARIABLES

Let

$$Q_n = \sum_{k=1}^n \sum_{j=1}^n f_{jk} M_j M_k, \quad n=1,2,\dots,$$

where (M_j) , $j=1,2,\dots$, are as above i.e. i.i.d. with $E \exp(itM_j) = \exp(-|t|^p)$, $0 < p < 2$. Denote the diagonal and off-diagonal parts of Q_n respectively by

$$D_n = \sum_{k=1}^n f_{kk} M_k^2 \quad \text{and} \quad R_n = \sum_{\substack{k,j=1 \\ k \neq j}}^n f_{jk} M_j M_k.$$

The diagonal part, being a series of independent random variables, is easy to handle.

THEOREM 4.1. *The sequence (D_n) converges a.s. as $n \rightarrow \infty$ if and only if*

$$\sum_{k=1}^{\infty} |f_{kk}|^{p/2} < \infty.$$

Proof. Let us observe that $\sum P(|f_{kk}| M_k^2 > 1) < \infty$ if and only if $\sum |f_{kk}|^{p/2} < \infty$.

Kolmogorov's three series theorem gives now the "only if" part of the claim. The proof of the "if" part follows directly from Proposition 5.2 because for $p < 2$ and $f_{kk} \neq 0$,

$$E |f_{kk}| M_k^2 I(|f_{kk}| M_k^2 \leq 1) = \int_0^1 P(M_k^2 > t |f_{kk}|^{-1}) dt \leq c |f_{kk}|^{p/2} \int_0^1 t^{-p/2} dt = \text{Const } |f_{kk}|^{p/2}. \quad \square$$

The above result and the results of previous sections give the following.

THEOREM 4.2. *The sequence of quadratic forms (Q_n) converges a.s. as $n \rightarrow \infty$ if and only if*

$$\sum_{k=1}^{\infty} |f_{kk}|^{p/2} < \infty$$

and

$$\sum_{k=2}^{\infty} \left\{ \left(\sum_{j=1}^{k-1} |f_{jk} + f_{kj}|^p \right) (1 + |\log \sum_{j=1}^{k-1} |f_{jk} + f_{kj}|^p|) \right\} < \infty.$$

Proof: The above result is a straightforward corollary to Theorems 4.1, 2.2, and 3.1 and to the fact that the sequence (Q_n) converges a.s. if and only if both

sequences (D_n) and (R_n) converge a.s. We prove the only if part of this latter assertion.

Let a, b be reals with $a \neq 0$, and set $\varepsilon = \text{sign}(b/a)$. Then for $M = M_1$, we have

$$\begin{aligned}
 (4.1) \quad P\{|aM^2 + bM| > 1\} &= P\{|a(\varepsilon M)^2 + b(\varepsilon M)| > 1\} = P\{|a|M^2 + |b|M| > 1\} \\
 &\geq P\{|a|M^2 + |b|M| > 1, M > 0\} \\
 &= P\{|a|M^2 + |b|M > 1, M > 0\} \\
 &\geq P\{M > |a|^{-1/2}\} = \frac{1}{2} P\{|M| > |a|^{-1/2}\}.
 \end{aligned}$$

Assume that (Q_n) converges a.s. We have $Q_n = \sum_{k=1}^n (V_k + f_{kk} M_k) M_k$ where $V_k = \sum_{j=1}^{k-1} (f_{jk} + f_{kj}) M_j$, $k \geq 2$, $V_1 = 0$. By the conditional Borel-Cantelli lemma (cf. Breiman (1968), p. 96), the a.s. convergence of (Q_n) implies that

$$\sum_{k=1}^{\infty} P\{|V_k M_k + f_{kk} M_k^2| > 1 | M_1, \dots, M_{k-1}\} < \infty \text{ a.s.}$$

It follows by (4.1) that $\sum P\{|f_{kk} M_k^2| > 1\} < \infty$, where the sum extends over all k for which $f_{kk} \neq 0$, which implies that $\sum_{k=1}^{\infty} |f_{kk}|^{p/2} < \infty$, in view of the tail behavior of the M_k 's. Now, by Theorem 4.1, D_n converges a.s. and by the assumption $R_n = Q_n - D_n$ converges a.s. as well. $\square$

In view of Theorem 5.1 below and of Theorem 2.1, we have the following corollary on the convergence in L^r , $r < p$, of the off-diagonal part of the quadratic form.

COROLLARY 4.1. *If $0 < r < p$ and*

$$\sum_{k=2}^{\infty} \left(\sum_{j=1}^{k-1} |f_{jk} + f_{kj}|^p \right)^{r/p} < \infty$$

then the sequence $R_n = \sum_{k \neq j}^n f_{jk} M_j M_k$ converges in L^r and a.s.

Proof: Here is a direct proof which does not use Theorems 2.1 and 5.1. By the martingale property of R_n we have, with $V_k = \sum_{j=1}^{k-1} (f_{jk} + f_{kj}) M_j$, $k \geq 2$, $V_1 = 0$,

$$\begin{aligned}
E|R_n - R_{m+1}|^r &= E\left|\sum_{k=m}^n V_k M_k\right|^r \leq \text{Const} \sum_{k=m}^n E|V_k M_k|^r \\
&= \text{Const} \sum_{k=m}^n E|V_k|^r E|M_k|^r \\
&= \text{Const} \sum_{k=m}^n \left(\sum_{j=1}^{k-1} |f_{jk} + f_{kj}|^p\right)^{r/p}
\end{aligned}$$

which proves the corollary. □

5. AUXILIARY RESULTS

The proof of Theorem 2.2 relies on the following technical property which, roughly speaking, justifies the change of the order of summation in the series

$$\sum_j (\sum_{k>j} f_{jk} e_k) M_j.$$

PROPOSITION 5.1. *Let E be a complete metric linear space, $x_{jk} \in E$, $n, j=1, 2, \dots$, and (Y_j) a sequence of non-zero independent symmetric real random variables. If*

$$S_n = \sum_{j=1}^n x_{jn} Y_j \rightarrow S$$

in probability as $n \rightarrow \infty$ then there exists a sequence $(x_j) \subset E$ such that for each j , $x_{jn} \rightarrow x_j$ as $n \rightarrow \infty$ and the series $\sum x_j Y_j$ converges a.s. to S .

Proof: Let $\|\cdot\|$ be a monotonic F-norm on E , i.e. $\|ax\| \leq \|x\|$ for $|a| \leq 1$ (which always exists by Rolewicz (1972), p. 16, Theorem I.2.2). Fix $j \geq 1$ and let $a, b > 0$ be such that $P(|Y_j| > a) > b$. Let $\epsilon > 0$. For $r \geq n \geq j$ we set $c(r, n) = 1$ if $\|a(x_{jr} - x_{jn})\| > \epsilon$ and $c(p, n) = 0$ otherwise. Since for a symmetric pair of r , vectors X, Y in E ,

$$P(\|X\| > \epsilon) \leq 2P(\|\frac{1}{2}(X+Y)\| > \frac{\epsilon}{2})$$

we obtain

$$\begin{aligned} b \cdot c(r, n) &\leq P(\|Y_j(x_{jr} - x_{jn})\| > \epsilon) \leq 2P(\|\frac{1}{2} \sum_{k=1}^n Y_k(x_{kr} - x_{kn})\| > \frac{\epsilon}{2}) \\ &\leq 4P(\|\frac{1}{4}(S_r - S_n)\| > \frac{\epsilon}{4}). \end{aligned}$$

It then follows from the assumption that $c(r, n) \rightarrow 0$ as $r, n \rightarrow \infty$, i.e. $\{x_{jr}\}$ as a Cauchy sequence for every j , and by the completeness of E there exists $x_j \in E$ such that $x_{jn} \rightarrow x_j$ as $n \rightarrow \infty$.

Let now $\epsilon > 0$ and let N be such that

$$P(\|\frac{1}{2}(S_r - S_n)\| > \frac{\epsilon}{2}) \leq \frac{\epsilon}{2}$$

for every $r \geq n \geq N$. By the symmetry argument used above

$$P\{\|\sum_{j=1}^n Y_j(x_{jr} - x_{jn})\| \geq \epsilon\} \leq \epsilon.$$

Keeping n fixed and letting $r \rightarrow \infty$ we get

$$P\{\|\sum_{j=1}^n x_j Y_j - S_n\| \geq \epsilon\} \leq \epsilon.$$

Thus $\sum_{j=1}^n x_j Y_j \rightarrow S$ in probability and, since the Y 's are independent, also a.s. $\square$

The following elementary proposition (cf. Szulga and Woyczynski (1983)) is used in the proof of Theorem 3.1.

PROPOSITION 5.2. *If for a sequence (W_k) of real random variables*

$$(5.1) \quad \sum_k P(|W_k| > 1) < \infty,$$

and

$$(5.2) \quad \sum_k E|W_k I(|W_k| \leq 1)| < \infty$$

then $\sum |W_k| < \infty$ a.s.

Proof: Indeed, let $Y_k = W_k I(|W_k| \leq 1)$, $Z_k = W_k - Y_k$. Then $|W_k| = |Y_k| + |Z_k|$. $\sum |Y_k|$ converges a.s. since $E(\sum |Y_k|) < \infty$ by (5.2), and $\sum |Z_k|$ converges a.s. by the Borel-Cantelli lemma. $\square$

We now establish a more precise criterion for summability of stable r.v.'s which is used in Section 4. For a symmetric p -stable r.v. X the quantity c_X is defined by $E \exp(itX) = \exp(-c_X |t|^p)$ and satisfies

$$\|X\|_{L^r} = (E|X|^r)^{1/r} = \text{Const}(r,p) c_X^{1/p}, \quad 0 < r < p,$$

and for independent symmetric p -stable r.v.'s (X_k) ,

$$(5.3) \quad c_{\sum_k a_k X_k} = \sum_k |a_k|^p c_{X_k}.$$

Definition 5.1. The r.v.'s (X_k) are jointly symmetric p -stable if for every sequence (a_k) with a finite number of nonzero elements the r.v. $\sum_k a_k X_k$ is symmetric p -stable.

THEOREM 5.1. Let (X_k) be jointly symmetric p -stable r.v.'s with $0 < p < 2$ and let $r > 0$. Then a necessary and sufficient condition for

$$\sum_{k=1}^{\infty} |X_k|^r < \infty \text{ a.s.}$$

is that for some $0 < s < p$,

$$E\left(\sum_{k=1}^{\infty} |X_k|^r\right)^s < \infty \quad \text{when } 0 < r \leq 1$$

and

$$E\left(\sum_{k=1}^{\infty} |X_k|^r\right)^{s/r} < \infty \quad \text{when } 1 < r.$$

Proof: Assume $X = (X_k)_k \in \ell^r$ a.s. and define $\phi: \Omega \rightarrow \ell^r$ by $\phi(\omega) = (X_k(\omega))_k = X(\omega)$ if $(X_k(\omega))_k \in \ell^r$ and $\phi(\omega) = 0$ otherwise. Then ϕ induces a symmetric p -stable measure $\mu = P \circ \phi^{-1}$ on ℓ^r . For $x = (x_k)_k \in \ell^r$ define $q(x) = \sum_k |x_k|^r$ when $0 < r \leq 1$ and $q(x) = (\sum_k |x_k|^r)^{1/r}$ when $r > 1$. Then q is a measurable seminorm on ℓ^r (a norm when $r \geq 1$) and by Theorem 3.2 in de Acosta (1975) we have for $0 < s < p$,

$$E\{q(X(\omega))\}^s = \int_{\Omega} q^s(X(\omega)) dP(\omega) = \int_{\ell^r} q^s(x) d\mu(x) < \infty.$$

The converse is clear. □

When $1 < r < p < 2$ we can take $s = r$ and the necessary and sufficient condition becomes

$$\sum_{k=1}^{\infty} E|X_k|^r < \infty \quad \text{or} \quad \sum_{k=1}^{\infty} c_{X_k}^{r/p} < \infty.$$

When $r = p$, Theorem 5.1 gives

$$(5.4) \quad \sum_{k=1}^{\infty} |X_k|^p < \infty \text{ a.s. if and only if } E\left(\sum_{k=1}^{\infty} |X_k|^p\right)^{s(1 \wedge \frac{1}{p})} < \infty \text{ for some } 0 < s < p.$$

This necessary and sufficient condition simplifies to

$$(5.5) \quad \sum_{k=1}^{\infty} c_{X_k} (1 + |\log c_{X_k}|) < \infty$$

when the X_k 's are independent, by Schwartz's theorem (cf. e.g. Woyczynski (1978), p. 277), and when the X_k 's are of the form $X_k = \sum_j f_{jk} M_j$, where the M_j 's are independent (cf. Corollary 3.1 and (5.3)). Since every sequence of jointly symmetric p -stable r.v.'s (X_k) is of the form $X_k = \int_0^1 f_k(t) dM(t)$, $k=1,2,\dots$, where $M(t)$, $0 \leq t \leq 1$, is a stable motion (i.e. has independent stationary symmetric p -stable increments) and $\int_0^1 |f_k(t)|^p dt < \infty$, $k=1,2,\dots$ (cf. Kuelbs (1973)), we conjecture that (5.5) is always a necessary and sufficient condition for (5.4).

REFERENCES

- DE ACOSTA, A. (1975). Stable measures and seminorms. *Ann. Probability* 3 865-875.
- DE ACOSTA, A. (1976). Quadratic zero-one laws for Gaussian measures and the distribution of quadratic forms. *Proc. Amer. Math. Soc.* 54 319-325.
- BREIMAN, L. (1968). *Probability*. Addison-Wesley, Reading, Mass.
- CHOW, Y.S., ROBBINS, H. and SIEGMUND, D. (1971). *Great Expectations: The Theory of Optimal Stopping*. Houghton Mifflin, Boston.
- HILL, T.P. (1982). Conditional generalizations of strong laws which conclude that partial sums converge almost surely. *Ann. Probability* 10 828-830.
- ITÔ, K. and NISIO, M. (1968). On the convergence of sums of independent Banach space valued random variables. *Osaka J. Math.* 5 35-48.
- KAHANE, J.P. (1968). *Some Random Series of Functions*. Heath, Lexington, Mass.
- KUELBS, J. (1973). A representation theorem for symmetric stable processes and stable measures on H . *Z. Wahrsch. verw. Geb.* 26 259-271.
- LE PAGE, R., WOODROOFE, M. and ZINN, J. (1981). Convergence to a stable distribution via order statistics. *Ann. Probability* 9 624-632.
- LINDE, W., MANDREKAR, V. and WERON, A. (1980). p -stable measures and p -absolutely summing operators. *Lecture Notes in Mathematics* No. 828, pp. 167-177, Springer-Verlag, Berlin.
- LINDE, W. and PIETSCH, A. (1974). Mappings of Gaussian cylindrical measures in Banach spaces. *Theor. Probability Appl.* 19 445-460.
- MARCUS, M. and PISIER, G. (1983). Characterizations of almost surely continuous p -stable random Fourier series and strongly stationary processes (to appear).
- MARCUS, M. and WOYCZYNSKI, W. (1979). Stable measures and central limit theorems in spaces of stable type. *Trans. Amer. Math. Soc.* 251 71-102.
- ROLEWICZ, S. (1972). *Metric Linear Spaces*. Polish Scientific Publ., Warszawa.
- ROSINSKI, J. and SZULGA, J. (1983). Product random measures and double stochastic integrals. In *Lecture Notes in Mathematics* No. 939 pp. 181-199. Springer-Verlag, Berlin.
- SZULGA, J. and WOYCZYNSKI, W.A. (1983). Existence of a double random integral with respect to stable measures. *J. Multivariate Anal.* 13 194-201.

THANG, D.H. and TIEN, N.D. (1982). Mappings of stable cylindrical measures in Banach spaces. Theor. Probability Appl. 27 492-501.

VARBERG, D. (1966). Convergence of quadratic forms in independent random variables. Ann. Math. Statist. 37 567-576.

WOYCZYNSKI, W.A. (1978). Geometry and martingales in Banach spaces, Part II. Adv. in Probability 4 267-517. Dekker, New York.

END

FILMED

1-84

DTIC