

MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

12

NAVENVPREDRSCHFAC
CONTRACTOR REPORT
CR 83-07



NAVENVPREDRSCHFAC CR 83-07

AD-A136146

TROPICAL CYCLONE WIND PROBABILITY FORECASTING FOR THE SOUTHERN HEMISPHERE (WINDPSH)


Prepared By:

Jerry D. Jarrell
Science Applications, Inc.
Monterey, CA 93940

Contract No. N00228-81-C-H361

OCTOBER 1983

APPROVED FOR PUBLIC RELEASE
DISTRIBUTION UNLIMITED



A
98

DTIC FILE COPY



Prepared For: 83

12 20

NAVAL ENVIRONMENTAL PREDICTION RESEARCH FACILITY
MONTEREY, CALIFORNIA 93943

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NAVENVPREDNSCHFAC Contractor Report CR 83-07	2. GOVT ACCESSION NO. AD-A136	3. RECIPIENT'S CATALOG NUMBER 146
4. TITLE (and Subtitle) Tropical Cyclone Wind Probability Forecasting for the Southern Hemisphere (WINDPSH)	5. TYPE OF REPORT & PERIOD COVERED Final	
7. AUTHOR(s) Jerry D. Jarrell	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Science Applications, Inc. Monterey, CA 93940	8. CONTRACT OR GRANT NUMBER(s) N00228-81-C-H361	
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Air Systems Command Department of the Navy Washington, DC 20361	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS PE 63207N PN 7W0513 TA CCOO NEPRF WU 3-14	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Naval Environmental Prediction Research Facility Monterey, CA 93943	12. REPORT DATE October 1983	
	13. NUMBER OF PAGES 20	
	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Tropical cyclone Strike probability Wind probability Southern Hemisphere		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) -> The development of a model to estimate 30 and 50 kt wind probabilities from Southern Hemisphere tropical cyclone forecasts is described. The model is based on position forecast errors, which are used to determine the probability of a tropical cyclone occupying a particular geographical position, and on wind profile errors. Wind profile errors consist of errors in the forecast maximum wind and errors in the forecast radius of 30 kt and 50 kt winds. The profile errors are used to estimate the probability of 30 and 50 kt winds occurring at (continued on reverse) <		

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

Block 20, Abstract, continued.

→ a point, given that the tropical cyclone occupies a particular position. These position and wind probability elements are combined by using an assumption of independence which was supported by correlation coefficients in an earlier work.

The model, which includes features of earlier strike and wind probability models, is tested on independent data. Test results show good agreement between forecast probability and the frequency of 30 and 50 kt winds.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

CONTENTS

1.0 Introduction 1

2.0 Model Description 5

 2.1 The Southern Ocean Model 5

3.0 Testing the Southern Hemisphere Wind
 Probability Program (WINDPSH) 7

4.0 Operational Products 11

5.0 Summary 15

Distribution 17

on For	
CR&I	<input checked="" type="checkbox"/>
TAB	<input type="checkbox"/>
Announced	<input type="checkbox"/>
Classification	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



TROPICAL CYCLONE WIND PROBABILITY FORECASTING
FOR THE SOUTHERN HEMISPHERE (WINDPSH)

1.0 Introduction

The concepts and development of tropical cyclone wind probability forecasting for the Southern Hemisphere follow closely those originally presented by Jarrell¹ (1981) for the western North Pacific. This is the fifth in a series of wind probability forecasting program reports covering different ocean basin areas. The initial report on the western North Pacific was followed by similar reports on the western North Atlantic, eastern North Pacific and North Indian Ocean. This and the previous reports are based upon an extension of the concepts developed for tropical cyclone strike probability forecasting (Jarrell², 1978). As these concepts have been previously presented and the programs are presently operational, a redevelopment of the concepts will not be presented in this report. A brief summary will be presented as a background for the new reader.

Differences between the development of Southern Hemisphere tropical cyclone wind probabilities and that of its closest previous counterpart, the model for the Atlantic Ocean is the crux of this report and these will be described in detail.

¹Jarrell, J.D., 1981: Tropical Cyclone Wind Probability Forecasting (WINDP), NAVENVPREDRSCHFAC Contractor Report CR 81-03.

²Jarrell, J.D., 1978: Tropical Cyclone Strike Probability Forecasting, NAVENVPREDRSCHFAC Contractor Report CR 78-01.

The tropical cyclone wind probability forecasting model generates estimates of the probability of 30 and 50 knot winds occurring at a point. The model includes many features of the strike probability model, which is based on an analysis of position forecast errors to determine the probability of a tropical cyclone occupying a particular geographic position. Wind profile errors based on the forecast of maximum wind and the forecast radius of 30 and 50 knot winds are similarly analyzed to determine the 30 and 50 knot probabilities of occurrence.

Jarrell² (1978), in the development of the strike probability model, based the theory of strike probability on three assumptions:

- 1) All tropical cyclone forecasts are subject to error;
- 2) Difficulty of the forecast and size of the forecast error are statistically related; and
- 3) The occurrence of errors is random and approximates a multimodal bivariate normal probability distribution.

Studies by independent investigators in three ocean basins frequented by tropical cyclones verified these assumptions. Nicklin³ (1977), Thompson et al⁴ (1981), and

³Nicklin, D.S., 1977: A Statistical Analysis of Western Pacific Tropical Cyclone Forecast Errors, Naval Postgraduate School, M.S. Thesis, June.

⁴Thompson, W.J., R.L. Elsberry and R.G. Read, 1981: An Analysis of Eastern North Pacific Tropical Cyclone Forecast Errors, Monthly Weather Review, V. 109, pp. 1930-1938.

Crutcher et al⁵ (1982) using data from the western Pacific, eastern Pacific and the Atlantic, respectively, also developed individual methods to group forecasts into three classes of forecast difficulty. A general relative classification evolved in each study to yield forecast error groups of below average, average, and above average errors. These groups were also referred to as Class I (easy forecasts), Class II (average forecasts) and Class III (difficult forecasts) in previous reports. Each investigator also utilized sufficient statistical data and prescribed sufficient parameters to describe the bivariate normal distributions for each of the three classes and for forecasts of 24, 48, and 72 hours.

Following the work of Crutcher et al⁵, southern hemisphere forecast errors were separated into groups using the NORMIX clustering model. The data set consisted of 789 forecasts issued by the Joint Typhoon Warning Center (JTWC) during the years 1976-1981. Unlike the other basins, only two significantly different clusters were identified. These two clusters are shown by 50% probability ellipses in Figure 1.

⁵Crutcher, H.L., C.J. Neumann and J.M. Pelissier, 1982: Tropical Storm Forecast Errors and the Multimodal Bivariate Normal Distribution, J. Appl. Meteor., V. 21, pp. 978-987.




Figure 1. Fifty percent error probability ellipses for two clusters of 24-hr forecast errors identified by the NORMIX clustering model in the Southern Hemisphere. Distances are actual position - forecast position in nautical miles (n mi). The small and large ellipses represent forecast classes referred to as Groups 1 and 2 respectively.

2.0 Model Description

2.1 The Southern Ocean Model

One version of the tropical cyclone wind probability model was developed for the Atlantic Ocean as a modification to the western Pacific wind probability program. The concepts of the strike portion of the Southern Ocean model are similar to those developed for the Atlantic. The Southern Ocean model uses a regression equation to estimate the probability of a forecast belonging to either of two forecast classes (associated with two forecast error clusters) and the Tsui, et al, model⁶ (used in the Atlantic wind probability model) to determine wind profiles and to handle asymmetrical wind distribution. Thus this model uses modified versions of both the wind probability and strike probability concepts used in the Atlantic.

The wind profiles and derived tropical cyclone asymmetry used in the Atlantic model were actually based on western Pacific data using methods produced by Tsui, et al, as indicated above. Tsui, et al, using wind radius data from tropical cyclone warnings over a 12-year period (1966 to 1977), determined that the profile of the tangential wind speed along the radial axis was exponential as a good approximation. He further determined that the size of the storm could be statistically related to the maximum wind and that the asymmetric shape of the storm's wind pattern was correlated to the forward speed of movement of the storm.

⁶Tsui, T.L., L.R. Brody and S. Brand, 1982: A Technique for Predicting Surface Wind Distributions of Tropical Cyclones in the Western North Pacific, NAVENVPREDRSCHFAC Technical Report TR 82-05.

A simple empirical relationship was derived to provide an estimation of wind at any radius.

$$V/V_{\max} = \exp (-0.693R),$$

where V is the wind speed of interest, V_{\max} is the maximum wind speed of the storm, and R is the ratio of the radius associated with V to the radius associated with one-half of V_{\max} (r_{half}). Radii are measured outward from the radius of maximum winds. Asymmetrical storm configuration is accommodated in the profile by an empirical adjustment of the one-half radius, r_{half} , dependent on bearing relative to direction of motion and translation speed. With due regard to the sparsity of data available in the southern hemisphere, 30 and 50 kt wind radii forecasts were treated as being without skill. Accordingly rather than predicting r_{half} as a function of these radii, a relationship based on Atlantic data was used relating r_{half} to maximum wind. While the Tsui model is considered entirely appropriate for this area, the Atlantic relationship may be somewhat suspect since Atlantic storms occur at more poleward latitudes and, all other things being equal, may be slightly larger than Southern Hemisphere counterparts. If true this would cause a slight over forecast of wind probabilities.

3.0 Testing the Southern Hemisphere Wind Probability Program (WINDPSH)

The methodology used in comparing WINDPSH predicted values against observed values is identical to that used in the North Indian Ocean. An array of 30 points in the Southern Hemisphere were selected (Figure 2). WINDPSH values for 30 and 50 knots were calculated at 12 hour intervals from the effective synoptic time of the Joint Typhoon Warning Center (JTWC) forecasts for the twenty 1982 season cyclones. Since most of these 30 points are not observing stations, actual verifying winds were not generally available. Consequently a verifying "warning time" probability of 50 or 30 winds greater than 50% constituted a verifying observation over 50 or 30 kt winds.

Tables 1 and 2 compare the expected to the observed occurrences of 30 and 50 knot winds. The test cases were 217 forecasts (12 hr intervals) on twenty 1982 cyclones. Of the 217 forecasts there were 129 and 88 verifiable at 24 and 48 hours respectively. Predictions are associated with percentage groups of increasing width, <1/2%, 1/2 to 1 1/2%, ..., etc. Time integrated probabilities were verified only if a continuous record was available over the entire time period. Significance of the differences between the expected and the observed, as discussed in previous reports (Jarrell⁷, 1981) is difficult to assess, but using a "t" test, agreement is excellent. The difference between the expected and observed occurrences was significant (5%) in only two of 72 comparisons and, although the sample is small, the overall results are considered statistically very sound.

⁷Jarrell, J.D., 1981: Atlantic Strike Probability Program, NAVENVPREDRSCHFAC Contractor Report CR 81-04.




Figure 2. Selected test points (30) in the Southern Hemisphere.

Table 1. Comparisons of expected occurrence (E) to observed occurrences (O) of 50 kt (a) and 30 kt (b) winds in a test sample of 3870 verifiable instantaneous wind probability computations.

a) 50 kt instantaneous wind probabilities

A<P<B	24 HR			48 HR		
	N	E	O	N	E	O
< 1/2*	3768	0	0	2446	0	0
1/2 - 1 1/2	36	0	0	96	1	0
1 1/2 - 3 1/2	33	1	0	47	1	1
3 1/2 - 7 1/2	18	1	1	48	2	3
7 1/2 - 15 1/2	9	1	1	3	0	0
15 1/2 - 31 1/2	6	1	2	0	0	0
31 1/2 - 63 1/2	0	0	0	0	0	0
> 63 1/2	0	0	0	0	0	0
ALL	3870	4	4	2640	4	4

b) 30 kt instantaneous wind probabilities

A<P<B	24 HR			48 HR		
	N	E	O	N	E	O
< 1/2*	3636	0	0	2329	0	0
1/2 - 1 1/2	69	1	0	93	1	0
1 1/2 - 3 1/2	37	1	1	64	2	1
3 1/2 - 7 1/2	37	2	2	63	3	1
7 1/2 - 15 1/2	37	4	3	54	6	8
15 1/2 - 31 1/2	30	7	6	37	7	11
31 1/2 - 63 1/2	24	13	11	0	0	0
> 63 1/2	0	0	0	0	0	0
ALL	3870	28	23	2640	19	21

Table 2. Comparisons of expected occurrence (E) to observed occurrences (O) of 50 kt (a) and 30 kt (b) winds in a test sample of 3840 time integrated wind probabilities.

a) 50 kt instantaneous wind probabilities

A<P<B	24 HR			48 HR		
	N	E	O	N	E	O
< 1/2%	3691	0	0	2319	0	0
1/2 - 1 1/2	48	0	0	73	1	0
1 1/2 - 3 1/2	22	1	0	61	1	0
3 1/2 - 7 1/2	30	2	0	41	2	0
7 1/2 - 15 1/2	20	2	1	43	5	1
15 1/2 - 31 1/2	22	5	3	35	7	5
31 1/2 - 63 1/2	6	3	3	7	3	3
> 63 1/2	1	1	1	1	1	1
ALL	3840	14	8	2580	20	10

b) 30 kt instantaneous wind probabilities

A<P<B	24 HR			48 HR		
	N	E	O	N	E	O
< 1/2%	3523	0	0	2204	0	0
1/2 - 1 1/2	71	1	0	71	1	0
1 1/2 - 3 1/2	57	1	0	65	2	0
3 1/2 - 7 1/2	37	2	1	61	3	1
7 1/2 - 15 1/2	42	5	2	66	7	2
15 1/2 - 31 1/2	56	12	8	65	14	11
31 1/2 - 63 1/2	33	15	16	31	14	17
> 63 1/2	21	17	16	17	13	13
ALL	3840	53	43	2580	54	44

4.0 Operational Products

The Southern Ocean wind probability program will be available for nine preselected points.* Probabilities will be given in two modes, instantaneous and time integrated, and at 0, 12, 24, 36 and 48 hours after the warning time. The instantaneous probability will be the probability at the stated time (i.e., 12 hr) and the time integrated probability will be summed from time 0 to the stated time for an estimate of the probability that the event will be observed within that period of time.

The greatest source of error for the WINDPSH program will be erroneous input data. An internal check for unusual motion (expected to occur only 5% of the time in nature) will be made and suspect motion flagged. The user should then recheck input data for accuracy.

When the forecast track approaches a land mass, the forecaster should be aware of program bias. This should be minor for seaward approach to low coastal areas or over smaller islands. However, in other cases, land influences will appear as rapid decreases in the instantaneous wind probabilities (especially 50 kt winds) near forecast landfall time. This will bias probabilities - overstate them for inland sites and understate them for coastal sites. Time integrated probabilities will be less biased. This problem is caused by wind forecasts being influenced by track forecasts where landfall is concerned. A bad track forecast

*Locations: Brisbane, Sydney, Auckland, Darwin, Perth, H.E. Holt, Learmonth, Fremantle and Diego Garcia.

may cause a bad wind forecast. This was not accounted for either in development or testing; hence the test results simulate expected actual operational results and some of the minor disparities between expected and observed occurrences no doubt stem from this.

Figure 3 depicts a standard output of WINDPSH for Cyclone Arthur, 14 January 1981. Probabilities are provided at 12 hour intervals for strike, 50 and 30 kt winds in both instantaneous and time integrated modes. The format is HHPIPS where HH is hours after forecast effective time, PI and PS are the instantaneous and time integrated probability rounded to the nearest whole percent. The letters "IN" mean less than one half percent while "THREAT NIL" means all probabilities were less than one half percent.

The other message that WINDPSH will generate is created for the Office of Foreign Disaster Assistance (OFDA), Agency for International Development. This message will be triggered when any of approximately 130 cities worldwide is threatened by a tropical cyclone. Of these cities 26 are in the Southern Hemisphere.

The message will state the storm's name and intensity category, which ocean the storm is in, the latitude and longitude of the storm, the center wind speed in knots, and estimates of the threat of hurricane force winds. The estimates are:

```

WINDP1      MESCON8090

STRIKE AND WIND PROBABILITY FORECAST

ARTHUR      141900Z

BRISBANE    THREAT NIL

SYDNEY      THREAT NIL

AUCKLAND    00ININ 12ININ 24ININ 36ININ 48ININ
           50 KNOT 00ININ 12ININ 24ININ 36ININ 48ININ
           30 KNOT 00ININ 12ININ 24ININ 36ININ 480102

DARWIN      THREAT NIL

PERTH       THREAT NIL

H E HOLT    THREAT NIL

LEARMONTH   THREAT NIL

DIEGO GARC  THREAT NIL

FOR METEOROLOGISTS:  FORECAST CONFIDENCE TABLE

           TIME  PROB  DIST  PROB  DIST  PROB
           12HR  22   50   20   75   58
           24HR  28  100   23  150   49
           48HR  26  200   22  300   52

PROBABILITIES BASED ON FOLLOWING FORECAST
001951787055 122131798045 242281785045 48260170040

```

Figure 3. Standard WINDPSH output, listing probabilities of a strike and 30 and 50 kt winds at Auckland for Cyclone Arthur, 14 January 1981. Forecast confidence table gives the probability of actual verification within 50 n mi, 50-75 n mi and outside 75 n mi for the 12 hour forecast and comparable values for radii twice as large for 24 hours and four times as large for 48 hours.

Danger (Red) if the probability of 65 kt winds or greater (P65) exceeds 20% within 18 hours, otherwise

Alert (Orange) if P65 exceeds 10% within 24 hours, otherwise

Caution (Yellow) if P65 exceeds 5% within 48 hours, otherwise

Notice (Green) if P65 exceeds 2 1/2% within 48 hours.

Figure 4 shows an example of the OFDA message.

A. FOR OFFICIAL USE ONLY

B. TROP STORM ARTHUR JAN 14, 1981 1900 GMT LOCATED IN S. PACIFIC OCEAN AT 19.5S 178.7E WITH CENTER WINDS OF 55 KTS. ESTIMATES OF THREAT OF HURRICANE FORCE WINDS FOLLOW: (REFER TO USERS MANUAL NAVENVPREDRSCHFAC DOCUMENT UM-06 8/82)

OFDA SELECTED POINTS	THREAT LEVEL
NANDI FIGI	CAUTION (YELLOW)
SUVA FIGI	ALERT (ORANGE)
NUKUALOFA TONGA IS.	NOTICE (GREEN)

C. THREAT TO ALL OTHER OFDA POINTS BELOW NOTICE (GREEN) THRESHOLD FROM THIS STORM

D. RECOMMEND READDRESS AS FOLLOWS:

 NIACT:
 SUVA

 IMMEDIATE:
 NONE

(Note: NIACT is a State Department Message handling Precedence .. Night Action)

Figure 4. OFDA message of threat estimates for cities near the path of Cyclone Arthur, 14 January 1981.

5.0 Summary

The wind probability model for the southern oceans is largely based on the wind probability program for the North Atlantic. Some features of wind probability programs for the eastern and western North Pacific and the North Indian oceans have also been incorporated. All of these programs are now operational. Test results for the Southern Hemisphere wind probability program (WINDPSH) demonstrated excellent agreement between expected and observed results. Operational products are available both in message form as described herein and through the Naval Environmental Display Station (NEDS) as plotted charts.

DISTRIBUTION

COMMANDER IN CHIEF
U.S. PACIFIC FLEET, CODE O2M
PEARL HARBOR, HI 96860

COMTHIRDFLT
CODE N33
PEARL HARBOR, HI 96860

COMSEVENTHFLT
ATTN: FLT METEOROLOGIST, N30W
FPO SAN FRANCISCO 96601

COMSEVENTHFLT
ATTN: NSAP SCIENCE ADVISOR
BOX 167
FPO SEATTLE 98762

COMMANDING OFFICER
USS AMERICA (CV-66)
ATTN: MET. OFFICER, OA DIV.
FPO NEW YORK 09531

COMMANDING OFFICER
USS FORRESTAL (CV-59)
ATTN: MET. OFFICER, OA DIV.
FPO MIAMI 34080

COMMANDING OFFICER
USS INDEPENDENCE (CV-62)
ATTN: MET. OFFICER, OA DIV.
FPO NEW YORK 09537

COMMANDING OFFICER
USS J. F. KENNEDY (CV-67)
ATTN: MET. OFFICER, OA DIV.
FPO NEW YORK 09538

COMMANDING OFFICER
USS NIMITZ (CVN-68)
ATTN: MET. OFFICER, OA DIV.
FPO NEW YORK 09542

COMMANDING OFFICER
USS D. D. EISENHOWER (CVN-69)
ATTN: MET. OFFICER, OA DIV.
FPO NEW YORK 09532

COMMANDING OFFICER
USS SARATOGA (CV-60)
ATTN: MET. OFFICER, OA DIV.
FPO MIAMI 34078

COMMANDING OFFICER
USS CARL VINSON (CVN-70)
ATTN: MET. OFFICER, OA DIV.
FPO NEW YORK 09558

COMMANDING OFFICER
USS CONSTELLATION (CV-64)
ATTN: MET. OFFICER, OA DIV.
FPO SAN FRANCISCO 96635

COMMANDING OFFICER
USS CORAL SEA (CV-43)
ATTN: MET. OFFICER, OA DIV.
FPO NEW YORK 09550

COMMANDING OFFICER
USS ENTERPRISE (CVN-65)
ATTN: MET. OFFICER, OA DIV.
FPO SAN FRANCISCO 96636

COMMANDING OFFICER
USS KITTY HAWK (CV-63)
ATTN: MET. OFFICER, OA DIV.
FPO SAN FRANCISCO 96634

COMMANDING OFFICER
USS MIDWAY (CV-41)
ATTN: MET. OFFICER, OA DIV.
FPO SAN FRANCISCO 96631

COMMANDING OFFICER
USS RANGER (CV-61)
ATTN: MET. OFFICER, OA DIV.
FPO SAN FRANCISCO 96633

COMMANDING OFFICER
USS NEW JERSEY (BB-62)
FPO SAN FRANCISCO 96688

PCO, IOWA (BB-61)
SUPERVISOR OF SHIPBUILDING
CONVERSION & REPAIR, USN
PASCAGOULA, MS 39567

COMMANDING OFFICER
USS MOUNT WHITNEY (LCC-20)
ATTN: MET. OFFICER
FPO NEW YORK 09517

COMMANDING OFFICER
USS BLUERIDGE (LCC-19)
ATTN: MET. OFFICER
FPO SAN FRANCISCO 96628

COMMANDING OFFICER
USS GUADALCANAL (LPH-7)
ATTN: MET. OFFICER
FPO NEW YORK 09562

COMMANDING OFFICER
USS GUAM (LPH-9)
ATTN: MET. OFFICER
FPO NEW YORK 09563

COMMANDING OFFICER
USS INCHON (LPH-12)
ATTN: MET. OFFICER
FPO NEW YORK 09529

COMMANDING OFFICER
USS IWO JIMA (LPH-2)
ATTN: MET. OFFICER
FPO NEW YORK 09561

COMMANDING OFFICER
USS NASSAU (LHA-4)
ATTN: MET. OFFICER
FPO NEW YORK 09557

COMMANDING OFFICER
USS SAIPAN (LHA-2)
ATTN: MET. OFFICER
FPO NEW YORK 09549

COMMANDING OFFICER
USS NEW ORLEANS (LPH-11)
ATTN: MET. OFFICER
FPO SAN FRANCISCO 96627

COMMANDING OFFICER
USS OKINAWA (LPH-3)
ATTN: MET. OFFICER
FPO SAN FRANCISCO 96625

COMMANDING OFFICER
USS TRIPOLI (LPH-10)
ATTN: METEOROLOGICAL OFFICER
FPO SAN FRANCISCO 96626

COMMANDING OFFICER
USS TARAWA (LHA-1)
FPO SAN FRANCISCO 96622

COMMANDING OFFICER
USS BELLEAU WOOD (LHA-3)
ATTN: METEOROLOGICAL OFFICER
FPO SAN FRANCISCO 96623

COMMANDING OFFICER
USS PELELIU (LHA-5)
FPO SAN FRANCISCO 96624

COMMANDING OFFICER
USS LASALLE (AGF-3)
ATTN: METEOROLOGICAL OFFICER
FPO NEW YORK 09577

CINCPAC
BOX 13
STAFF CINCPAC J37
CAMP SMITH, HI 96861

SPECIAL ASST. TO THE ASST.
SECRETARY OF THE NAVY (R&D)
ROOM 4E741, THE PENTAGON
WASHINGTON, DC 20350

OFFICE OF NAVAL RESEARCH
CODE 422AT
ARLINGTON, VA 22217

OFFICE OF NAVAL RESEARCH
COASTAL SCIENCES PROGRAM
CODE 422 CS
ARLINGTON, VA 22217

CHIEF OF NAVAL OPERATIONS
(OP-952)
U.S. NAVAL OBSERVATORY
WASHINGTON, DC 20390

CHIEF, ENV. SVCS. DIV.
OJCS (J-33)
RM. 2877K, THE PENTAGON
WASHINGTON, DC 20301

NAVAL DEPUTY TO THE
ADMINISTRATOR, NOAA
ROOM 200, PAGE BLDG. #1
3300 WHITEHAVEN ST. NW
WASHINGTON, DC 20235

OFFICER IN CHARGE
NAVOCEANCOMDET
BOX 81
U.S. NAVAL AIR STATION
FPO SAN FRANCISCO 96637

OFFICER IN CHARGE
NAVOCEANCOMDET
U.S. NAVAL AIR FACILITY
FPO SEATTLE 98767

OFFICER IN CHARGE
NAVOCEANCOMDET
MONTEREY, CA 93940

OFFICER IN CHARGE
U.S. NAVOCEANCOMDET
APO SAN FRANCISCO 96519

OFFICER IN CHARGE
U.S. NAVOCEANCOMDET
FPO SAN FRANCISCO 96685

COMMANDING OFFICER
FLEET INTELLIGENCE CENTER
(PACIFIC)
PEARL HARBOR, HI 96860

COMMANDER
NAVAL OCEANOGRAPHY COMMAND
NSTL, MS 39529

COMMANDING OFFICER
FLENUMOCEANCEN
MONTEREY, CA 93940

COMMANDING OFFICER
NAVWESTOCEANCEN
BOX 113
PEARL HARBOR, HI 96860

COMMANDING OFFICER
NAVEASTOCEANCEN
MCADIE BLDG. (U-117)
NAVAL AIR STATION
NORFOLK, VA 23511

COMMANDING OFFICER
U.S. NAVOCEANCOMCEN
BOX 12, COMNAV Marianas
FPO SAN FRANCISCO 96630

COMMANDING OFFICER
U.S. NAVOCEANCOMFAC
FPO SEATTLE 98762

COMMANDING OFFICER
U.S. NAVOCEANCOMFAC
BOX 63, NAS (CUBI PT)
FPO SAN FRANCISCO 96654

PRESIDENT
NAVAL WAR COLLEGE
ATTN: GEOPHYSICS OFFICER
NEWPORT, RI 02840

COMMANDER (2)
NAVAIRSYSCOM
ATTN: LIBRARY (AIR-00D4)
WASHINGTON, DC 20361

COMMANDER
NAVAIRSYSCOM (AIR-330)
WASHINGTON, DC 20361

COMMANDER
AWS/DN
SCOTT AFB, IL 62225

USAFETAC/TS
SCOTT AFB, IL 62225

DIRECTOR (12)
DEFENSE TECH. INFORMATION
CENTER, CAMERON STATION
ALEXANDRIA, VA 22314

DIRECTOR
OFFICE OF ENV. & LIFE SCI.
OFFICE OF THE UNDERSEC OF
DEFENSE FOR RSCH & ENG, E&LS
RM 3D129, THE PENTAGON
WASHINGTON, DC 20505

FEDERAL COORD. FOR METEORO.
SERVS. & SUP. RSCH. (OFCM)
11426 ROCKVILLE PIKE
SUITE 300
ROCKVILLE, MD 20852
ROCKVILLE, MD 20852

DIRECTOR
NATIONAL HURRICANE CENTER
NOAA, GABLES ONE TOWER
1320 S. DIXIE HWY
CORAL GABLES, FL 33146

DIRECTOR
INTERNATIONAL AFFAIRS OFFICE
NOAA (OAX4)
6010 EXECUTIVE BLVD.
ROCKVILLE, MD 20852

DIRECTOR (AOML)
NATIONAL HURRICANE RSCH. LAB.
1320 S. DIXIE HWY.
CORAL GABLES, FL 33146

THE EXECUTIVE DIRECTOR
AMERICAN METEORO. SOCIETY
45 BEACON ST.
BOSTON, MA 02108

AMERICAN METEORO. SOCIETY
METEOR. & GEOASTRO. ABSTRACTS
P.O. BOX 1736
WASHINGTON, DC 20013

MR. W. G. SCHRAMM/WWW
WORLD METEOROLOGICAL
ORGANIZATION
CASE POSTALE #5, CH-1211
GENEVA, SWITZERLAND

DIRECTOR, JTWC
BOX 17
FPO SAN FRANCISCO 96630

BUREAU OF METEOROLOGY
BOX 1289K, GPO
MELBOURNE, VIC, 3001
AUSTRALIA

LIBRARIAN, NEW ZEALAND
METEOROLOGICAL SERVICE
P.O. BOX 722
WELLINGTON, NEW ZEALAND

NAVAL POSTGRADUATE SCHOOL
METEOROLOGY DEPT., CODE 63
MONTEREY, CA 93943

**ATE
LME**