

AD-A139 415

SCALING OF THE BEAM PLASMA DISCHARGE FOR LOW MAGNETIC
FIELDS(U) MARYLAND UNIV COLLEGE PARK DEPT OF PHYSICS
AND ASTRONOMY K PAPADOPOULOS JAN 84 AP-84-026

1/1

UNCLASSIFIED

N00014-79-C-0665

F/G 12/1

NL


END

DATE

FILED

4-84

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963 A

AD A139415

12



DTIC
ELECTE
MAR 27 1984
S B

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

UNIVERSITY OF MARYLAND
DEPARTMENT OF PHYSICS AND ASTRONOMY
COLLEGE PARK, MARYLAND

DTIC FILE COPY

84 03 20 008

Scaling of the Beam Plasma
Discharge for Low Magnetic Fields

by
K. Papadopoulos

DTIC
ELECTE
S MAR 27 1984 D
B

DISTRIBUTION STATEMENT A

Approved for public release
Distribution Unlimited

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER University of Maryland Report AP84-026	2. GOVT ACCESSION NO. AD-A139415	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Scaling of the Beam Plasma Discharge for Low Magnetic Fields		5. TYPE OF REPORT & PERIOD COVERED Interim report on a continuing problem
7. AUTHOR(s) K. Papadopoulos		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Maryland Department of Physics and Astronomy College Park, Maryland 20742		8. CONTRACT OR GRANT NUMBER(s) N00014-79C-0665
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217 NASA Washington, DC		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE JANUARY 1984
		13. NUMBER OF PAGES 24
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES To be published in Journal of Geophysical Research		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Beam plasma, Discharge, Beam plasma instability, Electron beam in space <i>omega sub 2 cap to omega sub 2</i>		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A theoretical analysis of the scaling law and the value of the threshold current for beam plasma discharge (BPD) is presented, based on the requirement for an absolute instability near the plasma frequency. It is shown that both the scaling law as well as the numerical values of ω_c are consistent with the experimental data, both in the low and high pressure regimes for weak magnetic field experiments ($\omega_c \ll \omega_p$). The differences in scaling between the regimes are attributed to transition from Bohm to classical diffusion. It is found that the value of the pressure minimum for ignition increases linearly with the ambient magnetic field.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 68 IS OBSOLETE

S/N 0102-LF-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

AP 84
026

Scaling of the Beam Plasma
Discharge for Low Magnetic Fields

by

K Papadopoulos

University of Maryland
Department of Physics and Astronomy
College Park, Maryland 20742



Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By PER CALL JC	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Abstract

A theoretical analysis of the scaling law and the value of the threshold current for beam plasma discharge (BPD) is presented, based on the requirement for an absolute instability near the plasma frequency. It is shown that both the scaling law as well as the numerical values of I_c are consistent with the experimental data, both in the low and high pressure regimes for weak magnetic field experiments ($\omega_e > \Omega_e$). The differences in scaling between the two regimes are attributed to transition from Bohm to classical diffusion. It is found that the value of the pressure minimum for ignition increases linearly with the ambient magnetic field.

I. Introduction

Laboratory studies of energetic electron beam injection experiments in a neutral gas filled vacuum chamber carried out in the large vacuum facility at the Johnson Space Center (JSC) (Bernstein et al. 1978, 1979, 1980) have provided important informations for the interpretation of data from space based electron beam injection. Perhaps the most important aspect was the determination of an empirical relationship of the form

$$I_c \sim \frac{E_b^{3/2}}{B^\lambda L} f(p)$$

for the critical current I_c required for beam plasma discharge ignition and the values of beam energy (E_b), ambient magnetic field (B), system length L , and ambient pressure p . The value of $\lambda \approx .5-1$ and the pressure function $f(p)$ was a function with a minimum at $p_0 \approx 20 \mu T$ and varying roughly as $p^{\pm .5}$ to $p^{\pm 1}$ above and below this pressure (Kellogg et al. 1981). The minimum value of $I_c \approx 10$ mA for a system with $L \approx 20$ m, $E_b \approx 1$ keV and $B \approx 1$ G, typical of the experiment. The operating pressure range was between 1-50 μT , with most measurements on the 1-20 μT range. Similar results have been reproduced in a number of other experiments (Konrad et al. 1982, Lyakhov et al. 1982, Bernstein et al. 1983) in which the collisional ionization by the beam is sufficient to bring the plasma density of the system to the point that $\omega_e > \Omega_e$, where ω_e , Ω_e are the plasma and cyclotron frequency. The observed p , B scaling is shown graphically in Fig. 1.

Triggering of BPD has been long associated with a beam plasma instability between the electron beam and the beam generated plasma

(Kharchenko et al. 1962, Getty and Smullin 1963, Galeev et al. 1976, Linson and Papadopoulos 1980, Papadopoulos 1981, Galeev 1983). For $\omega_e > \Omega_e$ and finite size systems Rowland et al. (1981) and Papadopoulos (1982), have associated the triggering of the BPD with the threshold for an absolute beam plasma instability near ω_e . As explained in Papadopoulos (1981) for systems such as the JSC tank the system length ($L \lesssim 20$ m) is not long enough to allow convective modes to grow to sufficient amplitude. The requirement that the waves grow at frequencies near ω_e , is connected with the fast rate of non-linear energy transfer of the beam energy to ionizing suprathermal electron tails (Papadopoulos and Coffey 1974, Papadopoulos 1975, Papadopoulos and Rowland 1978, Rowland et al. 1980, Galeev 1983). It is the purpose of this paper to develop a model for BPD ignition and the expected scaling laws on the basis of the criterion for an absolute instability near ω_e . Notice that for systems with $\omega_e < \Omega_e$ the ω_e waves are in the lower hybrid branch (Manickam et al. 1975) which in the absence of internal wave reflections gives always convective amplification. Our analysis therefore applies only to situations where $\omega_e > \Omega_e$.

The plan of the paper is as follows. We discuss next the beam plasma equilibrium expected on the basis of collisional ionization. Section III presents the instability theory for the configuration determined in section II and derives the threshold criteria. Section IV presents a comparison of the model to the BPD ignition values determined in the JSC experiment. The final section summarizes the findings and discusses their applicability to other situations. The numerical values are given in MKS units except for the beam energy (keV), pressure (μ T), magnetic field (G), and density (cm^{-3}).

II. Pre-BPD Density Buildup

Before entering the instability analysis it is necessary to establish the equilibrium density profiles for the plasma during the collisional ionization stage. We assume that the beam density profile is given by

$$n_b(r) = n_b e^{-r^2/a^2} = \frac{I_b}{e v_b \pi a^2} e^{-r^2/a^2} \quad (1)$$

where I_b and v_b are the electron beam current and velocity parallel to the magnetic field and a the beam radius given by

$$a = \frac{v_b \sin \theta_d}{\Omega_e} \quad (1a)$$

where θ_d is the equivalent divergence injection angle (Linson and Papadopoulos 1980). The equation for the ionization at midplane (i.e. ignoring the z dependence) is

$$\frac{\partial}{\partial t} n(r) - \frac{1}{r} \frac{\partial}{\partial r} r D \frac{\partial}{\partial r} n(r) + \frac{\alpha n(r)}{L} = \frac{I_b N_o \sigma}{e \pi a^2} e^{-r^2/a^2} \quad (2)$$

where D is the diffusion coefficient, L the system length, N_o the ambient neutral density, and σ the ionization cross section. The term $\frac{\alpha}{L} n(r)$ describes the axial losses. Notice that if we average (2) over the volume we recover the zero dimensional description, in terms of the confinement time τ (Papadopoulos 1982), i.e.

$$\frac{d}{dt} n = \frac{I_b}{e \pi a^2} N_o \sigma - \frac{n}{\tau} \quad (3)$$

which gives the steady state value of the density as

$$n = \frac{I_b}{e\pi a^2} N_o \sigma \tau \quad (4)$$

The general solution of eq. (2) in terms of the first and second order Bessel functions I_o , K_o is

$$n(r) = A_1 I_o \left(\frac{r}{b} \right) + A_2 K_o \left(\frac{r}{b} \right) \quad (5)$$

with

$$b^2 = \frac{LD}{\alpha^2} \quad (6)$$

The constants A_1 , A_2 to be found from the boundary conditions. In the thin beam limit ($a \ll b$), in which we have a line source we find

$$n(r) = \frac{I_b N_o \sigma}{2\pi e D} K_o \left(\frac{r}{b} \right) \equiv n_o K_o \left(\frac{r}{b} \right) \quad (7)$$

The more general solution gives, for regions inside the source ($0 \leq r \leq a$)

$$n(r) = \frac{I_b N_o}{e D \pi a^2} \int_0^\infty K_o \left(\frac{|r-r'|}{b} \right) r' e^{-r'^2/a^2} dr' \quad (8)$$

while outside the source ($a \leq r \leq b$)

$$n(r) = \frac{I_b N_o \sigma}{e D \pi a^2} \left(\int_0^\infty I_o \left(\frac{r'}{b} \right) e^{-r'^2/a^2} r' dr' \right) K_o \left(\frac{r}{b} \right). \quad (9)$$

Guided by the experimental results we restrict ourselves here to the thin beam limit $a < b$, which implies that $\frac{v_b \sin \theta_d}{\Omega_e} < \frac{D^{1/2} L^{1/2}}{\alpha}$. In this case the input parameters to the instability analysis are n_b , n_o , a , and b given by eqs. (1), (6) and (7). The values of n_b and n_o in the system units discussed in section I can be found from eqs. (1) and (7) as

$$n_b = 1.9 \times 10^6 \frac{I_b B^2}{E_b^{3/2}} \frac{1}{\sin^2 \theta_d} \quad (9a)$$

$$n_o = 3.2 \times 10^8 \frac{I_b P}{E_b^{1/2}} \frac{1}{D} \quad (9b)$$

III. Instability Theory

The homogeneous interaction between the beam and the plasma is described by the dispersion relation

$$\epsilon(\underline{k}, \omega) = 1 + K_p(\underline{k}, \omega) + K_b(\underline{k}, \omega) = 0 \quad (10a)$$

where K_p and K_b are the longitudinal dielectric functions of the plasma and the beam. Both K_p and K_b can be calculated for any type of distribution functions including collisional and finite size geometry effects (Briggs 1964). We choose here models that allow us to emphasize the physics and avoid the mathematical complexity. Consistently with Rowland et al. 1981 and Szuszcwicz et al. 1982 we consider only the synchronous Cerenkov interaction of a slow beam wave with an upper hybrid wave ω_o of the cold plasma. In this case

$$K_p(k, \omega) = - \frac{\omega_o^2}{\omega^2} \quad (10b)$$

$$\omega_o^2 = \frac{1}{2} (\omega_e^2 + \Omega_e^2) + [\frac{1}{4} (\omega_e^2 + \Omega_e^2)^2 - \omega_e^2 \Omega_e^2 \cos^2 \theta]^{1/2} \quad (10c)$$

$$K_B(k, \omega) = - \frac{\omega_b^2 R \cos^2 \theta}{(\omega - k_z v_b)^2} \quad (10d)$$

$$\cos^2 \theta = \frac{k_z^2}{k_z^2 + k_\perp^2} \quad (10e)$$

ω_p , ω_b and Ω_e are the plasma frequency, beam plasma frequency, cyclotron frequency respectively, and $k^2 = k_z^2 + k_\perp^2$. The potential was assumed to have the form $J_0(k_\perp r) \exp[i(k_z z - \omega t)]$ and thus the propagation is axial, with $k_\perp$ determined by the transverse geometry. The finite size beam reduction factor R enters through the boundary conditions at the radii a and b is given in terms of Bessel functions by (Manickam et al. 1975)

$$R = \frac{\pi}{2} (k_\perp a) \frac{a}{b} \frac{Y_0(k_\perp b)}{Y_1(k_\perp b)} [J_0^2(k_\perp a) + J_1^2(k_\perp a)] \quad (11a)$$

$$J_0(k_\perp b) = 0 \quad (11b)$$

The value $\omega_b R^{1/2} \cos \theta$ serves as a reduced effective beam plasma frequency. An important aspect of the dispersion (10) is the absorption of the transverse wavenumber $k_\perp$ and geometry effects into a single parameter R . Therefore from eq. (11), R is fixed when the mode number and the ratio $\frac{b}{a}$ are fixed. Fig. 7 of Manickam et al. (1975) shows the values of R as function of $\frac{b}{a}$ for the fundamental mode. For large $\frac{b}{a} \gg 1$, it has a logarithmic dependence approaching the value $R = .1$.

In order to determine the conditions for absolute instability in our system we follow the techniques developed by Bers (1972), in the weak coupling approximation. For the beam waves the dispersion relation is

$$D_b(k, \omega) = 1 - \frac{\omega_b^2 R \cos^2 \theta}{(\omega - k_z v_b)^2} = 0 \quad (12)$$

From this we find the usual fast and slow waves given by

$$\omega - k_z v = \pm \omega_b R^{1/2} \cos \theta \quad (12a)$$

The slow wave is a negative energy wave while the fast is positive, i.e.

$$W_b = \pm \frac{2\omega}{\omega_b R^{1/2} \cos \theta} \frac{1}{4} \epsilon_0 |\epsilon_b|^2 \quad (12b)$$

where W_b is the wave energy, ϵ_b the beam wave amplitude and ϵ_0 the free space dielectric constant. The negative energy wave can couple in synchronous interaction with the backward positive energy upper hybrid wave to produce an instability. Notice that for the upper hybrid wave

$$D_p(k, \omega) = 1 - \frac{\omega_o^2(k_z)}{\omega^2} \quad (13a)$$

while the wave energy is

$$W_p = \frac{2\omega}{\omega_o(k_z)} \frac{1}{4} \epsilon_0 |\epsilon_p|^2 \quad (13b)$$

We examine now the situation where the slow wave of the beam interacts resonantly with the plasma wave ω_o . We find a set of coupled equations

$$\left(\frac{\partial}{\partial t} + v_b \frac{\partial}{\partial z} + \nu_b\right) U_b = C_{bp} U_p \quad (14)$$

$$\left(\frac{\partial}{\partial t} + v_p \frac{\partial}{\partial z} + \nu_p\right) U_p = C_{pb} U_b$$

where v_b, v_p are the group velocities of the beam and plasma waves and ν_b, ν_p are phenomenological damping coefficients of the two waves. $U_{p,b}$ are the usual normalized amplitudes defined as

$$|U_{p,b}|^2 = \frac{W_{p,b}}{\omega} = \left| \frac{\partial D_{p,b}}{\partial \omega} \right| \frac{\epsilon_0}{4} |\epsilon_{p,b}|^2 \quad (15)$$

(Bers 1972, Davidson 1970, Weiland and Wilhelmson 1977) and the coupling coefficients C_{bp}, C_{pb} given by

$$C_{bp} = \frac{-\frac{1}{4} \epsilon_b^* J_{pb}}{W_b} \equiv \frac{P_{bp}}{W_b} \quad (16)$$

$$C_{pb} = \frac{-\frac{1}{4} \epsilon_p^* J_{bp}}{W_p} \equiv \frac{P_{pb}}{W_p}$$

where J_{pb} is the perturbed plasma current that interacts with the beam and J_{bp} is the perturbed beam current that interacts with the plasma. For conservative interactions

$$P_{bp} = -P_{pb}^* = -P_{pb} \quad (17)$$

From eqs. (12-16) we find the growth rate

$$\gamma = \frac{|P_{bp}|}{|W_b W_p|^{1/2}} = \frac{1}{2} \left(\frac{\omega_b^2 R \cos^2 \theta}{\omega_0^2} \right)^{1/2} \omega_0 \quad (18)$$

Absolute instability requires (Bers 1972)

$$v_b v_p < 0 \quad (19a)$$

$$\gamma^2 > v_b v_p \quad (19b)$$

$$L > \frac{(|v_b v_p|^{1/2})}{\gamma} \equiv L_c \quad (19c)$$

where L is the system size in the z -direction. The first condition enters through the requirement that the unstable pulse encompasses the origin at all times. The second from the requirement that the pulse growth exceeds the dissipation. The last is equivalent to the breakdown length L_c of an oscillator and implies that the feedback is stronger than convective losses. Notice that in the absence of wave reflecting boundaries only the upper hybrid branch can be absolutely unstable, since the lower hybrid branch corresponds to a forward wave (i.e. $v_b v_p > 0$). In a plasma with $\frac{\omega_e}{\Omega_e} < 1$, waves near the plasma frequency will be convectively unstable.

As mentioned in the introduction we associate the threshold of BPD in the Johnson chamber with an absolute instability near ω_e . For our parameters the collisional ionization generates a plasma with $\frac{\omega_e}{\Omega_e} > 2$, and the collisionality is such that condition (19b) is trivially satisfied. We therefore concentrate on eq. (19c), for a backward wave in the upper hybrid range with $\frac{\omega_e}{\Omega_e} \gg 1$. The group velocity of the waves v_b, v_p are

$$v_b = V_b \quad (20)$$

$$v_p = -2V_b \frac{\Omega_e^2}{\omega_e^2} \cos^2 \theta \sin^2 \theta$$

The value of $\sin \theta$ can be computed using (10e) and the first root of eq. (11b), i.e. $k_{\perp} b = 2.4$, giving

$$\sin^2 \theta = \frac{1}{1 + g^2} \quad (21a)$$

$$g^2 = \frac{b^2}{(2.4)^2} \frac{\omega_e^2}{V_b^2} \quad (21b)$$

From eqs. (18), (19c), (20) and (21) we find the criterion for absolute instability as

$$\omega_b^2 \geq \frac{2\sqrt{2}}{L} \left(\frac{n_b}{n_o}\right)^{1/2} \frac{\Omega_e V_b}{R^{1/2}} \sin \theta \quad (22)$$

IV. BPD Ignition Scaling

Eq. (22) is the threshold condition for an absolute instability near the plasma frequency in terms of the plasma parameters of the system. Using eq. (9) we find the current threshold condition as

$$I \geq I_c = 1.2 \times 10^{-2} \frac{E_b^{3/2}}{p^{1/2}} \frac{D^{1/2}}{L} \left(\frac{\sin \theta \sin \theta_d}{R^{1/2}} \right) \quad (23)$$

In order to make further process we have to specify the value and scaling of the diffusion coefficient D . In the low pressure regime it

was experimentally determined (Szuszczewicz et al. 1979) that the diffusion coefficient obeyed the Bohm diffusion law. Therefore

$$D \approx D_B = 6.25 \times 10^2 \left(\frac{T_e}{\text{eV}} \right) \frac{1}{B} \quad (24)$$

Using this expression in eq. (23) with $T_e \approx 2.5$ eV we find

$$I_c = .5 \frac{E_b^{3/2}}{p^{1/2}} \frac{1}{B^{1/2}} \frac{1}{L} \left(\frac{\sin \theta \sin \theta_d}{R^{1/2}} \right) A \quad (25a)$$

For values of $g \lesssim 1$, the factor in parenthesis is between 2-3, so that to within a factor of two (25a) reads

$$I_c = \frac{E_b^{3/2}}{p^{1/2}} \frac{1}{B^{1/2}} \frac{1}{L} A \quad (25b)$$

For the standard parameters ($E_b \approx 1$ keV, $B \approx 1$ G, $L = 20$ m), we find a at the transition point between the high and low density regime

($p_0 \approx 15$ μ T) the minimum current required for BPD triggering as $I_c = 11$ mA. The case analyzed in detail by Kellog et al. (1982), i.e. $p = 5$ μ T, gives $I_c = 22$ mA. Both values compare favorably with the observed values. While Bohm diffusion is independent of pressure, classical diffusion is pressure dependent, i.e.

$$D_{cl} \approx 1.8 \left(\frac{T_e}{\text{eV}} \right)^{3/2} \frac{p}{B^2} \quad (26)$$

We therefore expect that at some value $D_{cl} \geq D_B$. From eqs. (23) and (26) we find for $T_e \approx 2.5$ eV

$$I_c \approx 3 \times 10^{-2} \frac{E_b^{3/2} p^{1/2}}{B} \frac{1}{L} \left(\frac{\sin \theta \sin \theta_d}{R^{1/2}} \right) A \quad (27a)$$

or

$$I_c \approx 6 \times 10^{-2} \frac{E_b^{3/2} p^{1/2}}{B} \frac{1}{L} A \quad (27b)$$

For the inflection point (i.e. $p_0 \approx 15 \mu T$) this will give $I_c \approx 14$ mA.

Notice, however, that $I_c \sim p^{1/2}$, while $I_c \sim \frac{1}{B}$ rather than $\frac{1}{B^{1/2}}$ found for the low pressure regime.

An interesting scaling results from the above for the dependence of the minimum pressure p_0 for BPD as a function of B . Namely $p_0 \sim B$. Since $p_0 \approx 15 \mu T$ for $B \approx 1G$, we find p_0 in μT as

$$p_0 = 15 B \quad (28)$$

Namely the pressure range of the low pressure scaling increases with B . This is consistent not only with Fig. 1, but also with the small chamber results (Konradi et al. 1983, Bernstein et al. 1983) in which

$B \approx 38G$ and the low pressure scaling was consistent with eq. 27a. Let me finally note that eqs. (27a,b) are consistent with more recent experimental results (Lyakhov et al. 1982, Kawashima et al. 1982).

An approximate criterion for the ignition of BPD was given before (Rowland et al. 1981, Papadopoulos 1981) and was found consistent with the observations at the J.S.F.C. tank (Szuszczewicz et al. 1982) as $\frac{\omega_e}{\Omega_e} > 5$. It is appropriate to comment on its relationship to the present more detailed considerations. Referring to eq. (23) we note that $I_c \sim \sin \theta$ so that for $\sin \theta \ll 1$ I_c becomes very small,

independently of other considerations. From eq. (21) $\sin\theta \ll 1$ corresponds to

$$g = \frac{b}{2.4} \frac{\omega_e}{v_b} \gg 1 \quad (29)$$

which is the opposite limit from the one considered before. Taking $b \approx 0(a)$ and using eq. (1a) we recover the condition $\frac{\omega_e}{\Omega_e} > \frac{2.4}{\sin\theta_d} \approx 5$ as an approximate condition at which absolute instability develops. This criterion is extremely relevant for cases with preionized plasma such as the ionosphere at daylight conditions or high altitude (i.e. F peak). The $\frac{\omega_e}{\Omega_e} > 5$ criterion is a sufficient but not a necessary condition, and accounts in a natural fashion for the observed hysteresis during PBD extinction (Bernstein et al. 1979).

V. Summary and Conclusions

We presented a detailed physical analysis of BPD threshold scaling based on the conjecture proposed by Papadopoulos (1981), that the BPD threshold is associated with the triggering of an absolute instability near ω_e . This conjecture predicts a scaling

$$I_c \sim \frac{E_b^{3/2}}{p^{1/2} B^{1/2} L}$$

for low pressures ($p < p_0$) and

$$I_c \sim \frac{E_b^{3/2} p^{1/2}}{B L}$$

for high pressures ($p > p_0$). In deriving these scalings we assumed Bohm diffusion for $p < p_0$ and classical diffusion for $p > p_0$. The minimum threshold current at $p = p_0$ is associated with the transition from Bohm to classical diffusion. The predicted scaling, the numerical values of I_c , as well as the dominance of Bohm diffusion for $p < p_0$ are in good agreement with the data from JSC tank experiment (Bernstein et al. 1979, Szuszcwicz et al. 1979). It is important to restate some of the key assumptions of the theory and their consequences, since many were inspired from the JSC tank parameters and might not be applicable to other situations.

- (a) Weak magnetic field in the sense that at prebreakdown $\frac{\omega_e}{\Omega_e} \geq 2$. For situations where $\frac{\omega_e}{\Omega_e} < 1$, the ω_e waves lie in the lower hybrid branch of the dispersion curve which corresponds to forward waves and therefore produces convective instability. This seems to be the situation in Boswell and Kellog (1983) and the expected scaling should be derived from different considerations. The case $\Omega_e < \omega_e < 2\Omega_e$, requires special consideration due to the presence of strong cyclotron damping which was neglected here. Note also that end plate reflections can produce an absolute instability even in the lower hybrid branch.
- (b) Thin beam in the sense $a \ll b$. For situations where $a \approx b$, axial losses could be dominant and the more general eqs. (8) and (9) should be used instead of (7).
- (c) Weak coupling limit eqs. (14) are valid in the limit where the parameter $(\frac{\omega_b}{\omega_e})^{1/2} = (\frac{n_b}{n_0})^{1/2} \ll 1$. Otherwise the saddle

point method (Le Queau et al. 1980) should be used to determine the threshold condition given by eq. (22).

- (d) In associating the threshold for absolute instability with the BPD ignition we implicitly assumed that the energy deposition from the beam to the plasma will produce ambient electron fluxes whose ionization rate exceeds the beam ionization rate. This seems to be clearly satisfied in the low pressure regime. However as noted in Papadopoulos et al. (1983) by increasing the pressure there is a limit at which the energy deposition by the beam plasma instability is not sufficient to overcome line emission so that the electron energy stays below the ionization energy. This case will have many of the BPD signatures (i.e. broadening of the radiated region) but the electron plasma density will be controlled by beam ionization only (i.e. BPD without D). The small chamber experiment (Bernstein et al. 1983) is possibly indicative of such behavior.

Before closing we should comment on the applicability of the above concepts to electron beam injection in space. There are two fundamental differences between the laboratory and the ionospheric beam injection:

- (i) The existence of ambient plasma with long density gradient scales (i.e. $L \rightarrow \infty$) in the ionosphere.
- (ii) The laboratory experiments are steady state, while the vehicle motion across the magnetic field line can be thought as producing beam pulses with time length $\tau = \frac{U}{a}$ where U is the cross-field motion of the vehicle (i.e. $1-2 \frac{\text{km}}{\text{sec}}$ for rockets, $4-8 \frac{\text{km}}{\text{sec}}$ for the shuttle).

In assessing BPD for the ionospheric case we have to ask the following questions:

- (i) Is the plasma density produced by the collisional ionization due to the beam during $t \ll \tau$, larger than the ambient ionospheric density.
- (ii) Is there a beam instability at ω_e based on the ambient ionospheric plasma density.
- (iii) Is the ionization time due to the hot electrons produced by the instability shorter than the injection time τ (i.e. $\nu_{ion} \tau \gg 1$).

If the answer to question (i) is yes and to question (ii) no, the threshold condition is similar to eq. (23), with the length L given by the plasma density gradient due to collisional ionization by the beam. In view of the short injection time, however, eq. (23) is sufficient only for the beam plasma instability (BPI) but not BPD. In order to have BPD we need in addition to eq. (23) a positive answer to question (iii). The condition $\nu_{ion} \tau \gg 1$ is equivalent to the Townsend condition (Galeev et al. 1976, Papadopoulos 1981). If the answer to question (i) is no and to question (ii) is yes, the BPI criterion will be given by eq. (22) with n_0 the value of the ambient plasma density. BPD requires in addition $\nu_{ion} \tau > 1$. The above comments should be only taken as guidelines. More precise considerations require non-linear BPI computations for the evaluation of ν_{ion} as a function of the beam parameters and will be reported in the future.

Acknowledgements: This work has evolved from many years of association with the team performing the JSC experiments (Drs. H. Anderson, B. Bernstein, J. Jost, T. Hallinan, P. Kellog, H. Leinbach and E. Szuszczeicz) whose contribution to this work cannot be overemphasized. Special thanks belong to Dr. B. Bernstein for introducing me to the subject of BPD and for many heated but stimulating discussions. Last but not least I would like to thank my close collaborators in previous BPD work Dr. L. Linson for many valuable discussions and corrections. The work was supported in part by NASA and in part by ONR N00014-79-C-0665.

References

- Bernstein, W., H. Leinbach, P. Kellogg, S. Monson, T. Hallinan, O. K. Garriott, A. Konradi, J. McCoy, P. Daly, B. Baker, and H. R. Anderson, Electron beam injection experiments: The beam-plasma-discharge at low pressures and magnetic field strengths, Geophys. Res. Lett., 5, 127, 1978.
- Bernstein, W., H. Leinbach, P. J. Kellogg, S. J. Monson, and T. Hallinan, Further laboratory measurements of the beam-plasma discharge, J. Geophys. Res., 84, 7271, 1979.
- Bernstein, W., B. A. Whalen, F. R. Harris, A. G. McNamara, and A. Konradi, Laboratory studies of the charge neutralization of a rocket payload during electron beam emission, Geophys. Res. Lett., 7, 93, 1980.
- Bernstein, W., J. O. McGarity, and A. Konradi, Electron beam injection experiments: replication of flight observations in a laboratory beam plasma discharge, Geophys. Res. Lett., 10, 1124, 1983.
- Bers, A. Linear waves and instabilities, in "Plasma Physics, Les Houches 1973", p. 113, 1975.
- Briggs, R. J. Electron-stream interactive with plasma, M.I.T. Press, Cambridge, Mass, 1964.
- Davidson, R. C., Methods in non-linear plasma theory, Ch-6 Academic Press, N.Y., 1972.
- Galeev, A. A., E. V. Mishin, R. Z. Sagdeev, V. D. Shapiro, and V. I. Shevchenko, Discharge in the region around a rocket following the injection of electron beams in the ionosphere, Sov. Phys. Dokl., 21, 641, 1976.

- Galeev, A. A., R. Z. Sagdeev, V. D. Shapiro, and V. I. Shevchenko, Beam plasma discharge and suprathermal electron tails, in Active Experiments in Space ESA-SP-195, 151, 1983.
- Getty, W. D., and L. D. Smullin, Beam-plasma discharge: Buildup of oscillations, J. Appl. phys., 34, 3421, 1963.
- Kawashima, N., S. Sasaki, K. Takahashi, and T. Obayashi, BPD experiments in the laboratory, in Active Experiments in Space ESA-SP-195, 181, 1983.
- Kharchenko, I. F., Va. B. Feinberg, R. M. Nikolayev, E. A. Kornilov, E. Lutsenko, and N. S. Pedenko, The interaction of an electron beam with a plasma in a magnetic field, Sov. Phys. Tech. Phys., 6, 551, 1962.
- Konradi, A., W. Bernstein, D. L. Bulgher, J. O. Mc Garrity, and J. L. Winkler, Initial results from a laboratory size beam plasma discharge, in Active Experiments in Space ESA-SP-195, 185, 1983.
- Le Queau, D., R. Pellat, and A. Saint-Marc, An investigation of electrostatic linear instabilities of a radially limited electron beam, Annales de Géophysique 36, 423, 1980.
- Linson, L. M., and K. Papadopoulos, Review of the status of theory and experiment for injection of energetic electron beams in space, Re. PAPS-69/SAI-D23-459-LJ, Sci. Appl., Inc., LaJolla, Calif., April 1980.
- Lyakhov, S. B., A. D. Maiorov, G. G. Managadze, O. A. Povalyev, A. I. Chmil, L. Yu Kochmarev, E. G. Shustin, M. F. Friedrich, W. K. Riedler, A. N. Lalishvili, A. Kiraga, Z. Kloss, Z. Kravchic, and N. A. Leonov, Laboratory studies for beam plasma discharge in experiments in rockets, Planetary and Space Res., 30, 347, 1982.

- Manickam, J., W. Carr, B. Rosen, and M. Seidl, Convective and absolute instabilities in beam plasma systems, Phys. Fl., 18, 369, 1975.
- Papadopoulos, K., Non-linear stabilization of beam plasma instabilities by parametric effects, Phys. Fl., 18, 1769, 1975.
- Papadopoulos, K., Theory of beam plasma discharge, in Proceedings of NATO Advanced Research Institute on Artificial Particle Beams in Space Plasma Physics, (B. Grandal Ed.) Plenum, New York, p. 505, 1982.
- Papadopoulos, K., C. L. Chang, and K. Ko, Collective plasma effects on electron beam injection from rockets, SAI 83-1215, 1983.
- Papadopoulos, K., and T. Coffey, Nonthermal features of the auroral plasma due to the precipitating electrons, J. Geophys. Res., 79, 674, 1974.
- Papadopoulos, K., and H. L. Rowland, Collisionless effects on the spectrum of secondary auroral electrons at low altitudes, J. Geophys. Res., 83, 5768, 1978.
- Rowland, H.L., C. L. Change, and K. Papadopoulos, Scaling of the beam-plasma discharge, J. Geophys. Res., 86, 9215, 1981.
- Szuszcwicz, E. P., K. Papadopoulos, W. Bernstein, C. S. Liu, and D. N. Walker, Threshold criterion for a beam plasma discharge, J. Geophys. Res., 87, 1565, 1982.
- Szuszcwicz, E. P., D. N. Walker, J. C. Holmes, and H. Leinbach, Plasma diffusion in a space simulation beam plasma discharge, Geoph. Res. Lett., 6, 201, 1979.
- Weiland J, and H. Wilhelmson, Coherent non-linear interaction of waves in plasmas" Pergamon Press, 1977.

Figure Caption

Figure 1 Current threshold for BPD ignition as a function of
pressure for three magnetic field values ($E = 1.5$ kV and
 $L = 20\text{m}$).




FIGURE 1

DA
FIL