

AD-A141 429 ORGANOPHOSPHAZENES 17 THE SYNTHESIS OF
TRIMETHYLSILYLACETYLENE AND TERMIN..(U) VERMONT UNIV
BURLINGTON DEPT OF CHEMISTRY C W ALLEN ET AL.
UNCLASSIFIED 18 MAY 84 TR-14 N00014-77-C-0605

F/G 7/3

NL

1 /


END

DATE

FILMED

7 84

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963-A

AD-A141 429

DTIC FILE COPY

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 14	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Organophosphazenes. 17. The Synthesis of Trimethylsilylacetylene and Terminal Acetylene Derivatives of Hexafluorocyclotriphosphazene		5. TYPE OF REPORT & PERIOD COVERED Technical Report
7. AUTHOR(s) Christopher W. Allen, James L. Desorcie and Kolikkara Ramachandran		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry University of Vermont Burlington, Vermont		8. CONTRACT OR GRANT NUMBER(s) N001477C-0605
11. CONTROLLING OFFICE NAME AND ADDRESS Department of the Navy Office of Naval Research Arlington, VA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE 5/18/84
		13. NUMBER OF PAGES 12
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) JUL 25 1984 A		
18. SUPPLEMENTARY NOTES Accepted for publication in Journal of the Chemical Society, Dalton Transactions.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Cyclophosphazenes Acetylene derivatives nmr		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The reactions of lithio trimethylsilylacetylene, $\text{LiC}\equiv\text{CSiMe}_3$, with hexafluorocyclotriphosphazene, $\text{N}_3\text{P}_3\text{F}_6$, lead to the new alkynphosphazenes $\text{N}_3\text{P}_3\text{F}_6\text{-n}$ ($\text{C}\equiv\text{CSiMe}_3$) _n (n=1,2). Both the geminal and nongeminal isomers are observed in the ^{31}P and ^{19}F n.m.r. spectra. A reinvestigation of the reaction of lithio phenylacetylene, $\text{LiC}\equiv\text{CPh}$, with $\text{N}_3\text{P}_3\text{F}_6$ showed that, at the stage of bis substitution, in addition to the previously reported major product, gem- $\text{N}_3\text{P}_3\text{F}_4(\text{C}\equiv\text{CPh})_2$, small amounts of the non-geminal $\text{N}_3\text{P}_3\text{F}_4(\text{C}\equiv\text{CPh})_2$ isomers are formed. (Continued)		

DD FORM 1473

1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

S/N 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

84 05 24 008

20. Continued

The reactions of $N_3P_3F_{6-n}(C\equiv CSiMe_3)_n$ ($n=1,2$) with potassium fluoride in anhydrous ethanol leads to the terminal acetylene derivatives, $N_3P_3F_{4-n}(OC_2H_5)_2(C\equiv CH)_n$ ($n=1,2$).



Appetition For	
NAME	☒
DATE	☐
OR NAME	☐
DATE	☐
/	
Code	
and/or	
Serial	
A1	

Unclassified

OFFICE OF NAVAL RESEARCH
Contract N001477C-0605
Project NR 356-663
Technical Report No. ~~22~~ 14

Organophosphazenes. 17. The Synthesis of Trimethylsilylacetylene and Terminal Acetylene Derivatives of Hexafluorocyclotriphosphazene.

by

Christopher W. Allen, James L. Desorcie and Kolikkara Ramachandran

Accepted for Publication in the
Journal of the Chemical Society, Dalton Transactions

University of Vermont
Department of Chemistry
Burlington, Vermont 05405

May 18, 1984

Reproduction in whole or in part is permitted for any purposes of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

Proofs to: Christopher W. Allen
 Department of Chemistry
 Cook Science Building
 University of Vermont
 Burlington, Vermont 05405
 USA

Organophosphazenes. 17. The Synthesis of Trimethylsilylacetylene and
 Terminal Acetylene Derivatives of Hexafluorocyclotriphosphazene.^{1†}

Christopher W. Allen*, James L. Desorcie and Kolikkara Ramachandran
 Department of Chemistry, University of Vermont, Burlington, Vermont 05405,
 U.S.A.

The reactions of lithio trimethylsilylacetylene, $\text{LiC}\equiv\text{CSiMe}_3$, with hexa-
 fluorocyclotriphosphazene, $\text{N}_3\text{P}_3\text{F}_6$, lead to the new alkynphosphazenes
 $\text{N}_3\text{P}_3\text{F}_{6-n}(\text{C}\equiv\text{CSiMe}_3)_n$ ($n=1,2$). Both the geminal and non-
 geminal isomers are observed in the ^{31}P and ^{19}F n.m.r. spectra. A reinvestigation
 of the reaction of lithio phenylacetylene, $\text{LiC}\equiv\text{CPh}$, with $\text{N}_3\text{P}_3\text{F}_6$ showed that,
 at the stage of bis substitution, in addition to the previously reported
 major product, $\text{gem-N}_3\text{P}_3\text{F}_4(\text{C}\equiv\text{CPh})_2$, small amounts of the non-geminal
 $\text{N}_3\text{P}_3\text{F}_4(\text{C}\equiv\text{CPh})_2$ isomers are formed. The reactions of $\text{N}_3\text{P}_3\text{F}_{6-n}(\text{C}\equiv\text{CSiMe}_3)_n$
 ($n=1,2$) with potassium fluoride in anhydrous ethanol leads to the terminal
 acetylene derivatives, $\text{N}_3\text{P}_3\text{F}_{4-n}(\text{OC}_2\text{H}_5)_2(\text{C}\equiv\text{CH})_n$ ($n=1,2$).

†Non-S.I. unit employed: mm Hg $\approx$ 13.6 x 9.8 Pa.

The reactions of organolithium and Grignard reagents with halophosphazenes have been shown to be valuable routes to organophosphazenes.² Much of the effort in this area has been devoted to aryl²⁻⁴ and alkyl^{2,4-6} substituted species with alkenyl^{1,2,6-8} and especially alkynyl^{2,9,10} derivatives being less common. We are particularly interested in alkenyl and alkynyl phosphazenes because of the potential, as shown in our work on propenyl phosphazenes, for further synthetic transformations into new organophosphazenes^{2,7,12} and organophosphazene polymers.^{2,11,12} A previous study by Chivers involving phosphazene derivatives of phenylacetylene, $N_3P_3F_{6-n}(C\equiv CC_6H_5)_n$ ($n=1,2$)⁹ demonstrated the viability of synthesis of alkynylphosphazenes. Alkynylphosphazenes have also been observed in the reactions of copper or lithiophosphazene anions with prop-2-ynyl bromide leading to materials of the type $N_3P_3Cl_4(R)CH_2C\equiv CH$ ($R=alkyl., allyl$).¹⁰ Rearrangement from the prop-2-ynyl derivative to the prop-1-ynyl derivative, $N_3P_3Cl_4(R)C\equiv CCH_3$ was also observed.¹⁰ A prop-1-ynyl derivative, $N_3P_3F_5C\equiv CCH_3$, was previously reported as a byproduct in the reaction of the lithiopropene $LiCH=CHCH_3$ with $N_3P_3F_6$.⁷ In this paper we describe the synthesis of phosphazenes with trimethylsilylacetylene and terminal acetylene substituents.

Experimental

Materials and Methods. Hexafluorocyclotriphosphazene, $N_3P_3F_6$, was prepared from $N_3P_3Cl_6$ (Ethyl Corp.) by a previously reported procedure.¹³ Phenylacetylene and n-butyl lithium (1.55 M solution in hexane) were obtained from Aldrich Corp.. Trimethylsilylacetylene was obtained from Petrarch systems. Diethyl ether and tetrahydrofuran were distilled from sodium benzophenone

ketyl. Petroleum ether (b.p. 35-55°C) was distilled over sodium. NMR spectra (in CHCl_3) were recorded on a Bruker WM250 spectrometer operating at 250.1 MHz (^1H), 62.9 MHz (^{13}C), 235.2 MHz (^{19}F) and 101.2 MHz (^{31}P). For ^{31}P NMR, 85% H_3PO_4 was used as external reference. For ^{19}F NMR, CFC_3 was used as the reference. ^{13}C , ^{19}F , and ^{31}P NMR spectra were recorded under conditions of broad band decoupling. Due to the complex second-order nature of the ^{19}F and ^{31}P spectra, parameters reported for these systems are approximate. Chemical shifts upfield to the reference are assigned a negative sign. Infrared (IR) spectra were obtained as their thin films (NaCl disks) on a Beckman IR 20A spectrometer. Mass spectra were recorded on a Perkin Elmer RMU-6D spectrometer operating at 80 eV. Elemental analysis were performed by Integral Microanalytical laboratories.

Preparation of $\text{N}_3\text{P}_3\text{F}_5\text{C}\equiv\text{CSiMe}_3$ (1). n-Butyl lithium (45 mL, 70.0 mmoles) was reacted with trimethylsilylacetylene (6.7 g, 66 mmoles) in diethyl ether (150 mL at -78°C using a previously described apparatus.¹⁴ The $\text{LiC}\equiv\text{CSiMe}_3$ thus obtained was added slowly (30 min.) to a cold (-78°C) stirred solution of $\text{N}_3\text{P}_3\text{F}_6$ (17.0g, 68 mmoles) in diethyl ether (200 mL). After 12 hr, the solvent was removed and petroleum ether (300 mL) was added to the residue. Filtration followed by the removal of petroleum ether from the filtrate gave a reddish brown liquid which was distilled under reduced pressure to yield a colorless liquid (10.2 g, 45.6% of theory), b.p. 30°C at 0.05 mm Hg. Anal. Calcd. for $\text{C}_5\text{H}_9\text{F}_5\text{N}_3\text{P}_3\text{Si}$: C, 18.35; H, 2.75; mol. wt. 327. Found: C, 18.48; H, 2.79; mol. wt. 327 (mass spectrum).

^1H NMR: $\delta_{\equiv\text{PF}(\text{C}\equiv\text{CSiMe}_3)} = -0.5$ (s) ^{31}P NMR: $\delta_{\equiv\text{PF}(\text{C}\equiv\text{CSiMe}_3)} = 3.6$
 (complex multiplet), $J_{\text{PF}} = 916.0$; $\delta_{\equiv\text{PF}_2} = 8.2$ (complex multiplet), $J_{\text{PF}} = 950.0$.
 ^{19}F NMR: $\delta_{\equiv\text{PF}(\text{C}\equiv\text{CSiMe}_3)} = -45.5$, $J_{\text{PF}} = 900.0$; $\delta_{\equiv\text{PF}_2} = -68.4$, $J_{\text{PF}} = 920.0$;
 $\delta_{\equiv\text{PF}_2} = -69.7$, $J_{\text{PF}} = 880$. ^{13}C NMR: $\delta_{\equiv\text{PF}(\text{C}\equiv\text{CSiMe}_3)} = 92.6$ (complex multiplet),
 $J_{\text{PC}} = 332.8$, $^2J_{\text{FC}} = 56.0$; $\delta_{\equiv\text{PF}(\text{C}\equiv\text{CSiMe}_e)} = 112.7$ (d of d), $^2J_{\text{PC}} = 51.5$;
 $\delta_{\equiv\text{PF}(\text{C}\equiv\text{CSiMe}_3)} = -1.5$ (s). IR: 2960 (w, $\equiv\text{C-H}$), 2145 (m, $-\text{C}\equiv\text{C}-$ Str.),
 1240 (s, P=N Str.), 950 (s, Si-C), 850 (s, P-F), 810 (s, P-F), 760 (m, P-F).

The same product can be obtained, in lower yields, from $\text{Me}_3\text{SiC}\equiv\text{CLi}$ prepared from ¹⁵ the reaction of methyl lithium with bis (trimethylsilyl)acetylene.

Preparation of $\text{N}_3\text{P}_3\text{F}_3(\text{OC}_2\text{H}_5)_2\text{C}\equiv\text{CH}$ (2). To a refluxing suspension of KF (0.2g) in absolute ethanol (25 mL), compound (1) (5.0 g, 15.3 mmoles) was added from a syringe. After 3 hr, the solvent was removed

and the residue obtained was extracted with petroleum ether (150 mL). The mixture was filtered and petroleum ether was removed from the filtrate. The dark brown liquid obtained was distilled under reduced pressure to give a colorless liquid (2.2 g, 46.8% of theory), b.p.

40°C at 0.05 mm Hg. Anal. Calcd. for $\text{C}_6\text{H}_{11}\text{F}_3\text{N}_3\text{O}_2\text{P}_3$: C, 23.45; H, 3.58; mol. wt. 307. Found: C, 23.21; H, 3.46; mol. wt. 307 (mass spectrum).

^1H NMR: $\delta_{-\text{C}\equiv\text{CH}} = 3.09$ (d), $^3J_{\text{PH}} = 13.56$, $\delta_{-\text{C}\equiv\text{CH}} = 3.08$ (d), $^3J_{\text{PH}} = 13.75$;
 $\delta_{-\text{OCH}_2\text{CH}_3} = 4.16$ (m), $\delta_{-\text{OCH}_2\text{CH}_3} = 1.38$ (m). ^{31}P NMR: very complex spectrum; approximate $\delta_{\equiv\text{PF}(\text{C}\equiv\text{CSiMe}_3)}$ values are 11.8 and 3.2. ^{19}F NMR:
 $\delta_{\equiv\text{PF}(\text{C}\equiv\text{CSiMe}_3)} = -65.5$; $\delta_{\equiv\text{PF}(\text{C}\equiv\text{SiMe}_3)} = -69.2$; $\delta_{\equiv\text{PF}_2} = -42.5$, $J_{\text{PF}} = 895.0$
 IR: 3280 (m, $\equiv\text{C-H}$), 2990 (m, CH Str.), 2065 (s, $-\text{C}\equiv\text{C}-$ Str.), 1260 (s, P=N Str.),

1165 (m), 1045 (s, P-O Str.), 975 (m, P-F), 915 (m, P-F), 845 (m, P-F), 785 (m, P-F).

Preparation of $N_3P_3F_4(C\equiv C-SiMe_3)_2$ (3). Lithio trimethylsilylacetylene, $LiC\equiv C-SiMe_3$, prepared by the reaction of n-butyl Li (64 mL, 100 mmoles) and trimethylsilylacetylene (9.8 g 100 mmoles) was allowed to react with $N_3P_3F_6$ (12.5 g, 50 mmoles) in diethyl ether (200 mL) as before. The reaction mixture was stirred for 2 days and worked up to give a viscous liquid (7.25 g, 35.8% of theory), b.p. 65°C at 0.05 mm Hg. Anal. Calcd. for $C_{10}H_{18}F_4N_3P_3Si_2$: C, 29.63; H, 4.44; mol. wt. 405. Found: C, 30.12; H, 4.38; mol. wt. 405 (mass spectrum).

1H NMR: $\delta_{-SiMe_3} = -0.45$ (s). ^{31}P NMR: $\delta_{\equiv P(C\equiv SiMe_3)_2} = -27.0$, $^2J_{PP} = 80.0$; $\delta_{\equiv PF_2} = 7.5$, $J_{PF} = 900.0$; $\delta_{\equiv PF(C\equiv CSiMe_3)} = 1.5$; $J_{PF} = 920.0$. ^{19}F NMR: $\delta_{\equiv PF_2} = -42.6$, $J_{PF} = 9200$; $\delta_{\equiv PF_2} = -44.5$, $J_{PF} = 890$; $\delta_{\equiv PF(C\equiv CSiMe_3)} = -68.6$, $J_{PF} = 915.0$. ^{13}C NMR: $\delta_{\equiv P(C\equiv CSiMe_3)_2} = 112.2$ (c.m.), $\delta_{\equiv P(C\equiv CSiMe_3)} = 96.3$ (t,d), $J_{PC} = 256.0$; $\delta_{\equiv P(C\equiv CSiMe_3)_2} = -1.1$ (s); $\delta_{\equiv PF(C\equiv CSiMe_3)} = 112.2$ (c-m); $\delta_{\equiv PF(C\equiv CSiMe_3)} = 93.8$ (c.m.), $J_{PC} = 300$, $^2J_{PC} = 56.0$; $\delta_{\equiv PF(C\equiv CSiMe_3)} = -1.4$ (s). IR: 2965 (m, CH Str.), 2140 (m, $-C\equiv C-$), 1265 (s, P=N), 930 (m, Si-C), 850 (m, P-F), 815 (m, P-F) 795 (m, P-F).

Attempts at separation of the mixture by distillation or crystallization proved unsuccessful. The materials apparently bind to silica gel and hence can not be eluted from a chromatography column.

Preparation of $N_3P_3F_2(OC_2H_5)_2(C\equiv CH)_2$ (4). The reaction of $N_3P_3F_4(C\equiv CSiMe_3)_2$ (3) (3.2g, 8.0 m moles) with KF (3.2 g) in absolute ethanol (40 mL) was

carried out as before to yield a dark brown liquid. Reduced pressure distillation of this liquid gave yielded a viscous oil (0.9 g, 36.7% of theory), b.p. 105°C at 0.05 mm Hg. Anal. Calcd. for $C_8H_{12}F_2N_3O_2P_3$: C, 30.67; H, 3.83; mol. wt. 313. Found: C, 29.94; H, 3.78; mol. wt. 313 (mass spectrum).

1H NMR: $\delta_{-C\equiv CH} = 3.0$ (c.m.); $\delta_{-OCH_2CH_3} = 4.1$ (c.m.); $\delta_{-OCH_2CH_3} = 1.4$ (c.m.).

^{31}P NMR: $\delta_{\equiv P(C\equiv CH)_2} = 25.0$ (c.m.); $\delta_{\equiv PF_2} = 7.5$ (c.m.); $\delta_{\equiv PFR} = 2.5$.

IR: = 3280 (m, $-C\equiv CH$ Str.), 2460 (m, CH Str.), 2165 (s, $-C\equiv C$), 1255 (s, P=N) 1170 (m), 1055 (s, P-O Str.), 965 (m, P-F), 915 (s, P-F), 845 (m, P-F), 790 (m).

Preparation of $N_3P_3F_5C\equiv CPh(5)$. n-Butylli (32 mL, 50 mmoles) was allowed to react with phenylacetylene (5.4 g, 50 mmoles) in diethyl ether (150 mL).

After 24 hr, the above reaction mixture was added to a solution of $N_3P_3F_6$ (13.4 g, 50 mmoles) in diethyl ether (150 mL). The resultant reaction mixture was stirred for 24 hr and worked up as before. Reduced pressure distillation of the product obtained gave $C_8H_5N_3P_3F_5$ (7.7 g 46.5% of theory), b.p. 45°C at 0.1 mm Hg. (Lit. b.p. 82°C at 2.5 mm Hg).

1H NMR: $\delta_{-C_6H_5} = 7.3$ (m). ^{31}P NMR: $\delta_{\equiv PF_2} = 8.2$ (t), $J_{PF} = 941.2$;

$\delta_{\equiv PF(C\equiv CC_6H_5)} = 6.0$ (d), $J_{PF} = 910.8$. ^{19}F NMR: $\delta_{\equiv PF_2} = -69.5$, $J_{PF} = 870.3$;

$\delta_{\equiv PF_2} = -68.5$, $J_{PF} = 903.2$; $\delta_{\equiv PF(C\equiv CC_6H_5)} = -44.0$, $J_{PF} = 893.8$. ^{13}C NMR:

$\delta_{\equiv PF(C\equiv CSiMe_3)} = 102.7$ (complex d), $J_{PC} = 69.2$; $\delta_{\equiv PF(C\equiv CC_6H_5)} = 118.0$ (d),

$^2J_{PC} = 6.1$; $\delta_{C_6H_5} = 128.0, 131.9, 133.0$. IR: 2180 (s, $-C\equiv C$), 1250 (s, P=N Str.),

1060 (w), 1010 (m), 980 (m, P-F), 930 (s, P-F), 880 (s, P-F), 830 (s, P-F).

Preparation of $N_3P_3F_4(C\equiv CPh)_2$ (6). Lithio phenylacetylene, $LiC\equiv CPh$ prepared

by the reaction of phenylacetylene (2.25g, 22.0 m moles) and n-butyl lithium (15 mL, 23 m moles) was allowed to react with $N_3P_3F_6$ (2.7 g, 11.0 m moles) in Et_2O (200 mL). After 24 hr, the reaction mixture was worked as before to yield a dark brown oil which was dissolved in petroleum ether - methylene chloride (3:1) and cooled. The slightly yellowish crystals obtained was recrystallized from petroleum ether-methylene chloride (3:1) to obtain $2,2-N_3P_3F_4(C\equiv CPh)_2$, 1.2 g (27.3% of theory). 1H NMR: $\delta_{C_6H_5}$: 5.6 (m). ^{31}P NMR: $\delta_{\equiv PF_2}$: 6.9 (t), $J_{PF} = 910.8$; $\delta_{\equiv P(C\equiv CC_6H_5)_2} = -23.2$, $^2J_{PP} = 91.1$, $^3J_{PF} = 12.1$.

^{19}F NMR: $\delta_{\equiv PF_2}$: - 68.6 (d), $J_{PF} = 889.7$. ^{13}C NMR: $\delta_{\equiv P(C\equiv CC_6H_5)_2} = 101.8$ (d), $J_{PC} = 56.6$; $\delta_{\equiv P(C\equiv CC_6H_5)_2} = 119.1$ (d), $^2J_{PC} = 56.7$; $\delta_{C_6H_5} = 128.7, 131.3$ and 133.6. IR: 2140 (s, $-C\equiv C$), 1220 (s, $P=N$), 1060 (w), 1010 (w, PF), 920 (s, PF), 870 (s, PF), 800 (s, PF).

The solvent was removed from the mother liquor remaining after isolation of the geminal isomer and the residual oil was distilled under reduced pressure. The distillate was found to contain two components. The major component was the geminal bis derivative (6) obtained above. The mass spectrum of the mixture showed only parent peaks due to $N_3P_3F_4(C\equiv CPh)_2$, while the ^{19}F and ^{31}P nmr spectra showed a small amount of $\equiv PF(C\equiv CPh)$ centers in the mixture.

Results and Discussion

The reaction of $LiC\equiv CSiMe_3$ with $N_3P_3F_6$ proceeds smoothly to give the expected product, $N_3P_3F_5C\equiv CSiMe_3$ (1) in moderate yield. Elemental analyses,

mass spectrometric data and spectroscopic (n.m.r., i.r.) properties of 1 are in accord with the assigned structure. Of particular value are the ^{31}P and ^{19}F nmr spectra which show the presence of the $\equiv\text{PF}_2$ and $\equiv\text{PF}(\text{R})$ ($\text{R} = \text{C}\equiv\text{CSiMe}_3$) centers. The addition of two molar equivalents of $\text{LiC}\equiv\text{CSiMe}_3$ to $\text{N}_3\text{P}_3\text{F}_6$ produces a viscous liquid which was shown to be a mixture of the disubstituted isomers, $\text{N}_3\text{P}_3\text{F}_4(\text{C}\equiv\text{CSiMe}_3)_2$ (3). Further attempts at a separation of the components of the mixture proved unsuccessful. Some indication of the isomeric composition of 3 was obtained from an examination of the ^{31}P and ^{19}F nmr spectra. The ^{31}P spectrum clearly shows the presence of a $\equiv\text{P}(\text{C}\equiv\text{CSiMe}_3)_2$ center but the remaining portion of the spectrum is far too complex to be ascribed only to the $\equiv\text{PF}_2$ region of the geminal isomer. The observation of $\equiv\text{PF}(\text{C}\equiv\text{CSiMe}_3)$ and two $\equiv\text{PF}_2$ resonances in the ^{19}F nmr spectrum demonstrates the existence of both geminal and non-geminal isomers in the mixture. A comparison of the relative areas of the $\equiv\text{PF}_2$ resonances and of the $\equiv\text{PF}(\text{C}\equiv\text{CSiMe}_3)$ resonances indicates that significant amounts of both geminal and non-geminal isomers are present.

Since it has been reported that the reaction of $\text{LiC}\equiv\text{CPh}$ with $\text{N}_3\text{P}_3\text{F}_6$ follows a geminal pathway,⁹ we chose to reinvestigate the lithiophenylacetylene reaction. In large part, this reaction proceeds as previously reported leading to $\text{N}_3\text{P}_3\text{F}_5\text{C}\equiv\text{CPh}$ (5) and the geminal bis derivative, $2,2\text{-N}_3\text{P}_3\text{F}_4(\text{C}\equiv\text{CPh})_2$.⁹ We also observed a small amount of the previously unreported non-geminal $\text{N}_3\text{P}_3\text{F}_4(\text{C}\equiv\text{CPh})_2$ isomers. In a comparison of the isomer distribution at the stage of bis substitution in the reactions of $\text{N}_3\text{P}_3\text{F}_6$ with $\text{LiC}\equiv\text{CR}$ ($\text{R} = \text{SiMe}_3, \text{Ph}$), one observes a significant contribution from the non-geminal

pathway in the trimethylsilyl derivative while a predominantly geminal pathway is observed in the formation of phenyl derivative. The sensitivity of this reaction to the nature of the β -substituent on the acetylenic center is remarkable even in the context of the established complexity of organometallic reactions of the cyclophosphazenes.² Geminal substitution is the preferred pathway in the reactions of organometallic reagents with $N_3P_3F_6$.^{2,7} Formation of non-geminal isomers is usually ascribed to steric hinderance in the organometallic reagent.² It is not immediately obvious that this should be the case in alkynylphosphazenes. Although examination of the nmr data for 1 and 5 shows a striking variation in $^1J_{PC}$, the variation in ^{31}P shifts is less than is noted in various alkynylphosphazenes¹⁶ and the ^{19}F shifts are virtually identical. Thus, significant differences in electronic structure between 1 and 5 are not expected.

We have explored the fluoride ion induced cleavage of the carbon-silicon bond in the trimethylsilylacetylene derivatives 1 and 3. The reaction of 1 with anhydrous KF in absolute ethanol leads to an inseparable mixture of trisubstituted isomers of the type $N_3P_3F_3(OC_2H_5)_2C\equiv CH$ (2). The 1H nmr spectrum shows the presence of the terminal acetylene and ethoxy functions and the absence of the trimethylsilyl group. The ^{31}P nmr spectrum is complex but doesn't show any significant concentration of $\equiv PF_2$ or $\equiv PRR'$ centers. The ^{19}F spectrum confirms the predominance of $\equiv PFR$ ($R=C\equiv CH$, OC_2H_5) environments and shows a trace amount of a $\equiv PF_2$ center. Thus, the material is essentially a mixture of the three possible

non-geminal isomers, $2,4,6\text{-N}_3\text{P}_3\text{F}_3(\text{OC}_2\text{H}_5)_2\text{C}\equiv\text{CH}$) with a trace of a geminal isomer. The tendency towards non-geminal substitution by the ethoxide ion is consistent with previous observations of the reactions of anionic oxygen nucleophiles with $\text{N}_3\text{P}_3\text{Cl}_6$.¹⁷ The ethoxy function can be seen as arising from a variety of routes. The cleavage of the carbon-silicon bond results in the formation of trimethylsilyl fluoride and a carbanion which abstracts a proton from the solvent, thereby generating an ethoxide ion. The second ethoxy function comes directly from the solvent or from the ethanol-KF mixture. The fact that the reaction stops at the tri-substituted stage can be ascribed to the higher energy barrier associated with geminal substitution by alkoxides.¹⁷

The reaction of the mixture of bis trimethylsilylacetylene derivatives, 3, with KF in absolute ethanol leads to the formation of a mixture of terminal acetylene derivatives having the composition $\text{N}_3\text{P}_3\text{F}_2(\text{OC}_2\text{H}_5)_2(\text{C}\equiv\text{CH})_2$ (4). The ^{31}P and ^{19}F nmr spectra of 4 are very complex and outside of suggesting the presence of several environments, do not provide much useful information. These are nine possible isomers of 4 and most or all are presumed to be present.

Acknowledgement. This work was supported in part by the Office of Naval Research. The assistance of Ms. P. T. Malik in checking some of the synthetic results is appreciated.

References

1. Part 16: C.W. Allen and R.P. Bright, Inorg. Chem., 1983, 22, 1291.
2. C.W. Allen, Ind. Eng. Chem. Prod. Res. Dev., 1981, 20, 20.
3. C.W. Allen and P.L. Toch, Inorg. Chem., 1981, 20, 9; H.R. Allcock, T.L. Evans and D.B. Patterson, Macromolecules, 1980, 13, 201; T. Chivers and N.L. Paddock, Inorg. Chem., 1972, 11, 848; C.W. Allen, J. Chem. Soc., Chem. Commun., 1970, 152; C.W. Allen and T. Moeller, Inorg. Chem., 1968, 7, 2177.
4. P.J. Harris, J.L. Desorcie and H.R. Allcock, J. Chem. Soc., Chem. Comm., 1981, 852.
5. P.J. Harris and C.L. Fadeley, Inorg. Chem., 1983, 22, 561; K. Ramachandran and C.W. Allen, J. Amer. Chem. Soc., 1982, 104, 2396; H.R. Allcock and P.J. Harris, J. Amer. Chem. Soc., 1979, 101, 6221; J. Blumentritt and T. Moeller, Inorg. Nucl. Chem. Lett., 1978, 14, 263; T.N. Ranaganathan, S.M. Todd and N.L. Paddock, Inorg. Chem., 1973, 12, 316; N.L. Paddock, T.N. Ranaganathan and S.M. Todd, Can. J. Chem., 1971, 49, 164; T. Moeller, A. Failli and F.Y. Tsang, Inorg. Nucl. Chem. Lett., 1969, 1, 49.
6. E. Niecke, H. Thamm and O. Glemser, Z. Naturforsch., 1971, 26b, 366.
7. J.G. Dupont and C.W. Allen, Inorg. Chem., 1978, 17, 3093.
8. R.D. Sharma, S.J. Rettig, N.L. Paddock and J. Trotter, Can. J. Chem., 1982, 60, 535.
9. T. Chivers, Inorg. Nucl. Chem. Lett., 1971, 7, 827.
10. H.R. Allcock, P.J. Harris and R.A. Nissan, J. Amer. Chem. Soc., 1981, 103, 2256.
11. J.G. Dupont and C.W. Allen, Macromolecules, 1979, 12, 169.
12. C.W. Allen and J.G. Dupont, Ind. Eng. Chem. Prod. Res. Dev., 1979, 18, 81.
13. T. Moeller, K. John and F. Tsang, Chem. Ind. (London) 1961, 347.
14. C.W. Allen, R.P. Bright, J.L. Desorcie, J.A. MacKay and K. Ramachandran, J. Chem. Educ., 1980, 57, 565.
15. A.B. Holmes, C.L.D. Jennings-White, A.H. Schulthess, B. Akinde and D.R.M. Walton, J. Chem. Soc., Chem. Comm., 1979, 840.

16. P.J. Harris, Inorg. Chim. Acta., 1983, 71, 233.
17. K. Ramachandran and C.W. Allen, Inorg. Chem., 1983, 22, 1445; J.L. Schmutz and H.R. Allcock, Inorg. Chem., 1975, 14, 2433.

DL/413/83/01
GEN/413-2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
ONR Pasadena Detachment Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Weapons Center Attn: Dr. A. B. Amster Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	Dean William Tolles Naval Postgraduate School Monterey, California 93940	1
Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Mr. Vincent Schaper DTNSRDC Code 2830 Annapolis, Maryland 21402	1
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1	Mr. A. M. Anzalone Administrative Librarian PLASTEC/ARRADCOM Bldg 3401 Dover, New Jersey 07801	1

DL/413/83/01
356B/413-2

TECHNICAL REPORT DISTRIBUTION LIST, 356B

Dr. C. L. Schilling
Union Carbide Corporation
Chemical and Plastics
Tarrytown Technical Center
Tarrytown, New York

Dr. A. G. MacDiarmid
Department of Chemistry
University of Pennsylvania
Philadelphia, Pennsylvania 19174

Dr. E. Fischer, Code 2853
Naval Ship Research and
Development Center
Annapolis, Maryland 21402

Dr. H. Allcock
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. M. Kenney
Department of Chemistry
Case Western University
Cleveland, Ohio 44106

Dr. R. Lenz
Department of Chemistry
University of Massachusetts
Amherst, Massachusetts 01002

Dr. M. David Curtis
Department of Chemistry
University of Michigan
Ann Arbor, Michigan 48105

NASA-Lewis Research Center
Attn: Dr. T. T. Serafini, MS 49-1
21000 Brookpark Road
Cleveland, Ohio 44135

Dr. J. Griffith
Naval Research Laboratory
Chemistry Section, Code 6120
Washington, D.C. 20375

Professor G. Wnek
Department of Materials Science
and Engineering
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. R. Soulen
Contract Research Department
Pennwalt Corporation
900 First Avenue
King of Prussia, Pennsylvania 19406

Dr. G. Goodman
Globe-Union Incorporated
5757 North Green Bay Avenue
Milwaukee, Wisconsin 53201

Dr. Martin H. Kaufman
Code 38506
Naval Weapons Center
China Lake, California 93555

Professor R. Drago
Department of Chemistry
University of Florida
Gainesville, Florida 32611

Dr. D. L. Venezky
Code 6130
Naval Research Laboratory
Washington, D.C. 20375

Professor T. Katz
Department of Chemistry
Columbia University
New York, New York 10027

Professor James Chien
Department of Chemistry
University of Massachusetts
Amherst, Massachusetts 01002

Professor J. Salamone
Department of Chemistry
University of Lowell
Lowell, Massachusetts 01854

Dr. S. Cooper
Department of Chemistry
University of Wisconsin
750 University Avenue
Madison, Wisconsin 53706

DL/413/83/01
356B/413-2

TECHNICAL REPORT DISTRIBUTION LIST, 356B

Professor D. Grubb
Department of Materials Science
and Engineering
Cornell University
Ithaca, New York 14853

Professor T. Marks
Department of Chemistry
Northwestern University
Evanston, Illinois 60201

Professor C. Chung
Department of Materials Engineering
Rensselaer Polytechnic Institute
Troy, New York 12181

Professor Malcolm B. Polk
Department of Chemistry
Atlanta University
Atlanta, Georgia 30314

Dr. D. B. Cotts
SRI International
333 Ravenswood Avenue
Menlo Park, California 94205

Dr. Kurt Baum
Fluorochem, Inc.
680 S. Ayon Avenue
Azusa, California 91702

CAPT J. J. Auburn, USNR
AT&T Bell Laboratories
Room 6F-211
600 Mountain Avenue
Muray Hill, New Jersey 07974

Professor H. Hall
Department of Chemistry
University of Arizona
Tucson, Arizona 85721

Professor G. Whitesides
Department of Chemistry
Harvard University
Cambridge, Massachusetts 02138

Professor H. Ishida
Department of Macromolecular Science
Case Western University
Cleveland, Ohio 44106

Dr. K. Paciorek
Ultrasystems, Inc.
P.O. Box 19605
Irvine, California 92715

Professor D. Seyferth
Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. G. Bryan Street
IBM Research Laboratory, K32/281
San Jose, California 95193