

RD-A143 442

A COMPARISON OF NON-LINEAR REGRESSION AND WEIGHTED
LEAST SQUARES FOR PRED. (U) UNIVERSITY OF CENTRAL
FLORIDA ORLANDO DEPT OF MATHEMATICS AND.
L A FRANKLIN ET AL. 01 APR 84 SCIENTIFIC-6 F/B 12/1

1/1

UNCLASSIFIED


ML



END

FILMED

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A143 442

DTIC FILE COPY

AFGL-TR-84-0132

A Comparison of Non-Linear Regression
and Weighted Least Squares for
Predicting Visibility in Germany

LeRoy A. Franklin
Paul N. Somerville
Steven J. Bean

Department of Statistics
University of Central Florida
Orlando, FL 32816

Scientific Report No. 6

1 April 1984

Approved for Public Release, distribution unlimited

AIR FORCE GEOPHYSICS LABORATORY
AIR FORCE SYSTEMS COMMAND
UNITED STATES AIR FORCE
HANSCOM AFB, MASSACHUSETTS 01731




84 07 23 011

This report has been reviewed by the ESD Public Affairs Office (PA) and is releasable to the National Technical Information Service (NTIS).

This technical report has been reviewed and is approved for publication



CHARLES F. BURGER
Contract Monitor



DONALD D. GRANTHAM
Chief, Tropospheric Structure Branch

FOR THE COMMANDER



ROBERT A. McCLATCHEY
Director, Atmospheric Sciences Division

Qualified requestors may obtain additional copies from the Defense Technical Information Center. All others should apply to the National Technical Information Service.

If your address has changed, or if you wish to be removed from the mailing list, or if the addressee is no longer employed by your organization, please notify AFGL/DAA, Hanscom AFB, MA 01731. This will assist us in maintaining a current mailing list.

Do not return copies of this report unless contractual obligations or notices on a specific document requires that it be returned.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFGL-TR-84-0132	2. GOVT ACCESSION NO AD-A143 442	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A Comparison of Non-Linear Regression and Weighted Least Squares for Predicting Visibility in Germany		5. TYPE OF REPORT & PERIOD COVERED Scientific Report #6
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) LeRoy A. Franklin Paul N. Somerville Steven J. Bean		8. CONTRACT OR GRANT NUMBER(s) F19628-82-K-0001
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Central Florida Department of Statistics Orlando, Florida 32816		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 62101F 667009AK
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Geophysics Laboratory Hanscom AFB, MA 01731 Contract Monitor: C. F. Burger/LYT		12. REPORT DATE 1 April 1984
		13. NUMBER OF PAGES 15
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release, Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Non-Linear Regression Weighted Least Squares Sample Reuse Constants Model Variables Model		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Two methods of parameter estimation for the two parameter Weibull distribu- tion are investigated using data on visibility from Germany. They are: Non-Linear Regression and Weighted Least Squares. Non-Linear Regression gave superior results for both "constants" and "variables" models when measured either by overall RMS or by sample reuse. However, since the Weighted Least Squares gave similar results and utilized only about one- tenth the computer time, the authors recommend that it might be used to investigate other models of visibility for other countries.		

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ACKNOWLEDGEMENT

We would like to acknowledge the contribution of Mr. David Van Brackle in the accomplishment of the results of this paper. David was responsible for writing several computer programs and made all of the numerous runs required for this report.

DTIC
ELECTE
JUL 24 1984
B

Accession For	
NTIS	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

TABLE OF CONTENTS

	<u>Page</u>
1. INTRODUCTION	1
2. BACKGROUND OF PREVIOUS WORK ON VISIBILITY IN GERMANY	1
3. METHODOLOGY AND RESULTS	4
4. CONCLUSIONS	6
5. REFERENCES	7

A Comparison of Non-Linear Regression and Weighted Least Squares for Predicting Visibility in Germany

by

L. A. Franklin, P. N. Somerville, and S. J. Bean
University of Central Florida

1. INTRODUCTION

A goal of the Air Weather Service is to be able to state the probability that a weather element will have a value above a specified threshold for any location at any time. Many models have been developed for such weather elements as visibility, ceiling, sky cover, etc., for locations where records exist. Modeling these elements for locations where there are no records is a more difficult task. Some models have been developed by Bean and Somerville that require only knowledge of elevation of the location in question and the average of the elevations 20 kilometers from the location to estimate visibility probabilities for a specified month and hour period for Germany. These models were developed using the method of non-linear regression. The method of evaluating how well the models predict visibility at data-void locations (i.e., sample reuse) was also developed using non-linear regression. This paper compares the method of non-linear regression with a new method, "weighted least squares," in developing and evaluating these models.

2. BACKGROUND OF PREVIOUS WORK ON VISIBILITY IN GERMANY

Bean and Somerville in AFGL-TR-81-0144 "Some Models for Visibility for German Stations" showed that the Weibull distribution was able to fit visibility data for 30 stations in Germany. For any distance x , the probability $F(x)$ of visibility less than x miles is then given by the Weibull formula

$$F(x) = 1 - e^{-\alpha x^{\beta}}$$

where a different set of α and β values were derived for each month and each 3-hour period.

The values of α and β were estimated by choosing the values for which the Weibull cumulative distribution most closely fits the empirical cumulative distribution. Let $E_j(x_i)$ be the empirical probability (step function) that the visibility is less than x_i miles for the j^{th} station, using data from the RUSSWOs (Revised Uniform Summary of Surface Weather Observations). The values of x_i are the following distances in miles: $\frac{1}{4}, \frac{5}{16}, \frac{1}{2}, \frac{5}{8}, \frac{3}{4}, 1, \frac{5}{4}, \frac{3}{2}, 2, \frac{5}{2}, 3, 4, 5$ and 6 . Let

$$F_j(x; \alpha_j, \beta_j) = 1 - e^{-\alpha_j x^{\beta_j}}$$

be the Weibull distribution at station j . Then the values of α_j and β_j that are chosen are those that minimize the expression

$$\sum_j \sum_i [E_j(x_i) - F_j(x_i; \alpha_j, \beta_j)]^2 \quad (2.1)$$

That is, the values of α_j and β_j are those that minimize the sum of squares of the difference between the empirical and model probabilities over all stations and all distances for which data is available. This deviation is done for each of the 12 months and 8 three-hour periods of interest.

The coefficients α_j and β_j may themselves be dependent upon other variables. In fact, Bean and Somerville in AFGL-TR-82-0335 "Some New Practical Models for Visibility for Germany Locations Where No Records Exist" found that elevation of the location of interest and relative elevation at 20 kilometers around the location of interest provided an improvement in the model. The coefficients can be written as

$$\begin{aligned} \alpha_j &= \gamma_0 + \gamma_1 * EL_j + \gamma_2 * AE_j \\ \beta_j &= \delta_0 + \delta_1 * EL_j + \delta_2 * AE_j \end{aligned}$$

where EL_j and AE_j are transformed and scaled values for the elevation at station j and the average elevation at 20 equispaced locations on a circle of radius 20 kilometers and centered at the station. EL is the cube of the elevation in feet, divided by 10^9 . AE is the cube of the average elevation in feet, divided by 10^9 . The constants $\gamma_0, \gamma_1, \gamma_2$ and $\delta_0, \delta_1, \delta_2$ are determined by minimizing expression (2.1). The presence of all six constants comprises what has been called the "variables model" using non-linear regression. If we have only $\alpha_j = \gamma_0$ and $\beta_j = \delta_0$ the model has been called the "constants model" using non-linear regression and effectively fits only one model $\alpha = \delta_0$ and $\beta = \delta_0$ for all stations, ignoring any geographical features at the stations. The "constants model" has been evaluated and discussed for 30 stations in Germany in AFGL-TR-81-0313 "Modeling Visibility for Locations in Germany Where No Records Exist." The "variables model" has been evaluated and discussed for 60 stations in Germany in AFGL-TR-82-0335.

The method of minimizing expression (2.1) has been the method of non-linear regression and has been discussed in detail in AFGL-TR-80-0362 "Least Squares Fitting of Distribution Using Non-Linear Regression." The technique involves an iterative solution incorporating an initial estimate of the parameters. While the method has been extremely successful in fitting models and displays very robust features, it is time consuming to execute even on a mid-sized computer.

The method used to evaluate the models formulated is called sample reuse and has been discussed in AFGL-TR-82-0335. Briefly, sample reuse takes a single station and uses all the other stations to build the model and then calculates how well the model predicts visibility at the omitted station measured in the usual root mean square sense (i.e., RMS). This is done for each station for which you have data and hence, for n stations,

results in n times as many non-linear regressions as needed to fit the original model to all n stations at once.

It was in the interest of trying to find a new method of estimating the models that several methods were investigated in AFGL-TR-83-0248 "A Comparison of Several Alternatives to Maximum Likelihood for the Weibull Distribution." In that simulation study, the method of non-linear regression seemed more robust in that it was able to give better models when the data was contaminated or when the underlying true distribution was not the form of the distribution chosen to model it. However the method of weighted least squares showed promise in that it gave good results (although not as good as non-linear regression) and used only a small fraction of the time that non-linear regression needed. This method was initially suggested by Major Al Boehm, USAF, and it was the purpose of this report to compare this technique with non-linear regression on actual visibility data in Germany.

3. METHODOLOGY AND RESULTS

3.1 The Constants Model

The resulting RMS of fitting the "constants model" by both non-linear regression and weighted least squares to the first group of 30 stations in Germany is recorded in Exhibit 3.1 for all month and hour periods. In that exhibit for the "constant model," non-linear regression always produced smaller RMS than the weighted least squares. The difference was as small as .002 for the period 06-08 LST of October to as large as .112 for period 00-02 LST of January where RMS for weighted least squares was three times the value for non-linear regression.

In AFGL-TR-82-0187, "Evaluation of An Observation-Based Climatology Model for Predicting Visibility for Data-Void Locations in Germany," the constants model was expanded to a total of 60 stations in Germany and RMS from sample reuse based on non-linear regression was utilized to measure its

ability to predict visibility. The overall results of Exhibits 2.3 and 2.4 of that report are included in Exhibits 3.2 and 3.3 of this report along with the RMS from sample reuse based on weighted least squares. Hour periods 00-02 LST and 03-05 LST are omitted since data from these pre-dawn hours was frequently missing. For the stations, the weighted least squares method produced a larger RMS than non-linear regression at each station, usually between 2 and 3 times as large. The average RMS for non-linear regression was .098 while for weighted least squares, it was .177. Similar results hold for the monthly and hourly RMSs.

3.2 The Variables Model

The variables model was fit to all 60 stations in Germany and the coefficients for the α and β terms as well as overall fit RMS and sample reuse RMS were reported in AFGL-TR-82-0335 and displayed as Exhibit 2.4 of that report. It is included in Exhibit 3.4 along with comparable RMS results of overall fit and sample reuse using weighted least squares. Whether one compares RMS of the overall fitted "variables model" done by non-linear regression and by weighted least squares or whether one compares RMS from sample reuse from the two methods the results are the same. In virtually every instance non-linear regression gives a smaller RMS, usually by a factor of 2 to 3. Noteworthy exceptions occur at Grosser Falk and Bad Kreuznach where weighted least squares does marginally better. However both stations have previously presented problems of modeling and have unusually large RMS values under non-linear regression.

Exhibit 3.5 presents the overall RMS fit of the variables model to the complete set of 60 stations in Germany using both non-linear regression (the top entry in each cell) and weighted least squares (the bottom entry in each cell) for all 12 months and hour periods 3 through 8. Non-linear regression in every case produces a lower RMS than weighted least squares, sometimes as

little as 10 percent lower (for October and hour period 9-11) but usually almost 50 percent lower. The overall average RMS for all months and time periods for non-linear regression was .071 as compared to .095 for weighted least squares.

Exhibit 3.6 presents the RMS for the variables model using sample reuse on the 60 stations in Germany using non-linear regression (the top entry in each cell) and weighted least squares (the bottom entry in each cell). Again the RMS achieved by non-linear regression was always smaller than the RMS achieved by weighted least squares.

4. CONCLUSIONS

The results of investigating weighted least squares as a method to derive visibility models is clear and consistent with other results. Non-linear regression consistently provides better models than weighted least squares, usually dramatically better. This is "better" in the sense of better-fitting models (i.e., overall RMS) and "better" in the sense of better-predicting models (i.e., sample reuse).

However, results from weighted least squares is consistent with non-linear regression in the sense that relatively low RMS values derived by NLR correspond to relatively low RMS values derived by WLS. In particular, stations that are "difficult to fit" or may contain "bad data" remain so under either technique. Thus it seems reasonable that WLS can be used to construct many models of visibility and explore them for (relative) goodness of fit, since it is much faster to compute than NLR. Hence having eliminated a vast array of poor models by WLS, it would be possible to use NLR upon these few models which WLS points to as most promising. NLR can then choose the "best among the good" and give the lowest RMS and most accurate values for the coefficients of α and β in the Weibull Model.

It is hoped that such a two-step method can be employed by the authors on data from Norway to see if it is successful.

5. REFERENCES

- Bean, S. J., and P. N. Somerville, "Evaluation of an Observation-Based Climatology Model for Predicting Visibility for Data-Void Locations in Germany," AFGL-TR-82-0187, July 1982, AD A123998.
- Bean, S. J., P. N. Somerville and L. A. Franklin, "A Comparison of Several Alternatives to Maximum Likelihood for the Weibull Distribution," AFGL-TR-83-0248, September 1983, AD A139721.
- Heuser, M., P. N. Somerville, and S. J. Bean, "Least Squares Fitting of Distributions Using Non-Linear Regression," AFGL-TR-80-0362, September 1980, AD A097039.
- Somerville, P. N., and S. J. Bean, "Some Models for Visibility for German Stations," AFGL-TR-81-0144, April 1981, AD A104167.
- Somerville, P. N. and S. J. Bean, "Modeling Visibility for Locations in Germany Where No Records Exist," AFGL-TR-81-0313, December 1981, AD A111890.
- Somerville, P. N. and S. J. Bean, "Some New Practical Models for Visibility for German Locations Where No Records Exist," AFGL-TR-82-0335, 3 September 1982, AD A126339.

Month	Hour Period (LST)							
	00-02	03-05	06-08	9-11	12-14	15-17	18-20	21-23
January	.068 .180	.102 .151	.107 .144	.099 .104	.087 .096	.078 .086	.091 .114	.075 .165
February	.082 .178	.093 .135	.078 .081	.064 .067	.057 .060	.053 .057	.072 .088	.090 .164
March	.048 .105	.086 .120	.066 .069	.057 .061	.053 .062	.041 .045	.051 .064	.055 .091
April	.047 .086	.073 .106	.054 .059	.042 .047	.029 .033	.027 .036	.042 .058	.034 .057
May	.034 .064	.064 .090	.045 .051	.029 .032	.016 .019	.013 .018	.021 .029	.028 .050
June	.053 .082	.067 .097	.052 .059	.035 .042	.019 .024	.016 .020	.026 .042	.027 .048
July	.045 .078	.074 .104	.052 .059	.031 .036	.016 .022	.012 .017	.026 .047	.031 .052
August	.040 .072	.076 .113	.076 .091	.047 .058	.025 .034	.015 .019	.023 .030	.071 .460
September	.066 .115	.101 .144	.088 .097	.062 .072	.059 .063	.025 .031	.049 .066	.046 .079
October	.068 .162	.100 .151	.080 .082	.075 .081	.059 .069	.049 .059	.087 .115	.061 .129
November	.062 .163	.091 .136	.070 .073	.066 .070	.059 .065	.063 .069	.087 .103	.061 .147
December	.083 .198	.094 .146	.082 .088	.078 .083	.075 .084	.076 .083	.075 .100	.105 .192

Exhibit 3.1

Values of RMS for "constants Model" for 30 stations in Germany using non-linear regression (top entry) and weighted least square (bottom entry).

WMO	Station	Lat (North)	Long (East)	Non- Linear RMS	Weighted L.S. RMS
10501	Aachen, DL	50.78	6.12	.081	.169
10224	Bremen, GER	53.05	8.80	.048	.172
10305	Lingen, GER	52.52	7.33	.127	.168
10427	Kahler Asten, GER	51.18	8.50	.274	.178
11157	Aigen Ennstal, OS	47.53	14.15	.101	.202
11448	Plezen/Dobra, CZ	49.67	13.30	.145	.159
10900	Bremgarten, GER	47.90	7.63	.050	.164
10948	Oberstdorf, GER	47.40	10.30	.147	.198
10929	Konstanz, GER	47.68	9.20	.051	.170
11120	Innsbruck, OS	47.27	11.37	.081	.198
11150	Salzburg, OS	47.80	13.02	.059	.181
10893	Passau, GER	48.58	13.50	.066	.157
10908	Feldberg, GER	47.87	8.02	.321	.217
10921	Neuhausen, DL	47.98	8.92	.108	.146
10512	Nurburg, GER	50.33	6.97	.134	.108
10515	Koblenz, DL	50.35	7.60	.102	.191
10532	Giessen, GER	50.57	8.72	.076	.194
10542	Hersfeld, DL	50.87	9.72	.135	.180
10658	Kissinlen, DL	50.20	10.10	.126	.179
10671	Coburg, GER	50.27	10.97	.123	.171
10685	Hof, GER	50.32	11.90	.067	.140
10704	Berus, GER	49.27	6.70	.125	.160
10727	Karlsruhe, GER	49.02	8.40	.072	.178
10742	Ohringen, GER	49.20	9.53	.092	.174
10788	Staubing, GER	48.82	12.60	.112	.185
10791	Grosser Falk, GER	49.08	13.30	.275	.161
10805	Lahr, GER	48.37	7.85	.061	.157
10837	Laudheim, GER	48.22	9.93	.094	.159
10953	Kaufbeuren, DL	47.87	10.63	.065	.166
10875	Muhldorf, GER	48.25	12.55	.091	.164
10616	Hahn, AB	49.95	7.27	.048	.147
10610	Bitburg, AB	49.95	6.57	.056	.172
10614	Ramstein, AB	49.43	7.58	.064	.184
10607	Spangdahlem, AB	49.97	6.70	.067	.177
10384	Tempelhof, APRT	52.47	13.40	.074	.192
10755	Ansbach, AAF	49.32	10.63	.071	.174
10544	Fulda, AAF	50.53	9.63	.062	.171
10869	Erding, AS	48.32	11.93	.060	.177
10765	Feucht, AAF	49.38	11.18	.055	.171
10618	Baumholder, AAF	49.65	7.30	.176	.199
10626	Bad Kreuznach, AAF	49.87	7.88	.111	.189
10971	Bad Tolz, AAF	47.77	11.60	.123	.200
10714	Zweibrucken, AB	49.22	7.40	.046	.169
10633	Wiesbaden, AB	50.05	8.33	.050	.165
10633	Finthen, AAF	49.97	8.15	.057	.160
10763	Furth, AAF	49.50	10.95	.090	.187
10642	Hanau, AAF	50.17	8.95	.046	.171

(continued)

WMO	Station	Lat (North)	Long (East)	Non- Linear RMS	Weighted L.S. RMS
10852	Gablingen, AAF	48.45	10.87	.109	.192
10653	Giebelstadt, AUX AF	49.67	9.88	.082	.186
10687	Grafenwohr, AAF	49.70	11.95	.073	.169
10734	Heidelberg, AAF	49.40	8.64	.041	.167
10752	Illesheim, AAF	49.47	10.38	.149	.205
10659	Kitzingen, AAF	49.75	10.20	.066	.189
10763	Nurnberg	49.50	11.08	.066	.191
10729	Coleman, AAF	49.57	8.47	.114	.182
10657	Wertheim, AAF	49.77	9.48	.153	.177
10745	Schwaebisch Hall, AA	49.17	9.78	.097	.179
10712	Sembach, AB	49.52	7.87	.037	.169
10862	Siegenberg Gunnery	48.75	11.80	.177	.219
10738	Echterdingen, ARPT	48.68	9.22	.057	.183
	Overall			<u>.098</u>	<u>.177</u>

Exhibit 3.2

Overall RMS for Each of 60 Stations for the Constants Model Utilizing Sample
Reuse Based on Non-Linear Regression and Weighted Least Squares

	Hour Period (LST)						
	06-08	09-11	12-14	15-17	18-20	21-23	all
January	.126 .156	.119 .144	.114 .159	.113 .161	.124 .169	.166 .210	.128 .167
February	.117 .147	.105 .137	.099 .159	.099 .167	.111 .175	.152 .217	.115 .167
March	.103 .145	.097 .160	.087 .198	.083 .205	.094 .211	.116 .253	.097 .195
April	.101 .176	.089 .214	.074 .226	.072 .226	.080 .235	.089 .238	.085 .219
May	.093 .167	.076 .225	.060 .209	.051 .183	.060 .201	.077 .227	.071 .202
June	.094 .171	.073 .212	.054 .184	.045 .159	.056 .182	.070 .190	.067 .183
July	.088 .164	.066 .194	.049 .170	.039 .137	.047 .146	.065 .177	.061 .165
August	.093 .126	.070 .156	.051 .159	.046 .155	.055 .167	.080 .202	.068 .161
September	.106 .123	.085 .131	.064 .163	.053 .161	.074 .190	.097 .212	.082 .163
October	.100 .107	.097 .112	.082 .142	.078 .157	.100 .157	.134 .176	.100 .142
November	.112 .137	.108 .140	.104 .174	.106 .175	.121 .181	.151 .206	.118 .169
December	.113 .143	.107 .136	.106 .153	.108 .151	.115 .156	.167 .207	.121 .158
all	.104 .147	.092 .161	.082 .175	.079 .170	.090 .181	.119 .210	.096 .174

Exhibit 3.3

RMS Over All 60 Stations by Month and Hour Period for the Constants Model Utilizing Sample Reuse Based on Non-Linear Regression (top entry) and Weighted Least Squares (bottom entry).

WMO	Station	Lat (North)	Long (East)	Elev. (ft.)	Rel. Elev. (ft.)	Non-Linear			Weighted Least		
						Regression			Squares		
						Sample Reuse RMS	Overall Fit RMS	Overall Fit RMS	Sample Reuse RMS	Overall Fit RMS	Overall Fit RMS
10616	Hahn, AB	49.95	7.27	1650.	1267.	.062	.048		.172	.070	
10610	Bitburg, AB	49.95	6.57	1228.	1363.	.044	.042		.092	.091	
10614	Ramstein, AB	49.43	7.58	780.	710.	.051	.052		.109	.108	
10607	Spangdahlem, AB	49.97	6.70	1196.	1330.	.057	.055		.101	.100	
10384	Tempelhof, APRT	52.47	13.40	164.	164.	.058	.058		.115	.114	
10755	Ansbach, AAF	49.32	10.63	1542.	1181.	.068	.064		.100	.099	
10544	Fulda, AAF	50.53	9.63	1010.	1650.	.045	.046		.085	.085	
10869	Erding, AS	48.32	11.93	1522.	1525.	.064	.062		.098	.097	
10765	Feucht, AAF	49.38	11.18	1265.	1407.	.046	.045		.086	.086	
10618	Baumholder, AAF	49.65	7.30	1408.	1538.	.153	.154		.156	.153	
10626	Bad Kreuznach, AAF	49.87	7.88	355.	825.	.071	.071		.120	.119	
10971	Bad Tölz, AAF	47.77	11.60	2360.	2688.	.121	.102		.120	.117	
10714	Zweibrücken, AB	49.22	7.40	1132.	938.	.041	.040		.091	.090	
10633	Wiesbaden, AB	50.05	8.33	470.	617.	.036	.036		.083	.082	
10633	Finthen, AAF	49.97	8.15	769.	948.	.048	.047		.082	.081	
10763	Furth, AAF	49.50	10.95	1000.	1225.	.085	.085		.112	.111	
10642	Hanau, AAF	50.17	8.95	377.	408.	.037	.036		.089	.087	
10852	Gablingen, AAF	48.45	10.87	1530.	1550.	.084	.082		.120	.120	
10653	Giebelstadt, AUX AF	49.67	9.88	985.	950.	.060	.060		.112	.111	
10687	Grafenwohr, AAF	49.70	11.95	1370.	1510.	.063	.066		.088	.087	
10734	Heidelberg, AAF	49.40	8.64	369.	367.	.034	.034		.086	.084	
10752	Illesheim, AAF	49.47	10.38	1060.	1250.	.084	.084		.135	.135	
10659	Kitzingen, AAF	49.75	10.20	699.	899.	.051	.052		.112	.111	
10763	Nurnberg	49.50	11.08	1053.	1053.	.062	.063		.114	.113	
10729	Coleman, AAF	49.57	8.47	334.	350.	.078	.077		.111	.110	
10657	Wertheim, AAF	49.77	9.48	1120.	1075.	.095	.096		.167	.116	
10745	Schwaebisch Hall, AA	49.17	9.78	1303.	1632.	.046	.046		.093	.092	
10712	Sembach, AB	49.52	7.87	1052.	1315.	.037	.037		.082	.082	
10862	Siegenberg Gunnery	48.75	11.80	1325.	1450.	.140	.140		.164	.163	
10738	Echterdingen, ARPT	48.68	9.22	1306.	1200.	.056	.054		.105	.105	
10501	Aachen, DL	50.78	6.12	673.	755.	.081	.080		.099	.097	
10224	Bremen, GER	53.05	8.80	502.	250.	.043	.043		.091	.090	

(continued)

WMO	Station	Lat (North)	Long (East)	Elev. (ft.)	Rel. Elev. (ft.)	Non-Linear Regression			Weighted Least Squares		
						Sample Reuse RMS	Overall Fit RMS	Overall Fit RMS	Sample Reuse RMS	Overall Fit RMS	
10305	Lingen, GER	52.52	7.33	549.	250.	.040	.040	.040	.086	.085	
10427	Kahler Asten, GER	51.18	8.50	2822.	1563.	.204	.188	.188	.207	.174	
11157	Aigen Ennstal, OS	47.53	14.15	2139.	3350.	.056	.053	.053	.110	.097	
11448	Plezen/Dobra, CZ	49.67	13.30	1194.	1400.	.130	.131	.131	.109	.105	
10900	Bremgarten, GER	47.90	7.63	699.	1438.	.044	.041	.041	.077	.076	
10948	Oberstdorf, GER	47.40	10.30	2664.	3350.	.097	.080	.080	.157	.102	
10929	Konstanz, GER	47.68	9.20	1368.	1725.	.037	.036	.036	.082	.081	
11120	Innsbruck, OS	47.27	11.37	1962.	3400.	.069	.074	.074	.102	.089	
11150	Salzburg, OS	47.80	13.02	1463.	2338.	.048	.047	.047	.080	.078	
10893	Passau, GER	48.58	13.50	1335.	1613.	.056	.055	.055	.075	.074	
10908	Feldberg, GER	47.87	8.02	4898.	2550.	.280	.069	.069	.355	.078	
10921	Neuhausen, DL	47.98	8.92	2648.	2463.	.072	.063	.063	.087	.081	
10512	Nurburg, GER	50.33	6.97	2064.	1425.	.085	.077	.077	.059	.056	
10515	Koblentz, DL	50.35	7.60	318.	900.	.066	.068	.068	.116	.115	
10532	Giessen, GER	50.57	8.72	640.	825.	.059	.060	.060	.119	.118	
10542	Hersfeld, DL	50.87	9.72	738.	1200.	.059	.060	.060	.099	.098	
10658	Kissinlen, DL	50.20	10.10	704.	1215.	.039	.040	.040	.095	.094	
10671	Coburg, GER	50.27	10.97	1106.	1338.	.038	.039	.039	.085	.084	
10685	Hof, GER	50.32	11.90	1864.	1938.	.048	.057	.057	.064	.062	
10704	Berus, GER	49.27	6.70	1204.	1050.	.056	.054	.054	.088	.087	
10727	Karlsruhe, GER	49.02	8.40	394.	600.	.041	.041	.041	.099	.098	
10742	Ohringen, GER	49.20	9.53	909.	1075.	.049	.049	.049	.092	.091	
10788	Staubing, GER	48.82	12.60	1155.	1350.	.084	.085	.085	.117	.116	
10791	Grosser Falk, GER	49.08	13.30	4291.	2475.	.259	.064	.064	.098	.058	
10805	Lahr, GER	48.37	7.85	509.	800.	.074	.073	.073	.077	.075	
10837	Laudheim, GER	48.22	9.93	1765.	1938.	.053	.056	.056	.074	.073	
10953	Kaufbeuren, DL	47.87	10.63	2388.	2513.	.069	.062	.062	.080	.076	
10875	Muhlendorf, GER	48.25	12.55	1319.	1625.	.053	.053	.053	.077	.076	

Exhibit 3.4

RMS of Sample Re-use and Overall Fit RMS by Station Over all Month and Hour Periods 3 through 8 for the Variables Model Based on Non-Linear Regression and Weighted Least Squares

Hour Period
(LST)

Month	06-08	09-11	12-14	15-17	18-20	21-23	all
January	.106 .123	.089 .111	.086 .103	.083 .105	.100 .121	.140 .169	.104 .122
February	.088 .104	.074 .087	.061 .080	.060 .081	.079 .102	.120 .155	.083 .102
March	.075 .092	.066 .086	.056 .085	.049 .080	.060 .095	.085 .140	.066 .096
April	.069 .089	.058 .081	.041 .076	.035 .078	.042 .088	.053 .121	.051 .089
May	.068 .092	.048 .094	.031 .072	.026 .066	.030 .086	.048 .122	.044 .089
June	.070 .093	.048 .088	.031 .087	.026 .064	.032 .080	.047 .105	.095 .086
July	.071 .099	.046 .082	.029 .065	.023 .052	.030 .067	.047 .103	.044 .078
August	.082 .098	.053 .067	.036 .063	.028 .043	.035 .062	.063 .170	.053 .084
September	.095 .109	.063 .075	.050 .104	.036 .038	.054 .089	.073 .115	.065 .088
October	.087 .094	.071 .078	.057 .074	.052 .074	.081 .106	.111 .143	.079 .095
November	.082 .094	.076 .087	.064 .086	.065 .088	.087 .105	.115 .143	.083 .101
December	.089 .106	.087 .101	.079 .102	.080 .103	.090 .112	.142 .171	.097 .116
All	.083 .099	.068 .086	.055 .083	.051 .073	.065 .093	.094 .138	.071 .095

Exhibit 3.5

RMS by Month and Hour for Overall Fit of All 60 Germany Stations Using the Variables Model by Non-Linear Regression (top entry) and Weighted Least Squares (bottom entry).

Hour Period
(LST)

Month	06-08	09-11	12-14	15-17	18-20	21-23	a11
January	.118 .130	.111 .119	.100 .113	.099 .114	.115 .131	.150 .176	.117 .131
February	.102 .113	.089 .096	.081 .094	.081 .094	.097 .112	.132 .162	.097 .112
March	.092 .101	.086 .098	.079 .099	.074 .094	.084 .106	.098 .149	.086 .112
April	.089 .099	.078 .095	.066 .090	.064 .086	.072 .094	.080 .128	.075 .099
May	.085 .108	.060 .124	.040 .094	.038 .078	.052 .104	.069 .136	.060 .107
June	.085 .106	.058 .114	.041 .120	.046 .088	.057 .100	.074 .122	.062 .108
July	.087 .115	.055 .113	.036 .096	.041 .071	.056 .086	.074 .123	.061 .101
August	.099 .106	.069 .091	.056 .105	.050 .062	.062 .078	.088 .172	.073 .102
September	.105 .118	.079 .096	.067 .138	.060 .085	.079 .107	.097 .126	.083 .112
October	.096 .102	.085 .084	.078 .094	.078 .091	.102 .117	.126 .150	.096 .107
November	.094 .104	.090 .097	.074 .099	.083 .100	.102 .114	.125 .150	.096 .111
December	.104 .117	.102 .114	.096 .116	.097 .115	.107 .124	.151 .179	.111 .128
A11	.097 .110	.082 .104	.071 .105	.070 .090	.085 .106	.109 .148	.087 .111

Exhibit 3.6

RMS by Month and Hour for Sample Reuse of A11 60 German Stations Using the Variables Model by Non-Linear Regression (top entry) and Weighted Least Squares (bottom entry).

END

FILMED

8-84

DTIC