

AD-R143 995

MECHANISM OF ACTIVATION OF LYMPHOCYTES-SYNTHESIS OF
REGULATORY SUBSTANCES(U) NORTHEASTERN UNIV BOSTON MASS
DEPT OF BIOLOGY P R STRAUSS 01 OCT 83 N00014-82-K-0283


1/1

UNCLASSIFIED

F/G 6/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A



REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM				
1. REPORT NUMBER 1	2. GOVT ACCESSION NO. AD-A143995	3. RECIPIENT'S CATALOG NUMBER				
4. TITLE (and Subtitle) Mechanism of Activation of Lymphocytes- Synthesis of Regulatory Substances		5. TYPE OF REPORT & PERIOD COVERED Annual Report 6/1/83 - 10/1/83				
		6. PERFORMING ORG. REPORT NUMBER				
7. AUTHOR(s) Phyllis R. Strauss		8. CONTRACT OR GRANT NUMBER(s) N00014-82-K-0283				
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Biology Northeastern University 360 Huntington Avenue, Boston, MA 02115		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 666-005				
11. CONTROLLING OFFICE NAME AND ADDRESS Jeannine A. Majde, Ph.D., Scientific Officer Immunology Code 441, Cellular Biosystems Group, Dept. of the Navy, ONR, Arlington, VA 22217		12. REPORT DATE 10/1/83				
		13. NUMBER OF PAGES 12				
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified				
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE				
16. DISTRIBUTION STATEMENT (of this Report) Unlimited						
<table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td colspan="2" style="text-align: center;">DISTRIBUTION STATEMENT</td> </tr> <tr> <td colspan="2" style="text-align: center;">Approved for public release; Distribution Unlimited</td> </tr> </table>			DISTRIBUTION STATEMENT		Approved for public release; Distribution Unlimited	
DISTRIBUTION STATEMENT						
Approved for public release; Distribution Unlimited						
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)						
18. SUPPLEMENTARY NOTES						
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) lymphocytes, DNA, thymidine, replication concanavalin A						
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This report describes further progress in characterizing detergent soluble DNA from activated murine splenocytes and lymphocyte cell lines. It also describes experiments implicating the involvement of DNA replication in detergent soluble DNA synthesis and disproves the assumption that detergent soluble DNA is a random degradation process.						

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Dist	Avail and/or Special



DTIC
ELECTE
S AUG 9 1984 D

FILE COPY

SUMMARY OF RESEARCH PROGRESS
June 1983-October 1983

Phyllis R. Strauss
021-30-7461

The purpose of this contract is to characterize cytoplasm-associated DNA from murine splenocytes and to determine its physiological function. Throughout the report, I will refer to this DNA as "MADS" DNA, metabolically active detergent soluble DNA.

I. Characterization

- A. The manuscript entitled "Characterization of rapidly-labeled detergent soluble DNA in murine lymphocytes" has been accepted for publication by Biochemistry (Washington). A copy is enclosed as Appendix I.

- B. The question of whether each size class in MADS DNA is a unique sequence or contains common sequences with other size classes has moved more slowly than anticipated. Our initial attempts to clone the material 'shotgun' (that is, try it and see what happens) were not successful. Therefore, we needed a restriction analysis before proceeding. When Dr. Harry Meade (Biogen Corp, Cambridge, MA) was not forthcoming with a restriction map, we decided to invest in the required restriction endonucleases. The experiment was designed to maximize sensitivity and minimize the amount of MADS DNA needed by labeling either whole MADS DNA or one of its fragments with ^{32}P -phosphate using the nick translation system. The ^{32}P labeled material was exposed to each of a battery of restriction endonucleases and size changes were observed by autoradiography

after electrophoresis on agarose gels. Although the nick translated starting products of MADS DNA had themselves become smaller than native MADS DNA or its fragment, no difficulty was encountered with ϕ X174RF/Hae III control DNA. Despite this problem, nick translated MADS DNA fragments were treated with Eco RI, Acc, HindIII, SmaI, HincII, BglI, SauIIIa, MboI, HpaII, ClaI, HaeIII, PstI, MspI, Kpn, AluI, AvaI, BalI and BclI. Only HaeIII resulted in cleavage. The result with HaeIII was expected as it had been observed earlier and reported in the manuscript appended. However, the lack of effect of the array of other restriction endonucleases was not expected, especially in light of the presence in MADS DNA of a variety of DNA sequences including those of globin β chain and immunoglobulin κ chain. In evaluating these results we decided to investigate three additional restriction endonucleases because these are known to cleave mouse satellite DNA. The three, AvaII, EcoRII, and Sau96I, cleaved MADS DNA, the cleavage pattern with AvaII^{being} similar to that of HaeIII and different from Sau96I and EcoRII. Although we could not draw conclusions about the similarity of MADS DNA and mouse satellite DNA, all the enzymes which cleaved MADS DNA contained GGCC sequence specificity. Thus, it is significant that we initially reported a slightly higher GC content in MADS DNA than in bulk chromosomal DNA. Now that we have cleavage sites, we can generate blunt ends on MADS DNA fragments, attach linkers and insert the DNA into cloning vehicles.

- C. Pulse labeled MADS DNA contains very rapidly labeled sequences smaller than 100 base pairs. These were end labeled at the 5'

terminus with ^{32}P -phosphate using polynucleotide kinase and analyzed on Maxam-Gilbert denaturing gels (Maniatis et al., Molecular Cloning, Cold Spring Harbor Press, 1982). They were a discrete size between 10 and 14 base pairs in length. Since they were so short, it should be possible to sequence them by chemical methods. Sequencing experiments are in progress.

End labeling with polynucleotide kinase did not result in labeling size classes larger than the 8-12 mer. This observation has held even when labeling was performed under conditions for recessed 5' ends (3' overhang) or exchange conditions. We have not yet exhausted all possibilities, however, since before a new 5' phosphate can be added, the original 5' unlabeled phosphate must be removed with either calf or bacterial alkaline phosphatase. The conditions for removing 5' phosphate in the presence of a 3' overhang are far more rigorous than those we have previously employed. Experiments are underway towards this end. If we are still unable to label at the 5' end, it will be necessary to explore alternative reasons such as covalent linkage of other molecules either protein or RNA. Because we have had difficulty in end labeling at the 5' terminus, we end labeled MADS DNA at the 3' terminus using terminal deoxytransferase and cordycepin 5' triphosphate. Although better results were obtained, not all species visualized by ethidium bromide staining patterns were labeled. We consider the possibility that a 3' overhang is folded back onto itself in most MADS species.

D. Electron microscopy is a useful tool in analyzing unusual DNAs because it allows the investigator to examine single pieces for irregularities. We are currently examining MADS DNA and its isolated fragments under native and denaturing conditions. Although quantitative measurements have not yet been made, it is clear that larger fragments in MADS DNA contain bubbles and forks in what appears to be a regular fashion. This would confirm the predictions made in I-B, that MADS DNA contains single stranded regions.

E. Early attempts to determine which lymphocyte class (T or B) contained MADS DNA by means of differential sensitivity to immune sera were unsuccessful because controls treated with complement in the presence of calcium were also depleted of ³H-thymidine labeled MADS DNA. This was not true if calcium was not present in the medium. Divalent cations are notorious requirements for both nucleases and some proteases. The nuclease problem is insoluble because there are no good inhibitors of nucleases other than chelating divalent cations with EDTA or EGTA. The protease problem could be explored directly because there are several good protease inhibitors such as phenylmethylsulfonylflouride (PMSF) available, and amino acids covalently linked to viruses and mitochondrial DNA have been recently reported.

To determined whether MADS DNA contains covalently linked protein, NP40 supernates were prepared in the standard fashion except that PMSF was added as a protease inhibitor. MADS DNA was isolated as usual except that the initial step employing proteinase K was omitted. The final preparation was divided in half and one half was

subjected to treatment with proteinase. Both halves were analyzed by agarose gels under native conditions. The portion which was not treated with proteinase K behaved exactly like the standard preparation. Therefore, we conclude that MADS DNA does not contain a long chain of covalently linked amino acids sufficient to interfere with its mobility on agarose gels.

II. Function

- A. In the previous progress report we described a series of pulse chase experiments in which a small fragment incorporated ^3H -thymidine rapidly (15min) and could be chased uniformly into larger size classes thereafter. The small fragment has been further characterized (see I-C). We have now shown that two inhibitors of replication, aphidicolin and hydroxyurea, did not inhibit the rapid labeling of the small fragment. (See Figures 1 and 2) Both inhibitors, however, prevented the transfer of radiolabeled thymidine to MADS DNA. Therefore, the transfer process was associated with replication. In order to interpret these results, we have examined what happens to ^3H -thymidine which is incorporated into the detergent insoluble fraction and which presumably represents chromatin. No rapidly labeled oligonucleotide was found in the detergent insoluble fraction at any time. (See Figure 3) Moreover, all incorporated ^3H -thymidine was associated with DNA of higher molecular weight than that found in MADS DNA. A manuscript describing these findings is in preparation.
- B. The presence of MADS DNA in several cell lines is under investigation. Jurkat human lymphoma cells continue to synthesize

and label MADS DNA after they have been exposed to Concanalin A or phorbol myristate acetate plus poke weed mitogen, which are reported to stimulate IL-2 production. While we have not measured IL-2 production ourselves, we have observed that the cells aggregated and were very difficult to handle within 12-24 hrs.

Studies on cell cycle dependence in Jurkat cells are also in progress. We have developed a method to separate Jurkat cells on the basis of size by means of Nycodenz gradients. Nycodenz is a new polymer similar to Ficoll but less toxic. The separation appeared to correlate with stages in the cell cycle as analyzed by cell cycle analysis. Cells which had been labeled with ^3H -thymidine prior to separation have been analyzed for the distribution of incorporated nucleoside into detergent soluble and insoluble fractions. A striking increase in the ratio of MADS DNA to chromosomal DNA (detergent insoluble DNA) is seen about one third of the way down the gradient. These experiments are being repeated now and a manuscript will be in preparation shortly.

Further studies on function require a well defined cell line where stimulators and modulators of replication and transcription are known (See below).




Figure 1: Cells from Con A stimulated mice were incubated in buffer, 0.5% (v/v) DMSO, or DMSO containing 10 μ g/ml aphidicolin for 15 min. ³H-TdR was added to 6 μ Ci/ml. After a 15 min pulse with ³H-TdR the cells were washed and resuspended in medium containing DMSO or DMSO with aphidicolin for a chase period of 60 min. At the indicated time they were washed and detergent lysed. MADS DNA was isolated from the detergent soluble fraction and analyzed by agarose gel electrophoresis (1.2% gel). -7-




Figure 2: Cells from Con A stimulated mice were incubated in the presence or absence of 2mM hydroxyurea for 30 min at which time $^3\text{H-TdR}$ was added to 5 $\mu\text{Ci}/\text{ml}$. After a 15 min pulse, cells were washed and resuspended in medium with or without hydroxyurea for a 60 min chase. They were then washed and detergent lysed. MADS DNA was purified from the detergent soluble fraction and analyzed by gel electrophoresis (1.2% agarose).




Figure 3: DNA from detergent insoluble pellets from the experiment described in Figure 1 was purified and analyzed by agarose gel electrophoresis.

PROPOSED EXPERIMENTS
March 1984-February 1985

The following experiments are proposed in addition to continuing the work described above.

- A. We intend to sequence several purified fragments of MADS DNA from lymphocytes obtained from conA stimulated mice to determine whether there are consensus sequences (rationale described above in IB)

- B. Whenever a person versed in the chromatin field sees the MADS DNA results he/she concludes that MADS DNA is nucleosomal in origin on the basis of the size distributions alone. If MADS DNA is nucleosomal, it must represent a special class of nucleosomes since nucleosomes are generally considered to be detergent insoluble. In fact detergent lysis is often used to prepare the nuclei from which nucleosomes are obtained (see discussion in Strauss et al, In press). I propose to centrifuge the NP40 supernate prepared in the presence of PMSF on linear sucrose gradients. The fractions will then be analyzed for MADS DNA and histones. If MADS DNA is part of a nucleosome complex, histones must be present in the correct molar ratios. I also propose to perform a thorough study of the effects of a graded series of detergents on nucleosomes and on "newly synthesized" and "old" chromatin. These studies should answer the question once and for all as to whether MADS DNA is nucleosomal in origin. At this time we have run preliminary sucrose gradients and worked out polyacrylamide gel systems for analyzing histones.

- C. The pulse chase studies described in IIA were performed at intervals no shorter than 15 min. In order to study replication intermediates we must reduce the pulse interval to 30 sec. Since a half minute pulse is insufficient to get enough radiolabel into intact cells, we intend to use permeabilized cells as described by Kowalski and Denhardt (Nucleic Acids Res. 5: 4355-4373). The pulsed cells will be followed to determine the fate of radiolabeled materials. We will be sure to collect and analyze the chase supernate because we would expect newly synthesized MADS DNA to be liberated into the supernate.
- D. As described in the PROGRESS II-B section, we need to capitalize on a cell line in which replication and transcription are already well studied and can be manipulated. A search for lymphocyte lines has proved fruitless. As noted by the Contract Officer and confirmed by Dr. Emil Unanue (Harvard Medical School Department of Pathology), lymphocyte lines are already committed to an immunological function or unable to be committed at all. The lack of a manipulatable line has seriously hampered the understanding of the immune process, according to Dr. Unanue. Friend's erythroleukemia line offers an attractive alternative. For a recent review see Ann. Rev. Biochem 47: 419-448, 1978. We propose to explore 1) whether MADS DNA is present in Friend's erythroleukemia cells and 2) whether MADS DNA is altered after the cell line is exposed to agents which promote division leading to differentiation to the globin producing state. If both 1 and 2 hold, we can use probes to determine whether MADS DNA is enriched (depleted?) in globin sequences upon stimulation.

We can also pull the globin sequences out of MADS DNA and sequence them in order to determine whether they are normal or scrambled. For instance, Mark Rush (personal communication) has shown that non-chromosomal small polydisperse circular DNA from African green monkey kidney cells contains at least one Alu sequence which is spliced end to end.

DISTRIBUTION LIST

Immunological Defense Program

Annual, Final and Technical Reports (one copy each except as noted)

Dr. John D. Clements
Department of Microbiology
and Immunology
Tulane University Medical Center
1430 Tulane Avenue
New Orleans, LA 70112

Francis A. Ennis, M.D.
University of Massachusetts
Medical School
55 Lake Avenue
Worcester, MA 01605

Dr. Edward A. Havell
Trudeau Institute
P.O. Box 59
Saranac Lake, NY 12983

Fred D. Finkelman, M.D.
Department of Medicine
Uniformed Services University
of the Health Sciences
4301 Jones Bridge Road
Bethesda, MD 20814

Dr. Arthur G. Johnson
Department of Medical
Microbiology and Immunology
University of Minnesota
School of Medicine
2205 East 5th Street
Duluth, MN 55812

Dr. Philip Lake
Immunologic Oncology Division
Lombardy Cancer Center
Georgetown University School of Med.
Washington, DC 20007

Dr. Janice Longstreth
Director of Immunology/Virology
Borrison Laboratories, Inc.
5050 Beech Place
Temple Hills, MD 20748

Dr. Ernest D. Marquez
Bioassay Systems Corporation
225 Wildwood Avenue
Woburn, MA 01801

James J. Mond, M.D.
Department of Medicine
Uniformed Services University
of the Health Sciences
4301 Jones Bridge Road
Bethesda, MD 20814

Dr. Page S. Morahan
Department of Microbiology
Medical College of Pennsylvania
3300 Henry Avenue
Philadelphia, PA 19129

Zoltan Ovary, M.D.
Department of Pathology
New York University
School of Medicine
550 First Avenue
New York, NY 10016

Dr. Donna G. Sieckmann
Infectious Diseases Program Ctr.
Naval Medical Research Institute
National Naval Medical Center
Bethesda, MD 20814

David A. Stevens, M.D.
Department of Medicine
Santa Clara Valley Medical Ctr.
751 S. Bascom Avenue
San Jose, CA 95128

Dr. Phyllis R. Strauss
Department of Biology
Northeastern University
360 Huntington Avenue
Boston, MA 02115

Dr. Alvin L. Winters
Department of Microbiology
University of Alabama
University, AL 35486

Lyn Yaffe, M.D.
Research Support Center
Naval Medical Research Institute
National Naval Medical Center
Bethesda, MD 20814

Annual, Final and Technical Reports (con't)

Dr. Jeannine A. Majde, Code 441CB
Office of Naval Research
800 N. Quincy Street
Arlington, VA 22217

Defense Technical Information Center (2 copies)
Building 5, Cameron Station
Alexandria, VA 22314

Annual and Final Reports Only (one copy each)

Scientific Library
Naval Biosciences Laboratory
Naval Supply Center
Oakland, CA 94625

Commanding Officer
Naval Medical Research & Development Command
National Naval Medical Center
Bethesda, MD 20814

Director
Infectious Disease Program Center
Naval Medical Research Institute
National Naval Medical Center
Bethesda, MD 20814

Commander
Chemical and Biological Sciences Division
Army Research Office
Research Triangle Park, NC 27709

Commander
U.S. Army Research and Development Command
Attn: SGRD-PLA
Fort Detrick
Frederick, MD 21701

Commander
USAMRIID
Fort Detrick
Frederick, MD 21701

Commander
Air Force Office of Scientific Research
Bolling Air Force Base
Washington, DC 20332

Administrative Contracting Officer
Office of Naval Research Residency
address varies - obtain from Business Office)

Final and Technical Reports Only

Director, Naval Research Laboratory (6 copies)
Attn: Code 2627
Washington, DC 20375

END

FILMED