

AD-A147 062

OCEAN ACOUSTICS(U) NEW YORK UNIV NY COURANT INST OF
MATHEMATICAL SCIENCES D STICKLER OCT 84
N00014-75-C-0218

1/3

UNCLASSIFIED

F/G 20/1

NL




END

DATE

FILED

11 84

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

12

REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS BEFORE COMPLETING FORM

1. REPORT NUMBER		2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Ocean Acoustics		5. TYPE OF REPORT & PERIOD COVERED Final Technical Report	
7. AUTHOR(s) Dr. David Stickler		6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS COURANT INSTITUTE, NEW YORK UNIVERSITY 251 Mercer Street New York, NY 10012		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0218	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy 800 N. Quincy Street (613 C:DAW), Arlington, VA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 386-968/11-26-82 (420)	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Office of Naval Research 715 Broadway (5th Floor) New York, NY 10003		12. REPORT DATE October 1984	
		13. NUMBER OF PAGES 5	
		15. SECURITY CLASS. (of this report) Unclassified	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public release-distribution unlimited			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Inverse scattering, ocean sediments and basement trace method, continuous spectrum, ray acoustics, smooth caustics.			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The main element of this final report is a discussion of the development of the trace method for determining the acoustic properties of the ocean bottom sediments and basement. In addition a method is reported for the uniform determination of the continuous modal contribution in a normal mode expansion for the second term in the geometrical acoustic expansion near a smooth caustic. Originator-supplied keywords include:			

DTIC ELECTE
NOV 5 1984
S D D

AD-A147 062

DTIC FILE COPY

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

(When Data Entered)

BLANK PAGE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

The work on this contract has focussed on developing a method to determine the acoustic parameters of the ocean sediments and basement. The methods are indirect in that they attempt to determine the properties (density and sound speeds as a function of depth) from measurement of the field reflected from the ocean bottom.

The initial paper^[1] in this series shows how to use the methods of one-dimensional inverse scattering theory, in particular the new approach of Deift and Trubowitz [Comm. Pure Appl. Math. XXXII, 121-251 (1979)], to recover the sound speed for the case of a stratified ocean with a pressure release surface driven by a harmonic point source at a finite depth from a measurement of the normal derivative at the pressure release surface. The method is not a perturbation technique. For comparison, this same problem is examined when the sound speed is slightly perturbed from unity using a technique of Cohen and Bleistein [SIAM J. App. Math. 32, 784-799 (1977) and Geophysics 44, 1077-1087 (1979)]. The trace method of Deift and Trubowitz was introduced to examine the quantum mechanical problem where it was reasonable to assume that the sound speed approached the same known value at depth and in the ocean. This is in general not the case, but their method was extended to the "geophysical" acoustic case in the next paper^[2].

In this analysis the following problem is considered: A point harmonic source at frequencies ω_1 and ω_2 is located in the region $z < 0$, where the assumption is made that both the sound speed and density profiles are known. For $z > 0$ the the sound speed and density profiles are unknown. These two profiles are recovered from a measurement of the pressure field for all r at some fixed depth $z < 0$ at the two frequencies. Trace formula methods are used. The following assumptions are needed: First, $c(z)$ and $\rho(z)$ approach c_1, ρ_1 as $z \rightarrow -\infty$, and c_2, ρ_2 as $z \rightarrow +\infty$, and although c_2, ρ_2 are not known, it must be known that $c_2 > c_1$. Second, the angular frequency of the source must be such that no trapped modes are excited. While the sound speed $c(z)$ can be complex, the limiting values c_1 and c_2 must be real. If $c(z)$ also depends on frequency and the form of the dependence is known, say, $c(z) = c_R(z) + i\omega^\lambda(z)c_I(z)$, then $c_R(z), c_I(z)$, and $\lambda(z)$ can be recovered.

Explicit expressions are obtained in the realistic case when $c_1(z)$ is not large. If a density is known and $c_1(z) = 0$, then only a measurement at one frequency is required. Two numerical examples are given; in both examples $\rho(z) \approx 1$ and $c(z)$ is real. The first example is for a monotonically increasing profile and the second has a low velocity zone. Typical sediment parameters are used.

When the ocean bottom is modeled as a fluid, then only the sound speed and density need to be recovered. But if the sediments and basement support shear waves; then they must be modeled elastically. This means that two wave speeds and the density must be recovered. Before this problem was examined the trace method was applied to a simpler set of equations which nonetheless supported N-different types of waves, rather than just two.[3]

Trace formulas are derived for the coefficients of a matrix differential equation representing wave propagation in a medium which supports N types of waves. In general these waves are coupled to each other but are assumed to become uncoupled as $|z| \rightarrow \infty$, but unknown for $z > 0$. A point source is located in the region $z < 0$ and the point source field is measured in the same region. These measurements yield a reflection coefficient which is then used in the trace formulas, derived in the paper, to recover the two matrix coefficients for $z > 0$. The unique features of this work are a derivation of the trace formula in which the Jost functions are not needed, an alternate measurement method based on an impedance concept and a trace equation based on an impedance concept.

These results have been presented as invited talks at the

(1) Spring 1983 meeting of the Acoustical Society of America.

(2) Spring 1983 meeting of the Society of Photo-Optical Instrumentation Engineers.

This talk also appeared as a paper in their journal^[4] and indicates that the method can be used in contexts other than acoustics.

(3) Spring 1983, American Mathematical Society and Society of Industrial and Applied Mathematics.

This talk also appeared as a chapter in Vol. 14 of their series [5].

This analysis is being extended to the elastic case and has been

partially presented as an invited talk at the Spring 1984, International Conference on Acoustic and Elastic Inverse Scattering held at Cornell University.

Building of some earlier work by Stickler and Ammicht^[6] (in an earlier phase of this contract) an expansion was found for the asymptotic evaluation of the continuous spectrum for a normal model expansion of the pressure field in a stratified ocean^[7].

The determination of the continuous spectral contribution for a stratified ocean acoustic model can be a difficult and expensive numerical task. A uniform asymptotic technique has been used to calculate the pressure field for the Pekeris model. It is uniform in the sense that it is valid as a mode passes through cutoff. The basic idea of the technique is to exploit the effect of the various singular points of the integrand representing the continuous spectrum. This idea has been further exploited to describe the continuous spectrum contribution for a general stratified ocean. Numerical examples show that this technique is both accurate and fast.

Again building on some work sponsored in an earlier phase of this contract^[8], where the leading term was calculated for the temporal response of the acoustic pressure near a smooth caustic, the initial phase of the necessary calculation to determine the second term was completed^[9].

One method to estimate the validity of the leading term in the geometrical acoustics expansion is to calculate the second term in the expansion. This second term plays an essential role in the construction of the second term in Ludwig's uniform smooth caustic ansatz and in the construction of the progressing wave smooth caustic ansatz. Therefore, the determination of this term is important in the assessment of each of these expansions. Expressions for this second term are derived valid for arbitrary index profiles near the source point and near a smooth caustic. Numerical examples are presented for the n^2 -linear profile. These results show that near the source the second term is not singular, but near the caustic it is more singular than the leading order term.

References

- [1] D.C. Stickler, P.A. Deift "Inverse problem for a stratified ocean and bottom" Jour. Acoust. Soc. Am. 1981(70) 1723-1737
- [2] D.C. Stickler "Inverse scattering in a stratified medium" Jour. Acoust. Soc. Am. 1983(74) 994-1005
- [3] D.C. Stocker "Application of the trace method for Inverse Scattering to a medium which supports N types of Waves" Accepted for publication SIAM Jour. of Applied Mathematics
- [4] D.C. Stickler "Electromagnetic Inverse Scattering for Stratified Media" SPIE vol. 413 (Inverse Optics) 115-123
- [5] D.C. Stickler "Application of the trace formula methods to inverse scattering for some geophysical problems" Proceedings of the Symposium in Applied Mathematics of the American Mathematical Society and the Society for Industrial and Applied Mathematics. vol. 13, American Math. Soc. 13-32
- [6] D.C. Sticker, E. Amnicht "Uniform asymptotic evaluation of the continuous spectrum contribution for the Pekeris Model" J. Acoust. Soc. Am. Vol. 67 (1980) 2018-2024
- [7] E. Amnicht, D.C. Sticker "Uniform asymptotic evaluation of the continuous spectrum contribution for a stratified ocean" J. Acoust. Soc. Amer. Vol. 76 (1984) 186-191
- [8] D.C. Stickler, D.S. Ahluwalia, L. Ting "Application of Ludwig's uniform progressing wave ansatz to a smooth caustic" J. Acous. Soc. Am. vol. 69 (1981), 1673-1681
- [9] D.C. Stickler, J. Tavantzis, E. Amnicht "Calculation of the second term in the geometrical acoustic expansion" J. Acous. Soc. Am. vol. 75, (1984), 1071-1076

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A/1	

