

AD-A148 210 IONOSPHERIC AND MAGNETOSPHERIC MODIFICATIONS CAUSED BY 1/1
VLF WAVES(U) REGIS COLL WESTON MA M C LEE ET AL.
01 JUN 84 AFGL-TR-84-0301 F19628-83-K-0024

UNCLASSIFIED


F/G 4/1

NL

END

FILMED

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

Unclassified

AD-A148 210

10

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS None	
2a. SECURITY CLASSIFICATION AUTHORITY N/A		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release, distribution unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE N/A		4. PERFORMING ORGANIZATION REPORT NUMBER(S)	
5. MONITORING ORGANIZATION REPORT NUMBER(S) AFGL-TR-84-0301		6a. NAME OF PERFORMING ORGANIZATION Regis College	
6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION Air Force Geophysics Laboratory/LIS	
6c. ADDRESS (City, State and ZIP Code) 235 Wellesley Street Weston, Mass. 02193		7b. ADDRESS (City, State and ZIP Code) Hanscom AFB, MA 01731	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Air Force Geophysics Lab		8b. OFFICE SYMBOL (If applicable)	
8c. ADDRESS (City, State and ZIP Code) Hanscom AFB, MA 01731 John A. Klobuchar		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F19628-83-K-0024	
10. SOURCE OF FUNDING NOS.		11. TITLE (Include Security Classification) Ionospheric and magnetospheric modifications caused by VLF waves	
PROGRAM ELEMENT NO. 62101F		PROJECT NO. 4643	
TASK NO. 09		WORK UNIT NO. AB	
12. PERSONAL AUTHOR(S) M.C. Lee; S.P. Kuo *			
13a. TYPE OF REPORT Scientific - Reprint		13b. TIME COVERED FROM _____ TO _____	
14. DATE OF REPORT (Yr., Mo., Day) June 1, 1984		15. PAGE COUNT 4	
16. SUPPLEMENTARY NOTATION*Polytechnic Institute of NY, Long Island Center, Farmingdale, NY 11735 Proceedings of 1984 International Conference on Plasma Physics, Lausanne, Switzerland, June 27 - July 3, 1984			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB. GR.	
		Thermal filamentation instability; lower hybrid waves; VLF wave injection experiments	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The ionosphere and the magnetosphere may be significantly modified by the injected VLF waves via the thermal filamentation instability and the excitation of lower hybrid waves. → 10 p 2			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☐ DTIC USERS ☐		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL John A. Klobuchar		22b. TELEPHONE NUMBER (Include Area Code) 617-861-3988	
		22c. OFFICE SYMBOL AFGL/LIS	

DTIC FILE COPY

DTIC
ELECTE
S DEC 5 1984
A

AFGL-TR-84-0301

Ionospheric and magnetospheric modifications caused by
the injected VLF waves

M.C. Lee

Regis College Research Center, Weston, Mass. 02193, U.S.A.

S.P. Kuo

Polytechnic Institute of New York, Long Island Center,
Farmingdale, N.Y. 11735, U.S.A.

Abstract. The ionosphere and the magnetosphere may be significantly modified by the injected VLF waves via the thermal filamentation instability and the excitation of lower hybrid waves.

1. Introduction

cont → In VLF wave injection experiments performed, for example, at Siple, Antarctica, the ground-based transmitters are usually operated in a pulsed-wave mode with durations of a few seconds. These VLF wave pulses ~~have been~~ ^{were} used to study coherent wave-particle interactions in the magnetosphere such as the wave amplification, the triggering of wave emissions, the induced particle precipitation etc. (see, e.g., Helliwell, 1983). We show in this paper that if the transmitters are operated continuously for a few minutes, significant ionospheric and magnetospheric disturbances can be caused by the following two processes. One is the thermal filamentation of the pump wave and the other one is the excitation of lower hybrid waves. ↗

2. Thermal filamentation of whistlers

Monochromatic VLF waves have been observed to change from linear into circular polarization (i.e., a whistler mode) on their path through the neutral atmosphere and into the ionosphere. If the pump waves are intense enough, the filamentation instability of whistlers can be excited. This instability yields a whistler sideband and zero-frequency modes that are associated with the simultaneous excitation of both plasma density fluctuations (δn) and magnetic field fluctuations (δB). The source of these magnetostatic fluctuations stems from the wave-induced quasi-DC electric current due to the electron $F \times B$ drift motion under the influence of the differential Ohmic heating force.

*Proceedings of 1984 International Conference on Plasma Physics,
Lausanne, Switzerland, June 27-July 3, 1984*

The plasma density fluctuations (δn) and the magnetostatic fluctuations (δB) are found to be related as

$$\delta n/n_0 \approx [1 + (v_e/\gamma)(c^2/\lambda^2 f_p^2)](\delta B/B_0) \quad (1)$$

where f_p , v_e , n_0 , B_0 , γ , and λ are, respectively, the electron plasma frequency, the electron-ion collision frequency, the background plasma density, the earth's magnetic field, the growth rate of the instability, and the scale lengths of the excited modes. It is clear from (1) that significant magnetostatic fluctuations are associated with the excitation of large scale modes.

Although the whistler waves have a broad propagation regime: $|\Omega_e| \ll \omega_0 < |\Omega_e|$, the excitation of the filamentation instability is restricted to whistlers with frequencies (ω_0) satisfying the following condition:

$$1 - \omega_0^2 \omega_p^2 (|\Omega_e| - \omega_0)^{-1} (\omega_p^2 + k^2 c^2)^{-1} - k^2 c^2 |\Omega_e|^{-1} (|\Omega_e| + \omega_0)^{-1} (\omega_0^2 + k^2 c^2)^{-1} < 0 \quad (2)$$

where ω_p , $|\Omega_e|$ ($|\Omega_i|$), ω_0 , and k are the angular electron plasma frequency, the electron (ion) gyrofrequency, the angular whistler wave frequency, and the wave number of the excited modes, respectively. The inequality (2) demands that $\omega_0 > |\Omega_e|/2$ for the excitation of large scale modes. Since $|\Omega_e| \sim 1.4$ MHz (13.6 KHz) in the ionospheric F region (in the magnetosphere at $L = 4.0$), this result indicates that the thermal filamentation of whistlers can occur in the ionosphere (in the magnetosphere at $L = 4.0$) when the wave frequencies are within the frequency range: $1.4 \text{ MHz} > \omega_0/2\pi > 0.7 \text{ MHz}$ ($13.6 \text{ KHz} > \omega_0/2\pi > 6.8 \text{ KHz}$).

The threshold of this instability has been found to be

$$|e\epsilon_{th}/mc|^2 \sim 1.5 \omega_0^{-1} \omega_p^{-2} (\omega_0 - |\Omega_e|)^3 k^2 v_{ti}^2 (4 - a - ab)^{-1} (1 + b - ab) \quad (3)$$

where $a = k^2 c^2 (\omega_0^2 - \Omega_e^2) / \omega_0 |\Omega_e| \omega_p^2$, $b = \omega_0 \omega_p^2 / (\omega_0 - |\Omega_e|) (\omega_p^2 + k^2 c^2)$, and v_{ti} is the ion thermal velocity. The threshold fields thus calculated are of the order of a few $\mu\text{V/m}$ (a few mV/m) for exciting modes with tens of kilometers (tens of meters) and larger scales in the magnetosphere at $L = 4.0$ (in the ionospheric F region), that are achievable whistler wave field intensities with available facilities. However,

whistler waves with much higher intensities are required to ensure the excitation of the thermal filamentation instability in the magnetosphere. This is because the growth rate of the instability is rather small if the whistler wave field intensities just barely exceed the threshold fields.

In terms of the threshold (ϵ_{th}), the growth rate of large scale modes (i.e., $f_p \gg c/\lambda$) is given by

$$\gamma_{\infty} \sim (2v_e k v_{ti} / |\Omega_e|) (\epsilon_0 / \epsilon_{th}) \quad (4)$$

that turns out to be independent of the scale lengths because as shown in (3), ϵ_{th} is proportional to k . If $\epsilon_0 / \epsilon_{th} \sim 0(1)$, γ_{∞} is of the order of 10^{-5} Hz (10^{-3} Hz) in the magnetosphere at $L = 4.0$ (in the ionospheric F region). The Siple signals, propagating in the non-ducted whistler mode, have $\epsilon_0 / \epsilon_{th} \sim 0(1)$ in the magnetosphere at $L = 4.0$. The probability for seeing the Siple signals in the ducted whistler mode is about 20%. The growth rate can be increased by two to three orders of magnitude in the ducted whistler propagation. Therefore, if the Siple transmitter is operated for a few minutes, the thermal filamentation instability can excite in the magnetosphere at $L = 4.0$ the plasma density fluctuations with scale lengths greater than tens of kilometers. This instability can also be excited in the ionospheric F region by the MF waves with frequencies close to but less than the local electron gyrofrequency.

3. Excitation of lower hybrid waves

The injected VLF waves can interact directly with the ionosphere through the excitation of lower hybrid waves and a field-aligned purely growing mode. This instability can be excited in a broad whistler frequency range in two domains. They are Domain 1: $\omega_{LH} [1 + (M/m)(v_t^2/c^2)(\omega_p^2/\Omega_e^2)]^{1/2} < \omega_0 < \omega_{p1}$ for the non-oscillatory beating current to be the dominant nonlinear effect, and Domain 2: $\omega_{p1} < \omega_0 \ll |\Omega_e|$ for the thermal pressure force to be the dominant nonlinear effect, where ω_{LH} defined by $\omega_{p1} / (1 + \omega_p^2/\Omega_e^2)^{1/2}$ is the lower hybrid resonance frequency; ω_{p1} (ω_p), $|\Omega_e|$, v_t , and (M/m) are the ion (electron) plasma frequency, the electron gyrofrequency, the electron thermal velocity, and the ratio of ion to electron masses, respectively.

The optimum threshold field for the instability is found to be

$E_m \sim 1.2 (m/e) v_e v_t |\eta|^{1/2}$ in frequency domain 1 and $E_m \sim 0.86 (k^2 v_t^3 / \Omega_e) (m/e) [1 + (1 + 4\Omega_e^2 v_t^2 \eta^2 / k_s^2 v_t^2)^{1/2}]^{1/2} / |\eta|^{1/2}$ in frequency domain 2, where v_e and k_s are the electron-ion collision frequency and the wave number of the excited field-aligned mode, respectively; $\eta = [1 + (M/m)(k_o/k_s)^2] / [1 - (M/m)(K_o/k_s)^2 (\Omega_e/\omega_p)^2]$, where k_o/k_s is the ratio of the perpendicular to the parallel scale lengths of the excited lower hybrid waves.

The growth rate of the instability has the following expressions, $\gamma \sim 0.5 (v_e k_s^2 v_t^2 / \Omega_e^2) (E_R^2 - 1)$ for $E_R^2 \ll 10$ and $\gamma \sim 1.4 (v_e k_s^2 v_t^2 / \Omega_e^2) E_R$ for $E_R^2 \gg 10$, where E_R is the ratio of the whistler field intensity (E_o) to the optimum threshold field intensity (E_m) of the instability. Since $\omega_{LH} \sim 6$ KHz in the ionosphere, the whistler wave frequencies have to be greater than 6 KHz for the excitation of lower hybrid waves. The threshold field has been estimated to be about 1 mv/v. If $E_R \sim 0(10)$, the instability with dominant scale lengths near 10 meters can be excited within a few seconds in the ionosphere. By contrast, kilometer-scale lower hybrid waves can be excited in the magnetosphere at $L = 4.0$ with much lower threshold field ($\sim$ a few μ v/m). The growth rate is, however, rather small for non-ducted whistler modes, whose $E_R \sim 0(1)$. Whereas, the ducted whistler modes can excite the instability with growth rates as large as 10^{-2} Hz. In other words, lower hybrid waves can be produced by ducted whistler waves in the magnetosphere within a few minutes. Electron precipitation is the ionospheric effects expected from the excitation of lower hybrid waves. Indeed, airglow effects have been observed in the Russian Juliana program (Chmyrev et al., 1976) to be associated with the VLF transmitter cycle.

References

- Chmyrev, V.M. et al., JETP Lett., 23, 409, 1976.
Helliwell, R.A., Radio Sci., 18, 801, 1983.

Acknowledgements. This work was supported by AFGL contract F19628-83-K-0024 at Regis College Research Center and jointly in part by NSF grant ATM-8315322 and in part by the Air Force Office of Scientific Research grant AFOSR-83-0001 at Polytechnic Institute of New York.



Distribution/	
Availability Codes	
Avail and/or	Special
A-1	

END

FILMED

1-85

DTIC