

AD-A152 927

ON STOCHASTIC INTEGRAL REPRESENTATION OF STABLE  
PROCESSES WITH SAMPLE PAT. (U) NORTH CAROLINA UNIV AT  
CHAPEL HILL CENTER FOR STOCHASTIC PROC. J ROSINSKI  
JAN 85 TR-88 AFOSR-TR-85-0296 F/G 12/1

1/1


UNCLASSIFIED

NL

END

FORM 1

10-1




MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS 1963-A

# CENTER FOR STOCHASTIC PROCESSES

AD-A152 927

Department of Statistics  
University of North Carolina  
Chapel Hill, North Carolina



ON STOCHASTIC INTEGRAL REPRESENTATION OF STABLE

PROCESSES WITH SAMPLE PATHS IN BANACH SPACES

by

Jan Rosinski

Technical Report No. 88

January 1985

DTIC  
ELECTE  
APR 25 1985  
S D  
B

DTIC FILE COPY

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |       |                                                                                |                                                                                                                         |                                              |                          |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|--------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------|----------------------------------------------|--------------------------|
| 1a. REPORT SECURITY CLASSIFICATION<br><b>UNCLASSIFIED</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |       |                                                                                | 1d. RESTRICTIVE MARKINGS                                                                                                |                                              |                          |
| 2a. SECURITY CLASSIFICATION AUTHORITY                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |       |                                                                                | 3. DISTRIBUTION/AVAILABILITY OF REPORT<br>Approved for public release; distribution unlimited.                          |                                              |                          |
| 2b. DECLASSIFICATION/DOWNGRADING SCHEDULE                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |       |                                                                                | 5. MONITORING ORGANIZATION REPORT NUMBER(S)<br><b>AFOSR-TR. 85-0296</b>                                                 |                                              |                          |
| 4. PERFORMING ORGANIZATION REPORT NUMBER(S)<br>TR No. 88                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |       |                                                                                | 7a. NAME OF MONITORING ORGANIZATION<br>Air Force Office of Scientific Research                                          |                                              |                          |
| 6a. NAME OF PERFORMING ORGANIZATION<br>University of North Carolina                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |       | 6b. OFFICE SYMBOL<br>(If applicable)                                           | 7b. ADDRESS (City, State and ZIP Code)<br>Directorate of Mathematical & Information Sciences, Bolling AFB DC 20332-6448 |                                              |                          |
| 6c. ADDRESS (City, State and ZIP Code)<br>Center for Stochastic Processes, Department of Statistics, Phillips Hall 039-A, Chapel Hill NC 27514                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |       | 9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER<br>F49620-82-C-0009            |                                                                                                                         |                                              |                          |
| 8a. NAME OF FUNDING/SPONSORING ORGANIZATION<br>AFOSR                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |       | 8b. OFFICE SYMBOL<br>(If applicable)<br>NM                                     | 10. SOURCE OF FUNDING NOS.                                                                                              |                                              |                          |
| 8c. ADDRESS (City, State and ZIP Code)<br>Bolling AFB DC 20332-6448                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |       | PROGRAM ELEMENT NO. 61102F<br>PROJECT NO. 2304<br>TASK NO. A5<br>WORK UNIT NO. |                                                                                                                         |                                              |                          |
| 11. TITLE (Include Security Classification)<br>ON STOCHASTIC INTEGRAL REPRESENTATION OF STABLE PROCESSES WITH SAMPLE PATHS IN BANACH SPACE                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |       |                                                                                |                                                                                                                         |                                              |                          |
| 12. PERSONAL AUTHOR(S)<br>Jan Rosinski                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |       |                                                                                |                                                                                                                         |                                              |                          |
| 13a. TYPE OF REPORT<br>Technical                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |       | 13b. TIME COVERED<br>FROM _____ TO _____                                       |                                                                                                                         | 14. DATE OF REPORT (Yr., Mo., Day)<br>JAN 85 |                          |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |       |                                                                                |                                                                                                                         | 15. PAGE COUNT<br>37                         |                          |
| 16. SUPPLEMENTARY NOTATION                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |       |                                                                                |                                                                                                                         |                                              |                          |
| 17. COSATI CODES                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |       |                                                                                | 18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)                                       |                                              |                          |
| FIELD                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  | GROUP | SUB. GR.                                                                       | Stable process; stochastic integral; stable distributions on Banach spaces.                                             |                                              |                          |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |       |                                                                                |                                                                                                                         |                                              |                          |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |       |                                                                                |                                                                                                                         |                                              |                          |
| 19. ABSTRACT (Continue on reverse if necessary and identify by block number)<br>Certain path properties of a symmetric $\alpha$ -stable process<br>$X(t) = \int_S h(t,s) dM(s), \quad t \in T,$<br>are studied in terms of the kernel $h$ . The existence of an appropriate modification of the kernel $h$ enables one to use results from stable measures on Banach spaces in studying of $X$ . Bounds for the moments of the norm of sample paths of $X$ are obtained. This yields definite bounds for the moments of a double $\alpha$ -stable integral. Also necessary and sufficient conditions for the absolute continuity of sample paths of $X$ are given. Along with the above stochastic integral representation of stable processes, the representation of stable random vectors due to LePage, Woodroffe and Zinn is extensively used and the relationship between these two representations is discussed. |       |                                                                                |                                                                                                                         |                                              |                          |
| 20. DISTRIBUTION/AVAILABILITY OF ABSTRACT<br>UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |       |                                                                                | 21. ABSTRACT SECURITY CLASSIFICATION<br>UNCLASSIFIED                                                                    |                                              |                          |
| 22a. NAME OF RESPONSIBLE INDIVIDUAL<br>MAJ Brian W. Woodruff                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |       |                                                                                | 22b. TELEPHONE NUMBER<br>(Include Area Code)<br>(202) 767- 5027                                                         |                                              | 22c. OFFICE SYMBOL<br>NM |

ON STOCHASTIC INTEGRAL REPRESENTATION OF STABLE  
PROCESSES WITH SAMPLE PATHS IN BANACH SPACES

by

Jan Rosinski\*  
Center for Stochastic Processes  
Department of Statistics  
University of North Carolina  
Chapel Hill, NC 27514

DTIC  
ELECTE  
APR 25 1985  
S B D

ABSTRACT. Certain path properties of a symmetric  $\alpha$ -stable process

$$X(t) = \int_S h(t,s) dM(s), \quad t \in T,$$

are studied in terms of the kernel  $h$ . The existence of an appropriate modification of the kernel  $h$  enables one to use results from stable measures on Banach spaces in studying of  $X$ . Bounds for the moments of the norm of sample paths of  $X$  are obtained. This yields definite bounds for the moments of a double  $\alpha$ -stable integral. Also necessary and sufficient conditions for the absolute continuity of sample paths of  $X$  are given. Along with the above stochastic integral representation of stable processes, the representation of stable random vectors due to LePage, Woodrooffe and Zinn is extensively used and the relationship between these two representations is discussed.

AMS 1980 Subject Classification: Primary: 60G17, 60H05, 60B12

Keywords and Phrases: Stable process, stochastic integral, stable distributions on Banach spaces

\*on leave from Wroclaw University, Poland

Research supported by AFOSR Grant No. F49620 82 C 0009

AIR FORCE OFFICE OF  
NOTICE  
THE  
DEPT.  
MATHEMATICS  
Chief, Technical Division

## 1. Introduction.

The Central Limit Theorem and the stability property provide the basic reasons for regarding stable processes as a natural generalization of Gaussian ones. As an analog to the well-known spectral representation of stationary Gaussian processes, every *symmetric  $\alpha$ -stable* (SaS) stochastic process with parameter set  $T$  has a version of the form

$$(1.1) \quad X(t) = \int_S h(t,s) dM(s), \quad t \in T$$

(cf. [1], [12], [29], [30], [11] and the discussion of the history of (1.1) in [9]) and in the stationary case one can choose  $t \rightarrow h(t, \cdot)$  as an orbit of a group of isometries on  $L^\alpha$  (see [9]). Here  $M$  is an independently scattered SaS random measure on an abstract measurable space  $(S, \mathcal{A})$ .

There are two special cases of (1.1) that have been extensively studied: *harmonizable processes* given by

$$(1.2) \quad X(t) = \int_{-\infty}^{\infty} e^{its} dM(s), \quad t \in \mathbb{R}$$

(with appropriate modifications if  $t$  runs over a group) (cf. [10], [19], [6], [17], [2], [23] and [32]) and *moving averages*

$$(1.3) \quad X(t) = \int_{-\infty}^{\infty} g(t-s) dM(s), \quad t \in \mathbb{R}$$

(cf. [6], [25], [4] and [2]).

In this paper we study general SaS processes given by (1.1). They are determined by two quantities: the kernel  $h$  and the control measure  $m$  of  $M$ . In contrast with the approach taken in [9] and [2], which relies on the properties of the mapping  $T \ni t \rightarrow h(t, \cdot) \in L^\alpha$ , we relate path properties of  $X$

with properties of the mapping  $S\alpha s \rightarrow h(.,s) \in \mathbb{R}^T(\mathbb{C}^T)$  which plays the crucial role here. More specifically, we are concerned with processes having sample paths in a separable Banach space  $V(T)$  of functions defined on  $T$ . We show that the kernel  $h$  in (1.1) admits a modification with all sections  $h(.,s)$  in  $V(T)$  (Section 5). Therefore we may always replace (1.1) by

$$(1.4) \quad X = \int_S h(.,s) dM(s),$$

where on the right-hand side we have a stochastic integral of the  $V(T)$ -valued function  $s \rightarrow h(.,s)$ . Such stochastic integrals of Banach space valued functions have been investigated in [26] for infinitely divisible random measures. In the present stable case the construction can be simplified and this is done at the beginning of Section 3. Then we establish the relationship between the stochastic integral representation of stable random vectors in Banach spaces and the series representation due to LePage, Woodrooffe and Zinn [13].

In Section 4 we use some ideas of Marcus and Pisier [17] and an adaptation of Hoffmann-Jørgensen's inequality due to Giné and Zinn [8] to obtain bounds for moments of the norm of  $X(.,)$ . We also introduce a complete norm on the space of all vector valued functions  $f$  for which  $\int f dM$  exists, similar to Pisier's norm for CLT [21]. Theorem 4.5 establishes the role of simple functions in the series representation of stable random vectors.

In Section 6 we apply the results of Sections 5 and 4 to characterize the absolute continuity of sample paths of  $S\alpha S$  processes. Earlier the absolute continuity of sample paths has been investigated by Cambanis and Miller [3] in terms the so-called covariation function, only for the case  $\alpha > 1$ .

Continuing the above approach, we obtain in Section 7 definitive bounds for moments of a double  $\alpha$ -stable stochastic integral. We also give a short and

new proof of a Fubini-type result which allows the interchanging of stochastic and usual integration (cf. [4] Theorem 4.6 and [20] Lemma 4.4).



|                    |                                     |
|--------------------|-------------------------------------|
| Accession For      |                                     |
| NTIS GNA&I         | <input checked="" type="checkbox"/> |
| DTIC TAB           | <input type="checkbox"/>            |
| Unannounced        | <input type="checkbox"/>            |
| Justification      |                                     |
| By                 |                                     |
| Distribution/      |                                     |
| Availability Codes |                                     |
| Dist               | Avail and/or<br>Special             |
| A-1                |                                     |

## 2. Preliminaries and notation.

A systematic treatment of stable measures on Banach spaces one can find in Linde [14] and we shall refer to this book for basic definitions and facts. The characteristic functional of an SsS p.m.  $\mu$  on a separable Banach space  $B$  can be written in the form

$$(2.1) \quad \hat{\mu}(x') = \exp\left(-\int_{\partial U} |\langle x, x' \rangle|^\alpha d\sigma(x)\right),$$

$x' \in B'$ , where  $\partial U$  is the unit sphere in  $B$  and  $\sigma$  is a finite symmetric measure on  $\partial U$ .  $\sigma$  is uniquely determined by  $\mu$  and is called the *spectral measure* of  $\mu$  (cf. Theorem 6.4.4 [14]). Further, for every  $p \in (0, \alpha)$   $\int_B ||x||^p d\mu(x) < \infty$  and for every  $p, q \in (0, \alpha)$

$$(2.2) \quad \left(\int_B ||x||^p d\mu(x)\right)^{1/p} \underset{C}{\sim} \left(\int_B ||x||^q d\mu(x)\right)^{1/q}$$

where  $C = C(\alpha, p, q)$  (we shall write  $L \underset{C}{\sim} R$  if  $C^{-1}R \leq L \leq CR$  and  $C = C(a, b, \dots)$  means that a positive constant  $C$  depends only on parameters  $a, b, \dots$ ). If  $\hat{\mu}$  is given by

$$\hat{\mu}(x') = \exp\left(-\int_B |\langle x, x' \rangle|^\alpha d\sigma_0(x)\right),$$

$x' \in B'$ , where  $\sigma_0$  is a (non-necessary symmetric) Borel measure on  $B$ , which is  $\sigma$ -finite on  $B \setminus \{0\}$ , then  $\int_B ||x||^\alpha d\sigma_0(x) < \infty$  and for every  $p \in (0, \alpha)$

$$(2.3) \quad \left(\int_B ||x||^\alpha d\sigma_0(x)\right)^{1/\alpha} \leq C \left(\int_B ||x||^p d\mu(x)\right)^{1/p},$$

where  $C = C(\alpha, p)$  (cf. Proposition 6.4.5 and Corollary 7.3.5 in [14]).

Let  $A$  be a  $\delta$ -ring of subsets of a non-empty set  $S$  (i.e. a ring that is

closed under countable intersections). A stochastic process  $\{M(A): A \in \mathcal{A}\}$  is said to be an *S $\alpha$ S random measure* if

- (i) for every sequence  $\{A_n\} \subset \mathcal{A}$  of pairwise disjoint sets with  $\bigcup A_n \in \mathcal{A}$  the series  $\sum M(A_n)$  converges in probability to  $M(\bigcup A_n)$ ;
- (ii)  $M(A_1), M(A_2), \dots$  are independent, provided  $A_n \in \mathcal{A}$  are pairwise disjoint;
- (iii)  $M(A)$  has an *S $\alpha$ S* distribution for every  $A \in \mathcal{A}$ .

It follows that the characteristic function of  $M(A)$  can be written in the form

$$E \exp(it M(A)) = \exp(-m(A)|t|^\alpha), \quad t \in \mathbb{R}, \quad A \in \mathcal{A},$$

where  $m$  is a non-negative measure on  $\mathcal{A}$ .  $m$  is called the *control measure* of  $M$ .

The existence of an *S $\alpha$ S* random measure  $M$  with a given control measure  $m$  follows by Kolmogorov's Consistency Theorem. In particular, if  $X(s)$ ,  $s \geq 0$  is an independent stationary increment process such that  $E \exp(itX(s)) = \exp(-s|t|^\alpha)$ , then  $M$  defined on intervals by  $M((a,b]) = X(b) - X(a)$  extends to an *S $\alpha$ S* random measure on the  $\delta$ -ring of all Borel bounded subsets of  $[0, \infty)$  with the Lebesgue measure as the control measure. (see Prekopa [22]).

Throughout this paper we shall assume that  $(S, \mathcal{A})$  satisfies the following condition: there exists a sequence  $\{A_n\} \subset \mathcal{A}$  such that  $\bigcup A_n = S$ . Then every countably additive finite measure on  $\mathcal{A}$  extends uniquely to a  $\sigma$ -finite measure on  $\sigma(\mathcal{A})$  and as a consequence every control measure of an *S $\alpha$ S* random measure  $M$  is the restriction to  $\mathcal{A}$  of a  $\sigma$ -finite measure  $m$  defined on  $\sigma(\mathcal{A})$ . To avoid trivialities we shall always assume that  $m$  is not zero.

For every real function  $f \in L^\alpha(S, \sigma(\mathcal{A}), m)$  the stochastic integral  $\int_S f dM$  is defined as the limit in  $L^p$  of integrals of simple functions and satisfies

the equality

$$(E \mid \int_S f dM \mid^p)^{1/p} = C \left( \int_S |f|^\alpha dM \right)^{1/\alpha},$$

where  $C = C(\alpha, p)$ ,  $p < \alpha$  (cf. [1], and [29]). If  $B$  is a finite dimensional Banach space, then for  $f \in L_B^\alpha(S, \sigma(A), m)$

$$(2.4) \quad (E \mid \mid \int_S f dM \mid \mid^p)^{1/p} \sim_C \left( \int_S \mid \mid f \mid \mid^\alpha dm \right)^{1/\alpha},$$

where  $C$  depends on  $\alpha$  and  $p$  and additionally on  $\dim B$  and  $\mid \mid \cdot \mid \mid$ . Banach spaces in which (2.4) holds for all simple functions with a universal constant  $C$  (and thus  $\int_S f dM$  can be defined for all  $f \in L_B^\alpha$ ) have been characterized by Marcus and Woyczynski [18]. This is the class of spaces of stable type  $\alpha$ , including in particular Hilbert spaces and  $L^p$ -spaces for  $p > \alpha$  (see [14] for further references).

### 3. Stochastic integral and series representation of SαS random vectors in Banach spaces.

Let  $M$  be an SαS random measure on  $(S, A)$  with the control measure  $m$ . Let  $L_B^{\text{simple}}$  be the space of all simple measurable functions  $f: (S, \sigma(A)) \rightarrow (B, \text{Borel}(B))$  such that  $\{s: f(s) \neq 0\} \in A$ . As usual functions equal  $m$ -a.e. are indistinguishable. For every  $f \in L_B^{\text{simple}}$ ,  $f = \sum x_j I_{A_j}$ ,  $x_j \neq 0$ ,  $A_j \in A$  we set

$$\int_S f dM = \sum x_j M(A_j)$$

and

$$(3.1) \quad \lambda_{\alpha, p}(f) = (E || \int_S f dM ||^p)^{1/p},$$

where  $p \in (0, \alpha)$ .  $\lambda_{\alpha, p}$  is a well-defined quasi norm on  $L_B^{\text{simple}}$ . Since  $\int f dM$  is a  $B$ -valued SαS random vector such that

$$(3.2) \quad E \exp(i \langle \int f dM, x' \rangle) = \exp(-\int |\langle f(s), x' \rangle|^\alpha dm(s)),$$

$x' \in B'$ , inequality (2.3) yields

$$(3.3) \quad \lambda_{\alpha, p}(f) \geq C \left( \int_S ||f(s)||^\alpha dm(s) \right)^{1/\alpha}$$

where  $C = C(\alpha, p)$ . Moreover, by (2.2)  $\lambda_{\alpha, p} \sim_C \lambda_{\alpha, q}$  for every  $p, q \in (0, \alpha)$ , where  $C = C(\alpha, p, q)$ .

Let  $S_B^\alpha$  be a completion of  $L_B^{\text{simple}}$  in  $\lambda_{\alpha, p}$ . In view of (3.3)  $S_B^\alpha$  can be realized as a linear subspace of  $L_B^\alpha = L_B(S, \sigma(A), m)$  as follows:

$$S_B^\alpha = \{f \in L_B^\alpha: \text{there exists } \{f_n\}_{n=1}^\infty \subset L_B^{\text{simple}} \text{ such that } f_n \rightarrow f \text{ in } L_B^\alpha \text{ and}$$

$$\lim_{n, m \rightarrow \infty} \lambda_{\alpha, p}(f_n - f_m) = 0 \text{ for some (each) } p \in (0, \alpha)\}.$$

By (3.1) the mapping  $f \mapsto \int_S f dM$  extends to an isometric injection of  $(S_B^\alpha, \lambda_{\alpha, p})$

into  $L_B^p(\Omega, \mathcal{F}, P)$ . Values of this extension are also denoted by  $\int fdM$  and called the stochastic integral of  $f$  with respect to  $M$ .

Because of the lack of (2.4) in general, the stable stochastic integral can not be defined for all  $f$ 's in  $L_B^\alpha(S, \sigma(A), m)$  and  $S_B^\alpha$  is the largest subspace of  $L_B^\alpha$  where  $\int fdM$  is defined by taking limits of stochastic integrals of simple functions. Although the dependence of  $\lambda_{\alpha, p}(f)$  on  $f$  is not given explicitly, this is a useful quasi-norm which can be effectively estimated in many concrete examples of  $B$  (see e.g. Sections 6 and 7).

We shall frequently use the following particular case of Ito-Nisio theorem for Banach space valued stochastic integrals which was proven in [26].

Proposition 3.1.  $f \in S_B^\alpha$  and  $\mu = L(\int fdM)$  if and only if  $\int |\langle f(s), x' \rangle|^\alpha dm(s) < \infty$  for every  $x' \in B'$  and the cylindrical measure  $\mu_0$  given by

$$\hat{\mu}_0(x') = \exp \left\{ - \int |\langle f(s), x' \rangle|^\alpha dm(s) \right\}, \quad x' \in B',$$

extends to a countably additive Borel measure  $\mu$  on  $B$ .

As a consequence of the above proposition every SaS p.m. on  $B$  has a stochastic integral representation which follows from the following (cf. [26], Theorem 6.7):

Proposition 3.2.  $\{\int fdM: f \in S_B^\alpha\}$  is a closed linear subspace of  $L_B^p(\Omega, \mathcal{F}, P)$  consisting of SaS random vectors. If  $(S, A, m)$  is atomless, then  $\{L(\int fdM): f \in S_B^\alpha\}$  coincides with the set of all SaS p.m.'s on  $B$ .

We shall discuss now a different transformation of  $f$  which leads to the same distribution as  $\int fdM$ . We shall consider a series representation of stable random vectors due to LePage, Woodroffe and Zinn [13] as it was developed by Marcus and Pisier [17].

Assume that  $m(S) < \infty$  and  $\Lambda = \sigma(A)$ . Let  $\{\eta_j\}$  be a sequence of positive

i.i.d. random variables such that  $P(\eta_n > \lambda) = e^{-\lambda}$ ,  $\lambda \geq 0$ , and put  $\Gamma_j = \eta_1 + \dots + \eta_j$ . Let  $\{\xi_j\}$  be an i.i.d. sequence of symmetric random variables such that  $E|\xi_j|^\alpha = 1$ . Let  $\{\tau_j\}$  be a sequence of independent uniformly distributed random elements in  $(S, A, m)$ , i.e.  $P(\tau_j \in A) = m(A)/m(S)$  for every  $A \in A$ . We assume that all the sequences  $\{\eta_j\}$ ,  $\{\xi_j\}$  and  $\{\tau_j\}$  are independent of the others.

Let  $f \in S_B^\alpha$  and  $\mu = L(f dM)$ . Then for every  $x' \in E'$

$$E|<f(\tau_j), x'>|^\alpha = [m(S)]^{-1} \int_S |<f(s), x'>|^\alpha dm(s) < \infty$$

and by Lemma 1.4 in [17] the series

$$c(\alpha)[m(S)]^{1/\alpha} \sum_{j=1}^{\infty} (\Gamma_j)^{-1/\alpha} \xi_j <f(\tau_j), x'>$$

converges a.s. to a real  $S_\alpha S$  random variable with the characteristic function

$$\phi(t) = \exp(-|t|^\alpha \int |<f(s), x'>|^\alpha dm(s)), \text{ where } c(\alpha) = (\int_0^\infty x^{-\alpha} \sin x dx)^{-1/\alpha}.$$

Since  $\phi(t) = \hat{u}(tx')$  and the sequence  $\{\Gamma_j \xi_j f(\tau_j)\}$  is sign-invariant, Ito-Nisio theorem (see e.g. [16], II.4.3. and II.4.4) yields the a.s. convergence of the series

$$(3.4) \quad \Sigma(f) = c(\alpha)[m(S)]^{1/\alpha} \sum_{j=1}^{\infty} (\Gamma_j)^{-1/\alpha} \xi_j f(\tau_j)$$

and  $L(\Sigma(f)) = \mu$ . Conversely, if (3.4) converges a.s. or in P, then the function

$x' \rightarrow \exp(-\int |<f, x'>|^\alpha dm)$  is the characteristic functional of  $\Sigma(f)$  and by

Proposition 3.1.  $f \in S_B^\alpha$ .

We summarize above in the following:

**Proposition 3.3.** Let  $m(S) < \infty$  and  $A = \sigma(A)$ . Then  $f \in S_B^\alpha$  if and only if  $\Sigma(f)$  converges a.s. or in P. Moreover,

$$L(f dM) = L(\Sigma(f))$$

#### 4. Bounds for moments of an SαS stochastic integral.

To obtain the first proposition we argue similarly to Marcus [15] and to Giné et al. [7].

Since  $j^{-1} \Gamma_j \rightarrow 0$  a.s. by Kolmogorov's SLLN, we get that  $\Sigma(f)$  converges a.s. if and only if

$$(4.1) \quad A(f) = \sum_{j=1}^{\infty} j^{-1/\alpha} \varepsilon_j f(\tau_j)$$

converges a.s. (see (3.4)). Moreover, by contraction principle we have

$$\begin{aligned} & 2^{-1/p} [E \inf_j (j/\Gamma_j)^{p/\alpha}]^{1/p} [m(S)]^{1/\alpha} (E \|A(f)\|^p)^{1/p} \leq \\ & \leq (E \|\Sigma(f)\|^p)^{1/p} \leq \\ & 2^{1/p} [E \sup_j (j/\Gamma_j)^{p/\alpha}]^{1/p} [m(S)]^{1/\alpha} (E \|A(f)\|^p)^{1/p}, \end{aligned}$$

where  $C = C(\alpha)$  and  $p \in (0, \alpha)$ . Since  $E \inf_j (j/\Gamma_j)^{p/\alpha} > 0$  and  $E \sup_j (j/\Gamma_j)^{p/\alpha} < \infty$

(see [15] and [7] respectively) we get

Proposition 4.1. Let  $m(S) < \infty$  and  $A = \sigma(A)$ . Then  $f \in S_B^\alpha$  if and only if  $A(f)$  converges a.s. and/or in  $L_B^p$  for some (each)  $p \in [0, \alpha)$ . Moreover,

$$(E \|f dM\|^p)^{1/p} \leq C [m(S)]^{1/\alpha} (E \|A(f)\|^p)^{1/p}$$

where  $C = C(\alpha, p)$ ,  $p \in (0, \alpha)$ .

Let us note that even on the real line  $A(f)$  need not have the  $\alpha$ -th moment finite. This is a simple corollary to Proposition 4.2 in [5]. We shall normalize  $f$  to ensure finiteness of all moments of  $A(f)$ .

Let  $x_0$  be a fixed point on the unit sphere of  $B$ . For every  $f \in L_B^\alpha$  we define  $\bar{f}(s) = f(s)/\|f(s)\|$  if  $f(s) \neq 0$  and  $\bar{f}(s) = x_0$  otherwise. We define also a finite measure  $m_f$  on  $(S, \sigma(A))$  by  $dm_f(s) = \|f(s)\|^\alpha dm(s)$  ( $m(S)$  can be infinite). Let  $M_f$  be an S $\alpha$ S random measure on  $(S, \sigma(A))$  with the control measure  $m_f$ . In view of Proposition 3.1  $f \in S_B(S, A, m)$  if and only if  $f \in L_B^\alpha$  and  $\bar{f} \in S_B^\alpha(S, \sigma(A), m_f)$ . Moreover

$$(4.2) \quad L\left(\int_S f dM\right) = L\left(\int_S \bar{f} dM_f\right).$$

Let  $\{\tau_j^f\}$  be a sequence of independent uniformly distributed random elements in  $(S, \sigma(A), m_f)$  i.e.  $P\{\tau_j^f \in A\} = m_f(A)/m_f(S)$ . Let  $\{\varepsilon_j\}$  be a sequence of i.i.d. random variables such that  $P\{\varepsilon_j = -1\} = P\{\varepsilon_j = 1\} = 1/2$  and independent of  $\{\tau_j^f\}$ . By Proposition 4.1 we get

$$(E \left\| \int_S \bar{f} dM_f \right\|^p)^{1/p} \sim_C \left( \int \|f\|^\alpha dm \right)^{1/\alpha} (E \|A(\bar{f})\|^p)^{1/p},$$

where  $C = C(\alpha, p)$  and  $A(\bar{f}) = \sum_{j=1}^{\infty} j^{-1/\alpha} \varepsilon_j \bar{f}(\tau_j^f)$ . Since  $\sup_j \|j^{-1/\alpha} \varepsilon_j \bar{f}(\tau_j^f)\| = 1$

$A(\bar{f})$  converges in  $L_B^r$  for every  $r > 0$ . Using Theorem 3.3 in Giné and Zinn [8] we obtain for every  $p \in (0, \alpha)$  and  $r > 0$

$$(E \|A(\bar{f})\|^p)^{1/p} \sim_C 1 + (E \left\| \sum_{j=j_0+1}^{\infty} j^{-1/\alpha} \varepsilon_j \bar{f}(\tau_j^f) \right\|^r)^{1/r},$$

where  $C = C(p, r)$  and  $j_0$  is the greatest integer not exceeding  $8^{-1} 3^{p/r}$ . By contraction principle we have

$$\begin{aligned} E \left\| \sum_{j=j_0+1}^{\infty} j^{-1/\alpha} \varepsilon_j \bar{f}(\tau_j^f) \right\|^r &= E \left\| \sum_{i=1}^{\infty} (j_0+i)^{-1/\alpha} \varepsilon_i \bar{f}(\tau_i^f) \right\|^r \\ &\leq 2^{-1} (1 + j_0)^{-r/\alpha} E \|A(\bar{f})\|^r \end{aligned}$$

and clearly

$$E \left\| \sum_{j=j_0+1}^{\infty} j^{-1/\alpha} \varepsilon_j \bar{f}(\tau_j^f) \right\|^r \leq 2 E \left\| A(\bar{f}) \right\|^r.$$

We have proven the following

Theorem 4.2.  $f \in S_B^\alpha$  if and only if  $f \in L_B^\alpha$  and the series

$\sum_{j=1}^{\infty} j^{-1/\alpha} \varepsilon_j \bar{f}(\tau_j^f)$  converges in  $L_B^r$  for some (each)  $r \geq 0$ . Moreover,

$$(E \left\| \int f dM \right\|^p)^{1/p} \sim_C \left( \int \|f\|^\alpha dm \right)^{1/\alpha} [1 + (E \left\| \sum_{j=1}^{\infty} j^{-1/\alpha} \varepsilon_j \bar{f}(\tau_j^f) \right\|^r)^{1/r}]$$

where  $C = C(\alpha, p, r)$ ,  $p \in (0, \alpha)$  and  $r > 0$ .

We shall study now the relationship between boundedness and convergence in (4.1). In view of Proposition 4.1 this will give us an additional information about  $S_B^\alpha$ . Let  $m(S) < \infty$ ,  $A = \sigma(A)$  and let

$$bS_B^\alpha = bS_B^\alpha(S, A, m) = \{f: S \rightarrow B: \sup_n \left\| \sum_{j=1}^n j^{-1/\alpha} \varepsilon_j f(\tau_j) \right\| < \infty \text{ a.s.}\}.$$

According to Proposition 4.1

$$S_B^\alpha = S_B^\alpha(S, A, m) = \{f: S \rightarrow B: \sum_{j=1}^{\infty} j^{-1/\alpha} \varepsilon_j f(\tau_j) \text{ conv. a.s.}\}.$$

Obviously  $S_B^\alpha \subset bS_B^\alpha$ . Let

$$\|f\|_{\alpha, p} = \sup_n (E \left\| \sum_{j=1}^n j^{-1/\alpha} \varepsilon_j f(\tau_j) \right\|^p)^{1/p} \quad \text{for } p \in (0, \alpha),$$

and let

$$||f||_{\alpha,0} = \sup_n E \left( || \sum_{j=1}^n j^{-1/\alpha} \xi_j f(\tau_j) || \wedge 1 \right).$$

It is standard to check that  $|| \cdot ||_{\alpha,0}$  is a complete F-norm on  $bS_B^\alpha$  as well as on  $S_B^\alpha$ . Moreover, by Proposition 4.1 all the F-norms  $|| \cdot ||_{\alpha,p}$  are equivalent on  $S_B^\alpha$ ,  $p \in [0, \alpha)$ .

**Lemma 4.3.** For every  $f \in bS_B^\alpha$  and  $p \in (0, \alpha)$   $||f||_{\alpha,p} < \infty$ . Moreover  $bS_B^\alpha \subset L_B^\alpha$ .

Proof. Without loss of generality we may assume that  $B = C[0,1]$ . Let  $\{P_k\}$  be a sequence of finite rank operators on  $B$  with  $||P_k|| \leq 1$  and such that  $P_k x \rightarrow x$  for every  $x \in B$ . Put  $f_k(s) = P_k f(s)$ ,  $s \in S$ . Clearly

$\sup_n || \sum_{j=1}^n j^{-1/\alpha} \xi_j f_k(\tau_j) || < \infty$  a.s. for every  $k$  and since  $\dim P_k B < \infty$  the series  $\sum_{j=1}^{\infty} j^{-1/\alpha} \xi_j f_k(\tau_j)$  converges a.s. By Proposition 4.1  $f_k \in S_B^\alpha$ .

Since  $|| \cdot ||_{\alpha,0}$  and  $|| \cdot ||_{\alpha,p}$  are equivalent on  $S_B^\alpha$  there exists  $\varepsilon > 0$  such that  $||g||_{\alpha,p} \leq 1$  for every  $g \in S_B^\alpha$  with  $||g||_{\alpha,0} \leq \varepsilon$ . Since  $f \in bS_B^\alpha$  there exists  $\delta > 0$  such that  $||\delta f||_{\alpha,0} \leq \varepsilon$ . Therefore  $||\delta f_k||_{\alpha,0} \leq \varepsilon$  for every  $k$  and because  $f_k \in S_B^\alpha$   $||\delta f_k||_{\alpha,p} \leq 1$ . By (3.3) and Proposition 4.1  $(\int ||\delta f_k||_{\alpha,p}^\alpha dm)^{1/\alpha} \leq \text{const } ||\delta f_k||_{\alpha,p} \leq \text{const}$ . Letting  $k \rightarrow \infty$  we get  $||f||_{\alpha,p} \leq \delta^{-1} < \infty$  as well as  $\int ||f||_{\alpha,p}^\alpha dm < \infty$ .

**Corollary 4.4.**  $S_B^\alpha \subset bS_B^\alpha \subset L_B^\alpha$ . For every  $p \in [0, \alpha)$   $|| \cdot ||_{\alpha,p}$  is a complete F-norm on  $bS_B^\alpha$  as well as on  $S_B^\alpha$ .

Proposition 4.1 and above Corollary give the following:

Theorem 4.5.  $S_B^\alpha$  is the smallest closed subspace of  $bs_B^\alpha$  containing all simple functions. In other words  $f \in S_B^\alpha$  if and only if for every  $\varepsilon > 0$  and  $p < \alpha$  there exists a simple function  $f_\varepsilon$  such that

$$E \left\| \sum_{j=1}^n j^{-1/\alpha} \xi_j (f - f_\varepsilon)(\tau_j) \right\|^p < \varepsilon$$

for all  $n \in \mathbb{N}$

Remarks:

(a) For the sufficiency  $f_\varepsilon$  need not be a simple function. It is enough to have  $f_\varepsilon \in S_B^\alpha$ .

(b)  $bs_B^\alpha = S_B^\alpha$  provided  $B$  does not contain a subspace isomorphic to  $c_0$ .

(c)  $bs_B^\alpha = S_B^\alpha = L_B^\alpha$  provided that  $B$  is of stable type  $\alpha$  (cf. [18] and Lemma 4.3). In particular these equalities hold for  $\alpha < 1$  and any Banach space  $B$ .

(d) In general  $S_B^\alpha \neq bs_B^\alpha \neq L_B^\alpha$ . Indeed, Sztencel [31] has showed that for every  $\alpha \geq 1$  there exists a Banach space  $B$  and a sequence  $\{x_n\} \subset B$  such that  $\sup_n E \left\| \sum_{j=1}^n \theta_j x_j \right\|^p < \infty$  for every  $p < \alpha$  and  $\sum_{j=1}^\infty \theta_j x_j$  diverges a.s., where  $\{\theta_j\}$  is a sequence of i.i.d. random variables with  $E \exp(it\theta_1) = \exp(-|t|^\alpha)$ . Let  $M$  be an  $S\alpha S$  random measure on Borel subsets of the unit interval and with the Lebesgue measure as the control measure. Put  $f_n = n^{1/\alpha} \sum_{j=1}^n x_j I_{[(j-1)/n, j/n]}$ . Since  $L(\int_0^1 f_n dM) = L(\sum_{j=1}^n \theta_j x_j)$ ,  $\{f_n\}$  is a bounded sequence in  $S_B^\alpha$ . Therefore  $\sup_n \int_0^1 \|f_n\|^\alpha dt < \infty$ . This yields  $c^{-1} = \sum_{j=1}^\infty \|x_j\|^\alpha < \infty$ . Let  $\{A_j\}$  be a partition of  $[0,1]$  such that  $|A_j| = c \|x_j\|^\alpha$  and define  $g_k = \sum_{j=1}^k (x_j / \|x_j\|) I_{A_j}$  and  $g = \lim_{k \rightarrow \infty} g_k$ . Since  $L(\int_0^1 g_k dM) = L(c^{1/\alpha} \sum_{j=1}^k \theta_j x_j)$ , by Proposition 3.1 and Ito-Nisio theorem  $g \notin S_B^\alpha$ . On the other hand  $g_k \in S_B^\alpha$  and by Proposition 4.1 for every  $n$  and  $p < \alpha$

$$\begin{aligned}
E \left\| \sum_{j=1}^n j^{-1/\alpha} \varepsilon_j g(\tau_j) \right\|^p &\leq \liminf_{k \rightarrow \infty} E \left\| \sum_{j=1}^n j^{-1/\alpha} \varepsilon_j g_k(\tau_j) \right\|^p \\
&\leq \liminf_{k \rightarrow \infty} \|g_k\|_{\alpha, p}^p \leq C \liminf_{k \rightarrow \infty} \left\| E \left\| \int_0^1 g_k dM \right\|^p \right\| \\
&\leq C c^{p/\alpha} \liminf_{k \rightarrow \infty} E \left\| \sum_{j=1}^k x_j \theta_j \right\|^p \\
&\leq \text{Const.}
\end{aligned}$$

This proves that  $g \in bS_B^\alpha$ .

To show that  $bS_B^\alpha \not\subset L_B^\alpha$  (in general) it is enough to choose  $\alpha \geq 1$  and Banach space  $B$  which is not of stable type  $\alpha$  and does not contain any subspace isomorphic to  $c_0$ . Then by (b)  $bS_B^\alpha = S_B^\alpha$  and  $S_B^\alpha \not\subset L_B^\alpha$  (cf. [18]).

### 5. Modification of the kernel of a stochastic integral process.

In this section we shall study processes  $X(t)$ ,  $t \in T$  which sample paths  $X(\cdot, \omega)$  belong to a separable Banach space  $V(T)$  of functions defined on  $T$ . Let  $C_T$  be the cylindrical  $\sigma$ -field of  $V(T)$  i.e. the smallest  $\sigma$ -field of subsets of  $V(T)$  such that all evaluations:  $\delta_t: V(T) \rightarrow \mathbb{R}$ , where  $\langle x, \delta_t \rangle = x(t)$ ,  $x \in V(T)$ ,  $t \in T$ , are measurable. The equality

$$(5.1) \quad C_T = \text{Borel } (V(T))$$

is necessary and sufficient for regarding stochastic processes with sample paths in  $V(T)$  as Borel measurable random elements in  $V(T)$ . Observe that the inclusion  $C_T \subset \text{Borel } (V(T))$  implies that every evaluation  $\delta_t$  is Borel measurable, and since  $\delta_t$  is linear, Banach theorem yields that  $\delta_t$  is continuous. Conversely, if all evaluations  $\delta_t$ ,  $t \in T$  are continuous, then (5.1) holds. Indeed, since evaluations separate points in  $V(T)$  one can easily deduce from Hahn-Banach theorem (see e.g. [24], Sec.2, Chap.2) that the set  $W = \{\sum_{j=1}^n a_j \delta_{t_j} : a_j \in \mathbb{R}, t_j \in T, n \geq 1\}$  is dense in  $[V(T)]'$  with respect to the weak-star topology. Since  $V(T)$  is separable,  $W$  is also sequentially weak-star dense in  $[V(T)]'$ , and consequently, every functional  $x' \in [V(T)]'$  is  $C_T$ -measurable. Again by separability of  $V(T)$  we get that  $\text{Borel } (V(T)) \subset C_T$ . Therefore (5.1) is equivalent to the assumption that all evaluations  $x \rightarrow x(t)$  are continuous.

**Theorem 5.1.** *Let  $V(T)$  be a separable Banach space of functions defined on  $T$  such that all evaluations  $x \rightarrow x(t)$  are continuous. Assume that the SoS stochastic process*

$$X(t) = \int_S h(t,s) dM(s), \quad t \in T,$$

has a modification  $X_0$  with sample paths in  $V(T)$ , where  $h: T \times S \rightarrow \mathbb{R}$  is such that  $h(t, \cdot) \in L^\alpha(S, \delta(A), m)$  for every  $t \in T$ .

Then there exists a function  $h_0: T \times S \rightarrow \mathbb{R}$  such that

- (i) for every  $s \in S$   $h_0(\cdot, s) \in V(T)$ ;
- (ii) for every  $t \in T$   $h_0(t, \cdot) = h(t, \cdot)$   $m$ -almost everywhere on  $S$ ;
- (iii) for every  $x' \in [V(T)]'$

$$\langle X_0(\cdot, \omega), x' \rangle = \int_S \langle h_0(\cdot, s), x' \rangle dM(s)(\omega),$$

for almost all  $\omega \in \Omega$ .

Proof. The proof is divided in three parts.

Claim 1. Assume that  $T$  is a compact metric space and  $V(T) = C(T)$  is the space of all continuous functions on  $T$  with the supremum norm. Then the conclusion of Theorem 5.1 is true.

Proof of claim 1. Let  $D$  be a finite subset of  $T \times T$ . First we shall show that

$$(5.2) \quad \int_S \max \{ |h(t_1, s) - h(t_2, s)|^\alpha : (t_1, t_2) \in D \} dm(s) \\ \leq C(E \max \{ |X(t_1) - X(t_2)|^p : (t_1, t_2) \in D \})^{\alpha/p},$$

where  $C = C(\alpha, p)$  and  $p \in (0, \alpha)$ .

Indeed, let us define an SaS random vector in  $\mathbb{R}^D$  by

$$Y = \{(X(t_1) - X(t_2))\}_{(t_1, t_2) \in D}$$

and consider  $\mathbb{R}^D$  as a Banach space with the norm  $\|a\| = \max\{|a(t_1, t_2)| : (t_1, t_2) \in D\}$ .

Then for every  $b \in \mathbb{R}^D$  we have

$$\begin{aligned}
E \exp(i \langle Y, b \rangle) &= E \exp \{i \int b(t_1, t_2) [X(t_1) - X(t_2)]\} \\
&= E \exp \{i \int_S [b(t_1, t_2) [h(t_1, s) - h(t_2, s)]] dM(s)\} \\
&= \exp \left( - \int_S |\langle f(s), b \rangle|^\alpha dm(s) \right),
\end{aligned}$$

where  $f: S \rightarrow \mathbb{R}^D$ ,  $f(s) = \{(h(t_1, s) - h(t_2, s))\}_{(t_1, t_2) \in D}$ . By (2.3) we get

$$\left( \int_S \|f(s)\|^\alpha dm(s) \right)^{1/\alpha} \leq C(E \|Y\|^p)^{1/p}$$

which yields (5.2).

Let  $d$  be a metric on  $T$ . For  $n=1, 2, \dots$  let  $T_n$  be a finite  $1/n$ -net in  $T$  and let  $T_\infty = \bigcup_{n=1}^\infty T_n$ . Clearly  $T_\infty$  is dense in  $T$ . Define for  $x: T \rightarrow \mathbb{R}$ ,  $\delta > 0$  and

$n \in \mathbb{N}$

$$\phi_{\delta, n}(x) = \max \{ |x(t_1) - x(t_2)| : t_1, t_2 \in T_n, d(t_1, t_2) < \delta \}$$

and

$$\phi_\delta(x) = \sup \{ |x(t_1) - x(t_2)| : t_1, t_2 \in T_\infty, d(t_1, t_2) < \delta \}.$$

Inequality (5.2) applied for  $D = \{(t_1, t_2) \in T_n \times T_n : d(t_1, t_2) < \delta\}$  yields

$$\int_S \{\phi_{\delta, n}[h(\cdot, s)]\}^\alpha dm(s) \leq C \left[ \int_\Omega \{\phi_{\delta, n}[X_0(\cdot, \omega)]\}^p dP(\omega) \right]^{\alpha/p}.$$

Since  $\phi_{\delta, n}[X_0(\cdot, \omega)] \leq 2 \sup_{t \in T} |X_0(t, \omega)| = 2 \|X_0(\cdot, \omega)\|$  letting  $n \rightarrow \infty$  we obtain

$$\int_S \{\phi_\delta[h(\cdot, s)]\}^\alpha dm(s) \leq C \left[ \int_\Omega \{\phi_\delta[X_0(\cdot, \omega)]\}^p dP(\omega) \right]^{\alpha/p}$$

and since sample paths of  $X_0$  are continuous we get

$$(5.3) \quad \lim_{\delta \downarrow 0} \phi_\delta[h(\cdot, s)] = 0$$

for  $m$ -almost all  $s \in S$ . Let  $s \in S$ . If (5.3) holds then  $h(\cdot, s)$  is uniformly continuous on  $T_\infty$  and there is unique continuous function  $h_0(\cdot, s)$  defined on  $T$  which is equal to  $h(\cdot, s)$  on  $T_\infty$ . If (5.3) fails, then we put  $h_0(t, s) = 0$  for all  $t \in T$ . Therefore  $h_0(\cdot, s) \in C(T)$  for all  $s \in S$  and  $h_0(t, \cdot) = h(t, \cdot)$   $m$ -almost everywhere for every  $t \in T_\infty$ . By stochastic continuity of  $X$  and continuity of  $t \rightarrow h_0(t, s)$  for every  $s \in S$  we get (ii) for every  $t \in T$ . Clearly (iii) is satisfied for all  $x'$ 's of the form  $x' = \sum_{j=1}^n a_j \delta_{t_j}$ . Since such functionals are sequentially weak-star dense in  $[V(T)]'$  (iii) follows.

Claim 2. Assume that  $V(T)$  is a closed subspace of  $C(T)$ , where  $T$  is a compact metric space. Then the conclusion of Theorem 5.1 holds.

Proof of claim 2. By claim 1 there exists a function  $h_0$  such that  $h_0(\cdot, s) \in C(T)$ , (ii) and (iii) hold ((iii) for every  $x' \in [C(T)]'$ ). Let

$$[V(T)]^\perp = \{x' \in [C(T)]' : \langle x, x' \rangle = 0 \text{ for all } x \in V(T)\}.$$

For every  $x' \in [V(T)]^\perp$  we have

$$\int_S \langle h_0(\cdot, s), x' \rangle dM(s) \stackrel{\text{a.s.}}{=} \langle X_0(\cdot, \cdot), x' \rangle = 0.$$

Therefore  $\langle h_0(\cdot, s), x' \rangle = 0$  for  $m$ -almost every  $s \in S$  and every  $x' \in [V(T)]^\perp$ .

Let  $\Gamma$  be a countable weak-star dense subset of  $[V(T)]^\perp$ . Define

$$S_0 = \{s \in S : \langle h_0(\cdot, s), x' \rangle = 0 \text{ for all } x' \in \Gamma\}.$$

Then  $m(S \setminus S_0) = 0$  and for every  $s \in S_0$   $h_0(\cdot, s) \in V(T)$ . Thus a function  $h_1$

defined by  $h_1(\cdot, s) = h_0(\cdot, s)$  for  $s \in S_0$  and  $h_1(\cdot, s) \equiv 0$  for  $s \notin S_0$  fulfills (i), (ii) and (iii) of Theorem 2.1.

Proof of the theorem in general. Let  $U' = \{x' \in [V(T)]' : \|x'\| \leq 1\}$ .

$U'$  equipped with the relative weak-star topology is a metrizable compact space. Let  $\phi: V(T) \rightarrow C(U')$  be defined by  $[\phi(x)](x') = \langle x, x' \rangle$ ,  $x \in V(T)$  and  $x' \in U'$ . It is easy to check that  $\phi$  is an isometric linear injection of  $V(T)$  into  $C(U')$ . Put  $V(U') = \phi[V(T)]$ . Since the mapping  $\Omega \ni \omega \rightarrow X_0(\cdot, \omega) \in V(T)$  is Borel measurable we obtain that  $Y: \Omega \rightarrow V(U')$  defined by  $Y(\cdot, \omega) = \phi[X_0(\cdot, \omega)]$  is also Borel measurable. Therefore  $Y(x', \cdot)$ ,  $x' \in U'$  is a stochastic process with continuous sample paths belonging to  $V(U')$ . Let  $W$  be the set of all linear combinations of  $\delta_t$ 's. By the discussion preceeding Theorem 5.1, for every  $x' \in U'$  there exists  $\{x'_n\} \subset W \cap U'$ ,  $x'_n = \sum_{j=1}^{k_n} a_{nj} \delta_{t_{nj}}$ , such that  $\langle x, x'_n \rangle \rightarrow \langle x, x' \rangle$

for every  $x \in V(T)$ . Therefore

$$\begin{aligned} Y(x', \cdot) &= \lim_{n \rightarrow \infty} Y(x'_n, \cdot) = \lim_{n \rightarrow \infty} \langle X_0(\cdot, \cdot), x'_n \rangle \\ &= \lim_{n \rightarrow \infty} \sum_{j=1}^{k_n} a_{nj} X(t_{nj}) \\ &= \lim_{n \rightarrow \infty} \int_S \left[ \sum_{j=1}^{k_n} a_{nj} h(t_{nj}, s) \right] dM(s), \end{aligned}$$

a.s., where the first equality holds point-wise by continuity of sample paths of  $Y$ . Thus  $\sum_{j=1}^{k_n} a_{nj} h(t_{nj}, \cdot)$  converge in  $L^\alpha(S, \sigma(A), m)$  to some function  $q(x', \cdot)$  and

we have

$$Y(x', \cdot) = \int_S q(x', s) dM(s) \quad \text{a.s.}$$

for every  $x' \in U'$ . Moreover  $g(a\delta_t, \cdot) = ah(t, \cdot)$   $m$ -almost everywhere on  $S$ , provided  $a\delta_t \in U'$ . According to claim 2 there exists  $g_0(x', s)$  such that  $g_0(\cdot, s) \in V(U')$  for all  $s \in S$  and  $g_0(x', \cdot) = g(x', \cdot)$   $m$ -almost everywhere for every  $x' \in U'$ .

Define a Borel measurable function  $G: S \rightarrow V(U')$  by  $[G(s)](x') = g_0(x', s)$  and  $H: S \rightarrow V(T)$  by  $H = \phi^{-1} \circ G$ . Let  $h_0(t, s) = \langle H(s), \delta_t \rangle$ . Clearly (i) is fulfilled. To show (ii) let  $t \in T$  and let  $a > 0$  be such that  $a\delta_t \in U'$ . Since  $H(s) = \phi^{-1}[G(s)]$  if and only if  $\langle H(s), x' \rangle = g_0(x', s)$  for all  $x' \in U'$  we obtain

$$ah_0(t, s) = \langle H(s), a\delta_t \rangle = g_0(a\delta_t, s) = g(a\delta_t, s) = ah(t, s),$$

where the last two equalities hold for  $m$ -almost all  $s \in S$ . (ii) is proved. (iii) follows by the weak-star density of  $W$  in  $[V(T)]'$ . The proof of Theorem 5.1 is complete.

Corollary 5.2. Let  $X_0$  and  $h_0$  be as in Theorem 5.1. Then the function  $f: S \rightarrow V(T)$  defined by  $[f(s)](t) = h_0(t, s)$  belongs to  $S_{V(T)}^\alpha$ . In particular,  $\int_S \|h_0(\cdot, s)\|_{V(T)}^\alpha dm(s) < \infty$ . Further, for every  $\varepsilon > 0$  and  $p \in (0, \alpha)$  there exist a finite sequence  $\{x_j\}_{j=1}^n \subset V(T)$  and pairwise disjoint sets  $A_1, \dots, A_n \in \mathcal{A}$  such that

$$E \|X_0(\cdot) - X_\varepsilon(\cdot)\|_{V(T)}^p < \varepsilon$$

where  $X_\varepsilon(t) = \sum_{j=1}^n x_j(t) M(A_j)$ ,  $t \in T$ .

Proof. Follows by (iii) of Theorem 5.1 and Proposition 3.1.

# 6. A characterization of S $\alpha$ S processes with absolute continuous trajectories.

A characterization of S $\alpha$ S processes ( $1 < \alpha < 2$ ) with absolute continuous sample paths in terms the so-called covariation function has been obtained by Cambanis and Miller in [3]. We shall characterize S $\alpha$ S processes with above sample path property using the representing function  $h$ . Moreover, in our case  $0 < \alpha < 2$ .

Let us recall that a function  $x: [a,b] \rightarrow \mathbb{R}$  is absolutely continuous if for every  $\epsilon > 0$  there exists a  $\delta > 0$  such that for every disjoint family  $\{(t_k, u_k)\}_{k=1}^n$  of subintervals of  $[a,b]$   $\sum_{k=1}^n |x(u_k) - x(t_k)| < \epsilon$ , provided  $\sum_{k=1}^n |u_k - t_k| < \delta$ . Then  $x$  is absolutely continuous if and only if there exists  $\dot{x} \in L^1[a,b]$  such that

$$x(u) = x(a) + \int_a^u \dot{x}(t) dt$$

for every  $a \leq u \leq b$ . Every absolutely continuous function  $x$  is differentiable almost everywhere and  $\frac{dx}{dt} = \dot{x}$  almost everywhere on  $[a,b]$ . Let  $AC^p[a,b]$  be the space of all absolutely continuous functions on  $[a,b]$  whose derivatives are integrable in the  $p$ -th power,  $p \geq 1$ .

Let

$$X(t) = \int_S h(t,s) dM(s), \quad t \in [a,b]$$

be an S $\alpha$ S process, where  $M$  is an S $\alpha$ S random measure defined on a  $\delta$ -ring of subsets of  $S$  and with the control measure  $m$ . As before  $h$  is a deterministic function such that  $h(t, \cdot) \in L^{\alpha}(S, \mathcal{A}, m)$  for every  $t \in [a,b]$ .

Theorem 6.1.  $X$  has a modification with sample paths in  $AC^p[a,b]$  if and

only if  $h$  admits a modification  $h_0$  such that

- (i) for every  $t \in [a, b]$   $h_0(t, \cdot) = h(t, \cdot)$   $m$ -almost everywhere,
- (ii) for every  $s \in S$   $h_0(\cdot, s) \in AC^p[a, b]$ ,
- (iii)  $Q_{\alpha, p}(\frac{\partial h_0}{\partial t}) < \infty$ ,

where

$$Q_{\alpha, p}(\alpha) = \begin{cases} \left( \int_S \left( \int_a^b |g(t, s)|^p dt \right)^{\alpha/p} dm(s) \right)^{1/\alpha} & \text{if } 0 < \alpha < p, \\ \left[ \int_S \int_a^b |g(t, s)|^\alpha \left[ 1 + \log_+ \frac{|g(t, s)|^\alpha \int_S \int_a^b |g(u, v)|^\alpha du \, dm(v)}{\int_a^b |g(u, s)|^\alpha du \int_S |g(t, v)|^\alpha dm(v)} \right] dt \, dm(s) \right]^{1/\alpha} & \text{if } \alpha = p, \\ \left( \int_a^b \left( \int_S |g(t, s)|^\alpha dm(s) \right)^{p/\alpha} dt \right)^{1/p} & \text{if } 1 \leq p < \alpha. \end{cases}$$

The proof of Theorem 6.1 is preceded by the lemma below which extends Proposition 4.2 in [5], where only the case  $p = \alpha$  has been considered. Since we use a similar argument in all cases of  $p$ , the case  $p = \alpha$  is also proven here.

**Lemma 6.2.** Let  $\varepsilon, \varepsilon_1, \varepsilon_2, \dots$  be i.i.d. symmetric random variables and  $0 < \alpha < 2$ . Then the series  $\sum_j j^{-1/\alpha} \varepsilon_j$  converges a.s. if and only if  $E|\varepsilon|^\alpha < \infty$ . Further, for every  $p > 0$

$$\left( E \left| \sum_j j^{-1/\alpha} \varepsilon_j \right|^p \right)^{1/p} \sim C \begin{cases} (E|\varepsilon|^p)^{1/p} & \text{if } \alpha < p \\ [E|\varepsilon|^\alpha (1 + \log_+ \frac{E|\varepsilon|^\alpha}{E|\varepsilon|^\alpha})]^{1/\alpha} & \text{if } \alpha = p \\ (E|\varepsilon|^\alpha)^{1/\alpha} & \text{if } p < \alpha \end{cases}$$

where  $C = C(\alpha, p)$ .

Proof. Since  $L(\xi_j) = L(\xi)$  we have for every  $t > 0$

$$(6.1) \quad t^{-\alpha} E |\xi|^\alpha - 1 \leq \sum_j P(|\xi_j| > tj^{1/\alpha}) \leq t^{-\alpha} E |\xi|^\alpha.$$

Therefore  $E |\xi|^\alpha < \infty$  is a necessary condition for the a.s. convergence of

$\sum_j j^{-1/\alpha} \xi_j$ . The sufficiency follows from (6.1) and the following estimates for every  $t > 0$

$$\begin{aligned} \sum_j E(j^{-1/\alpha} \xi_j I(|\xi_j| \leq tj^{1/\alpha}))^2 &= \sum_j j^{-2/\alpha} E |\xi|^2 I(|\xi| \leq tj^{1/\alpha}) \\ &= E |\xi|^2 \sum_{j \geq |\xi/t|^\alpha} j^{-2/\alpha} \\ (6.2) \quad &\leq E |\xi|^2 \left[ \left( \left| \frac{\xi}{t} \right|^\alpha \right)^{-2/\alpha} + \left( \left| \frac{\xi}{t} \right|^\alpha \right)^{1-2/\alpha} \right] \\ &= t^2 \left( 1 + \frac{\alpha}{2-\alpha} E \left| \frac{\xi}{t} \right|^\alpha \right). \end{aligned}$$

To estimate the moments of  $\sum j^{-1/\alpha} \xi_j$  we use Corollary 3.4 in Gine and Zinn [8] which gives

$$(6.3) \quad E \left| \sum_j j^{-1/\alpha} \xi_j \right|^p \underset{C}{\sim} E \sup_j |j^{-1/\alpha} \xi_j|^p + \left[ \sum_j E(j^{-1/\alpha} \xi_j I(|\xi_j| \leq \delta j^{1/\alpha}))^2 \right]^{p/2},$$

where  $\delta = \inf \{t > 0: \sum P(|\xi| > tj^{1/\alpha}) \leq 8^{-1} 3^{2vp}\}$  and  $C = C(p)$ ,  $p > 0$ .

By (6.1) we get

$$(6.4) \quad \delta \underset{C}{\sim} (E |\xi|^\alpha)^{1/\alpha} \quad \text{with } C = C(\alpha, p).$$

To obtain bounds for the first term on the right side of (6.3) we utilize

Lemma 3.2 in [8] which yields

$$(6.5) \quad E \sup_j |j^{-1/\alpha} \xi_j|^p \leq C \delta^p + R_{\alpha,p},$$

where  $C = C(p)$  and

$$\begin{aligned} R_{\alpha,p} &= \sum_j E |j^{-1/\alpha} \xi_j|^p I(|\xi_j| > \delta j^{1/\alpha}) \\ &= E |\xi|^p \sum_{j < |\xi/\delta|^\alpha} j^{-p/\alpha}. \end{aligned}$$

Since for every  $x > 0$   $(1-r)^{-1}(x^{1-r}-1) \leq \sum_{j < x} j^{-r} \leq 1 + (1-r)^{-1}(x^{1-r}-1)$

provided  $r \neq 1$  and  $r > 0$  we get for  $r = p/\alpha \neq 1$

$$\begin{aligned} R_{\alpha,p} &\leq E |\xi|^p [1 + (1 - p/\alpha)^{-1} ((|\xi|/\delta)^\alpha)^{1-p/\alpha} - 1] \\ &\leq \frac{p}{|\alpha - p|} E |\xi|^p + \frac{\alpha}{|\alpha - p|} \delta^{p-\alpha} E |\xi|^\alpha \end{aligned}$$

and

$$R_{\alpha,p} \geq \frac{\alpha}{|\alpha - p|} E |\xi|^p - \frac{\alpha}{|\alpha - p|} \delta^{p-\alpha} E |\xi|^\alpha$$

Above estimates in conjunction with (6.4) and (6.5) give

$$(6.6) \quad E \sup_j |j^{-1/\alpha} \xi_j|^p \leq C E |\xi|^p + (E |\xi|^\alpha)^{p/\alpha}$$

with  $C = C(\alpha, p)$  and  $p \neq \alpha$ . In the case  $p = \alpha$  elementary inequalities

$$\log_+ x \leq \sum_{j < x} j^{-1} \leq 1 + \log_+ x, \quad x > 0$$

$$E |\xi|^\alpha \log_+ \left| \frac{\xi}{\delta} \right|^\alpha \leq R_{\alpha, \alpha} \leq E |\xi|^\alpha (1 + \log_+ \left| \frac{\xi}{\delta} \right|^\alpha).$$

Using (6.4) and (6.5) we get

$$(6.7) \quad E \sup_j |j^{-1/\alpha} \xi_j|^\alpha \sim_C E |\xi|^\alpha (1 + \log_+ \frac{|\xi|^\alpha}{E |\xi|^\alpha}),$$

where  $C = C(\alpha)$ .

By (6.2) for  $t = \delta$  and (6.4) the second term on the right side of (6.3) is bounded from above by  $C(E |\xi|^\alpha)^{1/\alpha}$  with  $C = C(\alpha, p)$ . Therefore (6.6) for  $p \neq \alpha$  and (6.7) for  $p = \alpha$ , respectively, in conjunction with (6.3) conclude Lemma 6.2.

Proof of Theorem 6.1. Clearly  $AC^p[a, b]$  is a separable Banach space with the norm

$$||x|| = |x(a)| + \left( \int_a^b \left| \frac{dx}{dt} \right|^p dt \right)^{1/p},$$

and the evaluations  $x \rightarrow x(t)$  are continuous for every  $t \in [a, b]$ . By Theorem 5.1 there exists a function  $h_0$  satisfying (i) and (ii) provided  $X$  has a modification with sample paths in  $AC^p[a, b]$ . Then a function  $f: S \rightarrow AC^p[a, b]$  defined by  $[f(s)](t) = h_0(t, s)$  is Borel measurable and in view of Theorem 5.1 (iii) and Proposition 3.1  $f \in S_{AC^p[a, b]}^\alpha$ . Conversely, if  $f \in S_{AC^p[a, b]}^\alpha$

then  $\int_S f(s) dM(s)$  is a random element in  $AC^p[a, b]$  such that  $X_0(t) = \langle \int_S f dM, \delta_t \rangle = \int_S h_0(t, s) dM(s) = X(t)$  a.s., i.e.  $X_0$  gives a required modification of  $X$ .

Therefore  $X$  has a modification with sample paths in  $AC^p[a, b]$  if and only if (i), (ii) and  $f \in S_{AC^p[a, b]}^\alpha$ . Note also that without loss of generality we may

assume that  $h_0(a, \cdot) = 0$  (replacing  $X$  by  $X_a(t) = X(t) - X(a)$ ).

By Theorem 4.2  $f \in S_{AC^p[a, b]}^\alpha$  if and only if  $f \in L_{AC^p[a, b]}^\alpha$  and the series

$\sum j^{-1/\alpha} \epsilon_j \bar{f}(\tau_j^f)$  converges in  $L^q_{AC^p[a,b]}$  for some (each)  $q \geq 0$ . Here  $[\bar{f}(s)](t) =$

$h_0(t,s) ||f(s)||^{-1}$  and

$$||f(s)|| = |h_0(a,s)| + \left( \int_a^b \left| \frac{\partial h_0}{\partial t} \right| (t,s) |^p dt \right)^{1/p} = \left( \int_a^b |g(t,s)|^p dt \right)^{1/p},$$

where  $g(t,s) = \frac{\partial h_0}{\partial t}(t,s)$ . Moreover  $\tau_j^f$ 's are i.i.d. random elements in  $S$  such

$P(\tau_j^f \in A) = m_f(A)/m_f(S)$ ,  $A \in \sigma(A)$  and  $dm_f(s) = ||f(s)||^\alpha dm(s)$ . We have

$$(6.8) \quad m_f(S) = \int_S ||f(s)||^\alpha dm(s) = \int_S \left( \int_a^b |g(t,s)|^p dt \right)^{\alpha/p} dm(s)$$

and  $m_f(S) < \infty$  provided  $f \in S^\alpha_{AC^p[a,b]}$ . Further,

$$E \left| \left| \sum_{j=1}^n j^{-1/\alpha} \epsilon_j \bar{f}(\tau_j^f) \right| \right|^p = \int_a^b E \left| \sum_{j=1}^n j^{-1/\alpha} \epsilon_j g(t, \tau_j^f) ||f(\tau_j^f)||^{-1} \right|^p dt = \int_a^b u_n(t) dt.$$

Since  $\{u_n\}_{n=1}^\infty$  is a point-wise increasing sequence of functions and for every  $t$

$$u_n(t) \geq E \left| \sum_{j=n_0}^n j^{-1/\alpha} \epsilon_j g(t, \tau_j^f) ||f(\tau_j^f)||^{-1} \right|^p dt$$

as  $n, n_0 \rightarrow \infty$ , by the Monotone Convergence Theorem  $\sum j^{-1/\alpha} \epsilon_j \bar{f}(\tau_j^f)$  converges in

$L^p_{AC^p[a,b]}$  if and only if

$$\lim_{n \rightarrow \infty} E \left| \left| \sum_{j=1}^n j^{-1/\alpha} \epsilon_j \bar{f}(\tau_j^f) \right| \right|^p < \infty.$$

Case  $\alpha < p$ . By Lemma 6.2

$$\begin{aligned} \lim_{n \rightarrow \infty} E \left| \left| \sum_{j=1}^n j^{-1/\alpha} \epsilon_j \bar{f}(\tau_j^f) \right| \right|^p &\sim \int_a^b E |g(t, \tau_1^f)| |f(\tau_1^f)|^{-1} dt \\ &= [m_f(S)]^{-1} \int_a^b \int_S |g(t, s)|^p |f(s)|^{\alpha-p} dm(s) dt = [m_f(S)]^{-1} \int_S |f(s)|^\alpha dm(s) = 1. \end{aligned}$$

Therefore the condition  $f \in L_{AC^p[a,b]}^\alpha$  is also sufficient for  $f \in S_{AC^p[a,b]}^\alpha$ .

Case  $\alpha = p$ . By Lemma 6.2

$$\begin{aligned} \lim_{n \rightarrow \infty} E \left| \left| \sum_{j=1}^n j^{-1/\alpha} \epsilon_j \bar{f}(\tau_j^f) \right| \right|^p \\ \sim [m_f(S)]^{-1} \int_a^b \int_S |g(t, s)|^\alpha \left[ 1 + \log_+ \frac{|g(t, s)|^{\alpha m_f(S)}}{|f(s)|^\alpha |g(t, v)|^\alpha dm} \right] dm(s) dt \end{aligned}$$

which together with (6.8) ends this proof.

Case  $\alpha > p$ . By Lemma 6.2

$$\begin{aligned} \lim_{n \rightarrow \infty} E \left| \left| \sum_{j=1}^n j^{-1/\alpha} \epsilon_j \bar{f}(\tau_j^f) \right| \right|^p \\ \sim [m_f(S)]^{-p/\alpha} \left[ \int_a^b \int_S |g(t, s)|^\alpha dm(s) \right]^{p/\alpha} dt, \end{aligned}$$

which in conjunction with (6.8) completes the proof of Theorem 6.1.

Remarks:

Theorem 6.1 with appropriate modifications gives conditions for paths to have  $(n-1)$  continuous derivatives with the  $(n-1)$ th derivative in  $AC^p[a,b]$ .

An alternative proof of Theorem 6.1 can be obtained by an observation that  $AC^p[a,b]$  is isomorphic with  $\mathbb{R} \times L^p[a,b]$  ( $x \rightarrow (x(a), \frac{dx}{dt})$ ), and by the fact

that a full characterization of stable measures on  $L^p$ -spaces is known (see [18], [3] and [14] for  $p \neq \alpha$  and [27] for  $p = \alpha$ ). The proof given here, which is a straightforward application of Theorem 4.2, uses the same argument for all cases of  $p$  and  $\alpha$  and is self-contained.

The following result gives a full characterization of harmonizable  $S_{\alpha}S$  processes with absolutely continuous sample paths. Cambanis and Miller [3], using different methods, solved the case  $\alpha > 1$ .

Corollary 6.3. *Let  $M$  be an  $S_{\alpha}S$  random measure on the Borel  $\sigma$ -algebra of  $\mathbb{R}$  with the finite control measure  $m$ . Then  $X(t) = \int_{-\infty}^{\infty} e^{its} dM(s)$ ,  $t \in [a, b]$ , has a modification with sample paths in  $AC^p[a, b]$ ,  $1 \leq p < \infty$ , if and only if*

$$(6.9) \quad \int_{-\infty}^{\infty} |s|^{\alpha} dm(s) < \infty.$$

Proof. Since  $\mathbb{R}^2$  and  $\mathbb{C}$  are isomorphic Lemma 6.2 can be immediately extended to the case of complex valued random variables by considering two-dimensional random vectors instead of real random variables. Therefore Theorem 6.1 remains true when we replace a real valued function  $h$  by a complex one. In our case

$$h(t, s) = e^{its} \text{ and } \frac{\partial h}{\partial t} = ise^{its}.$$

It is elementary to check that in all cases of  $p$  and  $\alpha$  Theorem 6.1 yields the same condition (6.9).

Another important class of stable processes which is disjoint from the class of harmonizable ones (see Cambanis and Soltani [6]), is the class of  $S_{\alpha}S$  processes having the moving average representation.

**Corollary 6.4.** Let  $k: \mathbb{R} \rightarrow \mathbb{R}$  be an absolutely continuous function on every finite interval and such that  $\int_{-\infty}^{\infty} |k(s)|^{\alpha} ds < \infty$ . Let  $X(t) = \int_{-\infty}^{\infty} k(t-s) dM(s)$ ,  $t \in \mathbb{R}$ , where  $M$  is an SaS random measure defined on Borel bounded subsets of  $\mathbb{R}$  with the Lebesgue measure as the control measure.

Then  $X$  has a modification with sample paths in  $AC^p[a,b]$  for every  $-\infty < a < b < \infty$  if and only if

$$\int_{-\infty}^{\infty} (k_p(u))^{\alpha} du < \infty \quad \text{if } \alpha < p$$

$$\int_0^1 \int_{-\infty}^{\infty} \left| \frac{dk}{ds} \right|^{\alpha} \left( 1 + \log_+ \frac{\left| \frac{dk}{ds} \right|}{k_{\alpha}(s+t)} \right) ds dt < \infty \quad \text{if } \alpha = p$$

and

$$\int_{-\infty}^{\infty} \left| \frac{dk}{ds} \right|^{\alpha} ds < \infty \quad \text{if } \alpha > p,$$

$$\text{where } k_p(u) = \left( \int_u^{u+1} \left| \frac{dk}{ds} \right|^p ds \right)^{1/p}, \quad u \in \mathbb{R}.$$

**Proof.** Since  $X$  is a strictly stationary process it is enough to show that  $\{X(t): t \in [0,1]\}$  has a modification with sample paths in  $AC^p[a,b]$  if and only if the above conditions hold. Define  $h_0(t,s) = k(t-s)$ ,  $t,s \in \mathbb{R}$ . Then  $\frac{\partial h_0}{\partial t}(t,s) = \frac{dk}{ds}(t-s)$  and it is easy to check that the condition  $Q_{\alpha,p}(\frac{\partial h_0}{\partial t}) < \infty$  is equivalent to the above conditions for  $k$ .

### 7. Bounds for moments of a double $\alpha$ -stable stochastic integral.

Let  $h: [0,1] \times [0,1] \rightarrow \mathbb{R}$  be a jointly measurable function such that  $h(t,s) = 0$  for  $s \geq t$ . Let  $M$  be an S $\alpha$ S random measure on the Borel  $\sigma$ -algebra of  $[0,1]$ .

McConnell and Taqqu [20] have proved that a double stochastic integral

$$(7.1) \quad J(h) = \int_0^1 \int_0^1 h(t,s) dM(s) dM(t)$$

exists as the limit in  $L^p$  ( $p < \alpha$ ) of integrals of "dyadic" functions if and only if

$$(7.2) \quad P\left\{\int_0^1 \left|\int_0^1 h(t,s) dM(s)\right|^\alpha dt < \infty\right\} = 1$$

and in this case

$$(7.3) \quad (E|J(h)|^p)^{1/p} \underset{C}{\sim} \rho_{\alpha,p}(h),$$

$$\text{where } \rho_{\alpha,p}(h) = \{E[\int_0^1 \left|\int_0^1 h(t,s) dM(s)\right|^\alpha dt]^{p/\alpha}\}^{1/p},$$

$C = C(\alpha,p)$  and  $p < \alpha$ . Moreover,  $\rho_{\alpha,p}$  is a complete norm (quasi-norm if  $p < 1$ ) on the space of all functions  $h$  such that  $J(h)$  exists.

At the same time Rosinski and Woyczynski [27] studied double  $\alpha$ -stable integrals as iterated Ito-type stochastic integrals and proved that the finiteness of

$$(7.4) \quad N_\alpha^\alpha(h) = \int_0^1 \int_0^1 |h(t,s)|^\alpha \left[1 + \log_+ \frac{|h(t,s)|^\alpha \int_0^1 |h(u,v)|^\alpha du dv}{\int_0^1 |h(t,v)|^\alpha dv \int_0^1 |h(u,s)|^\alpha du}\right] dt ds$$

is necessary and sufficient for the existence of  $J(h)$  in this sense. They have proved also the equivalence of (7.2) and  $N_\alpha(h) < \infty$ . This shows, in

particular, that both approaches to define a double  $\alpha$ -stable integral are equivalent.

A natural problem which arises here is the relation between the norm  $\rho_{\alpha,p}$  and the functional  $N_\alpha$ . We shall prove that  $\rho_{\alpha,p} \sim N_\alpha$ , where  $C = C(\alpha,p)$ .

This in conjunction with (7.3) yields definitive bounds for moments of  $J(h)$ .

Let now  $h: [0,1]^2 \rightarrow \mathbb{R}$  be a jointly measurable function such that for every  $t \in [0,1]$   $h(t, \cdot) \in L^\alpha[0,1]$ . By Proposition 6.1 in [27]

$$X(t) = \int_0^1 h(t,s) dM(s), \quad t \in [0,1]$$

can be defined as a measurable stochastic process and by (7.2)  $X(\cdot, \omega) \in L^\alpha[0,1]$  for almost all  $\omega$ . Therefore,  $\int_0^1 \int_0^1 |h(t,s)|^\alpha ds dt < \infty$  (cf. [28]).

The lemma below justifies the interchanging of stochastic and usual integration and for the case  $\alpha > 1$  has been proved in [4] (Theorem 4.6) and in [20], Lemma 4.4. We give here a simpler and shorter proof of this result.

Lemma 7.1. *Let  $\alpha \geq 1$  and let  $h$  and  $X$  be as above. Then for every  $\phi \in L^{\alpha'}[0,1]$ , ( $\frac{1}{\alpha} + \frac{1}{\alpha'} = 1$ )*

$$(7.5) \quad \int_0^1 \phi(t) X(t) dt = \int_0^1 \left[ \int_0^1 \phi(t) h(t,s) dt \right] dM(s) \quad \text{a.s.}$$

Proof. Let  $\{U_j\}$  be a sequence of i.i.d. random variables uniformly distributed in  $[0,1]$  and defined on an auxiliary probability space  $(\Omega', P')$ , so that  $\{U_j\}$  is independent of  $\{X(t): t \in [0,1]\}$ . For every fixed  $\omega \in \Omega$  such that  $X(\cdot, \omega) \in L^\alpha[0,1]$  random variables  $\Omega' \ni \omega' \rightarrow \phi(U_j(\omega')) X(U_j(\omega'), \omega)$  are i.i.d. and  $E' |\phi(U_j) X(U_j, \omega)| = \int_0^1 |\phi(t) X(t, \omega)| dt < \infty$ . Therefore Kolmogorov's

SLLN yields the  $P'$ -a.s. convergence:

$$\frac{1}{n} \sum_{j=1}^n \phi(U_j) X(U_j, \omega) \rightarrow E' \phi(U_1) X(U_1, \omega) = \int_0^1 \phi(t) X(t, \omega) dt.$$

By Fubini's theorem, for almost all  $\omega' \in \Omega'$

$$(7.6) \quad \frac{1}{n} \sum_{j=1}^n \phi(U_j(\omega')) X(U_j(\omega'), \cdot) \rightarrow \int_0^1 \phi(t) X(t) dt$$

$P$ -a.s. on  $\Omega$ .

Define now i.i.d. random elements  $Y_j: \Omega' \rightarrow L^\alpha[0,1]$  by  $[Y_j(\omega')](s) = \phi(U_j(\omega')) h(U_j(\omega'), s)$ . Then

$$\begin{aligned} E \|Y_j\|_{L^\alpha[0,1]} &= \int_0^1 \|\phi(t) h(t, \cdot)\|_{L^\alpha[0,1]} dt \\ &= \int_0^1 \left[ \int_0^1 |\phi(t)|^\alpha |h(t, s)|^\alpha ds \right]^{1/\alpha} dt \\ &= \int_0^1 |\phi(t)| \left[ \int_0^1 |h(t, s)|^\alpha ds \right]^{1/\alpha} dt \\ &\leq \left[ \int_0^1 |\phi(t)|^{\alpha'} dt \right]^{1/\alpha'} \left[ \int_0^1 \int_0^1 |h(t, s)|^\alpha ds dt \right]^{1/\alpha} < \infty. \end{aligned}$$

By SLLN in  $L^\alpha[0,1]$

$$\frac{1}{n} \sum_{j=1}^n Y_j \rightarrow E' Y_1 = \int_0^1 \phi(t) h(t, \cdot) dt$$

$P'$ -a.s. Therefore for almost all  $\omega' \in \Omega'$

$$\begin{aligned}
 (7.7) \quad \frac{1}{n} \sum_{j=1}^n \phi(U_j(\omega')) X(U_j(\omega'), \cdot) &= \int_0^1 \left[ \frac{1}{n} \sum_{j=1}^n \phi(U_j(\omega')) h(U_j(\omega'), s) \right] dM(s) = \\
 &= \int_0^1 \left[ \frac{1}{n} \sum_{j=1}^n Y_j(\omega') \right] dM \xrightarrow{P} \int_0^1 \left[ \int_0^1 \phi(t) h(t, s) dt \right] dM(s).
 \end{aligned}$$

Since there exists  $\omega' \in \Omega'$  for which both (7.6) and (7.7) hold the proof is complete.

Corollary 7.2. Let  $h: [0,1]^2 \rightarrow \mathbb{R}$  and  $X$  be as in Lemma 7.1. Then the function  $f: [0,1] \rightarrow L^\alpha[0,1]$  defined by  $[f(s)](t) = h(t, s)$  belongs to  $S^\alpha_{L^\alpha[0,1]}$  and

$$X = \int_0^1 f(s) dM(s), \text{ a.s.}$$

Proof. By (7.5) for every  $\phi \in (L^\alpha[0,1])'$   $\langle X, \phi \rangle = \int_0^1 \langle f, \phi \rangle dM$  a.s.

Thus Proposition 3.1 completes the proof.

Theorem 7.3. Let  $1 \leq \alpha < 2$  and  $p < \alpha$ . Then there exists  $C = C(\alpha, p)$  such that for every  $h$

$$(E|J(h)|^p)^{1/p} \sim C N_\alpha(h).$$

Proof. In view of (7.3) it is sufficient to prove  $\rho_{\alpha,p} \sim C N_\alpha$ . By Corollary 7.2 and Theorem 4.2

$$\begin{aligned}
 \rho_{\alpha,p}(h) &= [E(\int_0^1 |X(t)|^\alpha dt)^{p/\alpha}]^{1/p} = (E\|X\|_{L^\alpha[0,1]}^p)^{1/p} = (E\|\int_0^1 f dM\|_{L^\alpha[0,1]}^p)^{1/p} \\
 &\leq C \left( \int_0^1 \|f(s)\|_{L^\alpha[0,1]}^\alpha ds \right)^{1/\alpha} [1 + (E\|\sum_{j=1}^n j^{-1/\alpha} f_j\|_{L^\alpha[0,1]}^\alpha)^{1/\alpha}].
 \end{aligned}$$

Here  $\int_0^1 ||f(s)||_{L^\alpha[0,1]}^\alpha ds = \int_0^1 \int_0^1 |h(t,s)|^\alpha dt ds$  and by Lemma 6.2

$$\begin{aligned}
 E || \sum_{j=1}^{\infty} j^{-1/\alpha} \epsilon_j \bar{f}(\tau_j^f) ||_{L^\alpha[0,1]}^\alpha &= E \int_0^1 || \sum_{j=1}^{\infty} j^{-1/\alpha} \epsilon_j ||f(\tau_j^f)||_{L^\alpha[0,1]}^{-1} |h(t, \tau_j^f)|^\alpha dt \\
 &\leq \int_0^1 E ||f(\tau_1^f)||_{L^\alpha[0,1]}^{-\alpha} |h(t, \tau_1^f)|^\alpha [1 + \log_+ \frac{||f(\tau_1^f)||_{L^\alpha[0,1]}^{-\alpha} |h(t, \tau_1^f)|^\alpha}{E ||f(\tau_1^f)||_{L^\alpha[0,1]}^{-\alpha} |h(t, \tau_1^f)|^\alpha}] dt \\
 &= (\int_0^1 \int_0^1 |h(t,s)|^\alpha dt ds)^{-1} N_\alpha^\alpha(h)
 \end{aligned}$$

which finishes the proof.

Acknowledgements. I would like to thank Stamatis Cambanis for fruitful discussions and helpful remarks during preparation of this paper.

## REFERENCES

1. Bretagnolle, J., Dacunha-Castelle, D., and Krivine, J.-L. (1966), Lois stable et espaces  $L^p$ , Ann. Inst. H. Poincaré II 231-259.
2. Cambanis, S., Hardin, D.C., and Weron, A. (1984), Ergodic properties of stationary stable processes, Center for Stochastic Processes Tech. Rept. No. 32. Univ. of North Carolina, Chapel Hill.
3. Cambanis, S. and Miller, G. (1980), Some path properties of  $p$ th order and symmetric stable processes, Ann. Prob. 8, 1148-1156.
4. Cambanis, S., and Miller, G. (1981), Linear problems in  $p$ th order and stable processes, SIAM J. Appl. Math. 41, 43-69.
5. Cambanis, S., Rosinski, J. and Woyczynski, W.A. (1984), Convergence of quadratic forms in  $p$ -stable random variables and  $p$ -radonifying operators, Ann. Probability, to appear.
6. Cambanis, S., and Soltani, A.R. (1984), Prediction of stable processes: spectral and moving average representations, Z. Wahrsch. Verw. Gebiete 66, 593-612.
7. Gine, E., Marcus, M.B. and Zinn, J. (1984), A version of Chevet's theorem for stable processes, Preprint.
8. Gine, E., and Zinn, J. (1983) Central limit theorems and weak laws of large numbers in certain Banach spaces, Z. Wahrsch. verw. Gebiete 62, 323-354.
9. Hardin, C.D. (1982), On the spectral representation of symmetric stable processes, J. Multivariate Anal. 12, 385-401.
10. Hosoya, Y. (1982), Harmonizable stable processes, Z. Wahrsch. verw. Gebiete 60, 517-533.
11. Kanter, M. (1972), A representation theorem for  $L^p$  spaces, Proc. Amer. Math. Soc. 31, 472-474.
12. Kuelbs, J. (1973), A representation theorem for symmetric stable processes and stable measures on  $H$ , Z. Wahrsch. Verw. Gebiete 26, 259-271.
13. LePage, R., Woodroffe, M., Zinn, J. (1981), Convergence to a stable distribution via order statistics, Ann. Prob. 9, 624-632.
14. Linde, W. (1983), Infinitely divisible and stable measures on Banach spaces, Teubner-Texte zur Mathematik. Band 58, Leipzig.
15. Marcus, M.B. (1983), Extreme values for sequences of stable random variables, Proceedings of NATO Conference on Statistical extremes, Vimeiro, Portugal, 1983.

16. Marcus, M.B. and Pisier, G. (1981), Random Fourier series with applications to harmonic analysis, Ann. Math. Studies, Vol. 101, Princeton Univ. Press., Princeton, N.J.
17. Marcus, M.B. and Pisier, G. (1984), Characterizations of almost surely continuous  $p$ -stable random Fourier series and strongly stationary processes, Acta Mathematica 152, 245-301.
18. Marcus, M.B. and Woyczynski, W.A. (1979), Stable measures and central limit theorem in spaces of stable type, Trans. Amer. Math. Soc. 251, 71-102.
19. Masry, E., and Cambanis, S. (1984), Spectral density estimation for stationary stable processes, Stochastic Proc. Appl. 18, 1-31.
20. McConnell, T.R., and Taqqu, M.S. (1984), Double integration with respect to symmetric stable processes, Cornell U. Dept. of Oper. Res., Tech. Rept. No. 618.
21. Pisier, G. (1975), Le theoreme limite central et la loi du logarithme iteree dans les espaces de Banach, Seminaire Maurey-Schwartz 1975-1976
22. Prékopa, A. (1956), On stochastic set functions I, Acta Math. Acad. Scient. Hung. 7, 215-263.
23. Pourahmadi, M. (1984). On minimality and interpolation of harmonizable stable processes. SIAM J. Appl. Math. 44, 1023-1030.
24. Robertson, A. and W. Robertson, Topological vector spaces, Cambridge 1964.
25. Rootzen, H. (1978), Extremes of moving averages of stable processes. Ann. Prob. 6, 847-869.
26. Rosinski, J. (1985), Random integrals of Banach space valued functions, Studia Math. 78, 15-38.
27. Rosinski, J. and Woyczynski, W.A. (1984), On Ito stochastic integration with respect to  $p$ -stable motion: inner clock, integrability of sample paths, double and multiple integrals, Ann. Probability, to appear.
28. Rosinski, J. and Woyczynski, W.A. (1984), Moment inequalities for real and vector  $p$ -stable stochastic integrals, Center for Stochastic Processes Technical Report No. 87, University of North Carolina, Chapel Hill.
29. Schilder, M. (1970), Some structure theorems for the symmetric stable laws, Ann. Math. Statist. 41, 412-421.
30. Schreiber, M. (1972), Quelques remarques sur les caracterisations des espaces  $L^p$ ,  $0 \leq p < 1$ , Ann. Inst. H. Poincare 8, 83-92.
31. Sztencel, R. (1981), On boundedness and convergence of some Banach space valued random series, Probability and Math. Statistics 2, 83-88.
32. Weron, A. (1983). Harmonizable stable processes on groups, Center for Stochastic Processes Tech. Rept. No. 32. Univ. of North Carolina, Chapel Hill.

**END**

**FILMED**

**6-85**

**DTIC**