

MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A154 229

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1 REPORT NUMBER NSWC TR 84-289	2 GOVT ACCESSION NO AD-A154229	3 RECIPIENT'S CATALOG NUMBER
4 TITLE (and Subtitle) AN ALGORITHM FOR THE INFLUENCE NUMBERS IN THE EXPANSION OF THE LUMPED GRAVITATIONAL POTENTIAL COEFFICIENTS IN THE PRESENCE OF RESONANT INCLINATION VARIATION		5 TYPE OF REPORT & PERIOD COVERED FINAL
		6 PERFORMING ORG REPORT NUMBER
7 AUTHOR(s) WALTER J. GROEGER		8 CONTRACT OR GRANT NUMBER(s)
9 PERFORMING ORGANIZATION NAME AND ADDRESS NAVAL SURFACE WEAPONS CENTER (CODE K12) DAHLGREN, VA 22448		10 PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS O & M, DA
11 CONTROLLING OFFICE NAME AND ADDRESS DEFENSE MAPPING AGENCY WASHINGTON, DC 20305		12 REPORT DATE JULY 1984
		13 NUMBER OF PAGES 33
14 MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15 SECURITY CLASS (of this report) UNCLASSIFIED
		15a DECLASSIFICATION DOWNGRADING SCHEDULE
16 DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17 DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18 SUPPLEMENTARY NOTES		
19 KEY WORDS (Continue on separate page if necessary and identify by block number) Satellite geodesy Satellite orbit analysis Orbital inclination resonance Lumped gravitational potential coefficients		
20 ABSTRACT (Continue on separate page if necessary and identify by block number) Near resonance, the rate of change of the orbital inclination of an earth satellite may be expressed as an aggregate of several more or less complicated functions involving the parameters characterizing the resonance type, the orbital elements, certain quantities named "lumped coefficients," and time. The lumped coefficients are linear expressions consisting of the scalar products of sets of expansion coefficients of the earth's gravitational potential (in the sense that each lumped coefficient is associated		

DD FORM 13 SEP 73 473

EDITION OF NOV 65 IS OBSOLETE
S. N. 0102 15 014 6001

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

20. (Continued)

with a specific potential term order) times a second set of parameters customarily called "influence numbers." The latter are essentially functions of orbital semi-major axis, inclination, and eccentricity. The present report specifies a computer algorithm for the influence numbers. Included are numerical trial data and a mathematical derivation of the influence number algorithm.


FOREWORD

This technical report documents a computer algorithm for the "influence numbers" that govern the relationship between certain subsets of the earth's gravitational potential expansion coefficients and "lumped coefficients," each of which summarizes the gravitational effect of an entire group of potential expansion coefficients on the inclination resonance of earth satellites.

The work was performed in the Space and Surface Systems Division as part of a larger effort, with the purpose of establishing equations of condition for groups of gravitational field parameters as a contribution to the validation of the WGS-84 Geodetic Parameter Solution.

The present report was reviewed by C. W. Duke, Jr., Head of the Space and Surface Systems Division; R. L. Kulp, Head of the Space and Ocean Geodesy Branch; and Dr. C. Oesterwinter, Research Associate.

Released by:



THOMAS A. CLARE, Head
Strategic Systems Department

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



DTIC ELECTE
S APR 25 1985 D
B

CONTENTS

	<u>Page</u>
INTRODUCTION.....	1
ALGORITHM FOR THE INFLUENCE NUMBERS.....	3
INPUT DATA.....	3
THE INFLUENCE NUMBERS.....	4
INCLINATION FUNCTION SUBROUTINE.....	5
ECCENTRICITY FUNCTION SUBROUTINE.....	6
ALGORITHM FOR THE LUMPED COEFFICIENTS.....	7
REFERENCES.....	9
APPENDIXES	
A--NUMERICAL TRIAL DATA FOR THE INFLUENCE NUMBER ALGORITHM.....	A-1
B--MATHEMATICAL BACKGROUND.....	B-1
DISTRIBUTION.....	(1)

INTRODUCTION

An artificial earth satellite is said to be in β/a resonance* with the earth's gravitational field when it performs β nodal periods while the earth revolves a times relative to the satellite's precessing orbit plane. Near resonance, the perturbations caused by those tesseral components of the earth's gravitational potential with which the satellite happens to be in resonance generally exceed in magnitude all the remaining tesseral perturbations.

Near resonance, the rate of change of the orbital inclination of the satellite may be expressed as an aggregate of several more or less complicated functions involving the parameters characterizing the resonance type, the orbital elements, certain quantities named "lumped coefficients," and time. The lumped coefficients are linear expressions consisting of the scalar products of sets of expansion coefficients of the earth's gravitational potential (in the sense that each lumped coefficient is associated with a specific potential term order) times a second set of parameters customarily called "influence numbers." The latter are essentially functions of orbital semimajor axis, inclination, and eccentricity.

Satellites with sufficiently short orbital periods will, as their orbits decay, pass through resonance or, frequently, through a succession of different resonances. While orbital resonance effects are not restricted to the orbit inclination, conditions for acquiring useful inclination resonance data have been especially favorable, giving rise to a large body of investigations (References 1 through 78). Most of these are undertaking to evaluate the lumped coefficients for empirical time sequences of resonant inclination values. There is almost always an attempt made to subsequently infer numerical values for the geopotential expansion coefficients related to the lumped coefficients found. Most of the investigations seen are confined to specific resonance cases and to specific satellites. Some however attempt a synopsis (see especially Reference 62). An effort was recently made at the Naval Surface Weapons Center (NSWC) to systematically review the entire body of published research work and to compile all lumped coefficients listed. A system of linear conditional equations complete with the customary statistical information was generated for the geopotential expansion

*For details, see the discussions of resonance in References 1 and 33.

coefficients, the latter being the system unknowns, and ranging up to degree 42, each identified lumped coefficient contributing one equation. A general, numerical solution for those geopotential parameters contributing to the inclination resonance cases treated in the just-mentioned collection of literature will in due course be executed.

As already stated, a necessary ingredient of the conditional equations are the influence numbers. These are, essentially, geometry factors reflecting the orbital parameters at resonance. As a generally valid theory of the influence numbers could not be located in any of our sources, it was derived starting from the Lagrange planetary equation for the inclination variation. To document the resulting computer algorithm is the purpose of the present report.

ALGORITHM FOR THE INFLUENCE NUMBERS

INPUT DATA

R "Radius of the Earth" (semimajor axis of reference ellipsoid)
in km. Default value: $R = 6378.140$ km.

a Semimajor axis of satellite, in km.

e Orbital eccentricity

i Orbital inclination

β } Integers, characterizing β/α orbital resonance.

α } Normally, $\alpha = 1$ or 2 . In most cases, β will be larger than
5 and smaller than 50.

γ Integer, denoting magnitude of the resonance term (order of
"overtone"). Generally, γ will be equal to 1 or 2. Generally,
it will be a positive integer.

q Integer (positive, negative or zero) associated with the magnitude
of the orbital eccentricity. For the present purpose, the
following values of q may occur: $0, \pm 1, \pm 2, \pm 3$.

THE INFLUENCE NUMBERS

$$k = \alpha\gamma - q \quad (1)$$

$$m = \beta\gamma \quad (2)$$

$$K = (\beta - \alpha)\gamma + q \quad (3)$$

If K is even

$$l_r = \beta\gamma + 2r = m + 2r \quad (4)$$

$$m_r = \beta\gamma \quad (5)$$

$$p_r = \frac{1}{2}(\beta - \alpha)\gamma + \frac{1}{2}q + r \quad (6)$$

$$= \frac{1}{2}K + r \quad (6a)$$

If K is odd

$$l_r = \beta\gamma + (2r + 1) = m + 1 + 2r \quad (7)$$

$$m_r = \beta\gamma \quad (8)$$

$$p_r = \frac{1}{2}(\beta - \alpha)\gamma + \frac{1}{2}q + \frac{1}{2} + r \quad (9)$$

$$= \frac{1}{2}(K + 1) + r \quad (9a)$$

$$r = 0, 1, 2, 3, 4, \dots, r_{\max} \quad (10)$$

Expected maximum value for r_{\max}

$$r_{\max} \leq 50 \quad (11)$$

The influence numbers are then

$$Q_{\ell}^{q,k} = (-1)^r \left(\frac{R}{a}\right)^{2r} \frac{G_{\ell pq}}{G_{\ell_0 p_0 q}} \frac{FAN_{\ell mp}}{FAN_{\ell_0 m p_0}} \quad (12)$$

where

$$\ell = \ell_r \quad (13a)$$

$$p = p_r \quad (13b)$$

$$\ell_0 = \ell(r=0) \quad (13c)$$

$$p_0 = p(r=0) \quad (13d)$$

Note that the influence numbers are dimensionless quantities.

INCLINATION FUNCTION SUBROUTINE

$$FA_{\ell mp} \equiv FA_{\ell mp}(i) \equiv \frac{(\ell+m)!}{2^{\ell} p! (\ell-p)!} * \sum_{\eta_{\min}}^{\max} \left[(-1)^{\eta} \binom{2\ell-2p}{\eta} \binom{2p}{\ell-m-\eta} \right. \\ \left. * C^{3\ell-m-2p-2\eta} * S^{m-\ell+2p+2\eta} \right] \quad (14)$$

$$C = \cos \left(\frac{1}{2} i\right) \quad (15)$$

$$S = \sin \left(\frac{1}{2} i\right) \quad (16)$$

$$\eta_{\min} = \max[0, (\ell-m-2p)] \quad (17)$$

$$\eta_{\max} = \min[(\ell-m), (2\ell-2p)] \quad (18)$$

$$N_{\ell m} = + \sqrt{\frac{(\ell - m)!(2\ell + 1)(2 - \delta_m^0)}{(\ell + m)!}} \quad (19)$$

$$\delta_m^0 = 0 \quad \text{for } m \neq 0 \quad (20a)$$

$$= 1 \quad \text{for } m = 0 \quad (20b)$$

$$FAN_{\ell mp} = N_{\ell m} * FA_{\ell mp} \quad (21)$$

ECCENTRICITY FUNCTION SUBROUTINE

$$G_{\ell pq} \equiv G_{\ell pq}(e) \equiv (-1)^{|q|} (1 + \beta^2)^{\ell} \beta^{|q|} * \sum_{k=0}^N V_{\ell pqk} W_{\ell pqk} \beta^{2k} \quad (22)$$

$$\beta = \frac{e}{1 + \sqrt{1 - e^2}} \quad (23)$$

Stop summation over k when $[G]_{N+1}$ and $[G]_N$ agree to six significant figures.

$$\text{For } p \leq \ell/2, \quad p' = p \\ q' = q \quad (24)$$

$$\text{For } p > \ell/2, \quad p' = \ell - p \\ q' = -q \quad (25)$$

$$V_{\ell p q k} \equiv \sum_{r=0}^h \binom{2p' - 2\ell}{h - r} \frac{(-1)^r}{r!} \left(\frac{(\ell - 2p' + q')e}{2\beta} \right)^r \quad (26)$$

$$\text{For } q' > 0, \quad h = k + q' \quad (27)$$

$$\text{For } q' \leq 0, \quad h = k \quad (28)$$

$$W_{\ell p q k} \equiv \sum_{r=0}^h \binom{-2p'}{h - r} \frac{1}{r!} \left(\frac{(\ell - 2p' + q')e}{2\beta} \right)^r \quad (29)$$

$$\text{For } q' \geq 0, \quad h = k \quad (30)$$

$$\text{For } q' < 0, \quad h = k - q' \quad (31)$$

To avoid difficulties arising from use of negative arguments, calculate the binomial coefficients occurring in Equations 26 and 29 as follows:

$$\binom{A}{B} = \frac{1}{B!} \left\{ [A - 0] * [A - 1] * [A - 2] * \dots * [A - (B - 1)] \right\} \quad (32)$$

B may be expected to be a positive integer always. A may assume negative values.

ALGORITHM FOR THE LUMPED COEFFICIENTS

Let $C_{\ell m}$ and $\bar{S}_{\ell m}$ be the normalized expansion coefficients of the geopotential and $C_{\ell m}$ and $S_{\ell m}$ their unnormalized counterparts, then

$$\bar{C}_{\ell m} = C_{\ell m} / N_{\ell m} \quad (33)$$

$$\bar{S}_{\ell m} = S_{\ell m} / N_{\ell m} \quad (34)$$

NSM 1R 84-289

APPENDIX B

MATHEMATICAL BACKGROUND

NSWC TR 84-289

R (km)	6378.135	6378.135
i (deg)	38.92	74.02
a (km)	7403.314	8095
e	.082	.06
r	13	29
r	1	2
r	2	1
q	0	0
r	1	1
K	24	27
m	26	29
\bar{G}_0	26	30
\bar{P}_0	12	14
c	28	32
p	13	15
s	2	2
EA ₀	0.000415812	-0.021323873
EA _{N₀}	0.000022206133	0.434536005
G	2.966430720	2.218928773
G_0	2.613511626	2.039722371
Q(C, q, k)	-15.5991083	0.93315660

NSWC TR 84-289

R (km)	6378.135	6378.135	6378.135
i (deg)	32.97	51.6	51.6
a (km)	6902.251	9714.537	9714.537
e	.003	.180	.180
ω	15	9	9
τ	1	1	1
ρ	1	1	1
q	0	-1	-3
r	6	3	3
κ	14	7	5
m	15	9	9
ζ_0	15	10	10
P_0	7	4	3
ζ	27	16	16
p	13	7	6
k	1	2	4
FAN	0.396980759	-0.312034989	0.742922385
FAN_0	0.000215953	0.591373731	0.643760579
γ	1.001692824	3.610028758	0.35180026
β	1.000531091	1.070359599	0.998985059
ζ_{total}	3.41966	0.17123417	0.1711901

APPENDIX A

NUMERICAL TRIAL DATA FOR THE INFLUENCE NUMBER ALGORITHM

76. D. M. C. Walker, *Analysis of the U.S. Navy Orbits of 1973-78 and 1974-78 at 15th-Order Resonance*, Royal Aircraft Establishment Technical Report 80080, (Farnborough, Hants., England, July 1980).
77. D. M. C. Walker, "Analysis of the U.S. Navy Orbits of 1973-78 and 1974-78 at 15th-Order Resonance," *Geophys. J. R. Astr. Soc.* (1981), 67, pp. 1-18.
78. S. N. Yashkin, "Satellite Resonance Orbits and Their Use in Dynamic Problems of Space Geodesy," *Izvestiya VUZ: Geodeziya i aerofotogrammetriya*, No. 2, 1981, pp. 36-42. Technical Translation: DMAAC-TC-3163.
79. W. M. Kaula, "Theory of Satellite Geodesy," Blaisdell Publishing Company, Waltham, Massachusetts, 1966. See especially Page 34.

67. S. M. Klosko and C. A. Wagner, "Spherical Harmonic Representation of the Gravity Field From Dynamic Satellite Data," Paper presented at XVII General Assembly of the International Union of Geodesy and Geophysics, Canberra, Australia, December 1979.
68. S. M. Klosko and C. A. Wagner, "Spherical Harmonic Representation of the Gravity Field from Dynamic Satellite Data," *Planet. Space Sci.* 1982, Vol. 30, pp. 5-28.
69. C. A. Wagner, "The Accuracy of the Low Degree Geopotential: Implications for Ocean Dynamics," Undated Manuscript, NOAA/NOS/National Geodetic Survey, Rockville, MD.
70. D. M. C. Walker, "Analysis of the Orbit of Ariel 1, 1962-15A, Near 15th-Order Resonance," *Planet. Space Sci.* 1975, Vol. 23, pp. 565-574.
71. D. M. C. Walker, *29th-Order Harmonics in the Geopotential From the Orbit of Ariel 1 at 29:2 Resonance*, Royal Aircraft Establishment Technical Report 76110, (Farnborough, Hants., England, August 1976).
72. D. M. C. Walker, "29th-Order Harmonics in the Geopotential From the Orbit of Ariel 1 at 29:2 Resonance," *Planet Space Sci.* 1977, Vol. 25, pp. 337-342.
73. D. M. C. Walker, *COSMOS 462 (1971-106A): Orbit Determination and Analysis*, Royal Aircraft Establishment Technical Report 78089, (Farnborough, Hants., England, August 1978).
74. D. M. C. Walker, *SAMOS 2 (1961 ALPHA 1): Orbit Determination and Analysis at 31:2 Resonance*, Royal Aircraft Establishment Technical Report 80022, (Farnborough, Hants., England, February 1980).
75. D. M. C. Walker, "SAMOS 2 (1961 ALPHA 1): Orbit Determination and Analysis at 31:2 Resonance," *Planet. Space Sci.*, 1980, Vol. 28, pp. 1059-1072.

56. C. A. Wagner and S. M. Klosko, *15th-Order Resonance on the Decaying Orbit of TETR-3*, GSFC Technical Report X-592-73-279, Greenbelt, MD, September 1973.
57. C. A. Wagner, *Effect of Resonance-Ublateness Coupling on a Satellite Orbit*, GSFC Technical Report X-920-74-334, Greenbelt, MD, November 1974.
58. C. A. Wagner, *14 Years of Resonance on Vanguard Orbits*, GSFC Technical Report X-921-75-179, Greenbelt, MD, June 1975.
59. C. A. Wagner, "Thirteenth-Order Orbital Resonance on a Diademe 2 Fragment," *J. Geophys. Res.*, Vol. 80, No. 26, 10 September 1975, p. 3791.
60. C. A. Wagner and S. M. Kloski, "15th-Order Resonance Terms Using the Decaying Orbit of TETR-3," *Planet. Space Sci.* 1975, Vol. 23, pp. 541-549.
61. Smith, Lerch, Marsh, Wagner, Kolenkiewicz, Khan, "Contributions to the National Geodetic Satellite Program by Goddard Space Flight Center," *J. Geophys. Res.*, Vol. 81, No. 5, 10 February 1976, pp. 1006-1026.
62. C. A. Wagner, *The Accuracy of Goddard Earth Models*, GSFC Technical Report X-921-76-187, June 1976.
63. C. A. Wagner and S. M. Klosko, "Gravitational Harmonics From Shallow Resonant Orbits," *Celestial Mechanics*, Vol. 16, (1977), pp. 143-163.
64. C. A. Wagner and F. J. Lerch, "The Accuracy of Geopotential Models," *Planet. Space Sci.* 1978, Vol. 26, pp. 1081-1140.
65. C. A. Wagner, *Gravitational Spectra From the Tracking of Planetary Spacecraft in Eccentric Orbits*, GSFC NASA Technical Memorandum 79721, Greenbelt, MD, March 1979.
66. Lerch, Klosko, Laubscher, Wagner, "Gravity Model Improvement Using GEOS-3 (GEM-9 and 10)," *J. Geophys. Res.*, Vol. 84, No. B8, 30 July 1979, pp. 3897-3915.

46. D. E. Smith, "A Determination of the Odd Harmonics in the Geopotential Function," *Planet Space Sci.* 1963, Vol. 11, pp. 789-795.
47. D. E. Smith, "The Zonal Harmonics in the Geopotential: Our Present Knowledge and Future Requirements." Presented at NASA Special Projects Branch Astrodynamics Conference, GSFC, Greenbelt, MD, October 1968.
48. A. S. Sochilina, "On the Motion of a Satellite in Resonance With its Rotating Planet," *Celestial Mechanics* 26, (1982), pp. 337-352.
49. C. A. Wagner, "Determination of Low Order Resonant Gravity Harmonics From the Drift of Two Russian 12-Hour Satellites," AAS National Specialist Symposium "Orbital Resonance," 15-16 January 1968.
50. C. A. Wagner, "A Survey of Existing Satellites in Resonant Orbits for Geodetic Purposes," GSFC, Greenbelt, Md, July 1968.
51. C. A. Wagner, *Resonant Perturbations of Earth Satellites in 2-Day Commensurable Orbits*, GSFC Technical Report X-643-68-373, Greenbelt, MD, September 1968.
52. Bruce C. Douglas, Chesterton F. Martin, Donald G. Williamson, and Carl A. Wagner, "Error Analysis of Resonant Orbits for Geodesy," *Celestial Mechanics* 1, (1969), pp. 252-270.
53. C. A. Wagner, *Earth Zonal Harmonics From Rapid Numerical Analysis of Long Satellite Arcs*, GSFC Technical Report X-553-72-341, Greenbelt, MD, August 1972.
54. C. A. Wagner, *11th-Order Resonance Terms in the Geopotential From the Orbit of Vanguard 3*, GSFC Technical Report X-592-73-130, Greenbelt, MD, May 1973.
55. C. A. Wagner, "Zonal Gravity Harmonics From Long Satellite Arcs by a Semi-numeric Method," *J. Geophys. Res.*, Vol. 75, No. 32, 10 June 1973, pp. 3271-3280.

37. J. Klokocnik, "Some Characteristics of the Earth Gravitational Field From Orbital Resonances of the Interkosmos Satellites," Proceedings of the International Symposium on the Use of Artificial Satellites for Geodesy and Geodynamics, Athens, Greece, 1979.
38. J. Klokocnik and J. Kostecky, "On the Possibility of Determining Variations in the Geocenter Position Using Geostationary Satellite Observations," Bull. Astron. Inst. Czechosl. 31, (1980), pp. 123-126.
39. J. Klokocnik, "A Possible Contribution to the Earth Model 'Grim-3,'" Zpravni a Pozorovani Geodeticke Observatore Pecny, No. 6, 1981.
40. J. Klokocnik, "A Review of Methods of Comparison of 'Resonant' Lumped Geopotential Coefficients," Bull. Astron. Inst. Czechosl. 33, (1982), pp. 89-104.
41. J. Klokocnik, "A Readjustment of Lumped Coefficients From Inclination of 1974-70A," Bull. Astron. Inst. Czechosl. 33, (1982), pp. 259-262.
42. C. Oesterwinter, "Importance of Gravitational Harmonics of High Order for the Orbits of Earth Satellites," Paper proposed to COSPAR for the May 1965 Symposium.
43. C. H. Reigber and G. Balmino, "Even and Odd Degree 13th-Order Harmonics From Analysis of Stable Near Resonant Satellite Orbits," Groupe de Recherches de Geodesie Spatiale Bulletin Nr. 15, March 1976.
44. C. H. Reigber and G. Balmino, "14th-Order Harmonics From Analysis of Mean Longitude Variations of Resonant Satellites," presented at XIXth COSPAR Meeting, Philadelphia, PA, June 1976.
45. C. Reigber and R. Rummel, "12th-Order Harmonics From Resonant Perturbations of Satellites Using Non-Homogeneous Weight Functions," Proceedings of the International Symposium on the Use of Artificial Satellites for Geodesy and Geodynamics, Athens, Greece, 1979.

27. D. G. King-Hele, C. J. Brookes, and G. E. Cook, "The Pear-Shaped Section of the Earth," *Nature*, Vol. 286, No. 5771, pp. 377-378, 24 July 1980.
28. D. G. King-Hele, *Lumped Harmonics of 15th and 30th-Order in the Geopotential From the Resonant Orbit of 1971-54A*, Royal Aircraft Establishment Technical Report 80088, (Farnborough, Hants., England, July 1980).
29. D. G. King-Hele, C. J. Brookes, and G. E. Cook, "Odd Zonal Harmonics in the Geopotential From Analysis of 28 Satellite Orbits," *Geophys. J. R. Astr. Soc* 1981, 64, pp. 3-30.
30. D. G. King-Hele, "Lumped Harmonics of 15th and 30th-Order in the Geopotential From the Resonant Orbit of the Satellite 1971-54A," *Proc. R. Soc. London, A* 375, pp. 327-350 (1981).
31. D. G. King-Hele and D. M. C. Walker, "Evaluation of 15th-Order Harmonics in the Geopotential From Analysis of Resonant Orbits," *Proc. R. Soc. London, A* 379, pp. 247-288, (1982).
32. D. G. King-Hele and D. M. C. Walker, "Geopotential Harmonics of Order 29, 30 and 31 From Analysis of Resonant Orbits," *Planet. Space Sci.* 1982, Vol. 30, pp. 411-425.
33. J. Klokocnik, "Changes in the Inclination of a Close Earth Satellite Due to Orbital Resonances," *Bull. Astron. Inst. Czechosl.* 27, (1976), No. 5.
34. J. Klokocnik and J. Kostelecky, "Evaluation of the 14th-Order Harmonics From Resonant Inclination Variations," *Bull. Astron. Inst. Czechosl.* 29, (1978), pp. 10-14.
35. J. Kostelecky and J. Klokocnik, "30th-Order Harmonics From Resonant Inclination Variations of Ten Satellites," *Bull. Astron. Inst. Czechosl.* 30. (1979), pp. 45-51.
36. J. Klokocnik, "15th-Order Resonance of Interkosmos 11 - Analysis of the Inclination," *Bull. Astron. Inst. Czechosl.* 30. (1979), pp. 214-219.

18. D. G. King-Hele, D. M. C. Walker, and R. H. Gooding, "Evaluation of Harmonics in the Geopotential of Order 15 and Odd Degree," *Planet. Space Sci.* 1974, Vol. 22, No. 10, p. 1349.
19. D. G. King-Hele, D. M. C. Walker, and R. H. Gooding, "Geopotential Harmonics of Order 15 and Even Degree From Changes in Orbital Eccentricity at Resonance," *Planet. Space Sci.* 1975, Vol. 23, pp. 229-246.
20. D. G. King-Hele, D. M. C. Walker, and R. H. Gooding, "Geopotential Harmonics of Order 15 and Odd Degree From Analysis of Resonant Orbits," *Planet. Space Sci.* 1975, Vol. 23, pp. 1239-1256.
21. H. Hiller and D. G. King-Hele, "The Orbit of Proton 4 Redetermined, With Geophysical Implications," *Planet. Space Sci.* 1977, vol. 25, pp. 511-520.
22. D. G. King-Hele, D. M. C. Walker, and R. H. Gooding, *Evaluation of 14th-Order Harmonics in the Geopotential*, Royal Aircraft Establishment Technical Report 78015, (Farnborough, Hants., England, February 1978).
23. D. G. King-Hele, D. M. C. Walker, and R. H. Gooding, "Evaluation of 14th-Order Harmonics in the Geopotential," *Planet. Space Sci.* 1979, Vol. 27, pp. 1-18.
24. D. G. King-Hele, *Skylab 1 Rocket, 1973-27B: Orbit Determination and Analysis*, Royal Aircraft Establishment Technical Report 79044, (Farnborough, Hants., England, April 1979).
25. D. G. King-Hele, D. M. C. Walker, and R. H. Gooding, "Evaluation of 14th-Order Harmonics in the Geopotential," Proceedings of the International Symposium on the Use of Artificial Satellites for Geodesy and Geodynamics, Athens, Greece, 1979.
26. D. G. King-Hele, *Odd Zonal Harmonics in the Geopotential From Analysis of 25 Satellite Orbits*, Royal Aircraft Establishment Technical Report 80023, (Farnborough, Hants., England, February 1980).

8. P. Dunn, "Geopotential Resonances in a Landsat Orbit," *Bulletin Geodesique* 55, 1981, pp. 143-158.
9. R. H. Gooding, *Lumped Geopotential Coefficients $C_{15,15}$ and $S_{15,15}$ Obtained From Resonant Variation in the Orbit of Ariel 3*, Royal Aircraft Establishment Technical Report 71068, (Farnborough, Hants., England, April 1971).
10. R. H. Gooding, "Lumped Fifteenth-Order Harmonics in the Geopotential," *Nature Physical Science*, Vol. 231, No. 25, pp. 168-169, 21 June 1971.
11. R. H. Gooding, "A Recurrence Relation for Inclination Functions," *Celestial Mechanics* 4, 1971, pp. 91-98.
12. H. Hiller, *The Effects of 15th-Order Resonance on the Orbit of Nimbus 1 Rocket, 1964-52B*, Royal Aircraft Establishment Technical Report 76001, (Farnborough, Hants., England, January 1976).
13. H. Hiller, "Lumped Geopotential Harmonics of Order 29 From Analysis of the Orbit of Cosmos 837 Rocket," *Planet. Space Sci.* 1982, Vol. 30, pp. 73-80.
14. D. G. King-Hele, "15th-Order Harmonics in the Geopotential From Analysis of Decaying Satellite Orbits," Proceedings of COSPAR "Space Research XIII," Madrid, Spain, 10-24 May 1972, Akademie Verlag, Berlin, 1973.
15. H. Hiller and D. G. King-Hele, "Equations for 15th-Order Geopotential Coefficients From the Orbit of Transit 1B," *Planet. Space Sci.* 1972, Vol. 20, pp. 1213-1228.
16. D. G. King-Hele, "Analysis of the Orbit of Cosmos 387 (1970-111A) Near 15th-Order Resonance," *Planet. Space Sci.* 1974, pp. 509-524.
17. D. G. King-Hele, "Analysis of the Orbit of Cosmos 72 (1965-53B) Near 15th-Order Resonance," *Planet. Space Sci.* 1974, Vol. 22, pp. 1269-1277.

REFERENCES

1. R. R. Allan, "Satellite Resonance with Longitude-Dependent Gravity - II, Effects Involving the Eccentricity," *Planet. Space Sci.* 1967, Vol. 15, pp. 1829-1845.
2. R. R. Allan, *Resonant Effect on Inclination for Close Satellites*, Royal Aircraft Establishment Technical Report 71245, (Farnborough, Hants., England, December 1971).
3. R. R. Allan, "Satellite Resonance with Longitude-Dependent Gravity - III, Inclination Changes for Close Satellites," *Planet. Space Sci.* 1973, Vol. 21, pp. 205-225.
4. R. J. Anderle, "Observations of Resonant Effects on Satellite Orbits Arising From the Thirteenth and Fourteenth Order Tesseral Gravitational Coefficients," *J. Geophys. Res.*, Vol. 70, No. 10, 15 May 1965.
5. R. J. Anderle and S. J. Smith, "Observation of Twenty-Seventh and Twenty-Eighth Order Gravity Coefficients Based on Doppler Observations," *Journal of Astronautical Sciences*, Vol. XV, No. 1, pp. 1-4, January/February 1968.
6. L. Blitzer, "Satellite Resonances and Librations Associated with Tesseral Harmonics of the Geopotential," *J. Geophys. Res.*, Vol. 71, No. 14, 15 July 1966.
7. B. R. Bowman, "Analysis of Mean Elements of Three U.S. Navy Navigation Satellites for the Period 1974-1976," *Celestial Mechanics* 19, 1979, pp. 203-211.

where $N_{\ell m}$ is the normalization factor specified by Equation 19. The relationship among the lumped coefficients, $\bar{C}_m^{q,k}$ and $\bar{S}_m^{q,k}$, the potential expansion coefficients and the influence numbers is

$$\bar{C}_m^{q,k} = \sum_{r=0}^{rMAX} Q_{\ell}^{q,k} \bar{C}_{\ell m} \quad (35)$$

$$\bar{S}_m^{q,k} = \sum_{r=0}^{rMAX} Q_{\ell}^{q,k} \bar{S}_{\ell m} \quad (36)$$

Note once more that the influence numbers as well as the potential expansion coefficients are dimensionless.

The following derivation should be viewed in context with the introduction to this report. Lagrange's planetary equation for the inclination variation was adopted from Pages 287 and 288 of Reference 33 and reads as follows:

$$\frac{di}{dt} = \frac{\dot{M}}{\sqrt{1-e^2}} * \sum_{\ell m p q} \left\{ J_{\ell m} \left(\frac{R}{a} \right)^\ell * \operatorname{Re} \left[j [(\ell - 2p) \cotg i - m \operatorname{cosec} i] \right. \right. \\ \left. \left. * F_{\ell m p} (i) C_{\ell p q} (e) \exp (j \psi_{\ell m p q}) \right] \right\} \quad (B01)$$

$$\psi_{\ell m p q} = (\ell - 2p)\omega + (\ell - 2p + q)M + m(\Omega - S) - m\lambda_{\ell m} \quad (B02)$$

$$J_{\ell m} = \sqrt{C_{\ell m}^2 + S_{\ell m}^2} \quad (B03)$$

$$m\lambda_{\ell m} = \tan^{-1} (S_{\ell m} / C_{\ell m}) \quad (B04)$$

where a , e , i , ω , Ω are the familiar orbital elements, \dot{M} is the mean motion, R is the equatorial "radius of the earth," M is the mean anomaly, S is the sidereal Greenwich time of the epoch of the elements, and $C_{\ell m}$ and $S_{\ell m}$ are the unnormalized expansion coefficients of the gravitational potential.

Also from Page 288 of Reference 33, for β/α resonance there hold the relationships

$$\ell = \beta \ell_0 + p, q = \alpha \ell_0 - q_0 - m\lambda_{\ell m} \quad (B05)$$

$$\ell = \ell_0 + N + \ell_0 (\ell - S) \quad (B06)$$

$$k = \ell - 2p + q_0 + q \quad (B07)$$

$$m = \ell_0 \quad (B08)$$

as well as Equations 3 through 11 from the main body of this report. For the meaning of beta, alpha, gamma, and q see the segment "Input Data" from the "Algorithm for the Influence Numbers." Also note that

$$0 \leq p \leq \ell \quad (B9)$$

$$\ell \geq m \quad (B10)$$

Again from Page 288 of Reference 53, the Inclination Function, $F_{\ell mp}(i)$, is

$$F_{\ell mp}(i) \equiv j^{(\ell-m)} \text{FA}(\ell, m, p, i) \quad (B11)$$

where the function FA is specified by Equations 14 through 16 above.

$$\bar{F}_{\ell mp}(i) \equiv j^{(\ell-m)} \text{FAN}(\ell, m, p, i) \quad (B12)$$

is the Normalized Inclination Function. The function FAN is the normalized version of FA and is defined by Equations 18 through 21.

$G_{\ell pq}(e)$ is the Eccentricity Function. It will be found defined on Page 37 of Reference 79 and is for our purposes specified by Equations 22 through 32 above. It shall already now be stated that the normalized expansion coefficients of the geopotential, $\bar{C}_{\ell m}$ and $\bar{S}_{\ell m}$, and the influence numbers, $C_{\ell m}^{(k)}$ and $S_{\ell m}^{(k)}$, are obtained by Equations 33 through 36 in the main part of this report.

Now consider the portion

$$\begin{aligned} C_{\ell m}^{(k)} * \text{Re}[\dots] &= \left(\frac{2\ell+1}{4\pi} \right)^{1/2} j^{(\ell-m)} * \text{FA} * G_{\ell pq} \\ &* C_{\ell m} * \text{Re} \left\{ 1 + j^{(q-m)} \frac{1}{\lambda^2} \left(\frac{1}{\lambda^2} \right) \right\} \end{aligned} \quad (B13)$$

in the equation that

For $(\ell - m)$ equal to an even number:

$$\begin{aligned}
 J_{\ell m} * \text{Re}[\dots] &= J_{\ell m}^{(\ell-m)} \text{Re}(j \exp(j\psi_{\ell mpq})) \\
 &= -J_{\ell m}^{(\ell-m)} \sin \psi_{\ell mpq} = -J_{\ell m}^{(\ell-m)} \sin (\gamma \psi_{\ell} - q\omega) \cos m\lambda_{\ell m} \\
 &\quad + J_{\ell m}^{(\ell-m)} \cos (\gamma \psi_{\ell} - q\omega) \sin m\lambda_{\ell m}
 \end{aligned}
 \tag{B14}$$

$$m\lambda_{\ell m} = \tan^{-1}(S_{\ell m}/C_{\ell m})
 \tag{B14}$$

$$\begin{pmatrix} \bar{C}_{\ell m} \\ \bar{S}_{\ell m} \end{pmatrix} = \frac{1}{N_{\ell m}} \begin{pmatrix} C_{\ell m} \\ S_{\ell m} \end{pmatrix}
 \tag{B15}$$

$$\cos m\lambda_{\ell m} = \frac{1}{\sqrt{1 + S_{\ell m}^2/C_{\ell m}^2}} = N_{\ell m} \bar{C}_{\ell m}/J_{\ell m}
 \tag{B16}$$

$$\sin m\lambda_{\ell m} = N_{\ell m} \bar{S}_{\ell m}/J_{\ell m}
 \tag{B17}$$

and, finally,

$$J_{\ell m} * \text{Re}[\dots] = -N_{\ell m} j^{(\ell-m)} [\bar{C}_{\ell m} \sin (\gamma \psi_{\ell} - q\omega) - \bar{S}_{\ell m} \cos (\gamma \psi_{\ell} - q\omega)]
 \tag{B18}$$

where

$$j^{(\ell-m)} = +1 \text{ or } -1
 \tag{B19}$$

or if $(\ell - m)$ equal to an odd number, one obtains in a similar fashion

$$J_{\ell m} * \text{Re}[\dots] = N_{\ell m} j^{(\ell-m+1)} [\bar{C}_{\ell m} \cos (\gamma \psi_{\ell} - q\omega) + \bar{S}_{\ell m} \sin (\gamma \psi_{\ell} - q\omega)]
 \tag{B20}$$

where

$$j^{(\ell-m+1)} = +1 \text{ or } -1
 \tag{B21}$$

If one now enters Equations B18 and B20 respectively into Equation B13 and subsequently Equation B13 into Lagrange's planetary equation (B6), observing that under the summations occurring in the latter equation m and q are constants while the summations over i and p are according to Equations 4 and 9a subject to the common summation index r , there result expressions for the inclination variation where the geopotential expansion coefficients appear in groups that may be conveniently "lumped together" into the lumped coefficients $\sum_m \bar{C}_m^{qk}$ and $\sum_m \bar{S}_m^{qk}$ in a linear fashion.

For even K

$$\frac{di}{dt} = \frac{\dot{M}}{\sqrt{1-e^2}} \left(\frac{R}{a}\right)^{k_0} A_{mk}^{(i)} C_{(0)0}^{(e)} FAN_{(0)0}^{(i)} \quad (B22)$$

$$* \left[\sum_m \bar{C}_m^{qk} \sin(\gamma\dot{\phi} - q\omega) - \sum_m \bar{S}_m^{qk} \cos(\gamma\dot{\phi} - q\omega) \right]$$

For odd K

$$\frac{di}{dt} = \frac{\dot{M}}{\sqrt{1-e^2}} \left(\frac{R}{a}\right)^{k_0} A_{mk}^{(i)} G_{(0)p}^{(e)} FAN_{(0)p}^{(i)} \quad (B23)$$

$$* \left[\sum_m \bar{C}_m^{qk} \cos(\gamma\dot{\phi} - q\omega) + \sum_m \bar{S}_m^{qk} \sin(\gamma\dot{\phi} - q\omega) \right]$$

and

$$\frac{d\omega}{dt} = \frac{\dot{M}}{\sqrt{1-e^2}} \left(\frac{R}{a}\right)^{k_0} A_{mk}^{(i)} G_{(0)p}^{(e)} FAN_{(0)p}^{(i)} \quad (B24)$$

$$\frac{d\Omega}{dt} = \frac{\dot{M}}{\sqrt{1-e^2}} \left(\frac{R}{a}\right)^{k_0} A_{mk}^{(i)} G_{(0)p}^{(e)} FAN_{(0)p}^{(i)} \quad (B25)$$

$$\frac{d\dot{\phi}}{dt} = \frac{\dot{M}}{\sqrt{1-e^2}} \left(\frac{R}{a}\right)^{k_0} A_{mk}^{(i)} G_{(0)p}^{(e)} FAN_{(0)p}^{(i)} \quad (B26)$$

$$Q_{\ell}^{pq} = j^{2r} \left(\frac{a}{b}\right)^{2r} \frac{G_{\ell pq}}{G_{\ell_0 p_0 q_0}} \frac{FAN_{\ell mp}}{FAN_{\ell_0 mp_0}} \quad (B.27)$$

$$j^{2r} = (-1)^r \quad (B.28)$$

where the Q_{ℓ}^{pq} are the influence numbers for both even and odd values of F .

It should finally be stated that Equation B11 represents the Inclination function as defined by Allan (Reference 1, p. 1830). Kaula's Inclination function (Reference 79) "FK" may be calculated as follows:

For even values of $(\ell - m)$, including $(\ell - m) = 0$,

$$FK = j^{(\ell-m)} FA \quad (B.29)$$

For odd values of $(\ell - m)$

$$FK = j^{(\ell-m)} FA / (-j) = j^{(\ell-m+1)} FA \quad (B.30)$$

DISTRIBUTION

	Copies		Copies
Headquarters Defense Mapping Agency Building 56 Washington, DC 20360 ATTN: Dr. C. F. Martin	1	National Aeronautics & Space Administration Washington, DC 20546 ATTN: Library	1
Defense Mapping Agency Aerospace Center 2nd & Arsenal Streets St. Louis, MO 63118	1	Aerospace Corporation 2350 East El Segundo Boulevard El Segundo, CA 90245 ATTN: Library	1
Defense Mapping Agency Hydrographic/Topographic Center 6500 Brooks Lane Washington, DC 20315	1	Applied Physics Laboratory The Johns Hopkins University 8621 Georgia Avenue Silver Spring, MD 20910 ATTN: Library	1
Chief of Naval Research Department of the Navy Arlington, VA 22217	1	Local:	
		E211 (Green)	1
		E231	10
		E31 (GIDEP)	1
		E05	1
		E10	1
		E11 (Green/White)	16
			1
			10
		K17 (Orange)	50
		K17 (Green/White)	1
		M1 (Green/White)	1
		M1	1
		N1	10
		N1 (Green/White)	1
National Oceanic & Atmospheric Administration National Ocean Survey Rockville, MD 20850 ATTN: Library	1		

END

FILMED

6-85

DTIC