



National
Defence

Défense
nationale

UNCLASSIFIED

DRES

SUFFIELD REPORT

NO. 412

AD-A156 381

THE IDENTIFICATION OF COMPOUNDS IN
MUSTARD HYDROLYSATE (U)

20030115229

by

P.A. D'Agostino and L.R. Provost

PCN No. 13E20

DTIC
ELECTE
JUL 11 1985

B

DTIC FILE COPY

April 1985

DISTRIBUTION STATEMENT A

Approved for public release
Distribution Unlimited



DEFENCE RESEARCH ESTABLISHMENT SUFFIELD, RALSTON, ALBERTA

WARNING

The use of this information is permitted subject to
recognition of proprietary and patent rights.

Canada

85 06 25 16

UNCLASSIFIED

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD
RALSTON ALBERTA

SUFFIELD REPORT NO. 412

THE IDENTIFICATION OF COMPOUNDS IN
MUSTARD HYDROLYSATE (U)

by

P.A. D'Agostino and L.R. Provost

PCN No. 13E20

WARNING
The use of this information is permitted subject to recognition
of proprietary and patent rights.

DTIC
ELECTE
JUL 11 1985
S D
B

UNCLASSIFIED

DISTRIBUTION STATEMENT A
Approved for public release
Distribution Unlimited

UNCLASSIFIED

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD
RALSTON ALBERTA

SUFFIELD REPORT NO. 412

**THE IDENTIFICATION OF COMPOUNDS IN
MUSTARD HYDROLYSATE (U)**

by

P.A. D'Agostino and L.R. Provost

ABSTRACT

About 700 tons of World War II mustard stored at the Defence Research Establishment Suffield was destroyed by hydrolysis during the 1970's. Samples of the liquid and sludge layers of the hydrolysate were retained and have now been analysed by gas chromatography using flame ionization and mass spectral detection. Hydrolysis was essentially complete since only trace levels of unreacted mustard were detected. Thiodiglycol, a hydrolysis product of mustard, was the major component identified in the hydrolysate. A number of other compounds were identified (or tentatively identified) in both the liquid and sludge hydrolysate samples.

UNCLASSIFIED

UNCLASSIFIED

ACKNOWLEDGEMENTS

The authors wish to thank the following DRES staff:

- a. Dr. P.A. Lockwood's group for providing a purified mustard sample,
- b. Mr. W.N. Lawson and the Decontamination Unit for collecting and storing the hydrolysate samples, and
- c. Mr. J.P. Bitz for making the glass columns used for packed column gas chromatographic analysis.

Thanks are also extended to Mr. B.G. Cameron for his advice during this study.

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By PER CALL JC	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

UNCLASSIFIED



UNCLASSIFIED

TABLE OF CONTENTS

	Page No.
ABSTRACT	
ACKNOWLEDGEMENTS	
TABLE OF CONTENTS	
INTRODUCTION	1
EXPERIMENTAL	3
Hydrolysate Samples	3
Instrumental Analysis	3
RESULTS AND DISCUSSION	3
Determination of Mustard and Thiodiglycol	3
Identification of other Hydrolysate Compounds	4
Aqueous Hydrolysate	5
Chloroform Extracts of the Hydrolysate	6
CONCLUSIONS	8
REFERENCES	9
TABLES	
FIGURES	
APPENDIX	

UNCLASSIFIED

UNCLASSIFIED

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD
RALSTON ALBERTA

SUFFIELD REPORT NO. 412

**THE IDENTIFICATION OF COMPOUNDS IN
MUSTARD HYDROLYSATE (U)**



by

P.A. D'Agostino and L.R. Provost

INTRODUCTION

1. During World War II over 700 tons of the chemical warfare agent mustard were shipped to the Defence Research Establishment Suffield (DRES) and stored in five lead-lined concrete vaults (1). In the early 1970's research was begun to find a safe, efficient, economical and environmentally acceptable method of destroying the DRES mustard. Laboratory studies indicated the feasibility of using batch hydrolysis provided the ratio of water to mustard was large, the temperature was elevated to 100°C and the pH was maintained above 7 (2). The mustard was hydrolysed in 1000 gallon batches using 500 pounds of lime ($\text{Ca}(\text{OH})_2$) and 2500 gallons of water (2, 3).

UNCLASSIFIED

$$\text{Cl}-\underset{\underset{1}{\text{CH}_2}}{\text{CH}_2}-\text{S}-\underset{\underset{2}{\text{CH}_2}}{\text{CH}_2}-\text{CH}_2-\text{Cl} + \text{H}_2\text{O} \rightarrow \text{HO}-\underset{\underset{2}{\text{CH}_2}}{\text{CH}_2}-\text{S}-\underset{\underset{1}{\text{CH}_2}}{\text{CH}_2}-\text{CH}_2-\text{Cl} + \text{HCl}$$


4. The objectives of this study were: a) to confirm the presence of thiodiglycol and mustard, b) to identify, where possible, other components in both the liquid and sludge hydrolysate samples, and c) to provide information that will aid in the identification of mustard decomposition products in environmental samples. Chloroform extracts of the liquid and sludge samples, were screened for the presence of mustard and other compounds by packed column gas chromatography with flame ionization detection (GC-FID). The liquid layer samples and water extracts of the sludge layer samples were analysed similarly for thiodiglycol and other hydrolysate products. Thiodiglycol, mustard and a number of other compounds were identified in the hydrolysate samples by combined gas chromatography-mass spectrometry (GC-MS).

EXPERIMENTAL

Hydrolysate Samples

6. Samples of the liquid and sludge hydrolysate layers were stored in polyethylene bottles. Figure 1 illustrates the analytical scheme followed during analysis of the samples. The method was described in detail in a prior publication (5).

Instrumental Analysis

7. A Varian 3700 (Varian Associates, Georgetown, Ont.) gas chromatograph was used for packed column GC-FID and GC-MS analyses. Packed column GC-MS analyses were performed using a VG-Micromass 70/70H double-focusing mass spectrometer (VG Analytical, Wythenshaw, UK) in the electron-impact mode. Operating condition for packed column GC-FID and GC-MS are presented in Tables I and II respectively.

RESULTS AND DISCUSSION

8. Both mustard and thiodiglycol were quantitated in the DRES mustard hydrolysate samples. A number of compounds were identified (or tentatively identified) based on the mass spectral and gas chromatographic data obtained during this study. In addition, a semi-quantitative estimate of concentration was made for the major components found in both the liquid and sludge hydrolysate.

Determination of Mustard and Thiodiglycol

9. Trace levels of mustard were detected in the chloroform extracts of the sludge from vaults 6 and 8. Mustard was confirmed by packed column GC-MS in the selective ion-monitoring mode at 2.9 and 4.2 $\mu\text{g/g}$ of sludge in these two vaults respectively (5). No mustard was detected in the chloroform extracts of the liquid hydrolysate.

10. Thiodiglycol was quantitated by packed column GC-FID using external standard calibration and confirmed by GC-MS. It was found in the 6.2 to 13.9 mg/g range in the water extracts of the sludge and in the 2.2 to 10.3 mg/mL range in the neat liquid hydrolysate (5). Table III summarizes this data.

Identification of other Hydrolysate Compounds

11. Packed column GC-MS was used to obtain electron-impact mass spectra for the major components isolated in the aqueous and chloroform extracts of the hydrolysate samples. A mass spectral and GC retention time match with standards was used for identification purposes. A number of compounds were identified in the hydrolysate using this method. Tentative identification of some of the other compounds was possible on the basis of mass spectral match alone.

12. Fundamental interpretation of the mass spectral data was required where a match was not found in the DRES or other mass spectral data bases (6, 7). Possible molecular formulas and in some cases plausible structures were advanced based on the acquired data. The molecular ion isotope ratio and characteristic fragmentation ions were most useful in the interpretation of the mass spectra of the compounds isolated from the hydrolysate.

13. It was possible to estimate the number of sulfur atoms in an unknown using molecular ion information since the ^{34}S isotope occurs naturally at 4.39%. The molecular ion ratio, M^+ to $(M+2)^+$, of known components in the hydrolysate was compared to the theoretical value calculated based on the natural occurrence of ^{32}S and ^{34}S . The ratio data for a large number of the hydrolysate components is presented in Tables IV and V. The experimental M^+ to $(M+2)^+$ ratio was always higher than theoretically expected. This is probably due to background noise contributions which are more significant for the less intense $(M+2)^+$ ion than for the M^+ ion.

14. Experimental M^+ to $(M+2)^+$ ratios of approximately 100:7, 100:12 and 100:19 were determined for compounds known to contain one, two and three sulfur atoms respectively. This compares to theoretical values of about 100:5, 100:9 and 100:13. Enhancement of the $(M+2)^+$ ion due to carbon and oxygen presence were taken into account for all calculations. Computer generated formulas were limited to compounds containing carbon, hydrogen, oxygen and sulfur. Possible molecular formulas for hydrolysate unknowns with measureable M^+ to $(M+2)^+$ ratios are listed in Tables IV and V.

15. Compounds containing chlorine were easily identified based on the characteristic molecular ion cluster associated with the occurrence of ^{35}Cl and ^{37}Cl . Possible molecular formulas, based on the presence of two chlorine atoms, are presented in Table III for an unknown compound found in the chloroform extract of the sludge hydrolysate.

16. The presence of sulfur in these compounds was substantiated by several characteristic low mass fragmentation ions. Fragmentation ions at m/z 59, m/z 60 and m/z 61 were found to be characteristic of this group of sulfur containing compounds. These ions were observed repeatedly throughout the study and probably correspond to $(S - CH = CH_2)^+$, $(S - CH_2CH_2)^+$ and $(H - S - CH_2CH_2)^+$ respectively. One or more of these ions was evident in all the mass spectra illustrated in Figures 3, 4 and 5 and tabulated in Appendix 1. The m/z 45, $(CH - S)^+$, and m/z 46, $(CH_2 - S)^+$ fragmentation ions are also characteristic of these compounds. Figure 3 illustrates the presence of these ions in five compounds isolated from the liquid hydrolysate and the water extracts of the sludge samples. The m/z 64 fragmentation ion, due primarily to $(S - S)^+$, suggests that the unknown contains a minimum of two sulfur atoms. Significant m/z 64 ions are evident in Figures 3, 4 and 5 for compounds containing two or more sulfur atoms.

17. Appendix 1 contains the mass spectral data for all the compounds not represented in Figures 3, 4 and 5. This data was included since identification of unknowns may be possible in the future with the availability of new standards or further research.

Aqueous Hydrolysate

18. Five compounds, including thiodiglycol, were identified in the neat liquid hydrolysate and the water extracts of the sludge samples. Figure 2 illustrates two chromatograms obtained during packed column GC-FID analysis. 1,4-Thioxane, 1,4-dithiane, hemisulfur mustard and thiodiglycol were identified in one or more of the samples studied. Identifications were based on a mass spectral and GC retention time match with a standard. Figure 3 illustrates the electron-impact mass spectra obtained for these compounds. Published mass spectra (6,7) were similar to those obtained for 1,4-thioxane, 1,4-dithiane and thiodiglycol.

19. The mass spectrum illustrated in Figure 3(b) was not similar to any alkyl-thiol ($C_5H_{12}S$), oxathiolane or oxathiane published (6,7). Fundamental interpretation of the mass spectral data suggested that it could be (2-vinylthio)ethanol. This compound could be formed by the dehydration of thiodiglycol. Dehydration may be possible under acidic conditions at elevated temperatures. pH values of less than 7 were noted for the liquid hydrolysate samples provided.

20. Table III summarizes the amounts of each major component in the neat liquid hydrolysate and the water extracts of the sludge samples. Thiodiglycol was quantitated by packed column GC-FID (in triplicate) using the external standard calibration method. A semi-quantitative estimate was made for the other compounds based on the FID peak height response of thiodiglycol. This was considered sufficient for these compounds since the emphasis was placed on qualitative identification.

Chloroform Extracts of the Hydrolysate

21. A large number of compounds were observed during the gas chromatographic study of the chloroform extracts of the liquid and sludge hydrolysate. Figures 6 and 7 illustrate the packed column GC chromatograms obtained during analysis of the sludge and liquid samples respectively. Many compounds were identified on the basis of their mass spectra and gas chromatographic data. Tentative identification of several compounds was possible by comparison of the mass spectrum with library (6,7) or literature spectra. As is often the case in broad spectrum analysis, a number of compounds remain unidentified. Possible molecular formulas, based on their mass spectral data, are presented for these unknowns in Tables IV and V.

22. Twenty-five major components were identified in the chloroform extracts of the sludge hydrolysate samples. 1,4-Thioxane, 1,4-dithiane and 1,2,5-trithiapane were positively identified using the DRES data base. The mass spectra of these compounds are illustrated in Figures 3(a), 3(c) and 4(e) respectively. The mass spectrum of 1,2,5-trithiapane obtained was similar to that in the literature (8). Two other ring structures, 2-methyl-1,3-oxathiolane (Figure 2(a)) and 1-oxa-4,5-dithiacycloheptane (Appendix 1 - 16) were tentatively identified using mass spectral data published in references (6) and (9) respectively.

23. The mass spectrum of the unknown illustrated in Figure 4(b) suggests that this compound may be similar in structure to 2-methyl-1,3-oxathiolane (Figure 4(a)). Both compounds show significant $(M-CH_3)^+$ fragmentation ions. The presence of an intense m/z 64, $(S_2)^+$, ion and a M^+ to $(M+2)^+$ ratio of 100:9 suggest a compound containing two sulfur atoms. This component probably only differs from 2-methyl-1,3-oxathiolane by the replacement of an oxygen with a sulfur atom. Thus the unknown in Figure 4(b) is probably a methyl substituted 1,3-dithiolane.

24. Both bis(2-chloroethyl)disulfide and bis(2-chloroethyl)trisulfide were identified using mass spectral and gas chromatographic data. The presence of these polysulfides is consistent with an earlier study (10) which indicated that Levenstein mustard (undistilled mustard) contained approximately 30% by weight unhydrolysable material (namely polysulfides and elemental sulfur). Figures 5(a) and 5(b) illustrate the mass spectral data acquired for these polysulfides. An additional compound containing two chlorine atoms remains unidentified. Possible molecular formulas for this compound (Figure 5(e)) are presented in Table IV.

25. The mass spectra of several compounds that remain unknown are presented in Figures 4(c), 4(d), 5(b) and 5(d). Possible molecular formulas and fragmentation ions based on their mass spectral data are presented in these figures. The compounds in Figures 4(c) and 4(d) appear to differ only in the replacement of an oxygen with a sulfur atom since their fragmentation pattern is similar.

26. Table IV lists the compounds and possible molecular formulas of unknowns identified in the sludge sample extracts. A semi-quantitative estimate of concentration was made based on the FID response of mustard. The major components account for 0.1 to 3.6% of the total weight of the sludge. It is interesting to note that the two sludge samples with the highest organic content, namely vaults 6 and 8, contained the only detectable traces of mustard.

27. Fourteen compounds were isolated during GC analysis of the chloroform extracts of the liquid samples. Tentative molecular formulas were advanced based on mass spectral evidence. The molecular formulas and an estimate of concentration, also based on the FID response of mustard, are presented in Table V.

28. Most of the compounds were of higher molecular weight and identification of a molecular ion was often difficult. Since these compounds were water soluble it is possible that some of the data accumulated may be due to the thermal degradation products of sulfonium salts. These salts have been isolated during the hydrolysis of mustard (4).

29. The organic content was less in the chloroform extracts of the liquid hydrolysate. Organic content, based on the major components identified in the liquid hydrolysis, was less than 0.1% for all the samples studied.

CONCLUSIONS

30. The hydrolysis of all the mustard destroyed at DRES was essentially complete since only trace levels of mustard were detected in two of the sample extracts. Thiodiglycol, the principle hydrolysis product of mustard, was found to be the major organic component in the aqueous sample.

31. A number of other components were identified in the hydrolysate samples based on their mass spectral and gas chromatographic data. Tentative identification was possible for several compounds based on comparison of their mass spectral data with published spectra. Mass spectral interpretation led to the assignment of possible molecular formulas for several unknown components.

32. As is often the case in broad spectrum analysis, some compounds remain as unknowns. The mass spectral data for these compounds are included to aid in possible future identification.

33. It could not be determined from these samples whether many of the compounds observed in the hydrolysate were present in the original mustard or were hydrolysis products. However, since hydrolysis represents a major chemical decomposition pathway, it would be possible to suggest the prior presence of mustard in environmental samples based on the identification of thiodiglycol and other hydrolysate components.

REFERENCES

1. Reichert, C. "Study of Mustard Destruction by Hydrolysis". Suffield Technical Note No. 329, July 1975. UNCLASSIFIED.
2. Reichert, C. "Hydrolysis of Mustard in Full-Scale Reaction Vessel". Suffield Technical Note No. 352, July 1975. UNCLASSIFIED.
3. Weaver, R.S., Reichert, C., Mellsen, S.B. "Destruction and Disposal of Canadian Stocks of World War II Mustard Gas". Defence Research Establishment Special Publication No. 67, December 1975. UNCLASSIFIED.
4. Stein, W.H., Moore, S., Bergmann, M. *J. Org. Chem.* **XI**, 664 - 674, 1946.
5. D'Agostino, P.A., Provost, L.R. "The Determination of Mustard and Thiodiglycol in Mustard Hydrolysate". Suffield Memorandum No. 1113, August 1984. UNCLASSIFIED.
6. Heller, S.R., Milne, G.W.A. "EPA/NIH Mass Spectral Data Base". Volumes 1 to 4 (1978) and Supplements 1 (1980) and 2 (1983), U.S. Government Printing Office, Washington, D.C.
7. Cornu, A., Massot, R. "Compilation of Mass Spectral Data". Volumes 1 and 2, (1975), Heyden Press, New York.
8. Field, L., Foster, C.H. *J. Org. Chem.* **35**, 749 - 752, 1970.
9. Goodrow, M.H., Muskev, W.K. *Synthesis* **6**, 457 - 459, 1981.
10. Fuson, R.C., Price, C.C., Burness, D.M., Foster, R.E., Hatchard, W.R., Lipscomb, R.D. *J. Org. Chem.* **XI**, 487 - 498, 1946.

UNCLASSIFIED

TABLE I
PACKED COLUMN GC-FID CONDITIONS

	AQUEOUS¹ SAMPLES	CHLOROFORM² SAMPLES
GC COLUMN:	1.22 m × 1.5 mm i.d. Tenax GC, 60/80 mesh (Alltech Assoc., Arlington, IL)	1.22 × 1.5 m i.d. 5% OV 101 on 80/100 mesh Chromosorb W (Chromatographic Specialties Ltd., Brockville, Ont)
TEMPERATURE PROGRAM:	150° for 1 min, then 10°/min to 250°C and held for 5 min	50° for 2 min, then 5°/min to 250°C and held for 10 min
INJECTION TEMPERATURE:	250°C	250°C
CARRIER GAS:	High purity helium ^{3,5} at 20 mL/min	High purity helium ^{3,5} at 25 mL/min
FID GASES:	Zero air ^{4,5} at 300 mL/min, ultra high purity hydrogen ^{4,5} at 30 mL/min	
FID TEMPERATURE:	250°C	

- 1 Liquid layer samples and aqueous extraction of sludge layer samples.
- 2 Chloroform extracts of both the liquid and sludge layer samples.
- 3 Helium is passed through Drierite, molecular sieve, dust and oxygen removal filters.
- 4 Air and hydrogen are passed through Drierite, molecular sieve and dust filters.
- 5 Gas supplier: Liquid Carbonic Canada Ltd. (Scarborough, Ontario).

UNCLASSIFIED

UNCLASSIFIED

TABLE II
PACKED COLUMN GC-MS CONDITIONS

OPERATING PARAMETERS	AQUEOUS ¹ SAMPLES	CHLOROFORM ² SAMPLES
GC COLUMN:	1.22 m × 1.5 mm i.d. Tenax GC, 60/80 mesh	1.83 × 1.5 mm i.d. OV 101 on 80/100 Chromosorb W
GC-MS INTERFACE:	Jet Separator (230°C)	
IONIZATION MODE:	Electron Impact	
ELECTRON ENERGY:	70 eV	
EMISSION:	200 μ A	
SOURCE TEMPERATURE:	190 - 200°C	
SOURCE PRESSURE:	ca. 2×10^{-6} torr	
SCAN FUNCTION AND RATE:	Scanning: 350 to 20 amu, exponential down scan, 3 sec/decade SIM: m/z 100 (PFK lock mass), 109, 111, 158 and 160 with 200 μ sec/ion dwell time (Mustard Determination)	
ACCELERATING VOLTAGE:	Scanning: 4 kV SIM: stepped from 4 kV downwards	
RESOLUTION (10% VALLEY DEFINITION):	Scanning: 500 SIM: 200 to 300	

¹ Liquid layer samples and aqueous extraction of sludge layer samples.

² Chloroform extracts of both the liquid and sludge layer samples.

UNCLASSIFIED

UNCLASSIFIED

Table III
COMPOUNDS IDENTIFIED IN LIQUID HYDROLYSATE AND WATER EXTRACTS OF SLUDGE HYDROLYSATE

CHROMATOGRAM PEAK NUMBER ^a	COMPOUND IDENTIFIED	MOL. WT.	VAULT CONCENTRATION								MASS SPECTRAL DATA	
			6	7	8	9	10					
			mg/mL ^b	mg/g ^c	mg/mL ^b	mg/g ^c	mg/mL ^b	mg/g ^c	mg/mL ^b	mg/g ^c		mg/mL ^b
1	1,4-Thioxane 	104	1.4	1.5	0.84	2.2	0.56	0.66	2.5	1.9	1.8	Figure 3a
2	(2-Vinylthio)ethanol ^d CH ₂ = CH — S HO — CH ₂ — CH ₂ —	104	—	1.7	0.74	2.6	—	0.56	—	—	—	Figure 3b
3	1,4-Dithiane 	120	1.3	1.4	0.59	3.1	0.56	0.31	1.9	2.5	1.3	Figure 3c
4	Hemisulfur mustard HO — CH ₂ CH ₂ — S Cl — CH ₂ CH ₂ —	140	—	—	—	—	—	—	1.9	3.1	—	Figure 3d
5	Thiodiglycol HO — CH ₂ CH ₂ — S HO — CH ₂ CH ₂ —	122	4.7 ± 1.1	11.8 ± 9	4.4 ± 2	7.7 ± 5	2.2 ± 1	6.2 ± 3	10.3 ± 5	13.9 ± 3	6.1 ± 4	Figure 3e
% Organic Content in Sample			0.7%	1.6%	0.7%	1.6%	0.3%	0.8%	1.7%	2.1%	0.9%	1.6%

^a Refer to Figure 2.

^b mg/mL of liquid hydrolysate.


^c mg/g of sludge hydrolysate.

^d Tentative identification based on mass spectral interpretation.

UNCLASSIFIED

UNCLASSIFIED

Table IV
MAJOR COMPONENTS IDENTIFIED IN THE CHLOROFORM EXTRACTS OF THE SLUDGE HYDROLYSATE

CHROMATOGRAM PEAK NUMBER ^a	COMPOUND IDENTIFIED (or tentatively identified)	MOL. WT.	VAULT CONCENTRATION (mg/g of Sludge) ^c					M:M+2 RATIO		NUMBER OF SULFUR ATOMS ^e	MASS SPECTRAL DATA ^f
			6	7	8	9	10	EXPERIMENTAL	THEORETICAL		
2-Methyl-1,3-oxathiolane											
1		104	2.8	—	3.4	—	—	100:6.9	100:4.7	1	Figure 4a
1,4-Thioxane											
1		104	0.94	0.07	0.87	0.34	0.02	100:7.1	100:4.7	1	Figure 3a
(2-Vinylthio)ethanol CH ₂ =CH-S-CH ₂ -CH ₂ -OH											
3		104	0.62	0.32	0.22	—	—	100:6.1	100:4.7	1	Figure 3b
Methyl-1,3-dithiolane											
4		120	2.4	0.27	2.6	0.40	0.03	100:10	100:8.9	2	Figure 4b
1,4-Dithiane											
5		120	1.0	—	1.5	0.30	—	100:10	100:8.9	2	Figure 3c
•C ₂ H ₄ S ₂ O C ₂ H ₄ SO ₂											
6		124	3.4	0.92	3.7	1.5	0.14	100:14	100:9.0 100:4.8	2	Figure 4c
C ₂ H ₄ SO C ₂ H ₄ SO ₂ C ₂ H ₄ S ₂											
7		148	0.28	0.04	0.31	—	—	m/z 150 not detected	—	—	Appendix 1-15

.../2

UNCLASSIFIED

UNCLASSIFIED

Table IV (Cont'd)

CHROMATOGRAM PEAK NUMBER ^a	COMPOUND IDENTIFIED (or tentatively identified)	MOL. WT.	VAULT CONCENTRATION (mg/g of Sludge) ^c					M:M+2 RATIO		NUMBER OF SULFUR ATOMS ^d	MASS SPECTRAL DATA
			6	7	8	9	10	EXPERIMENTAL	THEORETICAL		
8	1-Oxa-4,5-dithiacycloheptane 	136	0.06	0.02	0.12	—	—	100:14	100:9.1	2	Appendix 1-16
9	Unknown	140	0.24	0.09	0.27	0.27	—	m/z 142 not detected	—	—	Appendix 1-17
10	*C ₄ H ₁₀ S ₂ O C ₄ H ₁₀ SO ₂	138	1.4	0.49	0.93	0.94	0.11	100:13	100:9.1 100:5.1	2	Appendix 1-3
11	*C ₃ H ₈ S ₂ O C ₃ H ₈ S ₂ O ₂	156	0.82	0.23	1.1	0.34	—	100:19	100:13 100:9.4	3	Figure 4d
12	1,2,5-Trithiapane 	152	1.6	0.39	2.0	0.67	0.05	100:19	100:13	3	Figure 4e
13	bis(2-Chloroethyl) disulfide Cl-CH ₂ CH ₂ -S ₂ -CH ₂ CH ₂ -Cl	192	1.5	—	2.4	—	—	—	—	—	Figure 5a
14	*C ₇ H ₁₆ S ₂ *C ₄ H ₁₀ S ₂ O C ₇ H ₁₆ SO ₂ C ₄ H ₁₀ SO ₂	164	2.6	1.1	—	—	—	100:16	100:9.1 100:9.2 100:5.1 100:5.2	2	Appendix 1-6

.../3

UNCLASSIFIED

UNCLASSIFIED

Table IV (Cont'd)

CHROMATOGRAM PEAK NUMBER ^a	COMPOUND IDENTIFIED ^b (or tentatively identified)	MOL. WT.	VAULT CONCENTRATION (mg/g of Sludge) ^c					M:M+2 RATIO		NUMBER OF SULFUR ATOMS ^d	MASS SPECTRAL DATA
			6	7	8	9	10	EXPERIMENTAL	THEORETICAL		
15	*C ₆ H ₁₀ S ₂	168	—	—	1.9	—	—	100:23	100:13	3	Figure 5b
	*C ₆ H ₈ S ₂ O								100:13		
	C ₆ H ₁₀ S ₂ O ₂								100:9.3		
	C ₆ H ₈ S ₂ O ₂								100:9.5		
16	Unknown	172	0.63	0.07	0.87	0.13	—	m/z 174 not detected	—	—	Appendix 1-18
17	Unknown	152	—	0.25	—	—	—	m/z 154 not detected	—	—	Appendix 1-19
18	*C ₆ H ₁₀ S ₂	174	1.3	0.22	1.4	0.94	—	100:12	100:9.2	2	Appendix 1-20
	C ₆ H ₁₀ SO ₂								100:5.2		
	C ₆ H ₁₀ SO								100:5.1		
19	bis(C-Chloroethyl)trisulfide Cl-CH ₂ -CH ₂ -S ₃ -CH ₂ -CH ₂ -Cl	222	5.4	—	7.3	1.9	—	—	—	—	Figure 5c
20	*C ₆ H ₈ S ₄	184	2.4	0.28	2.6	0.88	0.09	100:33	100:18	4	Figure 5d
	C ₆ H ₈ S ₂ O ₂								100:14		
	C ₆ H ₁₀ S ₂								100:14		
	C ₆ H ₈ S ₂ O ₂								100:9.7		
21	Unknown	—	—	1.9	—	—	—	—	—	—	Appendix 1-21

.../4

UNCLASSIFIED

UNCLASSIFIED

Table IV (Cont'd)

CHROMATOGRAM PEAK NUMBER ^a	COMPOUND IDENTIFIED (or tentatively identified)	MOL. WT.	VAULT CONCENTRATION (mg/g of Sludge) ^c					M:M+2 RATIO		NUMBER OF SULFUR ATOMS ^d	MASS SPECTRAL DATA
			6	7	8	9	10	EXPERIMENTAL	THEORETICAL		
22	*C ₈ H ₁₀ S ₃								100:14		
	C ₈ H ₁₀ S ₂ O ₃	206	2.9	0.95	1.6	0.54	—	100:24	100:9.6	3	Appendix 1-22
	C ₈ H ₁₀ S ₂ O								100:9.5		
	C ₁₀ H ₁₂ S ₃								100:9.4		
23	*C ₈ H ₁₀ S ₃								100:14		
	C ₈ H ₁₀ S ₂ O ₃	208	—	0.28	—	0.67	0.07	100:24	100:9.6	3	Appendix 1-23
	C ₈ H ₁₀ S ₂ O								100:9.5		
24	C ₈ H ₁₀ S ₃								100:14		
	C ₈ H ₁₀ S ₂ O ₃	208	—	0.07	0.30	—	—	100:24	100:9.6	3	Appendix 1-24
	C ₈ H ₁₀ S ₂ O ₂								100:9.5		
25	C ₈ H ₁₀ SOCl ₂										
	C ₈ H ₁₀ SO ₂ Cl ₂	246	0.44	0.26	0.29	0.08	0.45	—	—	—	Figure 5e
	C ₈ H ₁₀ SO ₂ Cl ₂										
	C ₈ H ₁₀ S ₂ Cl ₂										
% Organic Content in Extract			3.3	0.9	3.6	1.0	0.1				

^a Refer to Figure 6.

^b Based on Mass Spectral Data.

^c Semi-quantitative estimate based on FID response.

^d Probable number of sulfur atoms in the compound based on the M:M+2 ratio observed.

* Most likely molecular formula(s) based on M:M+2 ratio observed.

UNCLASSIFIED

UNCLASSIFIED

Table V
MAJOR COMPOUNDS IDENTIFIED IN THE CHLOROFORM EXTRACTS OF THE LIQUID HYDROLYSATE

CHROMATOGRAM PEAK NUMBER ^a	MS SCAN NUMBER ^a	COMPOUND IDENTIFIED (or tentatively identified)	MOL. WT.	VAULT CONCENTRATION (µg/mL of Liquid) ^c					M:M+2 RATIO		NUMBER OF SULFUR ATOMS ^d	MASS SPECTRAL DATA
				6	7	8	9	10	EXPERIMENTAL	THEORETICAL		
1	52	1,4-Thioxane 	104	32	17	—	110	10	100:7.1	100:4.7	1	Figure 3a
2	75	(2-Vinylthio)ethanol $\text{CH}_2=\text{CH}-\text{S}-\text{CH}_2\text{CH}_2\text{OH}$	104	—	240	—	—	—	100:6.1	100:4.7	1	Figure 3b
Not in Figure	178	•C ₄ H ₈ S ₂ O C ₄ H ₈ SO ₂ C ₄ H ₁₀ SO ₂	136	—	—	2.0	4.4	—	100:13	100:9.1 100:5.1 100:4.9	2	Appendix 1-1
	215	Unknown	148	24	68	19	82	48	—	—	—	Appendix 1-2
4	232	•C ₄ H ₁₀ S ₂ O C ₄ H ₁₀ SO ₂	138	23	—	8.9	60	40	100:13	100:9.1 100:5.1	2	Appendix 1-3
5	262	Unknown	132	2.9	—	3.3	15	5.8	—	—	—	Appendix 1-4
6	300	•C ₄ H ₈ S ₂ O ₂ •C ₄ H ₁₀ S ₂ O ₂ C ₄ H ₈ S ₂ C ₄ H ₁₀ SO ₂ C ₄ H ₁₀ SO ₄	152	7.7	11	5.6	15	7.6	100:14	100:9.2 100:9.1 100:13 100:5.1 100:5.3	2	Appendix 1-5

.../2

UNCLASSIFIED

UNCLASSIFIED

Table V (Cont'd)

CHROMATOGRAM PEAK NUMBER	MS SCAN NUMBER ^a	COMPOUND IDENTIFIED (or tentatively identified)	MOLE WT. ^b	VAULT CONCENTRATION ($\mu\text{g/mL}$ of Liquid) ^c					M:M+2 RATIO		NUMBER OF SULFUR ATOMS ^d	MASS SPECTRAL DATA
				6	7	8	9	10	EXPERIMENTAL	THEORETICAL		
7	307	$\bullet\text{C}_7\text{H}_{10}\text{S}_2$	164	9.4	61	—	—	—	100:16	100:9.1	2	Appendix 1-6
		$\bullet\text{C}_8\text{H}_{10}\text{S}_2\text{O}$								100:9.2		
		$\text{C}_7\text{H}_{10}\text{SO}_2$								100:5.1		
		$\text{C}_8\text{H}_{10}\text{SO}_2$								100:5.2		
Not in Figure	313	$\bullet\text{C}_8\text{H}_{10}\text{S}_2$	154	—	—	5.8	16	12	100:22	100:13	3	Appendix 1-7
		$\bullet\text{C}_7\text{H}_8\text{S}_2\text{O}$								100:13		
		$\text{C}_8\text{H}_8\text{S}_2\text{O}_2$								100:9.4		
8	360	Unknown	164	15	11	7.8	27	16	m/z 166 not detected	—	—	Appendix 1-8
										—		
9	420	$\bullet\text{C}_7\text{H}_{10}\text{S}_2$	164	38	136	26	63	36	100:27	100:9.1	3	Appendix 1-9
		$\bullet\text{C}_8\text{H}_{10}\text{S}_2\text{O}$								100:9.2		
		$\text{C}_7\text{H}_{10}\text{SO}_2$								100:5.1		
		$\text{C}_8\text{H}_{10}\text{SO}_2$								100:5.2		
10	482	Unknown	152	19	11	27	11	13	m/z 154 detected at trace level	—	—	Appendix 1-10
										—		
11	531	$\text{C}_8\text{H}_{10}\text{S}_2\text{O}$	208	89	110	65	160	97	m/z 210 detected at trace level	—	—	Appendix 1-11
		$\text{C}_8\text{H}_{10}\text{S}_2\text{O}_2$								—		
		$\text{C}_8\text{H}_{10}\text{S}_2$								—		

.../3

UNCLASSIFIED

UNCLASSIFIED

Table V (Cont'd)

CHROMATOGRAM PEAK NUMBER ^a	MS SCAN NUMBER ^b	COMPOUND IDENTIFIED (or tentatively identified)	MOL. ^c WT.	VAULT CONCENTRATION ($\mu\text{g/mL}$ of Liquid) ^c						M:M+2 RATIO		NUMBER OF SULFUR ATOMS ^d	MASS SPECTRAL DATA
				6	7	8	9	10	EXPERIMENTAL	THEORETICAL			
12	598	Unknown	—	72	82	70	79	77	—	—	—	Appendix 1-12	
13	679	$\text{C}_8\text{H}_{10}\text{S}_2\text{O}$	208	47	63	32	86	46	m/z 210 detected at trace level	—	—	Appendix 1-13	
		$\text{C}_8\text{H}_{10}\text{S}_2\text{O}_2$											
		$\text{C}_8\text{H}_{10}\text{S}_3$											
14	766	$\text{C}_8\text{H}_{10}\text{S}_3\text{O}$	208	56	74	35	91	62	m/z 210 detected at trace level	—	—	Appendix 1-14	
		$\text{C}_8\text{H}_{10}\text{S}_3\text{O}_2$											
		$\text{C}_8\text{H}_{10}\text{S}_4$											
% Organic Content in the Extract				0.04	0.09	0.03	0.08	0.05					

^a Refer to Figure 7.^b Based on Mass Spectral Data.^c Semi-quantitative estimate based on FID response.^d Probable number of sulfur atoms in the compound based on the M:M+2 ratio observed.

• Most likely molecular formula(s) based on M:M+2 ratio observed.

UNCLASSIFIED




Figure 1: ANALYTICAL SCHEME USED FOR THE ANALYSIS OF DRES MUSTARD HYDROLYSATE




Figure 2

Packed Column GC-FID Chromatograms of: a) Water Extract of the Equivalent of 380 µg of Vault 7 Sludge Hydrolysate and b) 1.1 µL of Vault 9 Liquid Hydrolysate. Numbered Peaks are Identified in Table III.




Figure 3

Electron-Impact Mass Spectra of: a) 1,4-Thioxane, b) (2-Vinylthio)Ethanol (tentative identification), c) 1,4-Dithiane, d) Hemisulfur Mustard and e) Thiodiglycol Identified in the Aqueous Hydrolysate Samples.




Figure 4

Electron-Impact Mass Spectra of: a) 2-Methyl-1,3-Oxathiolane (tentative identification), b) a Methyl Substituted 1,3-Dithiolane (tentative identification), c) an Unknown (probably $C_3H_8OS_2$), d) an Unknown (probably $C_3H_8OS_3$) and e) 1,2,5-Trithiapane Identified in the Chloroform Extracts of the Mustard Hydrolysate.




Figure 5

Electron-Impact Mass Spectra of: a) bis(2-Chloroethyl)Disulfide, b) an Unknown (probably C₄H₈OS₃), c) bis(2-Chloroethyl)Trisulfide, d) an Unknown (probably C₄H₈S₄) and e) an Unknown Containing Two Chlorine Atoms Identified in the Chloroform Extracts of the Mustard Hydrolysate.




Figure 6

Packed Column GC-FID Chromatograms of the Chloroform Extracts of Two Sludge Hydrolysate Samples: a) the Equivalent of 410 μg of Vault 8 and b) the Equivalent of 390 μg of Vault 9. Numbered Peaks are Identified in Table IV.




Figure 7

Packed Column GC-MS Total Ion Current Chromatograms of the Chloroform Extracts of Two Liquid Hydrolysate Samples: a) the Equivalent of 60 μL of Vault 7 and b) the Equivalent of 100 μL of Vault 6. Numbered Peaks are Identified in Table V.
(One MS scan takes 3.5 sec).

UNCLASSIFIED

Appendix 1

ELECTRON-IMPACT MASS SPECTRAL DATA FOR COMPOUNDS IDENTIFIED IN THE
CHLOROFORM EXTRACTS OF THE MUSTARD HYDROLYSATE

APPENDIX NUMBER	CHROMATOGRAM		TENTATIVE IDENTIFICATION ^d	MOL. WT. ^e	PARTIAL MASS SPECTRUM: m/z (RELATIVE INTENSITY)																	
	LIQUID ^a	SLUDGE ^b																				
					(VAULT #) ^c	(VAULT #) ^c																
1-1	No Chrom. # (8,9)	—	*C ₆ H ₈ S ₂ O C ₆ H ₈ SO ₂ C ₆ H ₁₀ SO ₂	136	138 (13)	137 (10)	136 (100)	108 (10)	97 (10)	93 (13)	92 (28)	89 (23)	78 (35)	64 (75)	61 (33)	60 (100)	59 (53)					
1-2	3 (6,7,8,9,10)	—	Unknown	148	148 (1)	135 (3)	118 (4)	109 (48)	106 (23)	104 (28)	93 (13)	91 (20)	89 (50)	78 (53)	74 (38)	72 (68)	71 (100)	65 (73)	61 (88)	59 (83)		
1-3	4 (6,7,8,9,10)	10 (6,7,8,9,10)	*C ₆ H ₁₀ S ₂ O C ₆ H ₁₀ SO ₂	138	140 (1)	138 (8)	120 (8)	105 (10)	94 (5)	91 (8)	61 (50)	60 (100)	59 (75)									
1-4	5 (6,8,9,10)	—	Unknown	132	132 (1)	130 (1)	118 (1)	117 (1)	103 (1)	88 (25)	61 (10)	60 (100)	59 (30)									
1-5	6 (6,7,8,9,10)	—	*C ₆ H ₈ S ₂ O ₂ *C ₆ H ₁₀ S ₂ O C ₆ H ₈ S C ₆ H ₁₀ SO ₂ C ₆ H ₁₀ SO ₂	152	154 (2)	153 (3)	152 (14)	89 (5)	88 (5)	87 (100)	86 (8)	85 (8)	60 (40)	59 (58)								
1-6	7 (6,7)	14 (6,7)	*C ₆ H ₁₀ S ₂ *C ₆ H ₁₀ SO ₂ C ₆ H ₁₀ SO ₂ C ₆ H ₁₀ SO ₂	164	166 (2)	165 (2)	164 (7)	132 (2)	120 (10)	118 (18)	105 (68)	104 (20)	91 (15)	90 (20)	87 (38)	86 (15)	85 (28)	74 (18)	73 (20)	60 (80)	59 (75)	59 (100)

.../2

UNCLASSIFIED

UNCLASSIFIED

Appendix 1 (Cont'd)


APPENDIX NUMBER	CHROMATOGRAM		TENTATIVE IDENTIFICATION ^d	MOL. WT. ^d	PARTIAL MASS SPECTRUM:m/z (RELATIVE INTENSITY)															
	LIQUID ^a	SLUDGE ^b			(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c
1-7	No Chrom. # (8,9,10)	—	•C ₈ H ₁₀ S ₂ •C ₈ H ₈ S ₂ O C ₈ H ₈ S ₂ O ₂	154	156 (5)	155 (3)	154 (23)	120 (10)	92 (45)	87 (15)	79 (23)	78 (8)	76 (13)	74 (15)	64 (38)	61 (28)	60 (100)	59 (68)		
1-8	8 (6,7,8,9,10)	—	Unknown	164	164 (2)	149 (1)	136 (8)	123 (2)	120 (2)	115 (2)	105 (10)	104 (8)	103 (3)	102 (6)	89 (13)	74 (18)	61 (53)	60 (100)	59 (73)	
1-9	9 (6,7,8,9,10)	—	•C ₈ H ₁₀ S ₂ •C ₈ H ₈ S ₂ O C ₈ H ₈ SO ₂ C ₈ H ₁₀ SO ₂	164	166 (1)	165 (1)	164 (3)	138 (5)	120 (4)	105 (20)	104 (12)	91 (13)	76 (15)	74 (20)	73 (13)	61 (63)	60 (100)	59 (75)		
1-10	10 (6,7,8,9,10)	—	Unknown	152	152 (1)	136 (3)	124 (2)	120 (2)	105 (45)	76 (25)	74 (20)	64 (38)	61 (33)	60 (100)	59 (80)					
1-11	11 (6,7,8,9,10)	—	C ₈ H ₁₀ S ₂ O C ₈ H ₈ S ₂ O ₂ C ₈ H ₁₀ S ₂	208	208 (2)	182 (2)	164 (4)	136 (4)	120 (10)	113 (4)	105 (25)	104 (18)	87 (15)	86 (10)	76 (15)	74 (18)	61 (15)	60 (60)	59 (100)	75)
1-12	12 (6,7,8,9,10)	—	Unknown	—	181 (2)	164 (2)	152 (2)	138 (2)	136 (4)	134 (1)	120 (15)	118 (4)	105 (23)	92 (10)	76 (20)	74 (15)	64 (23)	61 (43)	60 (100)	59 (75)
1-13	13 (6,7,8,9,10)	—	C ₈ H ₁₀ S ₂ O C ₈ H ₈ S ₂ O ₂ C ₈ H ₁₀ S ₂	208	208 (2)	182 (2)	164 (4)	136 (4)	120 (10)	118 (4)	105 (25)	104 (18)	87 (15)	86 (10)	76 (15)	74 (18)	61 (15)	60 (60)	59 (100)	75)

.../3

UNCLASSIFIED

UNCLASSIFIED

Appendix 1 (Cont'd)

APPENDIX NUMBER	CHROMATOGRAM		TENTATIVE IDENTIFICATION ^d	MOL. WT. ^d	PARTIAL MASS SPECTRUM: m/z (RELATIVE INTENSITY)																	
	LIQUID ^a	PEAK NUMBER																				
		SLUDGE ^b																				
(VAULT #) ^c		(VAULT #) ^c																				
1 - 14	14	—	C ₈ H ₁₀ S ₂ O	208	208	181	164	149	136	120	118	105	104	89	87	86	76	74	64	61	60	59
	(6,7,8,9,10)	(2)	(3)		(3)	(2)	(4)	(5)	(3)	(25)	(15)	(18)	(15)	(25)	(15)	(73)	(100)	(75)				
1 - 15	—	7 (6,7,8)	C ₈ H ₁₀ S ₂ O ₂ C ₈ H ₁₀ S ₃		148	149	148	136	105	104	103	102	89	85	76	75	74	61	60	59	(30)	
	—		C ₇ H ₁₀ SO	(3)	(8)	(13)	(20)	(28)	(68)	(45)	(10)	(8)	(23)	(20)	(100)	(25)	(18)	(30)				
	—		C ₈ H ₁₀ SO ₂ C ₈ H ₁₀ S ₃	136	138	137	136	118	103	102	92	89	87	86	85	78	73	72	61	59	(50)	
1 - 16	—	8 (6,7,8)	1-Oxa-4,5-dithia- cycloheptane 	140	140	133	117	111	109	108	107	106	97	93	78	75	72	71	65	63	(25)	
	—		Unknown		(33)	(13)	(8)	(15)	(55)	(15)	(20)	(28)	(5)	(8)	(58)	(83)	(68)	(100)	(75)			
1 - 17	—	9 (6,7,8,9)	Unknown		172	172	127	112	104	102	96	87	86	85	73	64	61	59				
	—			Unknown	(18)	(8)	(8)	(6)	(6)	(13)	(100)	(40)	(20)	(15)	(13)	(18)	(40)					
1 - 18	—	16 (6,7,8,9)		Unknown	152	152	121	120	118	104	90	61	60	59								
	—		Unknown		(3)	(4)	(4)	(10)	(5)	(8)	(15)	(100)	(83)									
1 - 19	—	17 (7)	*C ₈ H ₁₀ S ₃ C ₈ H ₁₀ SO ₂ C ₈ H ₁₀ SO		174	176	175	174	159	146	136	124	123	104	102	89	76	74	61	60	59	(73)
	—			Unknown	(1)	(1)	(8)	(2)	(7)	(6)	(1)	(1)	(7)	(3)	(5)	(10)	(15)	(28)	(100)			

.../4

UNCLASSIFIED

UNCLASSIFIED

Appendix 1 (Cont'd)

APPENDIX NUMBER	CHROMATOGRAM PEAK NUMBER		TENTATIVE IDENTIFICATION ^d	MOL. WT. ^d	PARTIAL MASS SPECTRUM: m/z (RELATIVE INTENSITY)																		
	LIQUID ^a	SLUDGE ^b			(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c	(VAULT #) ^c
1-21	—	21 (7)	Unknown	—	136 (2)	134 (3)	118 (6)	105 (13)	104 (12)	103 (3)	102 (4)	90 (5)	89 (4)	88 (3)	87 (14)	86 (16)	85 (4)	76 (15)	61 (23)	60 (100)	59 (85)		
1-22	—	22 (6,7,8,9)	*C ₈ H ₁₀ S ₂ C ₈ H ₈ S ₂ O ₂ C ₈ H ₁₀ S ₂ O C ₁₀ H ₁₂ S ₂	206	208 (10)	207 (5)	206 (43)	180 (8)	178 (25)	146 (13)	136 (13)	134 (100)	102 (28)	88 (10)	78 (70)	76 (13)	69 (20)	64 (25)	61 (70)	60 (53)	59		
1-23	—	23 (7,9,10)	*C ₈ H ₁₀ S ₂ C ₈ H ₁₀ S ₂ O ₂ C ₈ H ₁₀ S ₂ O	208	208 (1)	206 (2)	181 (1)	178 (1)	136 (4)	120 (1)	105 (8)	104 (8)	102 (4)	89 (4)	76 (15)	74 (12)	61 (25)	60 (100)	59 (75)				
1-24	—	24 (7,8)	C ₈ H ₁₀ S ₂ C ₈ H ₁₀ S ₂ O ₂	208	210 (13)	209 (10)	208 (55)	180 (13)	150 (10)	148 (33)	120 (8)	111 (7)	104 (43)	88 (35)	78 (10)	76 (100)							

^a Refer to Figure 7.

^b Refer to Figure 6.

^c Vault numbers where compound was identified.

^d Based on Mass Spectral Data.

* Most likely molecular formula(s) based on M:M + 2 ratio observed.

UNCLASSIFIED

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall document is classified)

1. ORIGINATING ACTIVITY Defence Research Establishment Suffield		2a. DOCUMENT SECURITY CLASSIFICATION UNCLASSIFIED	
		2b. GROUP	
3. DOCUMENT TITLE The Identification of Compounds in Mustard Hydrolysate (U)			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Suffield Report			
5. AUTHOR(S) (Last name, first name, middle initial) D'Agostino, P.A. and Provost, L.R.			
6. DOCUMENT DATE APRIL May 1985		7a. TOTAL NO. OF PAGES	7b. NO. OF REFS 10
8a. PROJECT OR GRANT NO. 13E20		9a. ORIGINATOR'S DOCUMENT NUMBER(S) Suffield Report No. 412	
8b. CONTRACT NO.		9b. OTHER DOCUMENT NO.(S) (Any other numbers that may be assigned this document)	
10. DISTRIBUTION STATEMENT UNLIMITED			
11. SUPPLEMENTARY NOTES		12. SPONSORING ACTIVITY Defence Research Establishment Suffield	
13. ABSTRACT (U) The mustard stored at the Defence Research Establishment Suffield was disposed of by hydrolysis during the 1970's. Samples of the liquid and sludge hydrolysate were analysed by gas chromatography with flame ionization and mass spectral detection. Hydrolysis was essentially complete since only trace levels of mustard were detected. Thiodiglycol, a hydrolysis product of mustard, was found as a major component in the hydrolysate. A number of other compounds were identified (or tentatively identified) in both the liquid and sludge hydrolysate samples. <i>Originator</i> <i>Suffield keywords include:</i>			

Cont

B-5

UNCLASSIFIED

This Sheet Security Classification

KEY WORDS

Gas Chromatography,
 Mass Spectroscopy,
 Solvent Extraction,
 Chemical Agent Detection,
 Military Chemical Agent,
 Mustard Agents,
 Thiodiglycol,
 Mustard Hydrolysate.

INSTRUCTIONS

1. ORIGINATING ACTIVITY. Enter the name and address of the organization issuing the document.
- 2a. DOCUMENT SECURITY CLASSIFICATION. Enter the overall security classification of the document including special warning terms whenever applicable.
- 2b. GROUP. Enter security reclassification group number. The three groups are defined in Appendix 'M' of the DRB Security Regulations.
3. DOCUMENT TITLE. Enter the complete document title in all capital letters. Titles in all cases should be unclassified. If a sufficiently descriptive title cannot be selected without classification, show title classification with the usual one capital-letter abbreviation in parentheses immediately following the title.
4. DESCRIPTIVE NOTES. Enter the category of document, e.g. technical report, technical note or technical letter. If appropriate, enter the type of document, e.g. interim, progress, summary, annual or final. Give the inclusive dates when a specific reporting period is covered.
5. AUTHOR(S). Enter the name(s) of author(s) as shown on or in the document. Enter last name, first name, middle initial if necessary, show rank. The name of the principal author is an absolute minimum requirement.
6. DOCUMENT DATE. Enter the date (month, year) of Establishment approval for publication of the document.
- 7a. TOTAL NUMBER OF PAGES. The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- 7b. NUMBER OF REFERENCES. Enter the total number of references cited in the document.
- 8a. PROJECT OR GRANT NUMBER. If appropriate, enter the applicable research and development project or grant number under which the document was written.
- 8b. CONTRACT NUMBER. If appropriate, enter the applicable number under which the document was written.
- 9a. ORIGINATOR'S DOCUMENT NUMBER(S). Enter the official document number by which the document will be identified and controlled by the originating activity. This number must be unique to this document.
- 9b. OTHER DOCUMENT NUMBER(S). If the document has been assigned any other document numbers (either by the originator or by the sponsor), also enter this number(s).
10. DISTRIBUTION STATEMENT. Enter any limitations on further dissemination of the document, other than those imposed by security classification, using standard statements such as:
 - (1) "Qualified requesters may obtain copies of this document from their defence documentation center."
 - (2) "Announcement and dissemination of this document is not authorized without prior approval from originating activity."
11. SUPPLEMENTARY NOTES. Use for additional explanatory notes.
12. SPONSORING ACTIVITY. Enter the name of the departmental project office or laboratory sponsoring the research and development. Include address.
13. ABSTRACT. Enter an abstract giving a brief and factual summary of the document, even though it may also appear elsewhere in the body of the document itself. It is highly desirable that the abstract of classified documents be unclassified. Each paragraph of the abstract shall end with an indication of the security classification of the information in the paragraph (unless the document itself is unclassified) represented as (TS), (S), (C), (R), or (U).

The length of the abstract should be limited to 20 single-spaced standard typewritten lines 7 1/4 inches long.
14. KEY WORDS. Key words are technically meaningful terms or short phrases that characterize a document and could be helpful in cataloging the document. Key words should be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context.