

AD-A158 132


TWO RESULTS ON DENSE IMBEDDINGS(U) WISCONSIN
UNIV-MADISON MATHEMATICS RESEARCH CENTER M RENARDY
JUN 85 MRC-TSR-2833 DAAG29-80-C-0041


1/1

UNCLASSIFIED

F/G 12/1

NL





NATIONAL BUREAU OF STANDARDS
MICROCOPY RESOLUTION TEST CHART

AD-A158 132

MRC Technical Summary Report # 2833

TWO RESULTS ON DENSE IMBEDDINGS

Michael Renardy

**Mathematics Research Center
University of Wisconsin—Madison
610 Walnut Street
Madison, Wisconsin 53705**

June 1985

(Received May 10, 1985)

DTIC
ELECTE
AUG 21 1985
S
D

DTIC FILE COPY

Approved for public release
Distribution unlimited

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

National Science Foundation
Washington, DC 20550

SW

85 8 9 123

(A)

UNIVERSITY OF WISCONSIN - MADISON
MATHEMATICS RESEARCH CENTER

TWO RESULTS ON DENSE IMBEDDINGS

Michael Renardy

Technical Summary Report #2833

June 1985

ABSTRACT

In a recent paper [1], J. U. Kim ^{studied} the Cauchy problem for a Bingham fluid in $\mathbb{R}^2$. He points out that the extension of his results to $\mathbb{R}^3$ depends on two lemmas concerning dense imbedding of C_0^∞ -functions in certain spaces. In this ^{paper} ~~note~~, these lemmas are proved. _{sub 0 infinity}

Additional keywords: Sobolev spaces; approximation (mathematical).

AMS(MOS) Subject Classification: 46E35

Key Words: Sobolev spaces, approximation by test functions

Work Unit Number 1 - Applied Analysis

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



Sponsored by the United States Army under Contract No. DAAG29-80-C-0041 and the National Science Foundation under Grant No. MCS-8215064.

TWO RESULTS ON DENSE IMBEDDINGS

Michael Renardy

Following Kim's notation, we define

$$\tilde{F}_p = \{u \in W^{1,2}(\mathbb{R}^n) \mid \forall u \in L^p(\mathbb{R}^n)\} .$$

Our first result is the following.

Lemma 1:

$$C_0^\infty(\mathbb{R}^n) \text{ is dense in } \tilde{F}_p \text{ for } 1 \leq p < \infty .$$

Proof:

Clearly it suffices to show that functions of compact support are dense; C^∞ -regularity can easily be achieved by using the Friedrichs mollifier. If we know that $u \in L^p(\mathbb{R}^n)$ or even that $u \in L^{p+\epsilon}(\mathbb{R}^n)$ for sufficiently small $\epsilon > 0$, then we can use the standard cut-off procedure to approximate u by functions of compact support. I.e., if we set $u_m(x) = u(x)\psi_m(x)$, where, for example,

$$\psi_m(x) = \begin{cases} 1 & , |x| \leq m , \\ 2 - \frac{|x|}{m} & , m \leq |x| \leq 2m , \\ 0 & , |x| \geq 2m , \end{cases}$$

then it can easily be shown that $u_m \rightarrow u$ in $\tilde{F}_p$. Therefore it suffices to show that $\tilde{F}_p \cap L^{p+\epsilon}$ ($\epsilon \geq 0$ small) is dense in $\tilde{F}_p$. If $p \geq 2$, it follows from the Sobolev imbedding theorem that $\tilde{F}_p \subset L^p$, and there is nothing left to prove.

In the following, we assume $p < 2$. We set

$$\varphi_N(x) = \begin{cases} \frac{1}{N^n \Omega_n} & , \quad |x| \leq N \\ 0 & , \quad |x| > N \end{cases}$$

Here Ω_n is the volume of the unit ball in $\mathbb{R}^n$. For $u \in \tilde{F}_p$, let $u_N = u - \varphi_N * u$, where $*$ denotes convolution. Clearly, u_N is in $\tilde{F}_p$, and we want to show $u_N \rightarrow u$ in $\tilde{F}_p$. We have $\nabla u_N = \nabla u - \varphi_N * \nabla u$, and, if $p = 1$, then $\int_{\mathbb{R}^n} \nabla u = 0$, since $u \in L^2(\mathbb{R}^n)$. Therefore, it suffices to show the following: Let $1 \leq r < \infty$, and $v \in L^r$, for $r = 1$, assume in addition that $\int_{\mathbb{R}^n} v = 0$. Then $v_N = v - \varphi_N * v \rightarrow v$ in L^r .

To see this, note first that $\|\varphi_N\|_{L^1} = 1$, and hence $\|v_N\|_{L^r} \leq 2\|v\|_{L^r}$, hence it suffices to show $v_N \rightarrow v$ for v in a dense subset of L^r . If $r > 1$, take $v \in L^1 \cap L^r$. Then $\|\varphi_N * v\|_{L^r} \leq \|\varphi_N\|_{L^r} \|v\|_{L^1}$, which tends to zero. For $r = 1$, let v have compact support, contained in, say $\{|x| \leq R\}$, and assume $\int_{\mathbb{R}^n} v = 0$. Then

$$\begin{aligned} \|\varphi_N * v\|_{L^1} &= \int_{\mathbb{R}^n} \left| \int_{\mathbb{R}^n} \varphi_N(x-y)v(y)dy \right| dx \\ &= \int_{N-R \leq |x| \leq N+R} \left| \int_{|y| \leq R} \varphi_N(x-y)v(y)dy \right| dx \\ &\leq \int_{N-R \leq |x| \leq N+R} \int_{|y| \leq R} |\varphi_N(x-y)| |v(y)| dy dx \\ &\leq \int_{N-2R \leq |z| \leq N} |\varphi_N(z)| dz \cdot \int_{|y| \leq R} |v(y)| dy . \end{aligned}$$

This tends to zero as $N \rightarrow \infty$.

It remains to be shown that u_N lies in $L^{p+\epsilon}(\mathbb{R}^n)$ for small $\epsilon > 0$. Let g denote the fundamental solution of the Laplacian. Then an explicit calculation shows that $g - \varphi_N * g$ lies in $L^{1+\delta}(\mathbb{R}^n)$ for small positive δ , and so do its derivatives. Hence it follows that $w_N := g * \nabla u_N = (g - \varphi_N * g) * \nabla u$ lies in $L^{p+\epsilon}(\mathbb{R}^n)$ and so does $u_N = \operatorname{div} w_N$. This completes the proof.

Again following Kim's notation, we now define the following spaces of vector-valued functions.

$$G_1(\mathbb{R}^3) = \{ \underline{f} \in (W^{1,2}(\mathbb{R}^3))^3 \mid \epsilon_{ij}(\underline{f}) := \frac{\partial f_i}{\partial x_j} - \frac{\partial f_j}{\partial x_i} \in L^1(\mathbb{R}^3) \\ \text{for } i, j = 1, 2, 3, \operatorname{div} \underline{f} = 0 \}$$

$$S(\mathbb{R}^3) = \{ \underline{f} \in (C_0^\infty(\mathbb{R}^3))^3 \mid \operatorname{div} \underline{f} = 0 \}.$$

The lemma required for Kim's result is the following.

Lemma 2:

$S(\mathbb{R}^3)$ is dense in $G_1(\mathbb{R}^3)$.

Proof:

Obviously, G_1 is contained in

$$G_p = \{ \underline{f} \in (W^{1,2}(\mathbb{R}^3))^3 \mid \epsilon_{ij}(\underline{f}) \in L^p(\mathbb{R}^3), \operatorname{div} \underline{f} = 0 \}$$

for any $p \in [1, 2]$. Moreover, lemma 1.9 in [1] says that

$$G_p = F_p = \{ \underline{f} \in (W^{1,2}(\mathbb{R}^3))^3 \mid \nabla \underline{f} \in L^p(\mathbb{R}^3), \operatorname{div} \underline{f} = 0 \}.$$

for $p > 1$. Let $\underline{f}_N = \underline{f} - \varphi_N^* \underline{f}$ with φ_N as in the proof of lemma 1. Then it follows as before that $\underline{f}_N \rightarrow \underline{f}$ in G_1 . Let $\underline{a}$ be defined by

$\underline{a} = g^* \operatorname{curl} \underline{f}$, where g is again the fundamental solution of the Laplacian.

The convolution makes sense, since we can write $g = g_1 + g_2$, where

$g_1 \in L^1(\mathbb{R}^3)$, $\nabla g_2 \in L^2(\mathbb{R}^3)$, and we define $g_2^* \operatorname{curl} \underline{f}$ by putting the

derivative on g_2 . Clearly, we have $\operatorname{div} \underline{a} = 0$, $\operatorname{curl} \underline{a} = \underline{f}$. Now let

$\underline{a}_N = \underline{a} - \varphi_N^* \underline{a}$, so that $\operatorname{curl} \underline{a}_N = \underline{f}_N$, and $\Delta \underline{a}_N = \operatorname{curl} \underline{f}_N$. Since

$\underline{a}_N = (g - \varphi_N^* g)^* \operatorname{curl} \underline{f}$ and $\operatorname{curl} \underline{f} \in L^{1+\epsilon}(\mathbb{R}^3)$ for ϵ small, we conclude as

in the proof of lemma 1 that $\underline{a}_N \in L^{1+\epsilon}(\mathbb{R}^3)$. From $\underline{a}_N \in L^{1+\epsilon}(\mathbb{R}^3)$,

$\Delta \underline{a}_N \in L^{1+\epsilon}(\mathbb{R}^3)$, it follows that $\underline{a}_N \in W^{2,1+\epsilon}(\mathbb{R}^3)$.

It thus remain to be shown that every $\underline{f} \in G_1$ which has the form $\underline{f} = \text{curl } \underline{a}$ with $\underline{a} \in W^{2,1+\epsilon}(\mathbb{R}^3)$ can be approximated by functions of compact support. This is easily achieved by multiplying $\underline{a}$ with an appropriate cut-off function.

REFERENCES

- [1] J. U. Kim, On the Cauchy problem associated with the motion of a Bingham fluid in the plane, to appear.

MR/jvs

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER #2833	2. GOVT ACCESSION NO. A158132	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Two Results on Dense Imbeddings		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Michael Renardy		8. CONTRACT OR GRANT NUMBER(s) MCS-8215064. DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53705		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis
11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18 below		12. REPORT DATE June 1985
		13. NUMBER OF PAGES 5
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES U. S. Army Research Office National Science Foundation P. O. Box 12211 Washington, DC 20550 Research Triangle Park North Carolina 27709		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Sobolev spaces, approximation by test functions		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In a recent paper [1], J. U. Kim studies the Cauchy problem for a Bingham fluid in $\mathbb{R}^2$. He points out that the extension of his results to $\mathbb{R}^3$ depends on two lemmas concerning dense imbedding of C_0^∞ -functions in certain spaces. In this note, these lemmas are proved.		

END

FILMED

10-85

DTIC