

AD-A161 064

THE ASYMPTOTIC BEHAVIOUR NEAR THE CREST OF WAVES OF
EXTREME FORM(U) WISCONSIN UNIV-MADISON MATHEMATICS
RESEARCH CENTER J B MCLEOD AUG 85 MRC-TSR-2856

1/1


UNCLASSIFIED

DAG29-88-C-0041

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

① ~~⊗~~

AD-A161 064

MRC Technical Summary Report #2856

THE ASYMPTOTIC BEHAVIOUR NEAR THE CREST
OF WAVES OF EXTREME FORM

J. B. McLeod

Mathematics Research Center
University of Wisconsin—Madison
610 Walnut Street
Madison, Wisconsin 53705

August 1985

(Received July 19, 1985)

DTIC
ELECTE
NOV 12 1985
S D
A B

DTIC FILE COPY

Approved for public release
Distribution unlimited

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

85 11 06 060

UNIVERSITY OF WISCONSIN-MADISON
MATHEMATICS RESEARCH CENTER

THE ASYMPTOTIC BEHAVIOUR NEAR THE CREST
OF WAVES OF EXTREME FORM

J. B. McLeod

Technical Summary Report #2856
August 1985

ABSTRACT

The angle which the free boundary of an extreme wave makes with the horizontal is the solution of a singular, nonlinear integral equation. It has been proved only recently that solutions exist and that (as Stokes suggested in 1880) these solutions represent waves with sharp crests of included angle $2\pi/3$. Amick and Fraenkel have investigated the asymptotic behaviour of the free surface near the crest and obtained an asymptotic expansion for this behaviour, but are unable to say whether the leading term in this expansion has a non-zero coefficient (and so whether it is in fact the leading term or not). The present paper shows that the coefficient is non-zero and determines its sign.

AMS (MOS) Subject Classifications: 76B15, 45G05

Key Words: water waves, nonlinear integral equations, asymptotic analysis

Work Unit Number 1 (Applied Analysis)

SIGNIFICANCE AND EXPLANATION

This paper concerns waves of permanent form on the free surface of an ideal liquid which is in two-dimensional, irrotational motion under the action of gravity. ~~We consider~~ ^{It} only extreme waves, often called "waves of greatest height"; each of these is the end-member of a one-parameter family of waves and is distinguished from other "smaller" members of the family by a sharp crest. Although this corner is physically unrealistic, oceanographers have given such idealised, extreme waves a great deal of attention since Stokes postulated their existence in 1880.

~~The present paper~~ is a contribution to the strict mathematical theory of extreme waves, which has emerged only since 1978. An asymptotic series is known that describes the flow near the crest, but it has never been proved whether the leading term in this expansion has a non-zero coefficient or not (and so whether it is in fact the leading term or not). ^{author} ~~The present paper~~ shows that the coefficient is non-zero and determines its sign. The result should play a useful part in numerical computation of extreme waves.

*Summary of nonlinear integral equations;
triple integrals; water waves*



PER CALL JC



A-1

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

THE ASYMPTOTIC BEHAVIOUR NEAR THE CREST OF WAVES OF EXTREME FORM

J. B. McLeod

1. Introduction

This note is in the nature of a postscript to the paper [1], "Behaviour Near the Crest of Waves of Extreme Form," by Amick and Fraenkel. In it they discuss gravity waves, of permanent and extreme form, on the free surface of an ideal liquid, the flow being two-dimensional, irrotational and in a vertical plane. By a wave of extreme form is meant the "largest" member of a one-parameter family of such waves, and it is characterised by a sharp crest of included angle $2\pi/3$. The existence of such waves was conjectured by Stokes in 1880 and recently proved by Amick, Fraenkel and Toland [2], where a fuller account of the problem is given. (See also [3] for background information)

Amick and Fraenkel are concerned with the asymptotic behaviour of the free surface of the wave near the crest. As is described in [1], the notation of which we adopt, it is a matter of discussing the behaviour as $\zeta \rightarrow 0$ of a solution

$$\psi(\zeta) = \frac{1}{3\pi} \int_0^{\infty} \frac{\omega(\eta) \sin \psi(\eta)}{\int_0^{\infty} \omega(\tau) \sin \psi(\tau) d\tau} \log \left| \frac{\zeta + \eta}{\zeta - \eta} \right| d\eta, \quad (1.1)$$

where $0 < \zeta < \infty$ and

$$\omega(\eta) = (1 + \eta^2)^{-1/2} (1 + b\eta^2)^{-1/2},$$

the constant b satisfying $0 < b < 1$. (The value $b = 0$ corresponds to solitary waves, the values $0 < b < 1$ to periodic waves of finite depth, and the value $b = 1$ to periodic waves of infinite depth.) By a solution of (1.1) is meant a function ψ satisfying (1.1) pointwise and such that $0 < \psi < \frac{1}{2}\pi$, ψ is continuous on $(0, \infty)$, $\psi(\zeta) = O(\zeta^{-1})$ as $\zeta \rightarrow \infty$ and $\psi(\zeta) \rightarrow \frac{1}{6}\pi$ as $\zeta \rightarrow 0$. The existence of such solutions is known (see [1] for references).

It is a standard integral that

$$\frac{1}{3\pi} \int_0^{\infty} \frac{1}{\eta} \log \left| \frac{\zeta + \eta}{\zeta - \eta} \right| d\eta = \frac{1}{6}\pi, \quad (1.2)$$

and so (1.1) and (1.2) together yield

$$\psi(\zeta) - \frac{1}{6} \pi = \frac{1}{3\pi} \int_0^{\infty} \left(\frac{\omega(\eta) \sin \psi(\eta)}{\int_0^{\eta} \omega(\tau) \sin \psi(\tau) d\tau} - \frac{1}{\eta} \right) \log \left| \frac{\zeta + \eta}{\zeta - \eta} \right| d\eta \quad (1.3)$$

The basic result of [1] is that

$$\psi(\zeta) - \frac{1}{6} \pi = O(\zeta^{\beta_1}) \text{ as } \zeta \rightarrow 0, \quad (1.4)$$

where β_1 is the smallest positive root of the equation

$$\sqrt{3} (1 + \beta) = \tan \frac{1}{2} \pi \beta, \quad (1.5)$$

so that $0 < \beta_1 < 1$, but it remains possible in [1] that in fact $\psi(\zeta) - \frac{1}{6} \pi$ is of smaller order than ζ^{β_1} . It is the object of the present note to show that this is not so, and that

$$\psi(\zeta) - \frac{1}{6} \pi \sim -A \zeta^{\beta_1} \text{ as } \zeta \rightarrow 0, \quad (1.6)$$

where A is a strictly positive constant. The method, as in [1] is to use the Mellin transform on equation (1.3). The final result is therefore the following:

Theorem. If ψ is a solution of (1.1) in the sense described, then (1.6) holds.

2. Proof of the Theorem

We define the Mellin transform \hat{f} of a function f in the usual way by setting

$$\hat{f}(s) = \int_0^{\infty} \zeta^{s-1} f(\zeta) d\zeta.$$

Since from [1] we know that (1.4) holds, the transform $(\psi - \frac{1}{6} \pi)(s)$ exists and represents an analytic function of s for $-\beta_1 < \text{re}(s) < 0$. For such s , we may multiply (1.3) by ζ^{s-1} and integrate to obtain

$$\begin{aligned} (\psi - \frac{1}{6} \pi)(s) &= \frac{1}{3\pi} \int_0^{\infty} \left(\frac{\omega(\eta) \sin \psi(\eta)}{\int_0^{\eta} \omega(\tau) \sin \psi(\tau) d\tau} - \frac{1}{\eta} \right) \left(\int_0^{\infty} \zeta^{s-1} \log \left| \frac{\zeta + \eta}{\zeta - \eta} \right| d\zeta \right) d\eta \\ &= \frac{1}{3\pi} \left(\int_0^{\infty} x^{s-1} \log \left| \frac{1+x}{1-x} \right| dx \right) \int_0^{\infty} \eta^s \left(\frac{\omega(\eta) \sin \psi(\eta)}{\int_0^{\eta} \omega(\tau) \sin \psi(\tau) d\tau} - \frac{1}{\eta} \right) d\eta, \end{aligned} \quad (2.1)$$

by setting $\zeta = \eta x$. The interchange of order of integration is justified since both

integrals in (2.1) are absolutely convergent. (Note that, by the properties of ψ , the n -integrand is

$$O(n^{s-1}) \text{ as } n \rightarrow \infty \text{ and } O(n^{s+\beta_1-1}) \text{ as } n \rightarrow 0 .)$$

If we now use the standard result that the first integral in (2.1) is just $\pi \tan \frac{1}{2} \pi s / s$, and integrate the second integral by parts, we obtain, for $-\beta_1 < \operatorname{re}(s) < 0$,

$$\widehat{\left(\psi - \frac{1}{6} \pi\right)}(s) = -\frac{1}{3} \tan \frac{1}{2} \pi s \int_0^\infty n^{s-1} \log\left(\frac{2}{n} \int_0^n \omega(\tau) \sin \psi(\tau) d\tau\right) dn \quad (2.2)$$

$$= -\frac{1}{3} \tan \frac{1}{2} \pi s \int_0^\infty n^{s-1} \log\left(\frac{2}{n} \int_0^n \sin \psi(\tau) d\tau\right) dv - F(s) , \quad (2.3)$$

where

$$F(s) = \frac{1}{3} \tan \frac{1}{2} \pi s \int_0^\infty n^{s-1} \log\left(\frac{\int_0^n \omega(\tau) \sin \psi(\tau) d\tau}{\int_0^n \sin \psi(\tau) d\tau}\right) dn . \quad (2.4)$$

Since $\omega(\tau) = 1 + O(\tau^2)$ as $\tau \rightarrow 0$, we see that $F(s)$ is an analytic function of s for $-1 < \operatorname{re}(s) < 0$. But also, since $0 < \psi < \frac{1}{2} \pi$ and $0 < \omega < 1$, we see that $F(s)$ is strictly positive for $-1 < s < 0$. If we further write the logarithm in (2.3) in the form

$$\log\left(1 + \frac{2}{n} \int_0^n (\sin \psi(\tau) - \frac{1}{2}) d\tau\right) ,$$

and note that

$$\log(1+t) < t \text{ for } t > -1$$

and that

$$\sin \psi - \frac{1}{2} < \frac{1}{2} \sqrt{3} \left(\psi - \frac{1}{6} \pi\right) \text{ for } 0 < \psi < \frac{1}{2} \pi ,$$

we see that (2.3) becomes

$$\widehat{\left(\psi - \frac{1}{6} \pi\right)}(s) = -\frac{1}{\sqrt{3}} \tan \frac{1}{2} \pi s \int_0^\infty n^{s-2} \left(\int_0^n (\psi(\tau) - \frac{1}{6} \pi) d\tau\right) dn - F_1(s) , \quad (2.5)$$

where $F_1(s)$ is a function analytic for $\max(-1, -2\beta_1) < \operatorname{re}(s) < 0$ and strictly positive for $\max(-1, -2\beta_1) < s < 0$. (In fact, β_1 can be explicitly evaluated from (1.5), and $2\beta_1 > 1$.)

If finally we perform an integration by parts on the integral in (2.5), we obtain,

for $-\beta_1 < \text{re}(s) < 0$,

$$\left(1 - \frac{1}{\sqrt{3}} \frac{\tan \frac{1}{2} \pi s}{s-1}\right) \left(\psi - \frac{1}{6} \pi\right)(s) = -F_1(s),$$

or

$$\left(\psi - \frac{1}{6} \pi\right)(s) = -\frac{\sqrt{3} (1-s) F_1(s)}{\sqrt{3} (1-s) + \tan \frac{1}{2} \pi s},$$

and we use the inversion theorem on this. Thus

$$\psi(\zeta) - \frac{1}{6} \pi = \lim_{T \rightarrow \infty} \frac{1}{2\pi i} \int_{c-iT}^{c+iT} \frac{-\sqrt{3} (1-s) F_1(s)}{\sqrt{3} (1-s) + \tan \frac{1}{2} \pi s} \zeta^{-s} ds,$$

where c is any number satisfying $-\beta_1 < c < 0$.

Now move the path of integration from $\text{re}(s) = c$ to $\text{re}(s) = -l$, where $\beta_1 < l < 1$. Since the integrand is analytic for $-1 < \text{re}(s) < 0$ except for a simple pole at $s = -\beta_1$, where the residue is

$$-\frac{\sqrt{3} (1 + \beta_1) F_1(\beta_1)}{-\sqrt{3} + \frac{1}{2} \pi \sec^2 \frac{1}{2} \pi \beta_1} \zeta^{\beta_1} = -\Lambda \zeta^{\beta_1},$$

say, where $\Lambda > 0$, we have

$$\left(\psi - \frac{1}{6} \pi\right)(s) = -\Lambda \zeta^{\beta_1} + \lim_{T \rightarrow \infty} \frac{1}{2\pi i} \int_{-l-iT}^{-l+iT} \frac{-\sqrt{3} (1-s) F_1(s)}{\sqrt{3} (1-s) + \tan \frac{1}{2} \pi s} \zeta^{-s} ds. \quad (2.6)$$

The proof of the theorem is thus complete provided we can show

(i) that the contribution from the horizontal portions of the transformed contour, on $\text{im}(s) = \pm T$, are negligible as $T \rightarrow \infty$,

and

(ii) that the remaining integral on the right of (2.6) is $O(\zeta^l)$ as $\zeta \rightarrow 0$.

To prove (i), consider $F(s)$ as given by (2.4). Since the logarithm is $O(n^2)$ as $n \rightarrow 0$ and $O(\log n)$ as $n \rightarrow \infty$, we see that $|F(s)| \rightarrow 0$ as $\text{im}(s) \rightarrow \pm \infty$, by the Riemann-Lebesgue lemma in its Mellin form, provided that $-1 < \text{re}(s) < 0$. A similar remark can be made about $F_1(s)$, and this is sufficient to prove (i).

To prove (ii), we can integrate (2.4) by parts to obtain

$$F(s) = -\frac{1}{3s} \tan \frac{1}{2} \pi s \int_0^{\infty} n^s \left(\frac{w(n) \sin \psi(n)}{\int_0^n w(\tau) \sin \psi(\tau) dt} - \frac{\sin \psi(n)}{\int_0^n \sin \psi(\tau) dt} \right) dn .$$

With $\text{re}(s) = -1$, consider $F(s)$ as a function of $\text{im}(s)$. Since the integrand is $O(n^{s-1})$ as $n \rightarrow \infty$ and $O(n^{s+1})$ as $n \rightarrow 0$, Parseval's theorem in its Mellin form tells us that the integral is $L^2(-\infty, \infty)$ as a function of $\text{im}(s)$. In view of the factor s^{-1} in front of the integral, we see that $F \in L(-\infty, \infty)$. Similar remarks can be made about F_1 , and it is then clear that the integral in (2.6) is $O(\zeta^{\frac{1}{2}})$.

This completes the proof of the theorem.

REFERENCES

1. C.J. Amick and L.E. Fraenkel, Behaviour near the crest of waves of extreme form, Trans. Amer. Math. Soc., to appear; MRC Technical Summary Report No. 2687, University of Wisconsin.
2. C.J. Amick, L.E. Fraenkel and J.F. Toland, On the Stokes conjecture for the wave of extreme form, Acta Math. 148(1982), 193-214.
3. L.M. Milne-Thomson, Theoretical Hydrodynamics, (5th ed., Macmillan, 1968).

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2856	2. GOVT ACCESSION NO. AD-A161064	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) THE ASYMPTOTIC BEHAVIOUR NEAR THE CREST OF WAVES OF EXTREME FORM		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) J. B. McLeod		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Madison, Wisconsin 53705		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE August 1985
		13. NUMBER OF PAGES 6
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) water waves nonlinear integral equations asymptotic analysis		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The angle which the free boundary of an extreme wave makes with the horizontal is the solution of a singular, nonlinear integral equation. It has been proved only recently that solutions exist and that (as Stokes suggested in 1880) these solutions represent waves with sharp crests of included angle $2\pi/3$. Amick and Fraenkel have investigated the asymptotic behaviour of the free surface near the crest and obtained an asymptotic expansion for this behaviour, but are unable to say whether the leading term in this expansion has a non-zero coefficient (and so whether it is in fact the leading term or not). The present paper shows that the coefficient is non-zero and determines its sign.		

END

FILMED

12-85

DTIC