

AD-A161 863

INDEPENDENCE AND BAYESTAN UPDATING METHODS(U) NAVAL
RESEARCH LAB WASHINGTON DC R W JOHNSON 06 NOV 85
NRL-MR-5689


1/1

UNCLASSIFIED

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

2

NRL Memorandum Report 5689

Independence and Bayesian Updating Methods

R. W. JOHNSON

*Computer Science and Systems Branch
Information Technology Division*

November 6, 1985

AD-A161 863

DTIC FILE COPY

DTIC
ELECTE
DEC 3 1985
S B D



NAVAL RESEARCH LABORATORY
Washington, D.C.

Approved for public release; distribution unlimited.

85 11 25 069

AD-A161863

REPORT DOCUMENTATION PAGE				
1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited.	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE				
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NRL Memorandum Report 5689			5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION Naval Research Laboratory	6b. OFFICE SYMBOL (If applicable) Code 7591	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) Washington, DC 20375-5000		7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code) Arlington, VA 22217		10. SOURCE OF FUNDING NUMBERS PROGRAM ELEMENT NO. 61153N	PROJECT NO.	TASK NO. RR021-05-42 WORK UNIT ACCESSION NO. 75-2424-0-5
11. TITLE (Include Security Classification) Independence and Bayesian Updating Methods				
12. PERSONAL AUTHOR(S) Johnson, R.W.				
13a. TYPE OF REPORT Interim	13b. TIME COVERED FROM TO	14. DATE OF REPORT (Year, Month, Day) 1985 November 6	15. PAGE COUNT 9	
16. SUPPLEMENTARY NOTATION				
17. COSATI CODES FIELD GROUP SUB-GROUP			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Expert system ; Bayes ; Probability updating ; Rule-based system ; Bayesian ;	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Duda, Hart, and Nilsson have set forth a method for rule-based inference systems to use in updating the probabilities of hypotheses on the basis of multiple items of new evidence. Pednault, Zucker, and Muresan claimed to give conditions under which independence assumptions made by Duda et al, preclude updating — that is, prevent the evidence from altering the probabilities of the hypotheses. Glymour refutes Pednault et al.'s claim with a counterexample of a rather special form (one item of evidence is incompatible with all but one of the hypotheses); he raises, but leaves open, the question whether their result would be true with an added assumption to rule out such special cases. We show that their result does not hold even with the added assumption, but that it can nevertheless be largely salvaged. Namely, under the conditions assumed by Pednault et al., at most one of the items of evidence can alter the probability of any given hypothesis; thus, although updating is possible, multiple updating for any of the hypotheses is precluded. Keywords:				
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT. ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL R. W. Johnson			22b. TELEPHONE (Include Area Code) (202) 767-3490	22c. OFFICE SYMBOL Code 7591

CONTENTS

I. BACKGROUND	1
II. COUNTEREXAMPLES	3
III. IMPOSSIBILITY OF MULTIPLE UPDATING	4
IV. DISCUSSION	5
REFERENCES	5

Accession For	
NTIS GOMA	☒
DAIR Tag	☐
Unrec. copy	☐
By	
Distribution	
Availability Codes	
Dist.	Avail and/or Special
A-1	



INDEPENDENCE AND BAYESIAN UPDATING METHODS

I. BACKGROUND

Duda, Hart, and Nilsson [1] consider the problem of updating the probability of a hypothesis H with prior probability $P(H)$ when new evidence is obtained in the form of propositions E_j for which the conditional probabilities $P(E_j | H)$ and $P(E_j | \bar{H})$ are known. They assume that the E_j are conditionally independent, both on condition H and on condition $\bar{H}$, so that

$$P(E_1 \cdots E_m | H) = \prod_{j=1}^m P(E_j | H)$$

$$P(E_1 \cdots E_m | \bar{H}) = \prod_{j=1}^m P(E_j | \bar{H}).$$

They can then use the odds-likelihood form of Bayes's theorem and write an updating formula for the odds on H in terms of a product of likelihood ratios:

$$\frac{P(H | E_1 \cdots E_m)}{P(\bar{H} | E_1 \cdots E_m)} = \frac{P(H)}{P(\bar{H})} \prod_{j=1}^m \frac{P(E_j | H)}{P(E_j | \bar{H})}.$$

Pednault, Zucker, and Muresan [2], in analyzing this updating scheme, considered the consequences of imposing the independence assumptions for each hypothesis H_i of a jointly exhaustive, mutually exclusive set. They [2] and other writers (see [3] and other references therein) agree that the assumptions are unreasonably strong, but there has been some confusion over the exact extent of the undesirable consequences. Pednault *et al.* [2] concluded that if there were at least three hypotheses, then no updating could take place—that the assumptions are too strong to be satisfied unless

$$P(E_j | H_i) = P(E_j | \bar{H}_i) = P(E_j)$$

holds for all i and j , and consequently

$$P(H_i | E_1 \cdots E_m) = P(H_i).$$

However, Glymour [3] gives a counterexample to their conclusion—three jointly exhaustive, mutually exclusive hypotheses H_i and two evidence propositions E_j that satisfy the independence assumptions but allow updating to occur. He points out that Pednault *et*

al. had relied on an erroneous result claimed by Hussain [4], also refuted by his counterexample. Glymour notes that the evidence proposition E_2 of his counterexample has the special property that $P(E_2 | H_2) = P(E_2 | H_3) = 0$, so that E_2 determines a posterior probability of 1 for one hypothesis, H_1 , and 0 for the rest. He raises, and leaves open, the question whether Pednault *et al.*'s result would be true with the additional requirement that for all i ,

$$P(H_i | E_1 \cdots E_m) \neq 0. \quad (1)$$

In the next section we answer that question by giving a counterexample that satisfies (1).

Glymour's counterexample has another special property, one that is sufficient to make it a valid counterexample: $P(E_1 | H_i) = P(E_1 | \bar{H}_i) = P(E_1)$, so that E_1 produces no updating; only E_2 produces updating. We give a second counterexample that lacks this second special property. However, we can show that Pednault *et al.*'s assumptions imply that for every i there is at most one value of j for which

$$P(E_j | H_i) = P(E_j | \bar{H}_i) = P(E_j)$$

does not hold. We show this in the section following the counterexamples. It follows that for each hypothesis H_i , there is at most one evidence proposition E_j that produces updating of the probability of H_i ; there is no hypothesis for which multiple updating is possible.

We conclude this section by stating the assumptions used by Pednault *et al.* The hypotheses are $H_1, \dots, H_n$; it is assumed that

$$n > 2 \quad (2)$$

and that the H_i are jointly exhaustive

$$\sum_{i=1}^n P(H_i) = 1 \quad (3)$$

and mutually exclusive

$$P(H_i H_j) = 0 \quad (i \neq j). \quad (4)$$

The evidence propositions are $E_1, \dots, E_m$. Since a subset of an independent set is independent, we write the independence assumptions as

$$P(E_{j_1} \cdots E_{j_k} | H_i) = \prod_{j \in J} P(E_j | H_i), \quad (5)$$

$$P(E_{j_1} \cdots E_{j_k} | \bar{H}_i) = \prod_{j \in J} P(E_j | \bar{H}_i). \quad (6)$$

for every subset $J = \{j_1, \dots, j_k\}$ of the indices $1, \dots, m$.

II. COUNTEREXAMPLES

For comparison, here is Glymour's counterexample.

H_i	$P(E_1 E_2 H_i)$	$P(E_1 \bar{E}_2 H_i)$	$P(\bar{E}_1 E_2 H_i)$	$P(\bar{E}_1 \bar{E}_2 H_i)$
H_1	1/6	0	1/6	0
H_2	0	1/6	0	1/6
H_3	0	1/6	0	1/6

It is straightforward to verify that

$$\begin{aligned} P(E_2 | H_1) &= 1, \\ P(E_2 | H_2) &= P(E_2 | H_3) = 0, \end{aligned} \quad (7)$$

as noted by Glymour, and that

$$P(E_1 | H_i) = P(E_1 | \bar{H}_i) = P(E_1) \quad (8)$$

for each i .

To answer the question raised by Glymour, we modify the example so that (7) no longer holds. We retain the same values for $P(H_i)$ and $P(E_1 | H_i)$, choose new, nonzero values for $P(E_2 | H_i)$ (say 1/2, 1/3, 1/6 for $i = 1, 2, 3$), and define the remaining relevant conditional probabilities and probabilities with the help of (5). Here is the result.

H_i	$P(E_1 E_2 H_i)$	$P(E_1 \bar{E}_2 H_i)$	$P(\bar{E}_1 E_2 H_i)$	$P(\bar{E}_1 \bar{E}_2 H_i)$
H_1	1/12	1/12	1/12	1/12
H_2	1/18	1/9	1/18	1/9
H_3	1/36	5/36	1/36	5/36

Assumptions (2)–(6) can be verified. In fact, we can show that as long as E_1 satisfies (8), we can choose $P(E_2 | H_i)$ arbitrarily, and the procedure we have just used will lead to a probability distribution that satisfies (2)–(6).

Now (8) implies that E_1 is irrelevant for inference about the hypotheses. Only E_2 produces updating—multiple updating does not occur. But by going to four hypotheses, we can dispense with (8) and obtain a counterexample such that E_1 and E_2 can both produce updating.

H_i	$P(E_1 E_2 H_i)$	$P(E_1 \bar{E}_2 H_i)$	$P(\bar{E}_1 E_2 H_i)$	$P(\bar{E}_1 \bar{E}_2 H_i)$
H_1	1/24	1/24	1/12	1/12
H_2	1/12	1/12	1/24	1/24
H_3	1/24	1/12	1/24	1/12
H_4	1/12	1/24	1/12	1/24

Again (2)–(6) can be verified. Furthermore E_1 and E_2 can both produce updating since

we have, for example, $P(E_1 | H_1) \neq P(E_1)$ and $P(E_2 | H_3) \neq P(E_2)$. However, we have

$$P(E_1 | H_i) = P(E_1 | \bar{H}_i) = P(E_1) \quad (i = 3, 4),$$

$$P(E_2 | H_i) = P(E_2 | \bar{H}_i) = P(E_2) \quad (i = 1, 2).$$

Thus only E_1 can update the probability of H_1 or H_2 , and only E_2 can update the probability of H_3 or H_4 ; for no hypothesis is multiple updating possible. This illustrates the general case, as we show in the next section.

III. IMPOSSIBILITY OF MULTIPLE UPDATING

Theorem. If the assumptions (2)–(6) hold, then for every H_i there is at most one E_j that produces updating for H_i .

Proof. No updating for H_i is possible if either $P(H_i)$ or $P(\bar{H}_i)$ is 0; therefore we may assume that both are nonzero. First consider the case $m = 2$, where the evidence propositions are just E_1 and E_2 . We follow Pednault *et al.* (see [2], equations (6)–(9)) in deriving

$$\begin{aligned} P(E_1)P(E_2) - P(E_1)P(E_2H_i) - P(E_1H_i)P(E_2) \\ = P(E_1E_2)[1 - P(H_i)] - P(E_1E_2H_i) \end{aligned} \quad (9)$$

and summing over i to obtain

$$(n - 2)P(E_1)P(E_2) = (n - 2)P(E_1E_2),$$

or

$$P(E_1)P(E_2) = P(E_1E_2). \quad (10)$$

Using (9) and (10) with the help of (5), we obtain

$$\begin{aligned} P(E_1)P(E_2 | H_i)P(H_i) + P(E_1 | H_i)P(E_2)P(H_i) \\ = P(E_1)P(E_2)P(H_i) + P(E_1 | H_i)P(E_2 | H_i)P(H_i) \end{aligned}$$

from which it follows that

$$[P(E_1) - P(E_1 | H_i)][P(E_2) - P(E_2 | H_i)] = 0.$$

One of the bracketed factors vanishes. If $P(E_j | H_i) = P(E_j)$, then in fact

$$P(E_j | H_i) = P(E_j | \bar{H}_i) = P(E_j);$$

this therefore holds either with $j = 1$ or with $j = 2$. Consequently E_1 and E_2 do not both produce updating for H_i . Thus we have proved the theorem for the case $m = 2$. But in the general case, if one of the evidence propositions, say E_j , produces updating for H_i , the result for $m = 2$ implies that no other evidence proposition E_k produces updating for H_i .

IV. DISCUSSION

Assumptions (2)–(6) lead to unreasonably severe restrictions on the possibility of probabilistic updating. If we wish to make inferences about more than two jointly exhaustive, mutually exclusive hypotheses, and if we wish to allow more than one piece of evidence to bear on one hypothesis, then we must eliminate either (5) or (6). Duda *et al.* made the assumptions (5) and (6) in the context of a single pair of hypotheses $H, \bar{H}$. It is clearly a mistake to carry them both over into the context of several exhaustive, mutually exclusive hypotheses.

Cases where (5) is justified (at least as an approximation) are quite common, but (6) is much less plausible. Consider, for example, a physical quantity x that can take any of a considerable range of numerical values v_i . Let y and z be measurements of x made with instruments subject to independent errors; that is, suppose $y - x$ and $z - x$ are independent random variables but are fairly small with high probability. Then $y - z$ is small with high probability, and so y and z are highly dependent; but on condition of a given value of x , say $x = v_i$, the conditional distributions of y and z are independent. Define H_i to be $x = v_i$. Then y and z are independent on condition H_i for any i . However, on condition $\bar{H}_i$, we expect them to be dependent, for the same reason that their unconditional distributions are dependent. In that case, we can take E_1 and E_2 to be propositions about y and z , respectively, and it is easy to choose E_1 and E_2 so that they are conditionally independent on condition H_i but not on condition $\bar{H}_i$.

We must conclude that for inference about jointly exhaustive, mutually exclusive hypotheses, updating schemes based on the independence assumption (5) alone may be useful, but schemes based on both (5) and (6) are too restrictive to be useful.

REFERENCES

1. R. O. Duda, P. E. Hart, and N. J. Nilsson, "Subjective Bayesian methods for rule-based inference systems," pp. 1075-1082 in *Proc. 1976 National Computer Conf.* (1976).
2. E. P. D. Pednault, S. W. Zucker, and L. V. Muresan, "On the independence assumption underlying subjective Bayesian updating," *Artificial Intelligence* 16, pp. 213-222 (May 1981).
3. C. Glymour, "Independence assumptions and Bayesian updating," *Artificial Intelligence* 25, pp. 95-99 (1985).
4. A. B. S. Hussain, "On the correctness of some sequential classification schemes in pattern recognition," *IEEE Trans. Computers* 21, pp. 318-320 (March, 1972).

END

FILMED

1-86

DTIC