

AD-A161 989

STABILITY ANALYSIS OF DIFFERENCE SCHEMES FOR VARIABLE  
COEFFICIENT SCHRODINGER (U) YALE UNIV NEW HAVEN CT DEPT OF  
COMPUTER SCIENCE T F CHAN ET AL SEP 85


1/1


UNCLASSIFIED

VALEU/DCS/RR-413 N80014-82-K-0814

F/G 12/1


NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS 1963-A

AD-A161 989



**Stability Analysis of Difference Schemes for  
Variable Coefficient Schrödinger Type Equations**

Tony F. Chan and Longjun Shen  
Research Report YALEU/DCS/RR-413  
September 1985

DISTRIBUTION STATEMENT A

Approved for public release;  
Distribution Unlimited

YALE UNIVERSITY  
DEPARTMENT OF COMPUTER SCIENCE

85 10 7 013

ALL COPY

*Free d...*  
**Abstract.** We consider the stability of difference schemes for the solution of the initial boundary value problem for the equation

$$u_t = (A(x, t)u_x)_x + B(x, t)u_x + C(x, t)u + f(x, t),$$

where  $u$ ,  $A$ ,  $B$ ,  $C$  and  $f$  are complex valued functions. Using energy methods, <sup>*the authors*</sup> we establish the stability of a general two level scheme which includes Euler's method, Crank-Nicolson's method and the backward Euler method. If the coefficient  $A(x, t)$  is purely imaginary, the explicit Euler's method is unconditionally unstable. For this case, we propose a new scheme with appropriately chosen artificial dissipation, which we prove to be conditionally stable.

Contract N00014-82-K-0814

**Stability Analysis of Difference Schemes for  
Variable Coefficient Schrödinger Type Equations**

Tony F. Chan and Longjun Shen  
Research Report YALEU/DCS/RR-413  
September 1985

DTIC  
ELECTE  
S DEC 05 1985 D

The work was supported in part by the Department of Energy under contract DE-AC02-81ER10996, by the Army Research Office under contract DAAG-83-0177.

**Keywords:** *Schrödinger Equation, Parabolic Equation, Difference Schemes, Stability, Variable Coefficient, Energy Methods.*

**DISTRIBUTION STATEMENT A**

Approved for public release;  
Distribution Unlimited

## 1. Introduction

Finite difference methods for the solution of equations of Schrödinger type have been studied by many authors, and extensively applied to solve practical problems in many disciplines [1, 5, 6, 7, 8, 9, 10, 11]. Since conventional explicit schemes are unstable [2, 4], implicit schemes are usually used, especially the Crank-Nicolson scheme. In [7], D.F. Griffiths et al discussed a predictor-corrector scheme. In [8, 12] the existence and convergence of solution for this difference equation have been studied. Since the solution of Schrödinger equation possesses conservation laws, the schemes which satisfy discrete conservation laws have also been investigated extensively [6, 7].

We consider the equation

$$u_t = (A(x, t)u_x)_x + B(x, t)u_x + C(x, t)u + f(x, t), \quad (1.1)$$

in the domain  $Q_T(0 \leq x \leq l, 0 \leq t \leq T)$ , where  $u(x, t)$ ,  $A(x, t)$ ,  $B(x, t)$ ,  $C(x, t)$  and  $f(x, t)$  are complex functions, and  $\operatorname{Re} A(x, t) \geq 0$  and  $|A(x, t)| \neq 0$ . This kind of equation arises in plasma physics and acoustics [1, 11]. Clearly, (1.1) involves both equations of Schrödinger type and parabolic equations. In this paper, we consider the initial-boundary value problem for (1.1), with the conditions :

$$u|_{t=0} = \tilde{u}(x) \quad (1.2)$$

$$u|_{x=0} = u|_{x=l} = 0, \quad (1.3)$$

where  $\tilde{u}(x)$  is a complex function.

In Section 2, we analyse the stability of a general two level difference scheme for (1.1), (1.2) and (1.3), which is a direct generalization of a well-known scheme for parabolic equations, and includes the explicit Euler scheme and the implicit Crank-Nicolson scheme as special members. Some new results are obtained from which we can see the relationship between the Schrödinger equation and the parabolic equation.

If  $\operatorname{Re} A(x, t) \equiv 0$ , the above mentioned explicit Euler scheme is unstable. Several interesting stable explicit schemes have been presented in [3] for the case of the simplified equation  $u_t = iu_{xx}$ , and applied to some underwater acoustics problem in [2]. In these schemes, Euler's method is stabilized by appropriately chosen artificial dissipation. In Section 4, we extend these results to equation (1.1).

We let  $h$  denote the spatial mesh size and divide the finite interval  $[0, l]$  into the mesh intervals by the points  $x_j = jh$  ( $j = 0, 1, \dots, J$ ), where  $Jh = l$ . We let  $k_n$  denote the size of the time

step at the  $n$ -th step. For convenience, we shall denote  $u(x_j, t_n)$  by  $u_j^n$ . We also use the following difference operators :

$$\begin{aligned} u_{t_j}^{n+1} &= u_{t_j}^n = \frac{u_j^{n+1} - u_j^n}{k_n}, \\ D_+ u_j &= \frac{u_{j+1} - u_j}{h}, & D_- u_j &= \frac{u_j - u_{j-1}}{h}, \\ Du_j &= \frac{u_{j+1/2} - u_{j-1/2}}{h}, & \tilde{D}u_j &= \frac{u_{j+1} - u_{j-1}}{2h} = \frac{1}{2}(D_+ + D_-)u_j. \end{aligned}$$

Hence, we have

$$\begin{aligned} \frac{1}{h^2}(u_{j+1} - 2u_j + u_{j-1}) &= D_+ D_- u_j = D^2 u_j, \\ \frac{1}{h^2} [A_{j+1/2}(u_{j+1} - u_j) - A_{j-1/2}(u_j - u_{j-1})] &= D(ADu_j). \end{aligned}$$

Finally, for any function  $\phi$ , we use  $\phi^{n+\alpha}$  to denote  $\alpha\phi^{n+1} + (1-\alpha)\phi^n$ , for  $0 \leq \alpha \leq 1$ .

Next, we give our definition of stability. First we define the inner product for  $u$  and  $v$  :

$$(u, v) = \sum_{j=1}^{J-1} u_j \bar{v}_j h$$

where  $\bar{v}$  denotes the complex conjugate of  $v$  and the norm for  $u$  :

$$\|u\| = \sqrt{(u, u)}.$$

**Definition 1.1.** We call a scheme stable if the solution  $u_j^n$  satisfies :

$$\|u^n\| \leq C_1 \|u^0\| + C_2 \sum_{l=0}^{J-1} \|f^l\| k_l$$

where  $C_1$  and  $C_2$  are constants which are independent of  $n$  and  $h$ .

## 2. A General Two Level Scheme.

We are going to consider the following scheme for (1.1), (1.2) and (1.3) :

$$\begin{aligned} \frac{u_j^{n+1} - u_j^n}{k_n} - \frac{1}{h^2} [A_{j+1/2}^{n+\alpha}(u_{j+1}^{n+\alpha} - u_j^{n+\alpha}) - A_{j-1/2}^{n+\alpha}(u_j^{n+\alpha} - u_{j-1}^{n+\alpha})] \\ - B_j^{n+\alpha} \left( \frac{u_{j+1}^{n+\alpha} - u_{j-1}^{n+\alpha}}{2h} \right) - C_j^{n+\alpha} u_j^{n+\alpha} = f_j^{n+\alpha} \quad j = 1, 2, \dots, J-1 \end{aligned} \quad (2.1)$$

$$u_j^0 = \tilde{u}_j, \quad j = 1, 2, \dots, J-1. \quad (2.2)$$

$$u_0^n = u_J^n = 0, \quad n = 0, 1, \dots \quad (2.3)$$

We can also write (2.1) in this form :

$$u_{t_j}^{n+1} - DA^{n+\alpha} Du_j^{n+\alpha} - B_j^{n+\alpha} \tilde{D} u_j^{n+\alpha} - C_j^{n+\alpha} u_j^{n+\alpha} = f_j^{n+\alpha}. \quad (2.4)$$

Clearly, (2.1) or (2.4) is explicit, implicit and the Crank-Nicolson scheme when  $\alpha = 0$ ,  $\alpha \neq 0$  and  $\alpha = \frac{1}{2}$ , respectively.

We will see that this scheme is unstable if  $\operatorname{Re} A(x, t) = 0$  and  $\alpha \neq 0$ . In this case, we will give another conditionally stable explicit scheme in Section 3.

To facilitate the analysis, we first transform (2.1) by a change of variable to eliminate the first order term.

**Lemma 2.1.** Suppose  $A \in C^3$ ,  $B \in C^2$ ,  $C \in C^1$  and  $\operatorname{Re} A \geq 0$ ,  $|A| \geq a_0 > 0$ , then we can choose a function  $\phi$ , such that  $0 < M_0 \leq |\phi_j| \leq M_1$ ,  $j = 0, 1, \dots, J$  where  $M_0$  and  $M_1$  are constants and under the transformation

$$u_j^{n+\alpha} = \phi_j^{n+\alpha} v_j^{n+\alpha} \quad (2.5)$$

(2.1) becomes :

$$L(v^{n+\alpha}) \equiv \frac{v_j^{n+1} - v_j^n}{k_n} - \frac{1}{h^2} \left[ A_{j+1/2}^{n+\alpha} (v_{j+1}^{n+\alpha} - v_j^{n+\alpha}) - A_{j-1/2}^{n+\alpha} (v_j^{n+\alpha} - v_{j-1}^{n+\alpha}) \right] - G_{j-1}^{n+\alpha} v_{j-1}^{n+\alpha} - H_j^{n+\alpha} v_j^{n+\alpha} - K_{j+1}^{n+\alpha} v_{j+1}^{n+\alpha} - Q_j^{n+\alpha} v_j^{n+1-\alpha} = F_j^{n+\alpha} \quad (2.6)$$

where  $G_j, H_j, K_j, Q_j, F_j$  are bounded in  $Q_T$ .

*Proof.* See Appendix A. ■

We now state our main stability result for (2.6).

**Theorem 2.1.** Suppose the conditions in Lemma 2.1 are satisfied. If  $\frac{1}{2} \leq \alpha \leq 1$ , then scheme (2.1) is stable.

*Proof.* In order to establish the stability of (2.6), we are going to estimate  $\|v^{n+1}\|$ . We multiply (2.6) by  $\bar{v}_j^{n+\alpha}$  to obtain :

$$(L(v^{n+\alpha}), v^{n+\alpha}) = (F^{n+\alpha}, v^{n+\alpha}) \quad (2.7)$$

Then we estimate every term in (2.7). For the first term, we have :

$$\begin{aligned}
 \operatorname{Re}(v_i^{n+1}, v^{n+\alpha}) &= \alpha \operatorname{Re}(v_i^{n+1}, v^{n+1}) + (1-\alpha) \operatorname{Re}(v_i^{n+1}, v^n) \\
 &= \frac{\alpha}{k_n} \sum |v_j^{n+1}|^2 - \left( \frac{1-\alpha}{k_n} \right) \sum |\bar{v}_j^n|^2 \\
 &\quad - \frac{\alpha}{k_n} \operatorname{Re} \sum v_j^n \bar{v}_j^{n+1} + \frac{(1-\alpha)}{k_n} \operatorname{Re} \sum v_j^{n+1} \bar{v}_j^n \\
 &= \frac{1}{2k_n} (\sum |v_j^{n+1}|^2 - \sum |v_j^n|^2) - \frac{(1-2\alpha)}{2k_n} \sum |v_j^{n+1} - v_j^n|^2 \\
 &= \frac{1}{2k_n} (\|v^{n+1}\|^2 - \|v^n\|^2) - \frac{(1-2\alpha)}{2} k_n \|v_i^{n+1}\|^2.
 \end{aligned} \tag{2.8}$$

For the second term, we have

$$\begin{aligned}
 \operatorname{Re}(DA^{n+\alpha} Dv, v^{n+\alpha}) &= -\operatorname{Re}(A^{n+\alpha} D_- v^{n+\alpha}, D_- v^{n+\alpha}) \\
 &= -\|\sqrt{\operatorname{Re} A^{n+\alpha}} D_- v^{n+\alpha}\|^2.
 \end{aligned} \tag{2.9}$$

For the rest of the terms on the left hand side of (2.7), we have

$$\begin{aligned}
 |G_{j-1}^{n+\alpha} v_{j-1}^{n+\alpha} \bar{v}_j^{n+\alpha}| &\leq \frac{|G_{j-1}^{n+\alpha}|}{2} (|v_{j-1}^{n+\alpha}|^2 + |v_j^{n+\alpha}|^2) \\
 |H_j^{n+\alpha} v_j^{n+\alpha} \bar{v}_j^{n+\alpha}| &\leq \frac{|H_j^{n+\alpha}|}{2} (|v_j^{n+\alpha}|^2 + |\bar{v}_j^{n+\alpha}|^2) \\
 |K_{j+1}^{n+\alpha} v_{j+1}^{n+\alpha} \bar{v}_j^{n+\alpha}| &\leq \frac{|K_{j+1}^{n+\alpha}|}{2} (|v_{j+1}^{n+\alpha}|^2 + |\bar{v}_j^{n+\alpha}|^2) \\
 |Q_j^{n+\alpha} v_j^{n+1-\alpha} \bar{v}_j^{n+\alpha}| &\leq |Q_j^{n+\alpha}| [(1-\alpha)|v_j^{n+1}| + \alpha|v_j^n|] [\alpha|v_j^{n+1}| + (1-\alpha)|v_j^n|] \\
 &\leq \frac{|Q_j^{n+\alpha}|}{2} (|v_j^{n+1}|^2 + |v_j^n|^2).
 \end{aligned}$$

So we have

$$\begin{aligned}
 & \left| \operatorname{Re} \left[ \sum G_{j-1}^{n+\alpha} v_{j-1}^{n+\alpha} \bar{v}_j^{n+\alpha} + \sum H_j^{n+\alpha} v_j^{n+\alpha} \bar{v}_j^{n+\alpha} + \sum K_{j+1}^{n+\alpha} v_{j+1}^{n+\alpha} \bar{v}_j^{n+\alpha} + \sum Q_j^{n+\alpha} v_j^{n+1-\alpha} \bar{v}_j^{n+\alpha} \right] \right| \\
 & \leq M_2 \left[ \|v^{n+\alpha}\|^2 + \|v^{n+\alpha}\|^2 + \|v^{n+\alpha}\|^2 + \frac{1}{2} \|v^{n+1}\|^2 + \frac{1}{2} \|v^n\|^2 \right] \\
 & = 3M_2 \|v^{n+\alpha}\|^2 + \frac{M_2}{2} \|v^{n+1}\|^2 + \frac{M_2}{2} \|v^n\|^2,
 \end{aligned} \tag{2.10}$$

where  $M_2$  is a constant which is an upper bound of  $|G|$ ,  $|H|$ ,  $|K|$  and  $|Q|$ .

For the right hand side of (2.7), we get

$$|\operatorname{Re}(F^{n+\alpha}, v^{n+\alpha})| \leq \frac{1}{2} \|v^{n+\alpha}\|^2 + \frac{1}{2} \|F^{n+\alpha}\|^2. \tag{2.11}$$

Now we substitute expressions (2.8), (2.9), (2.10) and (2.11) into (2.7) and obtain

$$\begin{aligned}
 \frac{1}{2k_n} (\|v^{n+1}\|^2 - \|v^n\|^2) - \left( \frac{1-2\alpha}{2} \right) k_n \|v_t^{n+1}\|^2 + \|\sqrt{Re A^{n+\alpha}} D_- v^{n+\alpha}\|^2 \\
 \leq 3M_2 \|v^{n+\alpha}\|^2 + \frac{M_2}{2} \|v^{n+1}\|^2 + \frac{M_2}{2} \|v^n\|^2 + \frac{1}{2} \|v^{n+\alpha}\|^2 + \frac{1}{2} \|F^{n+\alpha}\|^2 \\
 \leq M_3 (\|v^{n+1}\|^2 + \|v^n\|^2) + \frac{1}{2} \|F^{n+\alpha}\|^2,
 \end{aligned} \tag{2.12}$$

where

$$M_3 = 6M_2 + \frac{M_2}{2} + 1.$$

When  $\frac{1}{2} \leq \alpha \leq 1$ , from expression (2.12) we have

$$\frac{1}{2k_n} (\|v^{n+1}\|^2 - \|v^n\|^2) \leq M_3 (\|v^{n+1}\|^2 + \|v^n\|^2) + \frac{1}{2} \|F^{n+\alpha}\|^2.$$

It is easy to see that when  $k_n \leq \frac{1}{4M_3}$  we have

$$\begin{aligned}
 \|v^{n+1}\|^2 &\leq \frac{1+2k_n M_3}{1-2k_n M_3} \|v^n\|^2 + 2k_n \|F^n\|^2 \\
 &\leq (1+8k_n M_3) \|v^n\|^2 + 2k_n \|F^n\|^2.
 \end{aligned}$$

According to Lemma 4.1 in Appendix B, we obtain

$$\|v^{n+1}\|^2 \leq 2e^{8M_3 t^{n+1}} \left[ \|v_0\|^2 + \sum_{l=0}^n \|F^l\|^2 k_l \right]. \tag{2.13}$$

From (2.5) and  $0 < M_0 \leq |\phi_j| \leq M_1$  we obtain

$$\|u^{n+1}\|^2 \leq 2 \frac{M_1^2}{M_0^2} e^{8M_3 t^{n+1}} \left[ \|u_0\|^2 + \sum_{l=0}^n \|f^l\|^2 k_l \right]. \tag{2.14}$$

Now we turn to discuss the case  $0 \leq \alpha < \frac{1}{2}$ .

**Theorem 2.2.** If  $0 \leq \alpha < \frac{1}{2}$  and

$$k_n \leq \frac{1}{2(1-2\alpha) \left[ \frac{1}{h^2} \max_x \frac{|A^{n+\alpha}|^2}{Re A^{n+\alpha}} + \frac{1}{\eta} \right]} \tag{2.15}$$

where  $\eta$  is an arbitrary positive constant then the scheme (2.1) is stable.

|                    |                                           |
|--------------------|-------------------------------------------|
| Accession For      |                                           |
| NTIS               | CRA&I <input checked="" type="checkbox"/> |
| DTIC               | TAB <input type="checkbox"/>              |
| Unannounced        | <input type="checkbox"/>                  |
| Justification      |                                           |
| By                 |                                           |
| Distribution/      |                                           |
| Availability Codes |                                           |
| Dist               | Avail and/or Special                      |
| A-1                |                                           |

*Proof.* First let

$$\begin{aligned} P_{1j} &= \frac{1}{h^2} |A_{j+1/2}^{n+\alpha}| |v_{j+1}^{n+\alpha} - v_j^{n+\alpha}| \\ P_{2j} &= \frac{1}{h^2} |A_{j-1/2}^{n+\alpha}| |v_j^{n+\alpha} - v_{j-1}^{n+\alpha}| \\ P_{3j} &= |G_{j-1}^{n+\alpha}| |v_{j-1}^{n+\alpha}| \\ P_{4j} &= |H_j^{n+\alpha}| |v_j^{n+\alpha}| \\ P_{5j} &= |K_{j+1}^{n+\alpha}| |v_{j+1}^{n+1-\alpha}| \\ P_{6j} &= |Q_j^{n+\alpha}| |v_j^{n+\alpha}| \\ P_{7j} &= |F_j^{n+\alpha}|. \end{aligned}$$

From (2.6) we obtain immediately

$$|v_{t_j}^{n+1}|^2 \leq \left( \sum_{l=1}^7 P_{lj} \right)^2. \quad (2.16)$$

Next we construct a quadratic form in  $P_{lj}$  ( $l = 1, \dots, 7$ ) by :

$$P_j = \frac{h^2 \operatorname{Re} A_{j+1/2}^{n+\alpha}}{|A_{j+1/2}^{n+\alpha}|^2} P_{1j}^2 + \frac{h^2 \operatorname{Re} A_{j-1/2}^{n+\alpha}}{|A_{j-1/2}^{n+\alpha}|^2} P_{2j}^2 + 5\eta \sum_{l=3}^7 P_{lj}^2 - (1 - 2\alpha)k_n \left( \sum_{l=1}^7 P_{lj} \right)^2 \quad (2.17)$$

where  $\eta$  is an arbitrary positive constant. According to Lemma 4.2 (in Appendix C),  $P_j$  is nonnegative if

$$k_n \leq \frac{1}{(1 - 2\alpha) \left[ \frac{|A_{j+1/2}^{n+\alpha}|^2}{h^2 \operatorname{Re} A_{j+1/2}^{n+\alpha}} + \frac{|A_{j-1/2}^{n+\alpha}|^2}{h^2 \operatorname{Re} A_{j-1/2}^{n+\alpha}} + \frac{1}{\eta} \right]}, \quad (2.18)$$

which is true because of (2.15). Making this assumption, we have from (2.16) :

$$(1 - 2\alpha)k_n |v_{t_j}^{n+1}|^2 \leq \left( \frac{h^2 \operatorname{Re} A_{j+1/2}^{n+\alpha}}{|A_{j+1/2}^{n+\alpha}|^2} \right) P_{1j}^2 + \left( \frac{h^2 \operatorname{Re} A_{j-1/2}^{n+\alpha}}{|A_{j-1/2}^{n+\alpha}|^2} \right) P_{2j}^2 + 5\eta \sum_{l=3}^7 P_{lj}^2$$

Summing up these expressions from  $j = 1$  to  $j = J - 1$ , we obtain

$$\begin{aligned} (1 - 2\alpha)k_n \|v_f^{n+1}\|^2 &\leq 2\|\sqrt{\operatorname{Re} A^{n+\alpha}} D_- v^{n+\alpha}\|^2 + 5\eta M_2^2 (3\|v^{n+\alpha}\|^2 + \|v^{n+1-\alpha}\|^2) + 5\eta \|F^{n+\alpha}\|^2 \\ &\leq 2\|\sqrt{\operatorname{Re} A^{n+\alpha}} D_- v^{n+\alpha}\|^2 + 40\eta M_2^2 (\|v^{n+1}\|^2 + \|v^n\|^2) + 5\eta \|F^{n+\alpha}\|^2 \end{aligned} \quad (2.19)$$

Substituting (2.19) into (2.12), we have

$$\frac{1}{k_n} (\|v^{n+1}\|^2 - \|v^n\|^2) \leq M_4 (\|v^{n+1}\|^2 + \|v^n\|^2) + M_5 \|F^{n+\alpha}\|^2,$$

where  $M_4 = 2M_3 + 40\eta M_2^2$  and  $M_5 = 1 + 5\eta$ . It is easy to see that when  $k_n \leq \frac{1}{2M_4}$ , then

$$\begin{aligned}\|v^{n+1}\| &\leq \frac{1 + k_n M_4}{1 - k_n M_4} \|v^n\|^2 + \frac{M_5 k_n}{1 - k_n M_4} \|F^{n+\alpha}\|^2 \\ &\leq (1 + 4k_n M_4) \|v^n\|^2 + 2M_5 k_n \|F^{n+\alpha}\|^2.\end{aligned}$$

According to Lemma 4.1, we obtain

$$\|v^{n+1}\|^2 \leq e^{4M_4 t^{n+1}} (\|u^0\|^2 + 2M_5 \sum_{l=0}^n \|F^{l+\alpha}\|^2 k_l) \quad (2.20)$$

and from (2.5) and  $0 < M_0 \leq |\phi| \leq M$ , we obtain

$$\|u^{n+1}\|^2 \leq \frac{M_1^2}{M_0^2} e^{4M_4 t^{n+1}} (\|u^0\|^2 + 2M_5 \sum_{l=0}^n \|F^{l+\alpha}\|^2 k_l) \quad (2.21)$$

which implies that the scheme (2.1) is stable. ■

### 3. A stable explicit scheme for the case $\operatorname{Re} A \equiv 0$ .

In this section, we assume  $\operatorname{Re} A \equiv 0$  and let  $a(x, t)$  denote the imaginary part of  $A$ . We see from the previous section that the scheme (2.1) with  $\alpha = 0$  is unstable. This is unfortunate because in many applications an explicit scheme is desirable because they tend to be easier to implement, especially in a vector or parallel computing environment. In [2], we construct a stable explicit scheme for the simple equation  $u_t = iu_{xx}$  by adding appropriately chosen artificial dissipative terms to (2.1). Here we consider an extension of this scheme for the more general equation (1.1).

We construct the following scheme :

$$\begin{aligned}\frac{u_j^{n+1} - u_j^n}{k_n} - ia_j^n \frac{u_{j+1}^n - 2u_j^n + u_{j-1}^n}{h^2} - \frac{(ia_j^n - |a_j^n|)}{4} (v_{j+1}^n - 2v_j^n + v_{j-1}^n) \\ - B_j^n \frac{u_{j+1}^n - u_{j-1}^n}{2h} - C_j^n u_j^n = f_j, \quad j = 1, \dots, J-1,\end{aligned} \quad (3.1)$$

where

$$v_j^n = \frac{1}{h^2} (u_{j+1}^n - 2u_j^n + u_{j-1}^n), \quad j = 1, \dots, J-1, \quad (3.2)$$

with

$$v_0 = v_J = 0. \quad (3.3)$$

We will prove that this scheme is conditionally stable. For convenience, we assume

1.

$$a_j > 0, \quad j = 0, 1, \dots, J. \quad (3.4)$$

2.

$$\max \left( |a|, \left| \frac{\partial a}{\partial x} \right|, \left| \frac{\partial^2 a}{\partial x^2} \right|, |B|, \left| \frac{\partial B}{\partial x} \right|, |C| \right) \leq M. \quad (3.5)$$

We shall need the following lemmas :

**Lemma 3.1.** (Discrete Sobolev inequality)

Given  $\epsilon > 0$ , there exists a constant  $c$  dependent on  $\epsilon$  and  $n$  such that

$$\|D^l u\|_{L_\infty} \leq \epsilon \|D^n u\|_{L_2} + c \|u\|_{L_2} \quad l < n, \quad (3.6)$$

$$\|D^l u\|_{L_2} \leq \epsilon \|D^n u\|_{L_2} + c \|u\|_{L_2} \quad l \leq n. \quad (3.7)$$

**Lemma 3.2.** (Estimate  $\|u_t^n\|$ ).

For any  $\epsilon > 0$  there exists a constant  $K$  dependent on  $\epsilon$ , such that the solution  $u_j^n$  of (3.1) satisfies :

$$\|u_t^{n+1}\|^2 \leq \sum_{j=1}^{J-1} (a_j^n)^2 |D^2 u_j^n|^2 + \epsilon \|D^2 u^n\|^2 + K(\epsilon) [\|u^n\|^2 + \|f^n\|^2]. \quad (3.8)$$

*Proof.* From (3.1), we have

$$\|u_t^{n+1}\|^2 = \sum_{j=1}^{J-1} \left\{ \left[ i a_j^n D^2 u_j^n + \frac{1}{4} (i-1) a_j^n D^2 v_j^n + G_j^n \right] \cdot \left[ -i a_j^n D^2 \bar{u}_j^n + \frac{1}{4} (-i-1) a_j^n D^2 \bar{v}_j^n + \bar{G}_j^n \right] \right\}$$

where

$$G_j = B_j D u_j + c_j u_j + f_j. \quad (3.9)$$

Expanding the above expression, we have

$$\begin{aligned} \|u_t^{n+1}\|^2 = & \sum_{j=1}^{J-1} \left\{ (a_j^n)^2 |D^2 u_j^n|^2 + \frac{1}{2} \operatorname{Re} [(1-i)(a_j^n)^2 h^2 D^2 u_j^n \cdot D^2 \bar{v}_j^n] + \frac{1}{8} (a_j^n)^2 h^4 |D^2 v_j^n|^2 \right\} \\ & + 2 \operatorname{Re} \sum_{j=1}^{J-1} \left[ i a_j^n D^2 u_j^n + \frac{1}{4} (i-1) a_j^n h^2 D^2 v_j^n \right] \bar{G}_j^n. \end{aligned}$$

Using (3.3), we obtain

$$\begin{aligned} & h^2 \sum_{j=1}^{J-1} (a_j^n)^2 D^2 u_j^n D^2 \bar{v}_j^n \\ &= \sum_{j=1}^{J-1} v_j^n [(a_{j+1}^n)^2 (\bar{v}_{j+1}^n - \bar{v}_j^n) - (a_j^n)^2 (\bar{v}_j^n - \bar{v}_{j-1}^n) - ((a_{j+1}^n)^2 - (a_j^n)^2) (\bar{v}_{j+1}^n - \bar{v}_j^n)] \\ &= \sum_{j=2}^{J-1} v_{j-1}^n a_j^n (\bar{v}_j^n - \bar{v}_{j-1}^n) - \sum_{j=1}^{J-1} v_j^n (a_j^n)^2 (\bar{v}_j^n - \bar{v}_{j-1}^n) - \sum_{j=1}^{J-1} v_j^n ((a_{j+1}^n)^2 - (a_j^n)^2) (\bar{v}_{j+1}^n - \bar{v}_j^n) \\ &= - \sum_{j=1}^J (a_j^n)^2 |v_j^n - v_{j-1}^n|^2 - \sum_{j=1}^{J-1} ((a_{j+1}^n)^2 - (a_j^n)^2) v_j^n (\bar{v}_{j+1}^n - \bar{v}_j^n). \end{aligned}$$

Hence

$$\begin{aligned}
& \frac{1}{2} \sum_{j=1}^{J-1} \operatorname{Re} [(1-i)(a_j^n)^2 h^2 D^2 u_j^n \cdot D^2 \bar{v}_j^n] + \frac{1}{8} \sum_{j=1}^{J-1} (a_j^n)^2 h^4 |D^2 v_j^n|^2 \\
& \leq -\frac{1}{2} \sum_{j=1}^J (a_j^n)^2 |v_j^n - v_{j-1}^n|^2 + \left| \sum_{j=1}^{J-1} ((a_{j+1}^n)^2 - (a_j^n)^2) v_j^n (\bar{v}_{j+1}^n - \bar{v}_j^n) \right| \\
& \quad + \frac{1}{4} \sum_{j=1}^{J-1} (a_j^n)^2 [|v_{j+1}^n - v_j^n|^2 + |v_j^n - v_{j-1}^n|^2] \\
& \leq \left| \sum_{j=1}^{J-1} ((a_{j+1}^n)^2 - (a_j^n)^2) v_j^n (\bar{v}_{j+1}^n - \bar{v}_j^n) \right| - \frac{1}{4} \sum_{j=1}^J ((a_j^n)^2 - a_{j-1}^n) |v_j^n - v_{j-1}^n|^2 \\
& \leq 2 \sum_{j=1}^{J-1} |(a_{j+1}^n)^2 - (a_j^n)^2| |v_j^n| |v_{j+1}^n - v_j^n|.
\end{aligned}$$

So

$$\begin{aligned}
\|u_t^{n+1}\|^2 & \leq \sum_{j=1}^{J-1} (a_j^n)^2 |D^2 u_j^n|^2 + 2 \sum_{j=1}^{J-1} |(a_{j+1}^n)^2 - (a_j^n)^2| |v_j^n| |v_{j+1}^n - v_j^n| \\
& \quad + 2 \operatorname{Re} \sum_{j=1}^{J-1} \left[ i a_j^n D^2 u_j^n + \frac{1}{4} (i-1) a_j^n h^2 D^2 v_j^n \right] \bar{G}_j^n.
\end{aligned}$$

For  $G_j$ , using Lemma 3.1, we have

$$\begin{aligned}
\|G^n\|^2 & \leq 3M^2 (\|Du^n\|^2 + \|u^n\|^2) + 3\|f^n\|^2 \\
& \leq 6M^2 \epsilon_1 \|D^2 u^n\|^2 + (6M^2 K(\epsilon_1) + 3M^2) \|u^n\|^2 + 3\|f^n\|^2,
\end{aligned}$$

where  $\epsilon_1 > 0$ . It follows that if we define

$$Q \equiv 2 \operatorname{Re} \sum_{j=1}^{J-1} \left[ i a_j^n D^2 u_j^n + \frac{1}{4} (i-1) a_j^n h^2 D^2 v_j^n \right] \bar{G}_j^n,$$

we have

$$\begin{aligned}
Q & \leq 6M \|D^2 u^n\| \|G^n\| \\
& \leq 3M \left[ \epsilon_2 \|D^2 u^n\|^2 + \frac{1}{\epsilon_2} \|G^n\|^2 \right] \\
& = \left[ 3M \epsilon_2 + \frac{18M^3 \epsilon_1}{\epsilon_2} \right] \|D^2 u^n\|^2 + \frac{1}{\epsilon_2} [18M^3 K(\epsilon_1) + 9M^3] \|u^n\|^2 + \frac{9M}{\epsilon_2} \|f^n\|^2,
\end{aligned}$$

where  $\epsilon_2$  is any positive constant. If we define  $\epsilon_1$  by

$$\epsilon_1 = \frac{\epsilon_2^2}{864M^4},$$

and  $\epsilon_2$  by

$$\epsilon_2 = \frac{\epsilon}{12M} \quad ,$$

then we have

$$Q \leq \frac{\epsilon}{2} \|D^2 u^n\|^2 + K_1(\epsilon) [\|u^n\|^2 + \|f^n\|^2] .$$

where  $K_1(\epsilon)$  is a constant dependent on  $\epsilon$ . On the other hand using Lemma 3.1,

$$\begin{aligned} 2 \sum_{j=1}^{J-1} |(a_{j+1}^n)^2 - (a_j^n)^2| |v_j^n| |v_{j+1}^n - v_j^n| &\leq 8M^2 h \|v^n\|^2 \\ &\leq 8M^2 h [\epsilon_3 \|D^2 u^n\|^2 + K_2(\epsilon_3) \|u^n\|^2] \\ &= 8M^2 h \epsilon_3 \|D^2 u^n\|^2 + 8M^2 h K_2(\epsilon_3) \|u^n\|^2 \quad , \end{aligned}$$

where  $\epsilon_3$  is an arbitrary positive constant. If we take

$$\epsilon_3 = \frac{\epsilon}{16M^2 h} \quad ,$$

then we have

$$2 \sum_{j=1}^{J-1} |(a_{j+1}^n)^2 - (a_j^n)^2| |v_j^n| |v_{j+1}^n - v_j^n| \leq \frac{\epsilon}{2} \|D^2 u^n\|^2 + K_3(\epsilon) \|u^n\|^2 \quad .$$

where  $K_3(\epsilon)$  is a constant dependent on  $\epsilon$ . Combining the above results, we finally have

$$\|u_i^{n+1}\|^2 \leq \sum_{j=1}^{J-1} (a_j^n)^2 |D^2 u_j^n|^2 + \epsilon \|D^2 u^n\|^2 + K_4(\epsilon) \|u^n\|^2 + K_1(\epsilon) \|f^n\|^2 ,$$

where

$$K_4(\epsilon) = K_1(\epsilon) + K_3(\epsilon) \quad ,$$

which completes the proof. ■

**Theorem 3.1.** *If (3.4) holds and*

$$\min |a(x)| > \beta > 0 \tag{3.10}$$

*then the scheme of (3.1) is stable if*

$$k_n < \frac{1}{2} \frac{h^2}{\max |a|} \quad . \tag{3.11}$$

*Proof.* Multiply (3.1) by  $\bar{u}_j^n$  and we obtain

$$(Lu^n, u^n) = (f^n, u^n). \tag{3.12}$$

Next we estimate each term in (3.12). For the first term we have

$$\operatorname{Re}(u_t^{n+1}, u^n) = \frac{1}{2k_n} (\|u^{n+1}\|^2 - \|u^n\|^2) - \frac{k_n}{2} \|u_t^{n+1}\|^2,$$

which is similar to (2.8). Applying Lemma 3.2, we obtain, for any  $\epsilon > 0$ ,

$$\operatorname{Re}(u_t^{n+1}, u^n) \geq \frac{1}{2k_n} (\|u^{n+1}\|^2 - \|u^n\|^2) - \frac{k_n}{2} \left[ \sum_{j=1}^{J-1} (a_j^n)^2 |D^2 u_j^n|^2 + \epsilon \|D^2 u^n\|^2 + k_4 \|u^n\|^2 + k_1 \|f^n\|^2 \right]. \quad (3.13)$$

The second term is given by

$$I_2 = \sum_{i=1}^{J-1} \left( -i a_j^n \frac{u_{j+1}^n - 2u_j^n + u_{j-1}^n}{h^2} \cdot \bar{u}_j^n \right).$$

We first employ the following expansion :

$$\begin{aligned} a_j^n \frac{u_{j+1}^n - 2u_j^n + u_{j-1}^n}{h^2} &= \frac{1}{h^2} \left[ a_{j+1/2}^n (u_{j+1}^n - u_j^n) - a_{j-1/2}^n (u_j^n - u_{j-1}^n) \right] \\ &\quad + \frac{1}{h^2} \left[ (a_j^n - a_{j+1/2}^n) (u_{j+1}^n - u_j^n) - (a_j^n - a_{j-1/2}^n) (u_j^n - u_{j-1}^n) \right]. \end{aligned}$$

Then, using the boundary conditions, we have :

$$\begin{aligned} &\sum_{j=1}^{J-1} \frac{1}{h^2} \left[ a_{j+1/2}^n (u_{j+1}^n - u_j^n) - a_{j-1/2}^n (u_j^n - u_{j-1}^n) \right] \bar{u}_j^n \\ &= \sum_{j=1}^{J-1} \frac{1}{h^2} a_{j+1/2}^n (u_{j+1}^n - u_j^n) \bar{u}_j^n - \sum_{j=0}^{J-2} a_{j+1/2}^n (u_{j+1}^n - u_j^n) \bar{u}_{j+1}^n \\ &= -\frac{1}{h^2} \sum_{j=0}^{J-1} a_{j+1/2}^n |u_{j+1}^n - u_j^n|^2, \end{aligned}$$

and

$$\begin{aligned} &\left| \sum_{j=1}^{J-1} \frac{1}{h^2} \left[ (a_j^n - a_{j+1/2}^n) (u_{j+1}^n - u_j^n) - (a_j^n - a_{j-1/2}^n) (u_j^n - u_{j-1}^n) \right] \bar{u}_j^n \right| \\ &\leq \frac{1}{2} \sum_{j=1}^{J-1} \left| \frac{(u_{j+1}^n - u_j^n)}{h} \right|^2 + \frac{1}{2} \sum_{j=1}^{J-1} \left| \frac{(a_j^n - a_{j+1/2}^n)}{h} \right|^2 |u_j^n|^2 + \frac{1}{2} \sum_{j=1}^{J-1} \left| \frac{(u_j^n - u_{j-1}^n)}{h} \right|^2 \\ &\quad + \frac{1}{2} \sum_{j=1}^{J-1} \left| \frac{(a_j^n - a_{j-1/2}^n)}{h} \right|^2 |u_j^n|^2 \\ &\leq 2[\epsilon \|D^2 u^n\|^2 + c_2 \|u^n\|^2] + M^2 \|u^n\|^2 \\ &\leq 2\epsilon \|D^2 u^n\|^2 + (2c_2 + M^2) \|u^n\|^2. \end{aligned}$$

Hence we have

$$|Re I_2| \leq 2\epsilon \|D^2 u^n\|^2 + (2c_2 + M^2) \|u^n\|^2.$$

The third term is given by

$$I_3 = \sum_{i=1}^{J-1} \left( -\frac{(ia_j^n - |a_j^n|)}{4} (v_{j+1}^n - 2v_j^n + v_{j-1}^n) \cdot \bar{u}_j^n \right).$$

Using the expression :

$$\begin{aligned} \frac{(1-i)}{4} a_j^n \frac{(v_{j+1}^n - 2v_j^n + v_{j-1}^n)}{h^2} &= \frac{(1-i)}{4} \frac{1}{h^2} [a_{j+1}^n v_{j+1}^n - 2a_j^n v_j^n + a_{j-1}^n v_{j-1}^n] \\ &\quad + \frac{(1-i)}{4} \frac{1}{h^2} [(a_j^n - a_{j+1}^n) v_{j+1}^n + (a_j^n - a_{j-1}^n) v_{j-1}^n], \end{aligned}$$

we have

$$\begin{aligned} h^2 \frac{(1-i)}{4} \sum_{j=1}^{J-1} \frac{1}{h^2} [a_{j+1}^n v_{j+1}^n - 2a_j^n v_j^n + a_{j-1}^n v_{j-1}^n] \bar{u}_j^n \\ = h^2 \frac{(1-i)}{4} \left[ \frac{1}{h^2} \sum_{j=1}^{J-1} a_j^n v_j^n (\bar{u}_{j+1}^n - \bar{u}_j^n + \bar{u}_{j-1}^n) \right] \\ = h^2 \frac{(1-i)}{4} \sum_{j=1}^{J-1} a_j^n |v_j^n|^2, \end{aligned}$$

and

$$\begin{aligned} \left| \frac{(1-i)}{4} h^2 \sum_{j=1}^{J-1} \frac{1}{h^2} [(a_j^n - a_{j+1}^n) v_{j+1}^n + (a_j^n - a_{j-1}^n) v_{j-1}^n] \bar{u}_j^n \right| \\ \leq \frac{h}{2} \sum_{j=1}^{J-1} \left| \frac{a_j^n - a_{j+1}^n}{h} \right| |v_{j+1}^n| |u_j^n| + \frac{h}{2} \sum_{j=1}^{J-1} \left| \frac{a_j^n - a_{j-1}^n}{h} \right| |v_{j-1}^n| |u_j^n| \\ \leq Mh[\epsilon \|v^n\|^2 + \frac{1}{\epsilon} \|u^n\|^2] \\ = h\epsilon M \|v^n\|^2 + \frac{Mh}{\epsilon} \|u^n\|^2. \end{aligned}$$

The rest of the terms are easily bounded :

$$\begin{aligned} \left| \sum_{j=1}^{J-1} B_j^n \frac{u_{j+1}^n - u_{j-1}^n}{2h} \cdot \bar{u}_j^n \right| &\leq M \|Du^n\| \|u^n\| \\ &\leq \frac{M}{2} [\epsilon \|D^2 u^n\|^2 + K_2(\epsilon) \|u^n\|^2], \\ |(cu, u)| &\leq M \|u\|^2 \end{aligned}$$

and

$$|(f, u)| \leq \frac{1}{2} \|f\|^2 + \frac{1}{2} \|u\|^2.$$

Summing all the above terms, we get

$$\begin{aligned} & \frac{1}{2k_n} (\|u^{n+1}\|^2 - \|u^n\|^2) \\ & \leq \sum_{j=1}^{J-1} \left( \frac{k_n}{2} (a_j^n)^2 - \frac{h^2}{4} a_j^n \right) |D^2 u_j^n|^2 + \left[ \frac{k_n}{2} \epsilon + 2\epsilon + h\epsilon M + \frac{M\epsilon}{2} \right] \|D^2 u^n\|^2 \\ & \quad + \left[ \frac{k_n}{2} K_4 + 2c_2 + M^2 + \frac{Mh}{\epsilon} + \frac{Mk_2}{2} + M + \frac{1}{2} \right] \|u^n\|^2 + \left[ \frac{k_n}{2} K_1 + \frac{1}{2} \right] \|f^n\|^2. \end{aligned}$$

Of course, when (3.11) holds, we have

$$\frac{k_n}{2} (a_j^n)^2 - \frac{h^2}{4} a_j^n \leq -\frac{\beta^2 \Delta}{2}.$$

where

$$\Delta = \frac{h^2}{2 \max |a|} - k_n > 0.$$

As long as  $\epsilon$  is small enough, we can make

$$\frac{k_n}{2} \epsilon + 2\epsilon + h\epsilon M + \frac{M\epsilon}{2} \leq \frac{\beta^2 \Delta}{2}.$$

For such an  $\epsilon$ , we have

$$\frac{1}{2k_n} (\|u^{n+1}\|^2 - \|u^n\|^2) \leq \widetilde{M}_1 \|u^n\|^2 + \widetilde{M}_2 \|f^n\|^2.$$

Here the constants  $\widetilde{M}_1, \widetilde{M}_2$  depend on  $\Delta$  and  $M$ . Using Lemma 2.1, we have the proof. ■

#### 4. Concluding Remarks

Using energy methods, we have established the stability properties of the two schemes considered in this paper. Since equation (1.1) includes both parabolic and Schrödinger type equations as special cases, our stability results provide a unified treatment for both types of equations. Finally, the results here agree with the stability results obtained for the constant coefficient Cauchy problem via Fourier analysis [3, 4].

**Acknowledgement** The authors would like to thank Prof. Martin Schultz of Yale University and Dr. Ding Lee of the Naval Underwater Systems Center, New London, CT., for their encouragement and helpful discussions throughout this work, and Mr. Herbert Beke for his help in the preparation of this manuscript.

## Appendix

## Appendix A (Proof of Lemma 2.1)

We substitute (2.5) into (2.1) and get, for the first term :

$$\begin{aligned}
 u_{i_j}^{n+1} &= (\phi v)_{i_j}^{n+1} \\
 &= \frac{1}{k} \left[ \phi_j^{n+\alpha} (v_j^{n+1} - v_j^n) + (\phi_j^{n+1} - \phi_j^{n+\alpha}) v_j^{n+1} + (\phi_j^{n+\alpha} - \phi_j^n) v_j^n \right] \\
 &= \frac{1}{k} \left[ \phi_j^{n+\alpha} (v_j^{n+1} - v_j^n) + (\phi_j^{n+1} - \phi_j^n) ((1+\alpha) v_j^{n+1} + \alpha v_j^n) \right] \\
 &= \phi_j^{n+\alpha} v_{i_j}^{n+1} + v_j^{n+1-\alpha} \phi_{i_j}^{n+1}.
 \end{aligned}$$

In the following, for simplicity, we shall drop the index " $n + \alpha$ ". For the second term using the expressions

$$\begin{aligned}
 \phi_{j+1} &= \phi_j + \tilde{D}\phi_j h + \frac{1}{2} D_+ D_- \phi_j h^2 \\
 \phi_{j-1} &= \phi_j - \tilde{D}\phi_j h + \frac{1}{2} D_+ D_- \phi_j h^2,
 \end{aligned}$$

we obtain :

$$\begin{aligned}
 D(ADu_j) &= D(AD(\phi v)_j) \\
 &= \frac{\phi_j}{h^2} [A_{j+1/2}(v_{j+1} - v_j) - A_{j-1/2}(v_j - v_{j-1})] \\
 &\quad + \frac{1}{2h^2} [A_{j+1/2}(\phi_{j+1} - \phi_{j-1})v_{j+1} - A_{j-1/2}(\phi_{j+1} - \phi_{j-1})v_{j-1}] \\
 &\quad + \frac{(\phi_{j+1} - 2\phi_j + \phi_{j-1})}{2h^2} (A_{j+1/2}v_{j+1} + A_{j-1/2}v_{j-1}) \\
 &= \frac{\phi_j}{h^2} [A_{j+1/2}(v_{j+1} - v_j) - A_{j-1/2}(v_j - v_{j-1})] + \frac{(\phi_{j+1} - \phi_{j-1})}{2h^2} A_j(v_{j+1} - v_{j-1}) \\
 &\quad + \frac{(\phi_{j+1} - \phi_{j-1})}{2h^2} [(A_{j+1/2} - A_j)v_{j+1} + (A_j - A_{j-1/2})v_{j-1}] \\
 &\quad + \frac{(\phi_{j+1} - 2\phi_j + \phi_{j-1})}{2h^2} (A_{j+1/2}v_{j+1} + A_{j-1/2}v_{j-1}) \\
 &= \phi_j DADv_j + 2A_j \tilde{D}\phi_j \tilde{D}v_j + \left[ \frac{A_{j+1/2} - A_j}{h} v_{j+1} + \frac{A_j - A_{j-1/2}}{h} v_{j-1} \right] \tilde{D}\phi_j \\
 &\quad + \frac{(A_{j+1/2}v_{j+1} + A_{j-1/2}v_{j-1})}{2} D^2\phi_j.
 \end{aligned}$$

For third term using the expressions

$$\phi_{j+1} = \phi_j + D_+\phi_j h, \quad \phi_{j-1} = \phi_j - D_-\phi_j h$$

we obtain

$$\begin{aligned}
 B_j \tilde{D}u_j &= \frac{B_j}{2h} (\phi_{j+1}v_{j+1} - \phi_{j-1}v_{j-1}) \\
 &= B_j \phi_j \tilde{D}v_j + \frac{B_j}{2} (D_+\phi_j v_{j+1} + D_-\phi_j v_{j-1}) .
 \end{aligned}$$

In the second and third term, the first order difference term for  $v$  is  $(2A_j \tilde{D}\phi_j + B_j \phi_j) \tilde{D}v_j$ . We can choose  $\phi_j$  to make this term vanish by setting

$$2A_j \tilde{D}\phi_j + B_j \phi_j = 0. \quad (1)$$

Since  $|A| \geq a_0 > 0$  and  $A, B$  are bounded, it is easy to see that we can choose  $\phi_j$  such that  $0 < M_0 \leq |\phi_j| \leq M_1$ , where  $M_0, M_1$  are constants. Moreover, from (1) and  $|\phi_j| \leq M_1$ , we have

$$|\tilde{D}\phi_j| \leq \text{constant} \quad \text{in } Q_T.$$

If we take the finite difference of (1), then because of the assumptions on  $A, B$ , we can obtain

$$|D_+ D_- \phi_j| \leq \text{constant} \quad \text{in } Q_T.$$

After some manipulation, we obtain (2.6), where

$$\begin{aligned} G_{j-1}^{n+\alpha} &= \frac{1}{\phi_j^{n+\alpha}} \left[ \tilde{D}\phi_j^{n+\alpha} \frac{(A_j^{n+\alpha} - A_{j-1/2}^{n+\alpha})}{h} + \frac{1}{2} D_+ D_- \phi_j^{n+\alpha} A_{j-1/2}^{n+\alpha} + \frac{b_j^{n+\alpha}}{2} \tilde{D}\phi_j^{n+\alpha} \right] \\ H_j^{n+\alpha} &= C_j^{n+\alpha} \\ K_{j+1}^{n+\alpha} &= \frac{1}{\phi_j^{n+\alpha}} \left[ \tilde{D}\phi_j^{n+\alpha} \frac{(A_{j+1/2}^{n+\alpha} - A_j^{n+\alpha})}{h} + \frac{1}{2} D_+ D_- \phi_j^{n+\alpha} A_{j+1/2}^{n+\alpha} + \frac{b_j^{n+\alpha}}{2} \tilde{D}\phi_j^{n+\alpha} \right] \\ Q_j^{n+\alpha} &= -\frac{\phi_j^{n+1}}{\phi_j^{n+\alpha}} \\ F_j^{n+\alpha} &= \frac{f_j^{n+\alpha}}{\phi_j^{n+\alpha}}. \end{aligned}$$

Because of the properties of  $A, B, C, F$  and  $\phi$ , the functions  $G, H, K, Q$  and  $F$  are all bounded.

## Appendix B

**Lemma 4.1.** (Duhamel's Principle).

Assume  $u^n \geq 0, v^n \geq 0$  for  $n = 1, 2, \dots$

If  $u^{n+1} \leq (1 + Mk_n)u^n + k_n v^n, \quad n = 0, 1, \dots,$

then

$$u^{n+1} \leq e^{Mt^{n+1}} \left( u^0 + \sum_{l=0}^n v^l k_l \right). \quad (2)$$

Here  $M$  is a positive constant and  $t^{n+1} = \sum_{l=0}^n k_l$ .

*Proof.*

We have

$$\begin{aligned}
 u^{n+1} &\leq (1 + Mk_n)u^n + k_nv^n \\
 &\leq (1 + Mk_n)[(1 + Mk_{n-1})u^{n-1} + k_{n-1}v^{n-1}] + k_nv^n \\
 &\leq \prod_{l=0}^n (1 + Mk_l)u^0 + \sum_{l=0}^{n-1} \prod_{s=l+1}^n (1 + Mk_s)k_lv^l + k_nv^n \\
 &\leq \prod_{l=0}^n (1 + Mk_l)u^0 + \sum_{l=0}^{n-1} \prod_{s=0}^n (1 + Mk_s)k_lv^l + k_nv^n \\
 &\leq \exp \left[ \sum_{l=0}^n \ln(1 + Mk_l) \right] \cdot \left[ u^0 + \sum_{l=0}^n v^l k_l \right].
 \end{aligned}$$

Since  $\ln(1 + Mk_l) \leq Mk_l$ , we obtain (2).

### Appendix C

**Lemma 4.2.** Assume  $m_i \geq 0$  ( $i = 1, \dots, n$ ) and  $k > 0$ . The necessary and sufficient condition for the polynomial in  $p_i$

$$\sum_{i=1}^n m_i p_i^2 - k \left( \sum_{i=1}^n p_i \right)^2 \quad (3)$$

to be nonnegative definite, is

$$k \leq \frac{1}{\sum_{i=1}^n \frac{1}{m_i}}. \quad (4)$$

*Proof.* Define  $p = (p_1, \dots, p_n)^T$ ,  $l = (1, 1, \dots, 1)^T$  and  $D = \text{diagonal}(m_1, m_2, \dots, m_n)$ . Then the polynomial in (3) can be rewritten as

$$p^T D p - k(p^T l)^2 = p^T (D - k l l^T) p$$

The condition for the matrix  $D - k l l^T$  to be nonnegative definite is  $1 - k l^T D^{-1} l \geq 0$  which is (4).

## References

- [1] K. Appert and J. Vaclavik, *Dynamics of Coupled Solitons*, Phys. Fluids, 20 (1977), pp. 1845.
- [2] T.F. Chan, Ding Lee, Longjun Shen, *A Stable Explicit Schemes for the Ocean Acoustic Wave Equation*, Technical Report YALEU/DC/TR-315, Dept. of Computer Science, Yale Univ., 1984. To appear in Comp. and Math Appl..
- [3] ———, *Stable Explicit Schemes for Equations of the Schrödinger Type*, Technical Report YALEU/DC/TR-305, Dept. of Computer Science, Yale Univ., 1984. To appear in Siam J. Numer. Anal..
- [4] T.F. Chan and Longjun Shen, *Difference Schemes for Equations of Schrödinger Type*, Technical Report YALEU/DC/TR-320, Dept. of Computer Science, Yale Univ., 1984.
- [5] L.M. Degtyarev and V.V. Krylov, *A method for the numerical solution of problems of the dynamics wave fields with singularities*, USSR Compt. Math. Math. Phys., 17/6 (1977), pp. 172.
- [6] M. Delfour, M. Fortin and G. Payre, *Finite Difference Solutions of a Nonlinear Schrodinger Equation*, J. Comp. Phys., 44 (1981), pp. 277-288.
- [7] D.F. Griffiths, A.R. Mitchell and J.U. Morris, *A Numerical Study of the Nonlinear Schrodinger Equation*, Technical Report Report NA/52, Univ. of Dundee, Dept. of Math., May 1982.
- [8] Boling Guo, *On the Problem of Numerical Calculation for a Class of Nonlinear Schrodinger Equations and Its Systems*, Math. Numer. Sinica, 3/3 (1981), pp. 211-223.
- [9] Yu.N. Karamzin, *Difference Schemes for Calculating the Three-Frequency Interactions of Quasi Monochromatic Waves Allowing for Diffraction*, USSR Comp. Math. Math. Phys., 15/2 (1975), pp. 142.
- [10] D. Lee and J.S. Papadakis, *Numerical Solutions of Underwater Acoustic Wave Propagation Problems*, Technical Report 5929, Naval Underwater Systems Center, New London, CT., February 1984.
- [11] G.J. Morales and Y.C. Lee, *Generation of Density Cavities and Localized Electric Fields in a Nonuniform Plasma*, Phys. Fluids, 20 (1977), pp. 1135.
- [12] Yulin Zhou, *Finite Difference Method of the Boundary Problems for the Systems of Generalized Schrodinger Type*, To appear.

**END**

**FILMED**

---

*1-86*

**DTIC**