

AD-A167 723

A LONG PERIOD RECORD OF BOTTOM CURRENTS ON THE TASMAN
ABYSSAL FLA(NCU) ROYAL AUSTRALIAN NAVY RESEARCH LAB
EDGECLIFF P J MULNEARN SEP 85 RANRL - IM-EXT-17/85


1/1

UNCLASSIFIED

F/G 8/3

NL

END
1985
6-18



12

UNCLASSIFIED

RANRL-TM-(EXT) 17/85

AR No: 003-440



DEPARTMENT OF DEFENCE
DEFENCE SCIENCE AND TECHNOLOGY ORGANISATION
WEAPONS SYSTEMS RESEARCH LABORATORY
RAN RESEARCH LABORATORY

AD-A167 723

RANRL TECHNICAL MEMORANDUM
(EXTERNAL) 17/85

A LONG PERIOD RECORD OF BOTTOM CURRENTS
ON THE TASMAN ABYSSAL PLAIN

P.J. Mulhearn

THE UNITED STATES NATIONAL
ARCHIVE INFORMATION SERVICE
IS AUTHORIZED TO
REPRODUCE AND SELL THIS REPORT

DTIC
ELECTE
MAY 01 1986
S E D
E

816 FILE COPY

COPY No. 4

Commonwealth of Australia
SEPT 1985

UNCLASSIFIED

This document has been approved
for public release and sales its
distribution is unlimited.

86 4 29 006

12

UNCLASSIFIED

(1)

AR No: 003-440

DEPARTMENT OF DEFENCE
DEFENCE SCIENCE AND TECHNOLOGY ORGANISATION
WEAPONS SYSTEMS RESEARCH LABORATORY
RAN RESEARCH LABORATORY

© Commonwealth of Australia 1985

RANRL TECHNICAL MEMORANDUM (EXTERNAL) No. 17/85

A LONG PERIOD RECORD OF BOTTOM CURRENTS
ON THE TASMAN ABYSSAL PLAIN

P.J. Mulhearn



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input checked="" type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

ABSTRACT

Results from a 99 day deployment of a recording current-meter 100 m above the bottom on the Tasman Abyssal Plain are presented. Strong southwards currents, up to 37.1 cm/s, were encountered. Kinetic energies are comparable with those found at abyssal depths under the Gulf Stream and the Kuroshio. Tidal currents are weak, of order 1 cm s⁻¹.

Technical memoranda are of a tentative nature, represent the views of the author(s), and do not necessarily carry the authority of the Laboratory.

POSTAL ADDRESS: The Superintendent, Maritime Systems Division, WSRL
RAN Research Laboratory,
PO Box 706, DARLINGHURST NSW 2010

JN0438
D23004

Introduction

To the author's knowledge no long-term current-meter measurements, obtained prior to those discussed here, have been reported from abyssal depths in the Tasman Basin. A times series, obtained concurrently with this one, of zonal bottom currents derived from measurements of ambient vertical electric field fluctuations is reported in Bindoff et al (1985). Laird and Ryan (1969) report current-meter measurements over only 70 minutes for a site at $30^{\circ}57.9'S$, $155^{\circ}00.8'E$ at 4572 m depth. Long term velocity measurements down the New south Wales continental slope to 2000 m depth were obtained during the Australian Coastal Experiment (Clarke and Thompson, 1984).

From 13 December 1983 to 21 March 1984 (99 days) an RCM-5 Aanderaa recording current-meter was deployed at $35^{\circ}54'S$, $151^{\circ}23'E$, 100 m above the bottom in 4850 m of water (see Fig.1). This meter was deployed as part of the Tasman Project of Sea Floor Magnetotelluric Exploration (Ferguson et al, 1985). The low frequency (sub-tidal) velocity fluctuations from the current-meter have been compared with other abyssal observations, and with the motion of the East Australian Current and an associated warm-core eddy in Mulhearn et al (1986). Figure 2 shows the time series of the zonal and meridional velocities, to which a Munk "tide-killer" filter have been applied. The present paper mainly considers fluctuations at tidal and higher frequencies. Bindoff et al (1985) concentrate on this frequency range in their discussion of results from vertical electric field measurements obtained farther east as another part of the Tasman Project of Sea Floor Magnetotelluric Exploration.





FIGURE 1. MAP SHOWING POSITION OF RECORDING CURRENT-METER (RCM)




FIGURE 2. VELOCITY TIME SERIES FROM RECORDING CURRENT-METER (TIDE-KILLER FILTER APPLIED).

Equipment and Analysis

The current meter was a standard RCM-5 Aanderaa. Every 15 minutes it recorded instantaneous direction and the average speed over that sampling period. The starting speed was 2 cm s^{-1} and for zero turns of the meter's rotor a speed of 1.5 cm s^{-1} was recorded. There were periods with flow less than 2 cm s^{-1} but such sections of the time series should not have a great effect on the overall statistics as they only represent 7% of records. Spectral and tidal analyses were restricted to the longest period (59 days) with currents above 2 cm s^{-1} . The meter was moored via an AMF Sea-Link Model 322 recoverable acoustic transponder to a number of cast iron weights, and had glass floats attached for buoyancy.

The record was analysed to produce standard statistics and histograms, power spectra and tidal components.

Results

Statistics from the whole record length are presented in Table I. The average current direction of 198° is within 9° of being parallel to the isobaths of the nearest part of the continental slope. The approximately southward mean direction is an indication of the influence of the East Australian Current system. The mean kinetic energy per unit mass is $8 \text{ cm}^2 / \text{sec}^2$, the fluctuation kinetic energy per unit mass is $60 \text{ cm}^2 / \text{sec}^2$ and their ratio is 7.5. These values are comparable with those under the Gulf Stream (Dickson, 1983) and the Kuroshio Extension (Schmitz 1984 a and b).

The current rose (Fig.3) shows that the strongest currents are predominately southwards and in this direction are usually over 5 cm s^{-1} . Away from the southwards direction speeds are most often less than 10 cm/sec . Detailed histograms show the median speed is 5 cm/sec , and the median direction is 183.6° . The speed histogram is bimodal with a second peak at 19 cm/sec .

Power spectra for the velocity components are shown in Fig. 4. Spectra were obtained in two separate runs. For the low frequency part they were obtained via 1024 point Fast Fourier Transforms on each component using hourly data. For the high frequency section ten spectra obtained via 512 point FFT's on 15 minute samples were averaged for each component. These spectra were obtained in the period after 11 January 1984 when currents were strong enough to keep the rotor turning.

Peaks at diurnal and semi-diurnal frequencies are prominent in the meridional component but are barely visible in the zonal component. There is some evidence of the presence of inertial oscillations ($f = .0489 \text{ cycles/hour}$) in the "shoulders" on the high frequency side of the diurnal peak. From approximately 0.1 to 0.7 cycles/hour the spectra fall off as $(\text{frequency})^{-2}$ and are approximately equal. These properties are as expected from the Garrett and Munk (1975, 1979) spectrum for internal waves.

The spectral levels in the internal wave frequency-band are typical of those found in deep moorings (e.g. Wunsch 1976). A direct comparison with the Garrett and Munk spectrum cannot be obtained with any accuracy because the buoyancy frequency, N , is not adequately known. From Nansen cast data, obtained when the current meter was deployed, a value for




FIGURE 3. CURRENT ROSE (NUMBERS ON AZIMUTHAL LINES ARE SPEED IN CM S^{-1})
 (DATA IS FROM 9507 FIFTEEN MINUTE AVERAGES).

between 0 and 0.9 cycles/hour was inferred. Taking the best linear fit through the in-situ density points gives $N = 0.6$ cycles/hour. It can be seen that this gives a curve within a factor of three of the data. Taking $N = N_0 e^{-z/b}$, as in Garrett and Munk (1979), with $N_0 = 3$ cycles/hour, $b = 1.3$ km and $z =$ depth (km) one obtains a curve approximately a factor of two below the data. However, this gives $N = .078$ cycles/hour, a 12.8 hour period! Considering the uncertainty in N and the possible errors from Doppler shifting in strong currents and from mooring motion, there is reasonable agreement between the data and the Garrett - Munk spectrum. Beyond 1 cycle/hour the data lie above the $(\text{frequency})^{-2}$ line and appear rather noisy.

Results of tidal analysis are shown in Table II. Where certain components, such as P_1 and K_1 , S_2 and K_2 , are too close together for them to be distinguished in this limited length of record only that associated with the larger tide generating force is presented. It can be seen that the M_2 component is the largest, followed in order of decreasing magnitude by K_1 , S_2 , O_1 , N_2 and Q_1 . However they all represent very small currents.

Acknowledgements

Thanks are extended to the officers and crews of HMAS COOK and RV SPRIGHTLY, the former for help in deploying and the latter for help in recovering the mooring. Mr Fred Boland of the Commonwealth Scientific Industrial and Research Organisation's Division of Oceanography is thanked for his advice on mooring designed and for having the current-meter results transcribed to computer compatible tape. Dr Jason Middleton is thanked for his advice on tidal analysis and the assistance of Mr Peter Tate and Mr Les Hamilton with data analysis is acknowledged.




FIGURE 4. POWER SPECTRA OF VELOCITY COMPONENTS: X, ZONAL COMPONENT; ●, MERIDIONAL COMPONENT. FULL LINE IS GARRETT - MUNK SPECTRUM WITH $N = 0.6$ CYCLES/HOUR. DASHED LINE IS GARRETT - MUNK SPECTRUM USING $N = N_0 e^{-z/b}$ WITH $N_0 = 3$ CYCLES/HOUR, $b = 1.3$ KM, AND Z IS DEPTH (KM).

References

- Bindoff N.L., J.H. Filloux, P.J. Mulhearn, F.E.M. Lilley and I.J. Ferguson (1985) Vertical electric field fluctuations at the floor of the Tasman Abyssal Plain. Deep-Sea Research (submitted)
- Clarke A.J. and R.O.R.Y. Thompson (1984) Large-scale wind-driven ocean response in the Australian Coastal Experiment region. Journal of Physical Oceanography, 14, 338-352.
- Dickson, R.R. (1983) Global Summaries and Intercomparisons: Flow Statistics from Long-Term Current Meter Moorings, in Eddies in Marine Science, ed. A.R. Robinson, Springer - Verlag, Berlin.
- Ferguson, I.J., J.H. Filloux, F.E.M. Lilley, N.L. Bindoff and P.J. Mulhearn (1985) A Seafloor Magnetotelluric Sounding in the Tasman Sea. Geophysical Research Letters (in press).
- Garrett, C. and W. Munk (1975) Space-time scales of internal waves: A progress report. Journal of Geophysical Research, 80, 291-297.
- Garrett, C. and W. Munk (1979) Internal waves in the ocean. Annual Review of Fluid Mechanics, 11, 339-369.
- Laird, M.P. and T.V. Ryan (1969) Bottom currents in the Tasman Sea. Journal of Geophysical Research, 74, 5433-5438.
- Mulhearn, P.J., J.H. Filloux, F.E.M. Lilley, N.L. Bindoff and I.J. Ferguson (1985) Abyssal currents during the formation and passage of an East Australian Current Warm Core Ring. Deep-Sea Research (submitted)
- Schmitz, W.J., Jr (1984a) Abyssal kinetic energy levels in the western North Pacific. Journal of physical Oceanography, 14, 198-201.
- Schmitz, W.J. Jr (1984b) Observations of the Vertical Structure of the Eddy Field on the Kuroshio Extension. Journal of Geophysical Research, 89, (C4), 6355-6364.
- Wunsch, C. (1976) Geographical Variability of the Internal Wave Field: A Search for Sources and Sinks. Journal of Physical Oceanography, 6, 471-485.

Table I

	u	v	Direction	Speed
Minimum	-16.9	-36.3	-	1.5
Maximum	25.6	11.6	-	37.1
Mean	- 0.7	- 3.9	197.6	9.5
Std. deviation	5.6	9.4	92.1	7.3
Skewness	1.3	- 1.3	- 0.2	1.5
Kurtosis	6.6	4.1	2.1	4.6

u is eastwards, v is northwards velocity component. u, v and speed are in cm s^{-1} ; direction is in degrees with 0° being true north

Table II

Component	Period (Hr)	ϵ ($^{\circ}$)	Major Axis (cm s^{-1})	Minor Axis (cm s^{-1})	Phase ($^{\circ}$)
Q ₁	26.868	6.88	.33	.09	- 57.15
O ₁	25.819	-68.18	.59	.01	107.41
K ₁	23.9	-75.53	1.10	-.11	142.17
N ₂	12.658	-13.76	.36	-.16	-144.28
M ₂	12.421	99.41	1.43	-.40	- 19.63
S ₂	12.000	-83.33	.66	-.28	175.46

ϵ is angle of major axis of tidal ellipse with respect to due east, measured anticlockwise.

Phase is lag of major axis current relative to tide generating force at Greenwich.

DISTRIBUTION LIST

	<u>Copy No.</u>
Chief Defence Scientist)	
Deputy Chief Defence Scientist)	1
Controller External Relations Projects and)	
Analytical Studies)	
Superintendent Science Programmes and Administration)	
Deputy Director, Scientific and Technical Intelligence, JIO	2
OIC, Document Exchange Centre, DISB, for National Library of Australia	3
US Defence Technical Information Center (DTIC)	4 - 15
UK Defence Research Information Centre (DRIC)	16 - 17
Director Scientific Information Services (Canada)	18
New Zealand Ministry of Defence	19
Microfiche copying then destruction	20
Librarian, Technical Reports Centre, Defence Central Library, Campbell Park	21
Librarian H Block, Victoria Barracks, Melbourne	22
Director RAN, Australian Joint Anti-Submarine School	23
Director RAN Tactical School HMAS WATSON	24
Naval Scientific Adviser	25
Air Force Scientific Adviser	26
DOM, HYDRO RAN	27
Senior Met. Officer, NAS Nowra	28
SOO HYDRO RAN	29
Librarian, CSIRO Marine Laboratories	30

Mr B.V. Hamon, Marine Studies Centre, Sydney University	31
Dr M. Tomczak, Marine Studies Centre, Sydney University	32
Principal Officer, Ocean Sciences Group RANRL	33
Author	34, 35
RANRL Master Copy	36
RANRL	37 - 44
Librarian, Defence Research Centre, Salisbury	45, 46
Librarian, Defence Signals Directorate	47

DOCUMENT CONTROL DATA

ADA 167723

1. a. AR No 003-440	1. Establishment No RANRL-TM-(EXT)-17/85	2. Document Date Sep 85	3. Task No 84/007
4. Title A Long Period Record of Bottom Currents on the Tasman Abyssal Plain.		5. Security a. document UNCLAS b. title c. abstract UNCLAS UNCLAS	6. No Pages 14 7. No Refs 11
4. Author(s) P.J. Mulhearn		9. Downgrading Instructions N/A	
10. Corporate Author and Address RAN Research Laboratory PO Box 706 DARLINGHURST NSW 2010		11. Authority (as appropriate) a. Sponsor b. Security c. Downgrading d. Approval a. Ocean Sciences Group b. PO/OSG c. UNCLASSIFIED d. DWSRL	
12. Secondary Distribution (of this document) Approved for Public Release			
Overseas enquiries outside stated limitations should be referred through ASDM, Defence Information Services Branch, Department of Defence, Campbell Park, CANBERRA ACT 2601			
13. a. This document may be ANNOUNCED in catalogues and awareness services available to ... No limitations			
13. b. Citation for other purposes (ie casual announcement) may be (select) unrestricted (or) as for 13 a.			
14. Descriptors Oceanography Current-meters Bottom currents Abyssal flow		15. COSATI Group 0803 0810	
16. Abstract Results from a 99 day deployment of a recording current-meter 100 m above the bottom on the Tasman Abyssal Plain are presented. Strong southwards currents, up to 37.1 cm/s, were encountered. Kinetic energies are comparable with those found at abyssal depths under the Gulf Stream and the Kuroshio. Tidal currents are weak, of order 1 cm s ⁻¹ .			

END

DATE
FILMED

6 - 86

DTI