

AD-A168 942

WEAK CONVERGENCE OF THE VARIATIONS ITERATED INTEGRALS
AND DOLEANS-DADE EX. (U) NORTH CAROLINA UNIV AT CHAPEL
HILL CENTER FOR STOCHASTIC PROC. F AVRAN MAR 86


1/1

UNCLASSIFIED

TR-135 AFOSR-TR-86-0327 F49620-85-C-0144 F/G 12/1

NL





MICROCOPY

101107

AD-A168 942


②

PORT DOCUMENTATION PAGE

1a. SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AND DATE UNCLASSIFIED JUN 06 1986		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for Public Release; Distribution Unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE UNCLASSIFIED		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR-86-0327	
4. PERFORMING ORGANIZATION REPORT NUMBER Technical Report No. 135		7a. NAME OF MONITORING ORGANIZATION AFOSR/NM	
6a. NAME OF PERFORMING ORGANIZATION University of North Carolina		7b. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448	
6b. ADDRESS (City, State and ZIP Code) Center for Stochastic Processes, Statistics Department, Phillips Hall 039-A, Chapel Hill, NC 27514		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620 85 C 0144	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR		10. SOURCE OF FUNDING NOS. PROGRAM ELEMENT NO. 6.1102F PROJECT NO. 2304 TASK NO. A5 WORK UNIT NO.	
8b. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC		11. TITLE (Include Security Classification) Weak convergence of the variations, iterated integrals, and Doleans-Dade exponentials of sequences of semimartingales"	
12. PERSONAL AUTHOR(S) Avram, F.		13a. TYPE OF REPORT Technical	
13b. TIME COVERED FROM 9/85 TO 8/86		14. DATE OF REPORT (Yr., Mo., Day) March 1986	
15. PAGE COUNT 6		16. SUPPLEMENTARY NOTATION	
17. COSATI CODES FIELD GROUP SUB GR. XXXXXXXXXXXXXXXXXXXX		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Keywords:	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) (n) If X is a sequence of semimartingales, converging to a semimartingale X , and such that $[X, X]^{(n)}$ converges to $[X, X]$, then all higher order variations and all the iterated integrals of $X^{(n)}$ converge jointly to the respective functionals of X .			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT ☐ DTIC USERS ☐		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL J. K. U. (202) 767-5027		22b. TELEPHONE NUMBER (Include Area Code) AFOSR/NM	
22c. OFFICE SYMBOL AFOSR/NM		UNCLASSIFIED	

CENTER FOR STOCHASTIC PROCESSES

Department of Statistics
University of North Carolina
Chapel Hill, North Carolina



WEAK CONVERGENCE OF THE VARIATIONS, ITERATED INTEGRALS,
AND DOLEANS-DADE EXPONENTIALS OF SEQUENCES OF SEMIMARTINGALES

by

Florin Avram

Technical Report No. 135

March 1986

Approved for public release;
distribution unlimited.

WEAK CONVERGENCE OF THE VARIATIONS, ITERATED INTEGRALS,
AND DOLEANS-DADE EXPONENTIALS OF SEQUENCES OF SEMIMARTINGALES

by

Florin Avram

University of North Carolina at Chapel Hill

Abstract

If $X^{(n)}$ is a sequence of semimartingales, converging to a semimartingale X , and such that $[X^{(n)}, X^{(n)}]$ converges to $[X, X]$, then all higher order variations and all the iterated integrals of $X^{(n)}$ converge jointly to the respective functionals of X .

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFOSR)
NOTES ON
THIS RESEARCH WAS SUPPORTED BY AFOSR AND IS
DISTRIBUTED UNDER THE TERMS OF AFOSR-12.
MATTHEW J. LAMAR
Chief, Technical Information Division

AMS 1980 Subject Classifications: Primary, 60F17; Secondary, 60H05.

Keywords and Phrases: Semimartingales, weak J_1 Skorohod topology, variations, multiple integrals, Doléans-Dade exponential.

This research supported by the Air Force Office of Scientific Research
Contract No. F49620 85C 0144.

$$I_k^{(n)}(X)_t = \sum_{1 \leq i_1 < \dots < i_k \leq [nt]} X_{i_1, n} \dots X_{i_k, n},$$

and

$$E(\lambda X)_t^{(n)} = \prod_{i=1}^{[nt]} (1 + \lambda X_{i, n}) = \sum_{k=0}^{[nt]} \lambda^k I_k^{(n)}(X)_t.$$

The problem of the convergence of these "moments", "symmetric statistics", and generating function of the symmetric statistics have been studied in [1], [3-5], [7], and [9].

C. From formula 41.1 of Meyer (1976), it follows that in the semimartingale context, just like in the discrete deterministic case, I_k , $k=1, \dots, m$ and V_k , $k=1, \dots, m$ can be represented as polynomials of n variables in one another (the Newton polynomials which relate sums of powers to the sums of products). Thus, the issue of the joint convergence of I_k , $k=1, \dots, m$, and that of the convergence of V_k , $k=1, \dots, m$, are equivalent.

D. $X \xrightarrow{w(J_1)} X$ does not imply in general $[X, X] \rightarrow [X, X]$, as the following deterministic example from Jacod (1983) shows:

$$X_t^{(n)} = \sum_{k=1}^{[n^2 t]} \frac{(-1)^k}{n} \text{ converges uniformly to 0, but } [X, X]_t^{(n)} = \sum_{k=1}^{[n^2 t]} \frac{1}{n^2} \rightarrow t.$$

E. However, the following result holds:

Theorem 1: The following three statements are equivalent.

$$(1.5) \quad (X, [X, X]) \xrightarrow[n \rightarrow \infty]{w(J_1)} (X, [X, X]),$$

$$(1.6) \quad (V_1^{(n)}(X), \dots, V_m^{(n)}(X)) \xrightarrow[n \rightarrow \infty]{w(J_1)} V_1(X), \dots, V_m(X), \quad \forall m \geq 2,$$

$$(1.7) \quad (I_1^{(n)}(X), \dots, I_m^{(n)}(X)) \xrightarrow[n \rightarrow \infty]{w(J_1)} I_1(X), \dots, I_m(X), \quad \forall m \geq 2.$$

They also imply:

$$(1.8) \quad E(\lambda X)_t^{(n)} \xrightarrow{w(J_1)} E(\lambda X), \quad \forall \lambda.$$

Corollary: If

$$(1.9) \quad \overset{(n)}{X} \xrightarrow{w(J_1)} X$$

and the condition of Jacod (1983) holds:

$$(1.10) \quad \lim_{b \rightarrow \infty} \sup_{n \rightarrow \infty} P\{\text{Var}(B^{h,n})_1 > b\} = 0$$

(where h is a truncation function and $(B^{h,n})_t$ is the previsible projection of the truncated semimartingale X), then (1.5), (1.6), (1.7) and (1.8) hold.

Proof: cf. Jacod (1983), Theorem 5.1.1, (1.9) and (1.10) imply (1.5).

2. Proofs

Introduce the following notation: For any real number x ,

$$x^{>a} := x \cdot 1_{\{|x| > a\}}$$

$$x^{\leq a} := x \cdot 1_{\{|x| \leq a\}}$$

We establish now the following:

Lemma 1: a) Suppose $\overset{(n)}{X}$ are semimartingales such that

$$(2.1) \quad \lim_{b \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} P\{[\overset{(n)}{X}, \overset{(n)}{X}]_1 > b\} = 0,$$

and let $f(x)$ be any real function such that $f(x) = o(x^2)$, as $x \rightarrow 0$. Then, for all ε ,

$$(2.2) \quad \lim_{a \rightarrow 0} \overline{\lim}_{n \rightarrow \infty} P\left\{\sum_{s \leq 1} |f(\Delta X_s^{\leq a})| \geq \varepsilon\right\} = 0.$$

b) If the assumptions of a) hold, $\overset{(n)}{X} \xrightarrow{w(J_1)} X$ and f is a continuous, vector valued function, then:

$$(2.3) \quad \sum_{s \leq t} f(\Delta \overset{(n)}{X}_s) \xrightarrow{w(J_1)} \sum_{s \leq t} f(\Delta X_s).$$

Proof: a) Note first that $\sum_{s \leq t} |f(\Delta X_s^{(n)})| < \infty$, since $\sum_{s \leq t} (\Delta X_s^{(n)})^2 < \infty$. Let now $g(a) = \sup_{|x| \leq a} |f(x)|/x^{-2}$. Then,

$$\begin{aligned} P\left\{\sum_{s \leq 1} |f(\Delta X_s^{(n)})| > \epsilon\right\} &\leq P\left\{\sum_{s \leq 1} (\Delta X_s^{(n)})^2 g(a) > \epsilon\right\} \\ &\leq P\left\{[X, X]_1^{(n)} > \epsilon/g(a)\right\}. \end{aligned}$$

Since $g(a) \rightarrow 0$, (2.2) follows from (2.1).

b) Let $U(X) = \{u > 0 : P\{|\Delta X_t| \neq u, \text{ for all } t\} = 0\}$. $U(X)$ is dense in $\mathbb{R}_+$. For any $a \in U(X)$, and f continuous, the functional

$$S_f^a(Z)_t = \sum_{s \leq t} f(\Delta Z_s^{>a})$$

is J_1 continuous a.s. (dist (X)). Thus, $X \xrightarrow{w(J_1)} X$ implies for $a \in U(X)$

$$S_f^a(X) \xrightarrow{w(J_1)} S_f^a(X).$$

Also,

$$S_f^a(X)_t \xrightarrow[a \rightarrow 0]{\text{a.s. } (J_1)} S_f(X)_t := \sum_{s \leq t} f(\Delta X_s).$$

The result follows now by (2.2) and Theorem 4.2 of Billingsley (1968).

Proof of Theorem 1:

By Lemma 1b, we have (1.5) $\Rightarrow$ (1.6), and in fact the same type of argument yields (1.5) $\Rightarrow$ (1.8), as follows: Assume for convenience $\lambda = 1$ and $1 \in U(X)$, let

$$f(x) = [\ell n(1+x) - x + \frac{x^2}{2}] \mathbf{1}_{\{|x| \leq 1\}},$$

and let $T: D_{[0,1]} \rightarrow D_{[0,1]}$ be defined by:

$$T(Z)_t := \prod_{s \leq t} \ell(\Delta Z_s^{>1}) = \prod_{s \leq t} (1 + \Delta Z_s^{>1}) \exp\{-\Delta Z_s^{>1} + \frac{1}{2}(\Delta Z_s^{>1})^2\}.$$

Since the Doléans-Dade exponential

$$E(X)_t = \exp\{X_t - \frac{1}{2}[X, X]_t + \sum_{s \leq t} f[\Delta X_s^{\leq 1}]\} \cdot T(X)_t,$$

it remains only to note that the functional:

$$X^a : D^{(2)}[0,1] \rightarrow D^{(4)}[0,1]$$

$$X(Z_1, Z_2) = (Z_1, Z_2, S_f^a(Z_1), T_{Z_1})$$

is continuous a.s., if both spaces are endowed with the respective J_1 topologies. Letting then $a \rightarrow 0$, as in the proof of Lemma 1, one gets:

$$((X_t^{(n)}, [X, X]_t^{(n)}, \sum_{s \leq t} f(\Delta X_s^{(n) \leq 1}), \prod_{s \leq t} \ell(\Delta X_s^{(n) > 1}))$$

$$\xrightarrow{w(J_1)} (X_t, [X, X]_t, \sum_{s \leq t} f(\Delta X_s^{\leq 1}), \prod_{s \leq t} \ell(\Delta X_s^{> 1})),$$

since $\ell_n(1+x) - x + \frac{x^2}{2} = o(x^2)$, and since (1.5) implies (2.1). Finally, applying the continuous functional

$$\rho : D^{(4)}[0,1] \rightarrow D[0,1],$$

$$\rho(Z_1, Z_2, Z_3, Z_4) = \exp[Z_1 - \frac{1}{2}Z_2 + Z_3] \cdot Z_4,$$

we get that

$$E(\lambda X^{(n)}) \xrightarrow{w(J_1)} E(\lambda X).$$

Since (1.6) is equivalent to (1.7) (by the use of the polynomial mapping), and (1.6) trivially implies (1.5), Theorem 1 is proved. $\square$

References

1. Avram, F., Taqqu, M.S.: Symmetric Polynomials of Random Variables Attracted to an Infinitely Divisible Law. To appear in Z. für Wahr.
2. Billingsley, P.: Convergence of probability measures. Wiley: New York, (1968).
3. Dynkin, E.B., Mandelbaum, A.: Symmetric statistics, Poisson point processes and multiple Wiener integrals. Ann. of Statistics 11, 739-745, (1983).
4. Denker, M., Grillenberger, Chr., Keller, G.: A note on invariance principles for von Mises' statistics. Metrika (to appear 1985).
5. Feinsilver, P.J.: Special functions, probability semigroups and Hamilton flows. Lect. Notes in Math., 696. Springer Verlag: New York (1978).
6. Jacod, J.: Théorèmes Limite Pour Les Processus. École d'Été de Probabilités de Saint-Flour XIII--1983. Lect. Notes in Math. 1117. Springer Verlag: New York (1983).
7. Mandelbaum, A., Taqqu, M.S.: Invariance principle for symmetric statistics. Ann. of Statistics 12, 483-496 (1984).
8. Meyer, P.-A.: Un cours sur les integrales stochastiques. Sem. Probl X. Lect. Notes Math. 511, 245-400. Springer Verlag: New York (1976).
9. Rubin, H., Vitale, R.A.: Asymptotic distribution of symmetric statistics. Ann. of Statistics 8, 163-170 (1980).

END

DTIC

7-86