

AD-A171 205

SLEPIAN'S INEQUALITY VIA THE CENTRAL LIMIT THEOREM(U)
STANFORD UNIV CA DEPT OF STATISTICS F W HUFFER
05 AUG 86 TR-378 N00014-86-K-0156

1/1

UNCLASSIFIED

F/G 12/1

NL





NATIONAL BUREAU OF STANDARDS-1963-A

AD-A171 205

12

SLEPIAN'S INEQUALITY VIA THE
CENTRAL LIMIT THEOREM

BY

FRED W. HUFFER

TECHNICAL REPORT NO. 378

AUGUST 5, 1986

PREPARED UNDER CONTRACT
N00014-86-K-0156 (NR-042-267)
FOR THE OFFICE OF NAVAL RESEARCH

Reproduction in Whole or in Part is Permitted
for any purpose of the United States Government

Approved for public release; distribution unlimited.


DEPARTMENT OF STATISTICS
STANFORD UNIVERSITY
STANFORD, CALIFORNIA



DTIC
ELECTE
AUG 25 1986
B

DTIC FILE COPY

86 6 25 098



SLEPIAN'S INEQUALITY VIA THE
CENTRAL LIMIT THEOREM

BY

FRED W. HUFFER

TECHNICAL REPORT NO. 378
AUGUST 5, 1986

Prepared Under Contract
N00014-86-K-0156 (NR-042-267)
For the Office of Naval Research

Herbert Solomon, Project Director


Reproduction in Whole or in Part is Permitted
for any purpose of the United States Government

Approved for public release; distribution unlimited.

DEPARTMENT OF STATISTICS
STANFORD UNIVERSITY
STANFORD, CALIFORNIA

DTIC
ELECTE
AUG 25 1986
S D
B

86 8 25 098



SLEPIAN'S INEQUALITY VIA THE
CENTRAL LIMIT THEOREM

By


Fred W. Huffer

We give a proof of Slepian's (1962) inequality which does not rely on Plackett's identity or geometric arguments. The proof uses a partial ordering of distributions which is preserved under convolutions and scale transformations. Slepian's inequality may be formulated in terms of such a partial ordering. The properties of this partial ordering allow us to obtain results for the multivariate normal distribution by using the central limit theorem.

Tchen (1980) also noted the preservation under convolution property and from this obtained Slepian's inequality in the bivariate case. Further information and references concerning partial orderings of probability distributions may be found in Eaton (1982) or Chapter 1 of Stoyan (1983).

Let F be a collection of bounded continuous real-valued functions defined on $\mathbb{R}^k$. Suppose that F is invariant under both translations and scaling, that is, for any $b \in \mathbb{R}^k$, $c \geq 0$ and $f \in F$, the function g defined by $g(x) = f(cx+b)$ also belongs to F . Define $X \ll Y$ if X and Y are random vectors satisfying $Ef(X) \leq Ef(Y)$ for all $f \in F$.

Our object will be to prove inequalities concerning multivariate normal distributions. Let Σ and Λ be any $k \times k$ covariance matrices. Let X^* and Y^* be k -dimensional normal random vectors with $EX^* = EY^* = 0$,



$\text{cov } X^* = \Sigma$, $\text{cov } Y^* = \Lambda$. We wish to determine if $X^* \ll Y^*$ in which case we also say that $\Sigma \ll \Lambda$. The following result can sometimes be used to make this determination.

Proposition 1: Let X^* and Y^* be as given above. Suppose the random vectors X and Y satisfy $EX = EY = 0$, $\text{cov } X = \Sigma$, $\text{cov } Y = \Lambda$. If $X \ll Y$, then $X^* \ll Y^*$.

In our application X and Y will be chosen to have simple discrete distributions so that the relationship $X \ll Y$ is easy to verify.

Proof: First note that

(a) If U, V, W are independent with $U \ll V$, then $U+W \ll V+W$.

This follows by conditioning on the value of W and using the translation invariance of F . Let $X_1, X_2, X_3, \dots$ be i.i.d. copies of X and $Y_1, Y_2, Y_3, \dots$ be i.i.d. copies of Y . Since $X \ll Y$, using (a) twice in succession gives $X_1 + X_2 \ll Y_1 + X_2 \ll Y_1 + Y_2$. By induction we obtain $X_1 + X_2 + \dots + X_n \ll Y_1 + Y_2 + \dots + Y_n$. By the scale invariance of F ,

(b) $U \ll V$ implies $cU \ll cV$ for all $c \geq 0$.

Thus $n^{-1/2}(X_1 + X_2 + \dots + X_n) \ll n^{-1/2}(Y_1 + Y_2 + \dots + Y_n)$. Now let $n \rightarrow \infty$ and use the Central Limit Theorem to obtain $X^* \ll Y^*$.

The next proposition is useful in extending the ordering $\ll$ to a broader class of covariance matrices.

Proposition 2: Suppose Λ_1 and Λ_2 are any $k \times k$ covariance matrices satisfying $\Lambda_1 \ll \Lambda_2$. Define $\Gamma = \Lambda_2 - \Lambda_1$. Choose $t > 0$. If Σ and $\Sigma + t\Gamma$ are both nonsingular covariance matrices, then $\Sigma \ll \Sigma + t\Gamma$.

Proof: Let ϕ_1, ϕ_2, ϕ_3 denote $k \times k$ covariance matrices. In terms of covariance matrices (a) and (b) become:

$$(c) \quad \phi_1 \ll \phi_2 \text{ implies } \phi_1 + \phi_3 \ll \phi_2 + \phi_3 .$$

$$(d) \quad \phi_1 \ll \phi_2 \text{ implies } c\phi_1 \ll c\phi_2 \text{ for all } c \geq 0 .$$

These are used implicitly in the following argument. Choose Δ small enough so that both $(\Sigma - \epsilon\Lambda_1)$ and $(\Sigma + t\Gamma - \epsilon\Lambda_1)$ are positive definite whenever $0 \leq \epsilon \leq \Delta$. By taking convex combinations of these matrices we find that $(\Sigma + s\Gamma - \epsilon\Lambda_1)$ is positive definite (and therefore a covariance matrix) when $0 \leq s \leq t$ and $0 \leq \epsilon \leq \Delta$. Now

$$\Sigma + s\Gamma = (\Sigma + s\Gamma - \epsilon\Lambda_1) + \epsilon\Lambda_1 \ll (\Sigma + s\Gamma - \epsilon\Lambda_1) + \epsilon\Lambda_2 = \Sigma + (s + \epsilon)\Gamma .$$

Here we have used $\Lambda_1 \ll \Lambda_2$. Thus $\Sigma + s\Gamma \ll \Sigma + (s + \epsilon)\Gamma$ for $0 \leq s \leq t$ and $0 \leq \epsilon \leq \Delta$. Since Δ does not depend on s , it is clear that $\Sigma \ll \Sigma + t\Gamma$ as desired.

For $x = (x_1, x_2, \dots, x_k)$ and $y = (y_1, y_2, \dots, y_k)$ define $x \vee y = (x_1 \vee y_1, x_2 \vee y_2, \dots, x_k \vee y_k)$ and $x \wedge y = (x_1 \wedge y_1, x_2 \wedge y_2, \dots, x_k \wedge y_k)$ where $\vee$ and $\wedge$ denote the maximum and minimum respectively. A function f defined on $\mathbb{R}^k$ is called L-superadditive if $f(x) + f(y) \leq f(x \vee y) + f(x \wedge y)$

for all x and y . This condition was introduced by Lorentz (1953) who also showed that when f has continuous second partial derivatives, f is L-superadditive if and only if

$$\frac{\partial^2 f(x)}{\partial x_i \partial x_j} \geq 0 \text{ for all } x \text{ and all } i \neq j .$$

See Marshall and Olkin (1979) for further information on L-superadditivity.

Proposition 3: Let F be the class of bounded, continuous, L-superadditive functions on $\mathbb{R}^k$. Suppose $\Sigma = (\sigma_{ij})$ and $\Pi = (\pi_{ij})$ are $k \times k$ non-singular covariance matrices. If $\sigma_{ii} = \pi_{ii}$ for all i and $\sigma_{ij} \leq \pi_{ij}$ for all $i \neq j$, then $\Sigma \ll \Pi$.

This result is very similar to Proposition 1 of Joag-dev, Perlman and Pitt (1983) and it easily implies Slepian's inequality as given in Slepian (1962). The argument needed to obtain Slepian's inequality is basically the same as that in Corollary 1 of Joag-dev, et al.

Proof: Let $\alpha, \beta, \theta, \phi$ be k -dimensional vectors defined by

$$\alpha_p = 1, \alpha_q = -1, \alpha_i = 0 \text{ for } i \neq p \text{ or } q ,$$

$$\beta_p = -1, \beta_q = 1, \beta_i = 0 \text{ for } i \neq p \text{ or } q ,$$

$$\theta_p = \theta_q = 1, \theta_i = 0 \text{ for } i \neq p \text{ or } q ,$$

$$\phi_p = \phi_q = -1, \phi_i = 0 \text{ for } i \neq p \text{ or } q ,$$

Define the random vectors X and Y by

$$P\{X=\alpha\} = P\{X=\beta\} = \frac{1}{2}, \quad P\{Y=\theta\} = P\{Y=\phi\} = \frac{1}{2}.$$

Since $\alpha \vee \beta = \theta$ and $\alpha \wedge \beta = \phi$, it is clear that $X \ll Y$. Now applying Proposition 1 leads to a corresponding ordering between normal random vectors with covariance matrices

$$\text{cov } X = S^{pq} \quad \text{and} \quad \text{cov } Y = T^{pq}$$

where the entries of S^{pq} and T^{pq} are given by $S_{pp}^{pq} = S_{qq}^{pq} = 1$, $S_{pq}^{pq} = S_{qp}^{pq} = -1$ and $S_{ij}^{pq} = 0$ otherwise, $T_{pp}^{pq} = T_{qq}^{pq} = T_{pq}^{pq} = T_{qp}^{pq} = 1$ and $T_{ij}^{pq} = 0$ otherwise. Since p and q are arbitrary, we have shown $S^{pq} \ll T^{pq}$ for all $p \neq q$. Now we can use (c) and (d) to deduce that

$\Lambda_1 \ll \Lambda_2$ where

$$\Lambda_1 = \sum_{\substack{p,q \\ p < q}} b_{pq} S^{pq}, \quad \Lambda_2 = \sum_{\substack{p,q \\ p < q}} b_{pq} T^{pq}$$

and b_{pq} are arbitrary nonnegative numbers. Choose $b_{ij} = (\pi_{ij} - \sigma_{ij})/2$ for all $i < j$. Now using Proposition 2 with $\Gamma = \Lambda_2 - \Lambda_1 = \Pi - \Sigma$ completes the proof.



Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution	
Availability Codes	
Dist	Avail and/or Special
A-1	

References

- Eaton, M.L. (1982). A review of selected topics in multivariate probability inequalities. Ann. Statistics, 10, 11-43.
- Joag-dev, K., Perlman, M.D., and Pitt, L.D. (1983). Association of normal random variables and Slepian's inequality. Ann. Probability, 11, 451-455.
- Lorentz, G.G. (1953). An inequality for rearrangements. Amer. Math. Monthly, 60, 176-179.
- Marshall, A.W., and Olkin, I. (1979). Inequalities: Theory of Majorization and Its Applications. Academic Press, New York.
- Slepian, D. (1962). The one-sided barrier problem for Gaussian Noise. Bell System Tech. J., 41, 463-501.
- Stoyan, D., and (edited with revisions by) Daley, D.J. (1983). Comparison Methods for Queues and Other Stochastic Models. John Wiley and Sons, New York.
- Tchen, A.H. (1980). Inequalities for distributions with given marginals. Ann. Probability, 8, 814-827.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 378	2. GOVT ACCESSION NO. AD-A171205	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Slepian's Inequality Via The Central Limit Theorem		5. TYPE OF REPORT & PERIOD COVERED TECHNICAL REPORT
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Fred W. Huffer		8. CONTRACT OR GRANT NUMBER(s) N00014-86-K-0156
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics Stanford University Stanford, CA 94305		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-042-267
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Statistics & Probability Program Code 1111.		12. REPORT DATE August 5, 1986
		13. NUMBER OF PAGES 7
14. MONITORING AGENCY NAME & ADDRESS (If different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Slepian's inequality, partial orderings of distributions, L-superadditive functions, multivariate normal distribution.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We define a partial ordering of distributions which is preserved under convolutions and scale transformations. Some properties of this partial ordering are developed and then used to give a new argument for Slepian's (1962) inequality.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

UNCLASSIFIED

7 SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

END

DTIC

9-86