

AD-A173 477


SPECIAL METHODS FOR PROBLEMS WHOSE OSCILLATORY SOLUTION 171  
IS DAMPED(U) NAVAL POSTGRADUATE SCHOOL MONTEREY CA  
B NETA SEP 86 NPS-53-86-812

UNCLASSIFIED

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

2

AD-A173 477

NPS-53-86-012

# NAVAL POSTGRADUATE SCHOOL

Monterey, California



DTIC  
ELECTE  
OCT 27 1986  
S D

DTIC FILE COPY

SPECIAL METHODS FOR PROBLEMS WHOSE  
OSCILLATORY SOLUTION IS DAMPED

by

Beny Neta

September 1986

Technical Report For Period

April 1986 - September 1986

Approved for public release; distribution unlimited

Prepared for: Naval Postgraduate School  
Monterey, CA 93943

NAVAL POSTGRADUATE SCHOOL  
MONTEREY CALIFORNIA 93943

R. C. AUSTIN  
Rear Admiral, U. S. Navy  
Superintendent

D. A. SCHRADY  
Provost


Reproduction of all or part of this report is authorized.


This report was prepared by:

  
BENY NETA  
Associate Professor of  
Mathematics

Reviewed by:

Released by:

  
HAROLD M. FREDRICKSEN, Chairman  
Department of Mathematics

  
KNEALE T. MARSHALL  
Dean of Information  
and Policy Sciences

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

| REPORT DOCUMENTATION PAGE                                                                                                                                                                                                                                                                                                                                                                                                                                                         |                                      | READ INSTRUCTIONS<br>BEFORE COMPLETING FORM                             |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------|-------------------------------------------------------------------------|
| 1. REPORT NUMBER<br>NPS-53-86-012                                                                                                                                                                                                                                                                                                                                                                                                                                                 | 2. GOVT ACCESSION NO.<br>AD-4173 477 | 3. RECIPIENT'S CATALOG NUMBER                                           |
| 4. TITLE (and Subtitle)<br>Special Methods for Problems Whose<br>Oscillatory Solution is Damped                                                                                                                                                                                                                                                                                                                                                                                   |                                      | 5. TYPE OF REPORT & PERIOD COVERED<br>Technical Report<br>04/86 - 09/86 |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |                                      | 6. PERFORMING ORG. REPORT NUMBER                                        |
| 7. AUTHOR(s)<br>Beny Neta                                                                                                                                                                                                                                                                                                                                                                                                                                                         |                                      | 8. CONTRACT OR GRANT NUMBER(s)                                          |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS<br>Naval Postgraduate School<br>Monterey, CA 93943                                                                                                                                                                                                                                                                                                                                                                                    |                                      | 10. PROGRAM ELEMENT, PROJECT, TASK<br>AREA & WORK UNIT NUMBERS          |
| 11. CONTROLLING OFFICE NAME AND ADDRESS                                                                                                                                                                                                                                                                                                                                                                                                                                           |                                      | 12. REPORT DATE<br>September 1986                                       |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)                                                                                                                                                                                                                                                                                                                                                                                                       |                                      | 13. NUMBER OF PAGES<br>19                                               |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |                                      | 15. SECURITY CLASS. (of this report)<br>Unclassified                    |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |                                      | 15a. DECLASSIFICATION/DOWNGRADING<br>SCHEDULE                           |
| 16. DISTRIBUTION STATEMENT (of this Report)<br><br>Approved for public release; distribution unlimited                                                                                                                                                                                                                                                                                                                                                                            |                                      |                                                                         |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)                                                                                                                                                                                                                                                                                                                                                                                        |                                      |                                                                         |
| 18. SUPPLEMENTARY NOTES                                                                                                                                                                                                                                                                                                                                                                                                                                                           |                                      |                                                                         |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number)<br>ordinary differential equations, exponentially-fitted methods, shallow water<br>equations, damped oscillations<br><i>exp (in x) data</i>                                                                                                                                                                                                                                                    |                                      |                                                                         |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number)<br>This paper introduces methods tailored especially for problems whose<br>solution behave like $e^{**}$ , where $x$ is complex. The shallow water<br>equations with topography admit such solution.<br>This paper complements the results of Pratt and others on exponential-<br>fitted methods and those of Gautschi, Neta, van der Houwen and others on<br>trigonometrically-fitted methods. |                                      |                                                                         |

DD FORM 1473  
1 JAN 73EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-LF-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

# ABSTRACT

This paper introduces methods tailored especially for problems whose solution behaves like  $e^{\lambda x}$ , where  $\lambda$  is complex. The shallow water equations with topography admit such solution.

This paper complements the results of Pratt and others on exponential-fitted methods and those of Gautschi, Neta, van der Houwen and others on trigonometrically-fitted methods.

## 1. Introduction

In this paper we consider linear multistep methods

$$\sum_{\lambda=0}^k a_{\lambda} y_{n+1-\lambda} = h \sum_{\lambda=0}^k b_{\lambda} f(x_{n+1-\lambda}, y_{n+1-\lambda}), \quad k \geq 1, \quad n \geq k-1 \quad (1)$$

for integrating the initial value problem

$$y'(x) = f(x, y(x)), \quad y(x_0) = y_0. \quad (2)$$

This linear multistep method is characterized by the polynomials

$$\rho(\zeta) = \sum_{\lambda=0}^k a_{\lambda} \zeta^{k-\lambda}, \quad \sigma(\zeta) = \sum_{\lambda=0}^k b_{\lambda} \zeta^{k-\lambda}. \quad (3)$$

The main assumption of this paper is that it is a priori known that the solution is approximately of the form

$$y(x) \sim c_0 + \sum_{j=1}^m c_j e^{i \lambda_j t} \quad (4)$$



|      |  |                      |  |
|------|--|----------------------|--|
| Dist |  | Avail and/or Special |  |
| A-1  |  |                      |  |

where  $\lambda_j = w_j + i\psi_j$ , and the frequencies  $w_j$  are in a given interval  $[w_\ell, w_u]$ .

The special case where  $\lambda_j = jw_0$  with  $w_0$  given was considered first by Gautschi [8]. His approach was the following. Let:

$$\phi(z) = \rho(e^z) - z\sigma(e^z) \quad (5)$$

then the local truncation error of (1) is given by Lambert [11]

$$T_{n+k} = \phi(h \frac{d}{dt}) y(t_n) . \quad (6)$$

Inserting (4) in (6) yields

$$T_{n+k} \sim \phi(0)c_0 + \sum_{j=1}^m c_j \phi(ih\lambda_j) e^{i\lambda_j t} , \quad \lambda_j = jw_0 . \quad (7)$$

The coefficients  $b_i$  are chosen in such a way that

$$\phi(ihw_0) = 0 , \quad j = 0, 1, \dots, q , \quad (8)$$

for the largest value of  $q$  possible.  $q$  is then called the trigonometric order of the method. Gautschi has chosen  $a_i$  such that the methods are of Adams and Stormer type. However, these methods are sensitive to changes in the frequency  $w_0$ . Neta and Ford [13] developed Nystrom and generalized Milne-Simpson type methods. These methods showed less sensitivity to perturbation in  $w_0$  but require the eigenvalues of the Jacobian to be purely imaginary. Neta [14] has developed families of backward differentiation methods that overcome the above-mentioned restriction. Salzer [17] has developed

predictor-corrector methods based on trigonometric polynomials. See also Steifel and Bettis [18] and Bettis [3]. Van der Houwen and Sommeijer [10] have developed an alternative approach. The conditions (8) were replaced by

$$\phi(0) = 0 , \tag{9}$$

$$\phi(ih\lambda^{(j)}) = 0 , \quad j = 1, 2, \dots, q ,$$

where the  $\lambda^{(j)}$  are appropriately chosen points in the interval  $[w_l, w_u]$ .

An advantage of this so-called minimax approach over the fitting approach is the increased accuracy in cases where no accurate estimate of  $w_0$  is available or when the frequency is varying in time.

The other special case considered in the literature is where  $\lambda_j = i\omega_j$ . Probably the first article on the subject is due to Brock and Murray [5]. They discuss the use of exponential sums in the integration of a system of first order ordinary differential equations. Dennis [7] also suggested special methods for problems whose solution is exponential. He suggested a transformation of variables. More recently, Carroll [6] has developed exponentially fitted one-step methods for the scalar Riccati equation. For the general first order system of equations, Pratt [16] suggests methods based on the three parameter exponential function

$$I(x) = A + Be^{zx} . \tag{10}$$

The parameters A, B are given in terms of values of  $y$  and  $f$ . Several possibilities for  $z$  are given based on results of Brandon [2] and Babcock et al. [1].

Lyche [12] analyzes multistep methods which exactly integrate the set  $\{x^m e^{w_n x}\}$ , where  $w_n$  is real or imaginary.

In this article we developed various methods fitting exponentials and methods obtained via the minimax approach.

## 2. Construction of Methods

### 2.1 Fitting Methods

In this subsection we discuss various fitting methods. To this end, we separate  $\phi(ihj\lambda) = 0$ ,  $j = 1, 2, \dots, q$ , into real and imaginary parts. This yields the following equations relating the coefficients  $a_i, b_i$ ,

$$\begin{aligned} \sum_{i=0}^k a_i e^{j\psi(k-i)} \cos jv(k-i) - \sum_{i=0}^k b_i e^{j\psi(k-i)} [jv \cos jv(k-i) \\ - jv \sin jv(k-i)] = 0, \\ \sum_{i=0}^k a_i e^{j\psi(k-i)} \sin jv(k-i) - \sum_{i=0}^k b_i e^{j\psi(k-i)} [jv \sin jv(k-i) \\ + jv \cos jv(k-i)] = 0, \end{aligned} \quad (11)$$

where  $\lambda = w + i\psi$ ,  $u = -h\psi$ ,  $v = hw$ ,  $j = 1, 2, \dots, q$ .

For explicit methods,  $b_0 = 0$ . For Adams type methods  $a_0 = 1$ ,  $a_1 = -1$ ,  $a_i = 0$  for  $i = 2, \dots, k$ . For Nystrom or generalized Milne-Simpson methods  $a_0 = 1$ ,  $a_2 = -1$  and other  $a_i = 0$ .

k = 1 Implicit

Adams

$$\begin{aligned} b_0 &= \frac{ve^{-u} + u \sin v - v \cos v}{(u^2 + v^2) \sin v}, \\ b_1 &= \frac{ve^u - u \sin v - v \cos v}{(u^2 + v^2) \sin v}. \end{aligned} \quad (12)$$

For  $\psi = 0$ , the coefficients become  $b_0 = \frac{1 - \cos v}{\sin v} = b_1$ , which agree with Gautschi [8] if the coefficients are expanded in Taylor series with respect to  $v$ .

k = 2 Explicit

Adams

$$\begin{aligned} b_1 &= \frac{(u \sin 2v - v \cos 2v)e^u + v \cos v - u \sin v}{(u^2 + v^2) \sin v}, \\ b_2 &= \frac{(v \cos v - u \sin v)e^{2u} - ve^u}{(u^2 + v^2) \sin v}. \end{aligned} \quad (13)$$

Nystrom

$$\begin{aligned} b_1 &= \frac{(u \sin 2v - v \cos 2v)e^u + ve^{-u}}{(u^2 + v^2) \sin v}, \\ b_2 &= \frac{(v \cos v - u \sin v)e^{2u} + ve^{-u}}{(u^2 + v^2) \sin v}. \end{aligned} \quad (14)$$

For  $\psi = 0$ , the coefficients become  $b_1 = 2 \frac{\sin v}{v}$ ,  $b_2 = 0$  which agree with Neta [14].

### k = 2 Implicit

In this case, one obtains a one-parameter family of (Adams, generalized Milne-Simpson) methods of trigonometric order 1. The free parameter can be used to increase the algebraic order of the method as in [13].

### Backward Differentiation

$$e^{2u} \cos 2v + a_1 e^u \cos v + a_2 - b_0 e^{2u} (u \cos 2v - v \sin 2v) = 0 ,$$

$$e^{2u} \sin 2v + a_1 e^u \sin v - b_0 e^{2u} (u \sin 2v + v \cos 2v) = 0 , \quad (15)$$

$$1 + a_1 + a_2 = 0 .$$

This system can be solved by MACSYMA (Project MAC's SYmbolic MANipulation system written in LISP and used for performing symbolic as well as numerical mathematical manipulation [4]) or by REDUCE [9]. The solution is

$$a_1 = \frac{ue^{2u} - u \sin 2v - v \cos 2v}{-e^u (v \cos v + u \sin v) + u \sin 2v + v \cos 2v} , \quad (16)$$

$$a_2 = -1 - a_1 ,$$

$$b_0 = \frac{-e^{2u} \sin v + e^u \sin 2v - \sin v}{-e^{2u} (v \cos v + u \sin v) + e^u (u \sin 2v + v \cos 2v)} .$$

For  $\psi = 0$ , the coefficients agree with those given in [14].

### k = 3 Explicit

Again here, one obtains a one-parameter family of methods of trigonometric order 1. In order to get methods of trigonometric order 2, one has to construct a 3 step implicit method of Adams or generalized Milne-Simpson type. In order to increase the trigonometric order without going to a higher step number, one can construct linear multistep methods for which the coefficients  $a_\ell$  are also functions of  $\lambda$ ,  $w$ . Some examples are given in the next subsection.

### 2.2 Generalized Fitting Methods

In this section, we construct some linear multistep methods of the form

$$\sum_{\ell=0}^k a_\ell(\lambda) y_{n+1-\ell} = h \sum_{\ell=0}^k b_\ell(\lambda) f_{n+1-\ell}, \quad k \geq 1, n \geq k-1. \quad (17)$$

Since  $a_\ell$  are functions of  $\lambda$  one has more free parameters for his disposal which can be used to obtain higher trigonometric order methods with relatively lower step number.

### k = 2 Implicit

In this case, one has to solve the following linear system of five equations for the parameters  $a_1, a_2, b_0, b_1, b_2$  to obtain a method of trigonometric order 2.

$$\begin{aligned}
& 1 + a_1 + a_2 = 0, \\
& e^{2u} \cos 2v + a_1 e^u \cos v + a_2 - b_0 e^{2u} (u \cos 2v - v \sin 2v) \\
& \quad - b_1 e^u (u \cos v - v \sin v) - u b_2 = 0, \\
& e^{2u} \sin 2v + a_1 e^u \sin v - b_0 e^{2u} (u \sin 2v + v \cos 2v) \\
& \quad - b_1 e^u (u \sin v + v \cos v) - v b_2 = 0, \quad (18) \\
& e^{4u} \cos 4v + a_1 e^{2u} \cos 2v + a_2 - b_0 e^{4u} (2u \cos 4v - 2v \sin 4v) \\
& \quad - b_1 e^{2u} (2u \cos 2v - 2v \sin 2v) - 2u b_2 = 0, \\
& e^{4u} \sin 4v + a_1 e^{2u} \sin 2v - b_0 e^{4u} (2u \sin 4v + 2v \cos 4v) \\
& \quad - b_1 e^{2u} (2u \sin 2v + 2v \cos 2v) - 2v b_2 = 0.
\end{aligned}$$

The system was solved by REDUCE [9]. The expressions for the coefficients are complicated but REDUCE produces an output in the form of Fortran statements that can be incorporated in a computer program for numerical experiments with such a method.

### 2.3 Minimax Methods

In this section we discuss minimax methods, i.e., methods obtained by satisfying conditions (9). These conditions can be written in terms of  $a_k, b_k$  as follows:

$$\begin{aligned}
& \sum_{k=0}^K a_k e^{(k-j)u(j)} \cos(k-j)v(j) \\
& = \sum_{k=0}^K b_k e^{(k-j)u(j)} \{-v(j) \sin(k-j)v(j) + u(j) \cos(k-j)v(j)\}, \quad (19) \\
& j = 1, 2, \dots, q,
\end{aligned}$$

$$\sum_{\ell=0}^k a_{\ell} e^{(k-\ell)\mu^{(j)}} \sin(k-\ell)v^{(j)}$$

$$= \sum_{\ell=0}^k b_{\ell} e^{(k-\ell)\mu^{(j)}} \{v^{(j)} \cos(k-\ell)v^{(j)} + \mu^{(j)} \sin(k-\ell)v^{(j)}\}, \quad (20)$$

$$j = 1, 2, \dots, q,$$

where  $\mu^{(j)} = h\psi^{(j)}$ ,  $v^{(j)} = hw^{(j)}$ .

$ih\lambda^{(j)}$  are the zeros of the function  $\phi(ih\lambda)$  such that it has a small maximum norm in the rectangle  $w_{\ell} \leq w \leq w_u$ ,  $\psi_{\ell} \leq \psi \leq \psi_u$ .

To obtain the best approximation in this case is certainly not easy; but we will assume that one can write  $\phi(ih\lambda^{(j)}) = \phi(\psi^{(j)}, w^{(j)})$  as a product of 2 one variable functions. Thus  $\psi^{(j)}$  and  $w^{(j)}$  can be taken as Chebyshev's points on the corresponding interval, i.e.

$$\psi^{(j)} = \frac{1}{2}(\psi_{\ell} + \psi_u) + \frac{1}{2}(\psi_u - \psi_{\ell}) \cos \frac{2j-1}{2q} \pi, \quad (21)$$

$$w^{(j)} = \frac{1}{2}(w_{\ell} + w_u) + \frac{1}{2}(w_u - w_{\ell}) \cos \frac{2j-1}{2q} \pi,$$

$$j = 1, 2, \dots, q.$$

For this choice of points, one can evaluate the coefficients  $a_{\ell}$ ,  $b_{\ell}$  by solving

$$\phi(0) = 0, \quad (22)$$

$$\phi(\psi^{(j)}, w^{(j)}) = 0, \quad j = 1, 2, \dots, q.$$

We call such methods product minimax ( $PM^2$ ).

The number of free parameters for implicit methods  $2k+1$  and the number of equations is  $2q+1$ , thus, the trigonometric order  $q$  is equal to the step number  $k$ .

$k = 1$  Implicit

In this case

$$\begin{aligned}\psi^{(1)} &= \frac{1}{2}(\psi_\ell + \psi_u) \\ w^{(1)} &= \frac{1}{2}(w_\ell + w_u) \\ a_0 &= 1 = -a_1\end{aligned}\tag{23}$$

and the system of equations can be solved for  $b_0, b_1$ . This yields the coefficients given by (12) where

$$u = -h\psi^{(1)}, \quad v = hw^{(1)}.\tag{24}$$

Thus, the product minimax method would suggest using the center of the rectangle  $[\psi_\ell, \psi_u] \times [w_\ell, w_u]$  as  $\lambda_0$ . To obtain a product minimax method of trigonometric order 2, one has to solve a system of 5 equations similar to equation (18) with the unknowns  $b_0, b_1, b_2, a_1, a_2$ . The difference is that in the last 2 equations one should replace  $2\mu$  by  $\mu^{(2)}$  and  $2v$  by  $v^{(2)}$ . In the second and third equations of (18), the  $\mu, v$ , should be replaced by  $\mu^{(1)}, v^{(1)}$  respectively. The resulting system can be solved by MACSYMA [4] or REDUCE [9].

In the next section we implement two methods of trigonometric order 1 and 2 and see how the product minimax methods compare with fitting methods.

### 3. Numerical Example

Both systems (18) and (19)-(21) were solved by REDUCE which produced a FORTRAN subroutine for the evaluation of the coefficients. This subroutine is called only once during the integration.

The methods were compared for the solution of the initial value problem

$$\begin{aligned}\dot{z} - \frac{\pi}{2} i(1+i)z &= 0, \quad 0 \leq t \leq 4, \\ z(0) &= 1\end{aligned}\tag{25}$$

whose exact solution

$$z(t) = e^{\frac{i\pi}{2}(1+i)t},\tag{26}$$

thus

$$\lambda = \frac{\pi}{2}(1+i), \quad \psi = -\frac{\pi}{2}, \quad w = \frac{\pi}{2}.\tag{27}$$

In order to avoid complex arithmetic, we rewrite the differential equation as a system of equations for the real and imaginary part of  $z = u + iv$ .

$$\begin{aligned}\dot{u} + \frac{\pi}{2}(u+v) &= 0, \\ \dot{v} - \frac{\pi}{2}(u-v) &= 0, \\ u(0) &= 1, \\ v(0) &= 0.\end{aligned}\tag{28}$$

The system is solved by fitting methods of trigonometric order 1 and 2 with  $h = .01$  and various values of  $\psi$  and  $w$ . In Table 1 we list the Euclidean norm of the error at  $t = 4$ . It is clear that the method is not sensitive to perturbations in the values of  $\psi$  and  $w$ .

| $\Delta\psi$ | $\Delta w$ | error       |              |
|--------------|------------|-------------|--------------|
|              |            | first order | second order |
| 0            | 0          | .3678(-12)  | .4433(-12)   |
| 0            | .1         | .4482(-7)   | .3508(-11)   |
| .1           | 0          | .4346(-7)   | .2544(-11)   |
| .1           | .1         | .6342(-7)   | .4421(-11)   |
| 0            | .2         | .9241(-7)   | .8126(-11)   |
| .2           | 0          | .8706(-7)   | .5095(-11)   |

Table 1

Using the product minimax methods of trigonometric order 1 and 2 with  $h = .01$  and various squares centered at  $\psi = -\pi/2$ ,  $w = \pi/2$ , the error is much larger but again is insensitive to small perturbations in the length of the sides of the squares. In Table 2 we list the Euclidean norm of the error at  $t = 4$ .

| length of side | error       |              |
|----------------|-------------|--------------|
|                | first order | second order |
| .4             | .3678(-12)  | .1469(-6)    |
| .8             | .3678(-12)  | .2348(-5)    |
| 1.2            | .3678(-12)  | .1186(-4)    |
| 1.6            | .3678(-12)  | .3728(-4)    |
| 2              | .3678(-12)  | .9001(-4)    |

Table 2

Note that the perturbations in the product minimax methods are larger than those allowed in the fitting methods. It is possible that the larger errors in the product minimax methods are due to the assumption that  $\phi$  can be written as a product of 2 one variable functions. Also note that for the first order method, one always gets a good result since  $PM^2$  always uses the center of the square.

#### Acknowledgment

The author would like to thank the NPS Foundation Research Program for its support of this research.

## REFERENCES

1. Babcock, P. D., Stutzman, L. F., Brandon, D. M., Improvements in a single-step integration algorithm, *Simulation* 28, 1 (1979), 1-10.
2. Brandon, D. M., A new single-step integration algorithm with A-stability and improved accuracy, *Simulation* 23, 1 (1974), 17.
3. Bettis, D. G., Numerical integration of products of Fourier and ordinary polynomials, *Numer. Math.*, 14 (1970), 421-434.
4. Bogen, R. et al., *MACSYMA Reference Manual*, MIT Press, Cambridge, MA, 1975.
5. Brock, P., Murray, F. J., The use of exponential sums in step by step integration, *Math. Tables Aids Comput.* 6 (1952), 63-78.
6. Carroll, J., Exponentially fitted one-step methods for the numerical solution of the scalar Riccati Equation, *IMA J. Numer. Anal.*, to appear.
7. Dennis, S. C. R., The numerical integration of ordinary differential equations possessing exponential type solutions, *Proc. Cambridge Phil. Soc.* 65 (1960), 240-246.
8. Gautschi, W., Numerical integration of ordinary differential equations based on trigonometric polynomials, *Numer. Math.*, 3 (1961), 381-397.
9. Hearn, A. C. (ed.), *REDUCE user's manual*, Rand Corporation, Santa Monica, CA, 1983.
10. van der Houwen, P. J., Sommeijer, B. P., Linear multistep methods with minimized truncation error for periodic initial value problem, *IMA J. Numer. Anal.*, 4 (1984), 479-489.
11. Lambert, J. D., *Computational Methods in Ordinary Differential Equations*, Wiley, London, 1977.
12. Lyche, T., Chebyshevian multistep methods for ordinary differential equations, *Numer. Math.*, 19 (1972), 65-75.
13. Neta, B., Ford, C. H., Families of methods for ordinary differential equations based on trigonometric polynomials, *J. Comp. Appl. Math.*, 10 (1984), 33-38.
14. Neta, B., Families of backward differentiation methods based on trigonometric polynomials, *Intern. J. Computer Math.*, to appear 1986.

15. Neta, B., Williams, R. T., Hinsman, D. E., Studies in a shallow water fluid model with topography, Proc. Eleventh IMACS World Congress, Oslo, Norway, August 5-9, 1985 (R. Vichnevetsky, ed.).
16. Pratt, D. T., Exponential-fitted methods for solving stiff systems of ordinary differential equations, Proc. Eleventh IMACS World Congress, Oslo, Norway, August 5-9, 1985 (R. Vichnevetsky, ed.).
17. Salzer, H. E., Trigonometric interpolation and predictor-corrector formulas for numerical integration, ZAMM, 42 (1962), 403-412.
18. Steifel, E., Bettis, D. G., Stabilization of Cowell's method, Numer. Math., 13 (1969), 154-175.

## APPENDIX

Here we show that the shallow water equations with topography have a solution of the form  $e^{\lambda x}$ , where  $\lambda$  is complex. This system of equations consists of three equations with three forecast variables  $u$ ,  $v$  and  $\phi$ . The equations are:

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} - fv + \frac{\partial \phi}{\partial x} = 0 , \quad (\text{A.1})$$

$$\frac{\partial v}{\partial t} + u \frac{\partial v}{\partial x} + v \frac{\partial v}{\partial y} + fu + \frac{\partial \phi}{\partial y} = 0 , \quad (\text{A.2})$$

$$\frac{\partial \phi}{\partial t} + \frac{\partial}{\partial x}[u(\phi - \phi_B)] + \frac{\partial}{\partial y}[v(\phi - \phi_B)] = 0 , \quad (\text{A.3})$$

where  $\phi = gh$  is the geopotential height ( $h$  = height of free surface),  $\phi_B$  is the bottom topography (assumed to be independent of time),  $u$ ,  $v$  are the components of the wind velocity in the  $x$ ,  $y$  direction, respectively, and  $f$  is the Coriolis parameter. Linearizing the equations by letting

$$u = U + u' , \quad v = V + v' , \quad \phi = \phi + \phi' ,$$

where  $U$ ,  $V$  are the constant mean flow and  $\phi$  is independent of time. Assuming that  $U$ ,  $V$  are related to  $\phi$  via the geostrophic relations

$$U = - \frac{1}{f} \frac{\partial \phi}{\partial y} , \quad V = \frac{1}{f} \frac{\partial \phi}{\partial x} \quad (\text{A.4})$$

one obtains the linear system (after dropping the primes):

$$\frac{\partial u}{\partial t} + U \frac{\partial u}{\partial x} + V \frac{\partial u}{\partial y} - fv + \frac{\partial \phi}{\partial x} = 0 , \quad (\text{A.5})$$

$$\frac{\partial v}{\partial t} + U \frac{\partial v}{\partial x} + V \frac{\partial v}{\partial y} + fu + \frac{\partial \phi}{\partial y} = 0 , \quad (\text{A.6})$$

$$\frac{\partial \psi}{\partial t} + U \frac{\partial \phi}{\partial x} + V \frac{\partial \phi}{\partial y} + \frac{\partial (u\gamma)}{\partial x} + \frac{\partial (v\gamma)}{\partial y} = U \frac{\partial \phi_B}{\partial x} + V \frac{\partial \phi_B}{\partial y} , \quad (\text{A.7})$$

where  $\gamma = \phi - \phi_B$ .

If the flow is assumed to be along the topography as in [15], then the right hand side of (A.7) is zero. In such a case, one can write the solution in the form

$$\begin{aligned} u &= u_0 e^{i(\xi x + \eta y - \sigma t)} , \\ v &= v_0 e^{i(\xi x + \eta y - \sigma t)} , \\ \psi &= \psi_0 e^{i(\xi x + \eta y - \sigma t)} , \end{aligned} \quad (\text{A.8})$$

where

$$\begin{aligned} \xi &= \mu - i\rho , \\ \eta &= \nu - i\theta . \end{aligned} \quad (\text{A.9})$$

In order for (A.8) to be a solution for (A.5)-(A.7), one must have

$$\lambda = -\sigma + \xi U + \eta V \quad (\text{A.10})$$

satisfying

$$\begin{aligned}
& i\lambda[(i\lambda)^2 - i\eta(i\eta\gamma + \frac{\partial\gamma}{\partial Y}) - f[-i\lambda f - i\xi(i\eta\gamma + \frac{\partial\gamma}{\partial Y})] \\
& + (i\xi\gamma + \frac{\partial\gamma}{\partial X})(-i\eta f + \xi\lambda) = 0 .
\end{aligned}
\tag{A.11}$$

The real and imaginary parts of

$$\lambda_r = -\sigma + \mu U + \nu V ,
\tag{A.12}$$

$$\lambda_i = -\rho U - \theta V ,$$

satisfy the following system of equations (after dropping nonlinear terms in  $\lambda$ )

$$\begin{aligned}
& \lambda_r[f^2 + \gamma(\xi^2 + \eta^2)] + \lambda_i(\xi \frac{\partial\gamma}{\partial X} + \eta \frac{\partial\gamma}{\partial Y}) = f(\eta \frac{\partial\gamma}{\partial X} - \xi \frac{\partial\gamma}{\partial Y}) , \\
& \lambda_r(\xi \frac{\partial\gamma}{\partial X} + \eta \frac{\partial\gamma}{\partial Y}) - \lambda_i[f^2 + \gamma(\xi^2 + \eta^2)] = 0 .
\end{aligned}
\tag{A.13}$$

In general,  $\lambda$  is complex and, thus, the shallow water equations have a solution in the class of problems to be discussed here.

DISTRIBUTION LIST

PROFESSOR B. NETA (10)  
Code 53Nd  
DEPARTMENT OF MATHEMATICS  
NAVAL POSTGRADUATE SCHOOL  
MONTEREY, CA 93943

PROFESSOR H.M. FREDRICKSEN  
Code 53Fs  
DEPARTMENT OF MATHEMATICS  
NAVAL POSTGRADUATE SCHOOL  
MONTEREY, CA 93943

DEPARTMENT OF MATHEMATICS  
Code 53  
NAVAL POSTGRADUATE SCHOOL  
MONTEREY, CA 93943

LIBRARY, Code 0142  
NAVAL POSTGRADUATE SCHOOL  
MONTEREY, CA 93943

PROFESSOR R. T. WILLIAMS  
Code 63Wu  
DEPARTMENT OF METEOROLOGY  
NAVAL POSTGRADUATE SCHOOL  
MONTEREY, CA 93943

PROFESSOR C. S. LIOU  
Code 63Lg  
DEPARTMENT OF METEOROLOGY  
NAVAL POSTGRADUATE SCHOOL  
MONTEREY, CA 93943

DR. I. N. NAVON  
FLORIDA STATE UNIVERSITY  
SUPERCOMPUTER COMPUTATIONS RESEARCH  
INSTITUTE  
TALLAHASSEE, FL 32306

PROFESSOR GEORGE J. FIX  
UNIVERSITY OF TEXAS AT ARLINGTON  
DEPARTMENT OF MATHEMATICS  
ARLINGTON, TX 76019

PROFESSOR PAUL NELSON  
DEPARTMENT OF MATHEMATICS  
TEXAS TECH UNIVERSITY  
LUBBOCK, TX 79409

PROFESSOR H. DEAN VICTORY, JR.  
DEPARTMENT OF MATHEMATICS  
TEXAS TECH UNIVERSITY  
LUBBOCK, TX 79409

RESEARCH ADMINISTRATION OFFICE  
Code 012  
NAVAL POSTGRADUATE SCHOOL  
MONTEREY, CA 93943

DEFENSE TECHNICAL INFORMATION  
CENTER (2)  
CAMERON STATION  
ALEXANDRIA, VA 22217

PROFESSOR P. J. VANDERHOVEN  
CENTRE FOR MATHEMATICS &  
COMPUTER SCIENCE  
DEPARTMENT OF NUMERICAL  
MATHEMATICS  
KRUISLAAN 413  
1098 SJ AMSTERDAM  
THE NETHERLANDS

END

12-86

DTIC