

AD-A174 438

THEORETICAL STUDY OF PULSED-LASER-INDUCED RESONANT
DESORPTION(U) STATE UNIV OF NEW YORK AT BUFFALO DEPT OF
CHEMISTRY S VAN SMALEN ET AL NOV 86


1/1


UNCLASSIFIED

UBUFFALO/DC/86/TR-19 N00014-86-K-0043

F/G 7/4

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963-A

12

AD-A174 438

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

R & T Code 413f001---01

TECHNICAL REPORT No. 19

Theoretical Study of Pulsed-Laser-Induced Resonant Desorption

by

Sander van Smaalen and Thomas F. George

Prepared for Publication

in

Surface Science

Departments of Chemistry and Physics
State University of New York at Buffalo
Buffalo, New York 14260

November 1986

Reproduction in whole or in part is permitted for any purpose of the
United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

DTIC FILE COPY

DTIC
ELECTE
NOV 26 1986
S E D

86 11 26 048

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/86/TR-19			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260			7b. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-86-K-0043	
8c. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia			10. SOURCE OF FUNDING NOS.		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
11. TITLE Theoretical Study of Pulsed-Laser-Induced Resonant Desorption					
12. PERSONAL AUTHOR(S) Sander van Smaalen and Thomas F. George					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Yr., Mo., Day) November 1986	
				15. PAGE COUNT 21	
16. SUPPLEMENTARY NOTATION Prepared for publication in Surface Science					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.	PULSED LASER		
			VIBRATIONAL LEVELS OF ADBOND		
			RESONANT DESORPTION		
			RESONANCE HEATING		
			HIGH-INTENSITY LIMIT		
			CONTINUOUS WAVE LASER		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Adsorbed atoms irradiated by an infrared laser in resonance with one pair of vibrational levels of the adbond are studied. A pulsed laser is considered which has pulse duration short compared to the relaxation times of the system. The equations governing the time evolution of the adbond under a series of δ -pulses are derived. Two criteria are defined to compare the effects of a series of π -pulses and of a continuous-wave laser. They are: (1) equal average energy in both lasers and (2) equal average resonant heating due to both lasers. These criteria are used to compare the effects of a pulsed laser and a continuous-wave laser on laser-induced resonant desorption. It is shown that a pulsed laser does not lead to a dramatic increase of the desorption. In the high-intensity limit, resonance heating and desorption reach the same saturation limit for a pulsed laser as for a continuous-wave laser.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☒ DTIC USERS ☐			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE NUMBER (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

Theoretical Study of Pulsed-Laser-Induced Resonant Desorption

Sander van Smaalen and Thomas F. George
Departments of Physics & Astronomy and Chemistry
239 Fronczak Hall
State University of New York at Buffalo
Buffalo, New York 14260

Abstract

Adsorbed atoms irradiated by an infrared laser in resonance with one pair of vibrational levels of the adbond are studied. A pulsed laser is considered which has pulse duration short compared to the relaxation times of the system. The equations governing the time evolution of the adbond under a series of π -pulses are derived. Two criteria are defined to compare the effects of a series of π -pulses and of a continuous-wave laser. They are: (1) equal average energy in both lasers and (2) equal average resonant heating due to both lasers. These criteria are used to compare the effects of a pulsed laser and a continuous-wave laser on laser-induced resonant desorption. It is shown that a pulsed laser does not lead to a dramatic increase of the desorption. In the high-intensity limit, resonance heating and desorption reach the same saturation limit for a pulsed laser as for a continuous-wave laser.

For	
I	☒
Unannounced	☐
Justification	☐
By _____	
Distribution/	
Availability Codes	
Avail and/or	
Dist	Special
A-1	



1. Introduction

When a coated surface is irradiated by a laser, the laser frequency can be chosen such that it is in resonance with one of the transitions of the adbond. If the substrate is transparent at that frequency, the primary effect of the laser is to bring the adsorbed particle into an excited state. In this way, the laser may influence desorption and other surface processes.

Due to the interaction of the adparticle with the substrate degrees of freedom, relaxation of the excited state will occur. This will limit the probability of finding the adparticle in an excited state of the adbond. Also, it means that there is an energy flow from the laser into the substrate, a process called resonant heating.¹ In a previous paper it was shown that this is in fact the most important process occurring, i.e., almost all of the absorbed laser energy is used for heating the substrate rather than for desorption.² If the heat conductivity of the substrate is not large enough to divert the absorbed energy immediately, resonant absorption will lead to elevated surface temperatures, and consequently will lead to enhanced desorption.³ In this way the laser indirectly influences the desorption process.

The resonant effect of laser absorption has been clearly established experimentally.³⁻⁸ However, it is yet an unresolved question whether laser-induced resonant desorption or laser-induced thermal desorption via resonant heating can explain these experiments,^{3,8} although there are strong indications that the experiments can be accounted for, at least partially, by resonant desorption. In this paper, we shall calculate both desorption and resonant heating. However, we will consider only direct resonant desorption, i.e., we assume that the thermal conductivity of the substrate is large enough to maintain a constant surface temperature.

Much theoretical effort has gone into studies of the effects of continuous-wave lasers on adspecies.^{2,9-15} Experimentally, either continuous-wave lasers have been used or pulsed lasers with a pulse duration long compared to the relaxation times.^{3,4,7,8,16,17} This means that for the laser-adsorbate dynamics, the theory for continuous-wave lasers will also apply for these pulsed lasers.

In this paper, we shall consider the effect of a short laser pulse on the dynamics of an adatom. A short pulse is defined as a pulse with a duration much shorter than the energy relaxation time (T_1) of the excited levels. Then, during the action of the pulse, the relaxation can be neglected, and the effect of the pulse is to change the occupation probabilities and coherences of the two coupled levels.¹⁸ Relaxation, desorption and other possible surface processes occur after the pulse as free evolution of the adspecies system under the initial conditions determined by the effect of the pulse. We shall derive the equations of motion describing the time evolution of the reduced density matrix of the adsorbate for a series of equally spaced π -pulses.

Of particular interest is the comparison of the effect of a series of short pulses with the effect of a continuous-wave laser. We shall do so by calculating the average desorption for both cases. The key point here is the criterion used for comparing both lasers. We shall use two criteria. The first is to require that both lasers have the same average energy. The second is that both lasers must lead to the same average energy flux into the substrate. It will follow that these two criteria lead to completely different requirements for the available power in both lasers.

2. Pulsed Laser Adbond Dynamics

Consider an adsorbed atom or molecule irradiated by a laser beam with frequency ω_L . Assume that the laser is in resonance with but one pair of vibrational levels of the adbond. Then, for pulses short compared to the energy relaxation times of the system, the time evolution is given by the optical Bloch equations:¹⁸

$$\begin{aligned}\frac{dR_1}{dt} &= \Delta R_2 \\ \frac{dR_2}{dt} &= -\Delta R_1 + \Omega_p(t) R_3 \\ \frac{dR_3}{dt} &= -\Omega_p(t) R_2.\end{aligned}\tag{2.1}$$

Here $\Omega_p(t) = \frac{1}{\hbar} \vec{\mu} \cdot \vec{E}_0(t)$ is the time-dependent Rabi frequency; $\vec{E}(t) = \vec{E}_0(t) \cos(\omega_L t)$ is the classical electric field amplitude of the laser, with slowly-varying envelope $\vec{E}_0(t)$; $\vec{\mu}$ is the transition dipole of the adbond; $\Delta = \omega_L - (\omega_e - \omega_g)$ is the detuning; and $\hbar\omega_g$ and $\hbar\omega_e$ are the energy of the ground level and the excited level, respectively. The real-valued quantities R_1 , R_2 and R_3 are defined by the populations of and coherences between the two coupled levels in the rotating frame, according to

$$\begin{aligned}R_1 &= P_{ge} + P_{eg} \\ R_2 &= -i(P_{ge} - P_{eg}) \\ R_3 &= P_e - P_g.\end{aligned}\tag{2.2}$$

For nonzero detuning, analytical solutions to Eq.(2.1) exist only for a few special pulse shapes $E_0(t)$.^{18,19} For zero detuning the solution is easily obtained. It does not depend on the pulse shape, but only on the integral,

$$\Theta(t) = \int_{-\infty}^t dt' \Omega_p(t'). \quad (2.3)$$

Since we are only interested in the populations after the pulse, the integral in Eq.(2.3) can be extended over the complete pulse to obtain the pulse area Θ , which is then independent of time. In this paper only zero detuning will be considered. The values of R_1 , R_2 and R_3 directly after the pulse are,¹⁸

$$\begin{aligned} R_1(\Delta t) &= R_1^0 \\ R_2(\Delta t) &= R_2^0 \cos(\Theta) + R_3^0 \sin(\Theta) \\ R_3(\Delta t) &= R_3^0 \cos(\Theta) - R_2^0 \sin(\Theta). \end{aligned} \quad (2.4)$$

The start of the pulse is taken as the zero point of time. The pulse duration is given by Δt , and R_1^0 , R_2^0 and R_3^0 are the values of R_1 , R_2 and R_3 directly before the pulse. From Eq.(2.4) it follows that maximum inversion R_3 is obtained for pulses with a value of Θ equal to π . Since we are interested in an optimal effect, that is, the maximal value for P_e , we shall consider only π -pulses in this paper.

Assume that the system is initially in thermal equilibrium, i.e., $R_1 = 0$, $R_2 = 0$ and $R_3 = R_3(\text{eq})$. Then, after a π -pulse, $R_1 = R_1(\Delta t)$ and $R_2 = R_2(\Delta t)$ are zero again, whereas R_3 is changed to

$$R_3(\Delta t) = -R_3(\text{eq}). \quad (2.5)$$

Between two consecutive pulses the adbond evolves freely through its relaxation against the substrate degrees of freedom. The off-diagonal elements of the reduced density matrix of the adbond decay exponentially to zero,¹⁵ and because they were initially zero, they remain zero throughout. The time evolution of the diagonal elements is described by the master

equation

$$\frac{dP_n(t)}{dt} = \sum_k \{W_{kn} P_k(t) - W_{nk} P_n(t)\}, \quad (2.6)$$

where the summation extends over all vibrational levels of the adbond, including those not involved in the coupling to the laser. W_{nk} is the rate constant for the transition from state n to state k , which can be obtained, for example, through second-order perturbation theory.^{15,20,21} The populations, $P_n(t)$, can be obtained by solving Eq. (2.5) under the initial conditions given by the result of the pulse, $P_n(\Delta t)$.

Consider the case where the adspecies is irradiated by a series of equally-spaced π -pulses, with interval time t_p . Then, after several pulses, a quasi steady state will be reached, wherein the time evolution in each interval t_p will be the same. The formal solution of Eq. (2.6) is ($\Delta t \leq t \leq t_p$),

$$\underline{P}(t) = \underline{P}(eq) + e^{-\underline{W}(t-\Delta t)} [\underline{P}(\Delta t) - \underline{P}(eq)], \quad (2.7)$$

where $\underline{W}$ is the matrix formed by the transition rates W_{nk} , and has elements

$$W_{nm} = + \sum_k W_{nk} \delta_{nm} - W_{mn}. \quad (2.8)$$

$\underline{P}(t)$ is the vector formed by the occupation probabilities $P_n(t)$, and $\underline{P}(eq)$ is the equilibrium solution of Eq. (2.6). Of course $\underline{P}(eq)$ can be omitted from Eq. (2.7). However, to clearly demonstrate the thermal and the laser effect, we have written it as it is. In the quasi steady state, Eq. (2.7) gives a relation between the $P_n(\Delta t)$ and $P_n(t_p)$ of

$$\underline{P}(t_p) = \underline{P}(eq) + e^{-\underline{W}t_p} [\underline{P}(\Delta t) - \underline{P}(eq)], \quad (2.9)$$

where we have used that $\exp[-\frac{W}{2}\Delta t] = 1$. Alternatively, the effect of the pulse is known from Eq. (2.5), which gives

$$\begin{aligned} P_n(t_p) &= P_n(\Delta t) = P_n^0 \quad (n \neq e, g) \\ P_g(t_p) &= P_g(\Delta t) = P_g^0 \\ P_e(t_p) &= P_e(\Delta t) = P_e^0. \end{aligned} \quad (2.10)$$

Equations (2.9) and (2.10) together define the quantities $P_n(t_p)$ and $P_n(\Delta t)$ in terms of the system parameters W_{nk} and the time t_p . They therefore provide the initial condition to be used in Eq. (2.7). A schematic drawing of the time evolution in the quasi steady state is given in Fig. 1.

3. Continuous-Wave Laser

Again, we consider a laser in resonance with only two vibrational levels of the adbond. For a continuous-wave laser, the equation of motion for the reduced density matrix of the adbond is²⁰

$$\begin{aligned} \frac{dP_n(t)}{dt} &= \sum_k \{W_{kn} P_k(t) - W_{nk} P_n(t)\} \quad (n \neq e, g) \\ \frac{dP_g(t)}{dt} &= \sum_k \{W_{kg} P_k(t) - W_{gk} P_g(t)\} + \frac{1}{2} \Omega_{CW} R_2(t) \\ \frac{dP_e(t)}{dt} &= \sum_k \{W_{ke} P_k(t) - W_{ek} P_e(t)\} - \frac{1}{2} \Omega_{CW} R_2(t) \\ \frac{dR_2(t)}{dt} &= - \sum_k \frac{1}{2} (W_{ek} + W_{gk}) R_2(t) + \Omega_{CW} (P_e(t) - P_g(t)) - \Delta R_1(t) \\ \frac{dR_1(t)}{dt} &= - \sum_k \frac{1}{2} (W_{ek} + W_{gk}) R_1(t) + \Delta R_2(t), \end{aligned} \quad (3.1)$$

where the symbols have the same meaning as in Eqs. (2.1), (2.2) and (2.6), and Ω_{CW} is the time independent Rabi frequency of the continuous wave laser. For the rate constants W_{nk} , the expressions derived by Efrima et al.²¹ or Arnoldus et al.¹⁵ can be used. Equations similar to Eq. (3.1) can be

obtained for the other coherences. They are not coupled to Eq. (3.1) and therefore need not be considered here.

We are interested in the long-time behavior of the system, i.e., only the steady-state solution of Eq. (3.1) is needed. Then the occupation probabilities follow from

$$\begin{aligned} \sum_k W_{nk} P_n(\infty) &= \sum_k W_{kn} P_k(\infty) \quad (n \neq e, g) \\ \sum_k W_{gk} P_g(\infty) &= \sum_k W_{kg} P_k(\infty) - \frac{1}{2} \frac{\Gamma_{eg}}{\frac{1}{2}\Gamma_{eg}^2 + 2\Delta^2} \Omega_{CW}^2 [P_g(\infty) - P_e(\infty)] \\ \sum_k W_{ek} P_e(\infty) &= \sum_k W_{ke} P_k(\infty) + \frac{1}{2} \frac{\Gamma_{eg}}{\frac{1}{2}\Gamma_{eg}^2 + 2\Delta^2} \Omega_{CW}^2 [P_g(\infty) - P_e(\infty)] \end{aligned} \quad (3.2)$$

with $\Gamma_{eg} = \sum_k (W_{ek} + W_{gk})$ and $P_n(\infty)$ denotes the steady-state value of $P_n(t)$.

4. Pulsed Laser and Continuous-Wave Laser

To be able to compare the effects of a pulsed laser and the effects of a continuous-wave laser, a criterion is necessary for comparing these two lasers. One possibility is to require that the total average energy in the continuous-wave laser and in the series of pulses be equal. Because the intensity of a laser is proportional to the square of the electric field amplitude, and the Rabi-frequency is proportional to the amplitude of the electric field itself, this leads to the condition:

$$\frac{1}{t_p} \int_0^{\Delta t} dt \Omega_p^2(t) = \Omega_{CW}^2, \quad (4.1)$$

where it is assumed that the constants of proportionality are the same in both cases. For a π -pulse we have also the condition,

$$\theta = \int_0^{\Delta t} dt \Omega_p(t) = \pi. \quad (4.2)$$

For a given average laser intensity, Ω_{CW}^2 , Eqs. (4.1) and (4.2) together give a relation between the pulse duration Δt and the pulse interval time t_p .

For a square pulse, that relation is

$$\Delta t \cdot t_p = \frac{\pi^2}{\Omega_{CW}^2}. \quad (4.3)$$

Further restrictions are $\Delta t \ll t_p$ and $\Delta t \ll \Gamma^{-1}$, with Γ^{-1} a typical relaxation time. Therefore, Eqs. (4.1)-(4.3) severely limit the number of available pulses for a given intensity.

A second criterion for comparing the effects of the two lasers is to require that the average energy flow into the substrate (resonant heating) is equal in both cases, which is almost equivalent to requiring that the total absorption from the lasers be equal.² For the continuous-wave laser the energy flow in the steady state is constant in time. Per unit of time it is given by,²

$$\frac{dE_{CW}}{dt} = \hbar \omega_L \frac{\frac{1}{2} \Gamma_{eg}}{\frac{1}{2} \Gamma_{eg}^2 + 2\Delta^2} \Omega_{CW}^2 (P_g(\infty) - P_e(\infty)). \quad (4.4)$$

For the pulsed laser the average absorption in the quasi steady state can simply be obtained from Eq. (2.10) as

$$\frac{\Delta E_p}{t_p} = \hbar \omega_L \frac{1}{t_p} (P_g^0 - P_e^0). \quad (4.5)$$

Combination of these two equations gives

$$\frac{1}{t_p} (P_g^0 - P_e^0) = \frac{\Omega_{CW}^2}{\Gamma_{eg}} [P_g(\infty) - P_e(\infty)], \quad (4.6)$$

where we have set $\Delta = 0$. The populations P_g^0 and P_e^0 can be expressed in terms of the rate constants and the interval time t_p by use of Eqs (2.9) and (2.10). The steady-state solutions $P_g(\infty)$ and $P_e(\infty)$ follow from Eq. (3.2) and are a function of the transition probabilities and the Rabi frequency Ω_{CW} .

5. Desorption

Desorption can be described by the addition of a loss term, $-W_{nc} P_n(t)$, to each of the master equations, Eqs. (2.6) and (3.1). Previously, the loss rates were defined as the rate constant for the transition from the bound state n to any of the continuum states of the adbond potential.²¹ However, the results presented here are not restricted to such an interpretation.

Here we consider only slow desorption, which allows the use of the populations for the steady-state and the quasi steady-state as determined by Eqs. (3.1) and (2.6), respectively. With these occupation probabilities the average rate of desorption is defined as,

$$D = \frac{1}{(t_p - \Delta t)} \int_{\Delta t}^{t_p} dt \sum_n W_{nc} P_n(t), \quad (5.1)$$

where by the nature of the assumptions made earlier the integral can be taken either from Δt to t_p or from 0 to t_p . To obtain the average desorption for the pulsed laser and for the continuous-wave laser, the respective functions $P_n(t)$ have to be substituted in Eq. (5.1).

The ratio between the pulsed-laser desorption and the continuous-wave-laser desorption is a measure of the effectiveness of the one over the other. This ration, μ , can be written as

$$\mu = \frac{\sum_n W_{nc} P_n(av)}{\sum_n W_{nc} P_n(\infty)}, \quad (5.2)$$

where $P_n(av)$ is the average occupation probability for the pulsed irradiated adbond given as

$$P_n(av) = \frac{1}{(t_p - \Delta t)} \int_{\Delta t}^t dt P_n(t), \quad (5.3)$$

and $P_n(\infty)$ is the steady state occupation probability for the continuous wave laser case.

For a continuous-wave laser, the high-intensity limit is defined by

$$\frac{\Omega_{CW}^2}{\Gamma_{eg}^2} \rightarrow \infty, \text{ for the pulsed laser the high-intensity limit corresponds to}$$

$\Gamma t_p \rightarrow 0$. Note that for the idea of a pulsed laser to be valid, the latter limit has to be taken under the condition $\Delta t \ll t_p$. It is easily shown that the occupation probabilities, the energy flux and desorption assume saturation values in the high-intensity limit. In the high-intensity limit the occupation probabilities for the pulsed laser and for the continuous-wave laser obey the same set of equations

$$\sum_k W_{nk} P_n = \sum_k W_{kn} P_k \quad (5.4)$$

$$P_g = P_e.$$

For a pulsed laser, Eq. (5.4) refers to $P_n(av) = P_n(\Delta t)$, and they are valid apart from a term of the order of Γt_p . For a continuous-wave laser,

Eq. (5.4) refers to $P_n(\infty)$, and they are valid apart from terms of the order

of $\left(\frac{\Omega_{CW}^2}{\Gamma_{eg}^2}\right)^{-1}$. It follows immediately that the desorption for both situations

is the same, i.e., $\mu = 1$ [Eq. (5.2)]. The saturation limit for the energy flux is also the same, and can be written as

$$\frac{dE_{CW}}{dt} = \frac{\Delta E_p}{t_p} = \gamma_{\omega_L} \sum_k \{W_{kg} P_k - W_{gk} P_g\}. \quad (5.5)$$

This means that use of criterion 2 leads to $\mu = 1$ in the high-intensity limit. Criterion 1 (equal average laser intensity) is defined by [Eq. (4.3)]

$$\frac{\gamma_{eg} \Delta t}{\pi^2} \gamma_{eg} t_p = \left(\frac{\Omega_{CW}^2}{\gamma_{eg}^2} \right)^{-1}, \quad (5.6)$$

One of our basic assumptions is $\frac{\gamma_{eg} \Delta t}{\pi^2} \ll 1$. Therefore, if we take $\frac{\Omega_{CW}^2}{\gamma_{eg}^2}$ $\approx \left(\frac{\gamma_{eg} \Delta t}{\pi^2} \right)^{-1} \gg 1$, $\gamma_{eg} t_p$ is about one. It follows that the saturation limit is reached for the continuous-wave laser at a much lower average intensity than for the pulsed laser. It can be concluded that in the high-intensity limit the pulsed laser is not more effective than the continuous-wave laser, whereas from an energetic point of view the continuous-wave laser will always be the best choice.

A weak continuous-wave laser is defined by $\frac{\Omega_{CW}^2}{\gamma_{eg}^2} \ll 1$. In that limit the populations are different from their equilibrium values by a term proportional to the laser intensity, i.e.,

$$P_n(\infty) = P_n(eq) + C_n \frac{\Omega_{CW}^2}{\gamma_{eg}^2}, \quad (5.7)$$

where C_n are constants. For $(\frac{\Omega_{CW}^2}{\Gamma_{eg}^2}) \ll 1$, we also have [Eqs. (4.3) and (4.6)] $\Gamma_{eg} t_p \gg 1$. Then, the pulsed-laser-induced desorption can approximately be evaluated from

$$P_n(av) = P_n(eq) + \frac{1}{\Gamma_{eg} t_p} \{P_n(\Delta t) - P_n(eq)\}, \quad (5.8)$$

where Γ is in the range of the nonzero eigenvalues of $\underline{W}$. The situation we are interested in is the situation where the laser gives a considerable enhancement of the desorption. This means that the pure thermal desorption can be neglected. An approximate expression for μ is then

$$\mu = \frac{\sum_n W_{nc} [P_n(\Delta t) - P_n(eq)]}{\sum_n W_{nc} C_n} \frac{1}{\frac{\Gamma_{eg}^2 t_p}{\Omega_{CW}^2}} \quad (5.9)$$

Since we do not know the constants C_n , an estimate for μ cannot be made. However, it does follow from Eq. (5.9) that both criteria for comparing the lasers [Eqs. (4.3) and (4.6)], lead to a μ which is independent of the laser power. For criterion 2 [Eq. (4.6)], μ becomes simply,

$$\mu = \frac{\sum_n W_{nc} [P_n(\Delta t) - P_n(eq)]}{\sum_n W_{nc} C_n}, \quad (5.10)$$

that is, μ is completely determined by system properties. For criterion 1 [Eq. (4.3)], a value less by a factor $\frac{\Gamma_{eg} \Delta t}{\pi^2} \ll 1$ is obtained. It is interesting to note that now μ becomes larger if the pulse duration becomes longer. Of course, the pulse duration is limited to values $(\Gamma_{eg} \Delta t) \ll 1$.

6. Two-Level System

A two-level system is of special interest because all the equations can be solved analytically, and values for the different quantities are obtained which depend only one relaxation time $\Gamma^{-1} = (W_{ge} + W_{eg})^{-1}$. There are only two states, the ground state $|g\rangle$ and the excited state $|e\rangle$, with the obvious relation between the populations of

$$P_e(t) + P_g(t) = 1. \quad (6.1)$$

For the pulsed-laser-irradiated adbond, it follows from Eqs. (2.7), (2.9) and (2.10) that

$$P_e(t) = P_e(\text{eq}) + \frac{[1 - 2P_e(\text{eq})]}{[1 + e^{-\Gamma t} P]} e^{-\Gamma t}, \quad (6.2)$$

with the equilibrium population given by

$$P_e(\text{eq}) = \frac{W_{ge}}{\Gamma}. \quad (6.3)$$

The average population of the upper level is [Eq. (5.3)]

$$P_e(\text{av}) = P_e(\text{eq}) + [1 - 2P_e(\text{eq})] \frac{1}{\Gamma t} \frac{[1 - e^{-\Gamma t} P]}{[1 + e^{-\Gamma t} P]}. \quad (6.4)$$

The average energy flow into the substrate is obtained as [Eq. (4.5)]

$$\frac{\Delta E}{t_p} = \hbar \omega_L \Gamma P_e(\text{av}) \quad (6.5)$$

For the continuous-wave laser irradiation, the excited level population is,

$$P_e(\infty) = P_e(\text{eq}) + [1 - 2P_e(\text{eq})] \frac{1}{\frac{1}{2}\Gamma^2 + \Omega_{cw}^2}, \quad (6.6)$$

which is constant in time. The energy flow into the substrate is given by,

$$\frac{dE_{cw}}{dt} = \hbar \omega_L \frac{\Omega^2}{\Gamma} [1 - 2P_e(\infty)]. \quad (6.7)$$

The situation is considered where pure thermal desorption can be neglected, i.e., the adspecies is stable without the laser. Then, for desorption the first term in Eq. (6.2) and in Eq. (6.6) can be neglected, and the ratio between pulsed-laser-induced desorption and continuous-wave-laser-induced desorption becomes [Eq. (5.2)]

$$\mu = \frac{2}{[1 + e^{-\Gamma t_p}]} \frac{[1 - e^{-\Gamma t_p}]}{\Gamma t_p} \frac{[\frac{1}{2}\Gamma^2 + \Omega_{CW}^2]}{\Omega_{CW}^2} \quad (6.8)$$

This implies that in the strong laser limit ($\Gamma t_p \rightarrow 0$; $\frac{\Omega_{CW}^2}{\Gamma^2} \rightarrow \infty$) $\mu = 1$, as it should be. Also, for $\frac{\Omega_{CW}^2}{\Gamma^2} \gg 1$ but $\Gamma t_p \neq 0$, it is easily shown that $\mu < 1$, with μ decreasing for increasing Γt_p . Using criterion 1, the ratio can be written as,

$$\mu = \frac{[1 - e^{-\Gamma t_p}]}{[1 + e^{-\Gamma t_p}]} \frac{\Gamma^2 + 2\Omega_{CW}^2}{\Gamma^2} \frac{\Gamma \Delta t}{\pi^2}, \quad (6.9)$$

which in the low-intensity limit leads to $\mu \approx \frac{\Gamma \Delta t}{\pi^2} \ll 1$. For Ω_{CW}^2/Γ^2 and/or Γt_p of the order unity, μ starts increasing, until it reaches its saturation value 1 in the high-intensity limit.

With the second criterion we obtain from Eqs. (6.5), (6.7) and (6.8) that μ equals unity, independent of the laser power used. Note that in this case μ is equal to the ratio of the quantum yield of desorption. Where the latter is defined as the number of photons used for desorption divided by the number of photons used for substrate heating.²

7. Conclusions

In this paper we have studied the dynamics of an adspecies irradiated by a series of equally-spaced π -laser pulses, in resonance with one pair of

vibrational levels of the adbond. The equations describing the time evolution of the reduced density matrix of the adbond are derived. A quasi steady state is defined as the state wherein the time evolution in each pulse interval is the same.

In particular, we have studied the laser-induced desorption and the resonant heating of the substrate. A comparison is made between the effects of a pulse train and of a continuous-wave laser. The essential point for making this comparison is how to relate both lasers. Two criteria are given which may serve for such a goal. The first is to require that both lasers have equal average energy. It is then found that the ratio between the pulsed-laser-induced desorption and the continuous-wave-laser-induced desorption [Eq. (5.2)] becomes proportional to $\Gamma\Delta t$ (low intensity limit), where Δt is the pulse duration. In other words, μ can be made arbitrarily small by decreasing the pulse duration. This result can be understood easily if one realizes that for one pulse the energy content is proportional to the intensity multiplied by the duration, whereas the total exciting power is only proportional to the product of Δt with the square root of the intensity [Eqs. (2.3) and (2.4)]. This should be compared with the continuous-wave laser case, where both the energy and the exciting power are proportional to the intensity [Eq. (3.2)]. This observation leads to an interesting prediction for the intensity dependence of the laser induced desorption. Experimentally, it has been found that the laser-induced desorption is proportional to some power, n , of the laser intensity, where n is interpreted as the number of photons necessary for providing the required desorption energy.⁷ From the analysis here, it follows that for short pulses an exponent of $\frac{1}{2}n$ will be found.

The situation is considered where pure thermal desorption can be neglected, i.e., the adspecies is stable without the laser. Then, for desorption the first term in Eq. (6.2) and in Eq. (6.6) can be neglected, and the ratio between pulsed-laser-induced desorption and continuous-wave-laser-induced desorption becomes [Eq. (5.2)]

$$\mu = \frac{2}{[1 + e^{-\Gamma t_p}]} \frac{[1 - e^{-\Gamma t_p}]}{\Gamma t_p} \frac{[\frac{1}{2}\Gamma^2 + \Omega_{CW}^2]}{\Omega_{CW}^2} \quad (6.8)$$

This implies that in the strong laser limit ($\Gamma t_p \rightarrow 0$; $\frac{\Omega_{CW}^2}{\Gamma^2} \rightarrow \infty$) $\mu = 1$, as it should be. Also, for $\frac{\Omega_{CW}^2}{\Gamma^2} \gg 1$ but $\Gamma t_p \neq 0$, it is easily shown that $\mu < 1$, with μ decreasing for increasing Γt_p . Using criterion 1, the ratio can be written as,

$$\mu = \frac{[1 - e^{-\Gamma t_p}]}{[1 + e^{-\Gamma t_p}]} \frac{\Gamma^2 + 2\Omega_{CW}^2}{\Gamma^2} \frac{\Gamma \Delta t}{\pi^2}, \quad (6.9)$$

which in the low-intensity limit leads to $\mu \approx \frac{\Gamma \Delta t}{\pi^2} \ll 1$. For Ω_{CW}^2/Γ^2 and/or Γt_p of the order unity, μ starts increasing, until it reaches its saturation value 1 in the high-intensity limit.

With the second criterion we obtain from Eqs. (6.5), (6.7) and (6.8) that μ equals unity, independent of the laser power used. Note that in this case μ is equal to the ratio of the quantum yield of desorption. Where the latter is defined as the number of photons used for desorption divided by the number of photons used for substrate heating.²

7. Conclusions

In this paper we have studied the dynamics of an adspecies irradiated by a series of equally-spaced π -laser pulses, in resonance with one pair of

vibrational levels of the adbond. The equations describing the time evolution of the reduced density matrix of the adbond are derived. A quasi steady state is defined as the state wherein the time evolution in each pulse interval is the same.

In particular, we have studied the laser-induced desorption and the resonant heating of the substrate. A comparison is made between the effects of a pulse train and of a continuous-wave laser. The essential point for making this comparison is how to relate both lasers. Two criteria are given which may serve for such a goal. The first is to require that both lasers have equal average energy. It is then found that the ratio between the pulsed-laser-induced desorption and the continuous-wave-laser-induced desorption [Eq. (5.2)] becomes proportional to $\Gamma\Delta t$ (low intensity limit), where Δt is the pulse duration. In other words, μ can be made arbitrarily small by decreasing the pulse duration. This result can be understood easily if one realizes that for one pulse the energy content is proportional to the intensity multiplied by the duration, whereas the total exciting power is only proportional to the product of Δt with the square root of the intensity [Eqs. (2.3) and (2.4)]. This should be compared with the continuous-wave laser case, where both the energy and the exciting power are proportional to the intensity [Eq. (3.2)]. This observation leads to an interesting prediction for the intensity dependence of the laser induced desorption. Experimentally, it has been found that the laser-induced desorption is proportional to some power, n , of the laser intensity, where n is interpreted as the number of photons necessary for providing the required desorption energy.⁷ From the analysis here, it follows that for short pulses an exponent of $\frac{1}{2}n$ will be found.

The second criterion used is to require that the energy flux from the laser into the substrate is the same in both cases. Then it is found that in the low-intensity limit μ becomes independent of the laser power. Of course, to achieve this, a pulsed laser is required which is much more intense than the continuous-wave laser.

In the high-intensity limit, all quantities assume saturation values. It is shown that $\mu = 1$ in this case. From criterion 1 it follows that the high-intensity limit is reached for much lower average power by the continuous-wave laser than by the pulsed laser.

For a two-level system the equations can be solved analytically. It is found that criterion 2 leads to $\mu = 1$ for all laser powers. In the low-intensity limit criterion 1 gives $\mu \ll 1$. For increasing intensity, μ grows gradually to one.

In this paper it has been shown that use of a pulsed laser instead of a continuous-wave laser does not lead to a dramatic increase of resonant desorption or resonant heating. Basically this can be understood from the fact that both processes depend in some way on the average populations of the excited vibrational levels. These cannot be enhanced by using a pulsed laser. Other processes, like laser-induced thermal desorption via resonant heating, depend on the average of some function (e.g., exponential) of the occupation probabilities. It can be expected that then quite different results will be obtained.

Acknowledgements

We thank H.F. Arnoldus for a critical reading of the manuscript. This research was supported by the Office of Naval Research, Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009, and the National Science Foundation under Grant CHE-8519053. The

United States Government is authorized to reproduce and distribute reprints notwithstanding any copyright notation hereon.



REFERENCES

1. Z.W. Gortel, H.J. Kreuzer, P. Piercy and R. Teshima, Phys. Rev. B 28 (1983) 2119.
2. S. van Smaalen, H.F. Arnoldus and T.F. George, Phys. Rev. B, submitted.
3. I. Hussla, H. Seki, T.J. Chuang, Z.W. Gortel, H.J. Kreuzer and P. Piercy, Phys. Rev. B 32 (1985) 3489.
4. J. Heidberg, H. Stein and E. Riehl, Phys. Rev. Lett. 49 (1982) 666.
5. T.J. Chuang and H. Seki, Phys. Rev. Lett. 49 (1982) 382.
6. T.J. Chuang, J. Chem. Phys. 76 (1982) 3828.
7. T.J. Chuang, Surf. Sci. Reports 3 (1983) 1.
8. K. Veeken, P.A.M. van der Heide, L.M. ten Dam, A.R. de Vroomen and J. Reuss, Surf. Sci. 166 (1986) 1.
9. M.S. Slutsky and T.F. George, Chem. Phys. Lett. 57 (1978) 474.
10. J.T. Lin and T.F. George, J. Phys. Chem. 84 (1980) 2957.
11. A.C. Beri and T.F. George, J. Vac. Sci Technol. B 3 (1985) 1529.
12. X.Y. Huang, T.F. George and J.M. Yuan, J. Opt. Soc. Am. B 2 (1985) 985.
13. Z.W. Gortel, H.J. Kreuzer, P. Piercy and R. Teshima, Phys. Rev. B 27 (1983) 5066.
14. B. Fain and S.H. Lin, Surf. Sci. 147 (1984) 497.
15. H.F. Arnoldus, S. van Smaalen and T.F. George, Phys. Rev. B, submitted.
16. R.B. Hall and S.J. Bares in: Chemistry of Structure at Interfaces: New Laser and Optical Techniques, Eds., R.B. Hall and A.B. Ellis (VCH Publishers, 1985).
17. R.B. Hall, A.M. DeSantolo and S.J. Bares, Surf. Sci. 161 (1985) L533.
18. L. Allen and J.H. Eberly, Optical Resonance and Two-Level Atoms, (Wiley, New York 1975).
19. A. Bambini and P.R. Berman, Phys. Rev. A 23 (1981) 2496.
20. W.H. Louisell, Quantum Statistical Properties of Radiation, (Wiley, New York, 1973) Chapt. 6.
21. S. Efrima, L. Jedrzejek, K.F. Freed, E. Hood and H. Metiu, J. Chem. Phys. 79 (1983) 2436.

FIGURE CAPTION

Fig. 1: Schematic drawing of the time evolution of the vibrational level populations for a pulse sequence in the quasi steady state. Curve (a) represents the inversion $R_3 = P_e - P_g$; curve (b) represents $P_e + P_g$; and curve (c) represents the population of any other level $n \neq e$ or g . The actual variation of the populations during the pulse ($0 < t < \Delta t$) depends on the pulse shape. For a two-level system there are no levels $n \neq e$ or g , and $P_e + P_g = 1$.

Figure 1.



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookeville, Tennessee 38501

Dr. A. Reisman
Microelectronics Center of North Carolina
Research Triangle Park, North Carolina
27709

Dr. D. DiLella
Chemistry Department
George Washington University
Washington D.C. 20052

Dr. M. Grunze
Laboratory for Surface Science and
Technology
University of Maine
Orono, Maine 04469

Dr. R. Reeves
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. J. Butler
Naval Research Laboratory
Code 6115
Washington D.C. 20375-5000

Dr. Steven M. George
Stanford University
Department of Chemistry
Stanford, CA 94305

Dr. L. Interante
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Mark Johnson
Yale University
Department of Chemistry
New Haven, CT 06511-8118

Dr. Irvin Heard
Chemistry and Physics Department
Lincoln University
Lincoln University, Pennsylvania 19352

Dr. W. Knauer
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. K.J. Klaubunde
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. Murday
Naval Research Laboratory
Code 6170
Washington, D.C. 20375-5000

Dr. J. B. Hudson
Materials Division
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne
Chemistry Department
Northwestern University
Evanston, Illinois 60637

Dr. J. M. White
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. D. E. Harrison
Department of Physics
Naval Postgraduate School
Monterey, California 93940

Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria
Electrical Engineering Department
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green
Quantum Surface Dynamics Branch
Code 3817
Naval Weapons Center
China Lake, California 93555

Dr. A. Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02912

Dr. S. L. Bernasek
Department of Chemistry
Princeton University
Princeton, New Jersey 08544

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Richard Colton
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Dan Pierce
National Bureau of Standards
Optical Physics Division
Washington, D.C. 20234

Dr. R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer
Materials Characterization Lab.
General Electric Company
Schenectady, New York 22217

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Ronald Lee
R301
Naval Surface Weapons Center
White Oak
Silver Spring, Maryland 20910

Dr. Paul Schoen
Code 6190
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. John T. Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene
Code 5230
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda
University of Pittsburgh
Chemistry Building
Pittsburg, PA 15260

Dr. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH
UNITED KINGDOM

Dr. H. Tachikawa
Chemistry Department
Jackson State University
Jackson, Mississippi 39217

Dr. John W. Wilkins
Cornell University
Laboratory of Atomic and
Solid State Physics
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis
Department of Physics
University of California
Irvine, California 92664

Dr. D. Ramaker
Chemistry Department
George Washington University
Washington, D.C. 20052

Dr. J. C. Hemminger
Chemistry Department
University of California
Irvine, California 92717

Dr. T. F. George
Chemistry Department
University of Rochester
Rochester, New York 14627

Dr. G. Rubloff
IBM
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. Horia Metiu
Chemistry Department
University of California
Santa Barbara, California 93106

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. P. Hansma
Department of Physics
University of California
Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser
Department of Chemistry
University of Richmond
Richmond, Virginia 23173

Dr. R. W. Plummer
Department of Physics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. Steckl
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. G.H. Morrison
Department of Chemistry
Cornell University
Ithaca, New York 14853

END

1-87

DTIC