

AD-A174 613

PARDAXIN'S ACTION IN SHARK(U) NEW YORK AQUARIUM
BROOKLYN OSBORN LABS OF MARINE SCIENCES
N PRIMOR ET AL 24 OCT 86 N00014-82-C-0435


1/1

UNCLASSIFIED

F/G 6/7

NL





MICROCOPY RESOLUTION TEST CHART



17

October 24, 1986

AD-A174 613

Dr. Eli D. Schmell
Code 1141MB
Office of Naval Research
800 North Quincy St.
Arlington, VA 22217-5000

Dear Dr. Schmell:

Enclosed please find a Progress Report. In addition, I am enclosing a paper which is being accepted for publication in J. Biol. Chem.. It is expected to be published in February-March, 1987. This is a major paper on the chemistry and the mechanism of ion-channel formation of pardaxin.

Sincerely,

Naftali Primor

Naftali Primor

NP:ms
Enc.

DTIC FILE COPY

DTIC
UNCLASSIFIED
10/13/1986
Ab

This document is not approved for public release and its distribution is restricted.

86 11 18 050

Title Project: Pardaxin's action in shark

Naftali Primor, Philip Lazarovici
Osborn Laboratories of Marine Sciences
Brooklyn, New York 11224

Contract: N 00014-82-C-0435

Objectives:

Structure and mode of action of the shark repellent pardaxin

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



Abstract

1. Amino Acid analysis

The amino acid composition of pardaxins I and II purified from P. marmoratus secretion is listed. -

Contd.

Amino acid	PXI	PXII
Asp	4.22 (4)	3.79 (4)
Thr	1.23 (1)	0.92 (1)
Ser	3.78 (4)	3.75 (4)
Glu	2.98 (3)	2.71 (3)
Pro	0.76 (1)	1.05 (1)
Gly	6.33 (6)	4.23 (4)
Ala	2.82 (3)	2.83 (3)
Val	1.77 (2)	1.13 (1)
Ile	1.93 (2)	1.85 (2)
Leu	2.25 (2)	0.62 (1)
Tyr	0	0
Phe	2.27 (2)	1.80 (2)
His	0.95 (1)	0.94 (1)
Lys	1.91 (2)	1.23 (1)
Arg	0.33 (0)	0.23 (0)
Trp	0	0
Cys	1.32 (2)	1.51 (2)

L.

1

Met	0	1.91 (1)
Total Residues	35	31
Molecular weight	3529	3188
Total hydrophylic ⁽³⁾	48.5%	45.2%
Total hydrophobic ⁽⁴⁾	51.5%	54.8%

2. Sequence determination of pardaxin I

Cont'd → Pardaxins were analyzed by automated Edman degradation *Cont'd* in a Beckman sequencer 890 C. The degradation was carried out using 0.25 M Quadrol, 5% phenylisothiocyanate with single cleavage. Conversion of the phenylthio derivatives was achieved with 1 N HCl for 10 min at 80°C. The phenylthiohydantoin (PTH) derivatives were identified by high pressure liquid chromatography.

The amino acid sequence of the NH₂-terminal of pardaxin I:

<u>Cycle</u>	<u>Amino</u>
<u>No.</u>	<u>acid</u>
1	Gly
2	Phe
3	Phe
4	Ala
5	Leu
6	Ile
7	Pro
8	Gly
9	Ile
10	Glu

Cont'd

3. Binding of Pardaxin to shark gill membranes ² is accomplished ^(were used)
Membranes from a blue shark (Prionace) gills. A
sucrose gradient was used for membrane purification.
Pardaxin was iodinated using the method of Bolton-Hunter
reagent (as described in the paper JBC, In press.). We are
in the process of evaluating this data.

Plans for next year:

1. Synthesis of N-terminal.
2. Test for activities of the synthesized N terminal:
 - a. hydrophobicity
 - b. Compative replacement with pardaxin using membranes
 - c. cytotoxicity
3. Binding of the iodinated pardaxin to shark gill membranes

Published work on the subject (1986)

Primor, N. Action on ileal smooth muscle of synthetic detergents and pardaxin. Gen. Pharmacol. 17, 413-418 (1986).

Lazarovici, P., Primor, N. and Loew, L. Purification and pore forming activity of two hydrophobic polypeptides from the Red Sea Moses sole. J. Biol. Chem. In press.

END

1-87

DTIC