

AD-A176 171

SLEEP DEPRIVATION AND EXERCISE TOLERANCE(U) INDIANA  
UNIV FOUNDATION INDIANAPOLIS RESEARCH AND SPONSORED  
PROGRAMS B J MARTIN 31 JAN 86 DAND17-81-C-1023


1/1

UNCLASSIFIED

F/G 6/19

NL





13

SE

AD-A176 171

DOCUMENTATION PAGE

Form Approved  
OMB No 0704-0188  
Exp Date Jun 30, 1986

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited.	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		7a. NAME OF MONITORING ORGANIZATION	
6a. NAME OF PERFORMING ORGANIZATION Indiana University Foundation	6b. OFFICE SYMBOL (if applicable)	7b. ADDRESS (City, State, and ZIP Code)	
6c. ADDRESS (City, State, and ZIP Code) 355 Lansing Street Indianapolis, Indiana 46202		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DAMD17-81-C-1023	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION U.S. Army Medical Research & Development Command	8b. OFFICE SYMBOL (if applicable) SGRD-RMI-S	10. SOURCE OF FUNDING NUMBERS	
8c. ADDRESS (City, State, and ZIP Code) Fort Detrick Frederick, Maryland 21701-5012		PROGRAM ELEMENT NO. 62777A	PROJECT NO. #3E1-62777A879
		TASK NO. BF	WORK UNIT ACCESSION NO. 084
11. TITLE (Include Security Classification) (U) Sleep Deprivation and Exercise Tolerance			
12. PERSONAL AUTHOR(S) Martin, Bruce J.			
13a. TYPE OF REPORT Annual	13b. TIME COVERED FROM 2/1/85 TO 1/31/86	14. DATE OF REPORT (Year, Month, Day) 1986 January 31	15. PAGE COUNT 8
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	stress diaphragm	
06	06	stress hormones	
06	16	exertion	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Sleep loss appears to influence subsequent exercise in a manner dependent upon the nature of that exercise. To investigate this, we examined the effect of 24 sleepless hours on the ability to produce and maintain maximal ventilation, a form of exercise involving a limited muscle mass. We found that sleep loss reduced maximal ventilatory output and also resulted in persistent mood changes. In addition, sleep loss worsened performance of both short- and long-term maximal ventilation, indicating that its influence was not limited to prolonged, boring tasks. We concluded that those forms of exercise utilizing the largest muscle mass are most resistant to the influence of sleep deprivation.			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED / UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			
22a. NAME OF RESPONSIBLE INDIVIDUAL Mary Frances Bostian		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22b. TELEPHONE (Include Area Code) 301-663-7325		22c. OFFICE SYMBOL SGRD-RMI-S	

DTIC  
SELECTED  
JAN 27 1987  
S. D.

DTIC FILE COPY

SLEEP DEPRIVATION AND EXERCISE TOLERANCE

Annual Report

Bruce J. Martin, Ph.D.

February 1, 1985 - January 31, 1986

Supported by

U.S. Army Medical Research and Development Command  
Fort Detrick, Frederick, Maryland 21701-5012

Contract No. DAMD17-81-C-1023

Indiana University Foundation  
Research and Sponsored Programs  
355 Lansing Street  
Indianapolis, Indiana 46202

Approved for public release; distribution unlimited.

The findings of this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

**SUMMARY**

A 24-hour sleepless period

- a) reduces short-term maximal ventilation
- b) reduces prolonged isocapnic maximal ventilation
- c) reduces short-term isocapnic submaximal ventilation
- d) increases sleepiness and worsens mood

We conclude that sleep loss has effects on small muscle mass performance at least as great as those evidenced in work requiring a larger muscle mass.



Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

**FOREWORD**

For the protection of human subjects this investigator has adhered to policies of applicable Federal Law 45 CFR 46.

### Statement of the Problem

Sleep deprivation is a common occurrence in both the military and civilian spheres. In many cases, severe or prolonged exercise follows sleep loss. In many instances, this exercise involves a small, not large muscle mass. Because we know nothing of how variation in active muscle mass influences sleep deprivation's effects, this study was undertaken.

### Background

Previous work under this contract identified several effects of sleep deprivation upon subsequent exercise. Sleep loss did not influence  $VO_2$ max or short-term work tolerance (1), while it did blunt the ability to carry out long-term heavy exercise (2, 3), in proportion to the "dosage" of sleeplessness. Measurements of physiological responses to exercise after sleep loss revealed no clear effects on oxygen uptake, heart rate, minute ventilation or blood lactate. In addition, prolonged mild exercise provoked no greater or lesser stress hormonal (B-endorphin, cortisol) response after sleep loss than it did after normal sleep (4). Most provocatively, this form of exercise significantly reduced mood disturbance and sleepiness when carried out for 3 hours.

### Approach to the Problem

These previous results provoked our inquiry into the relation of muscle mass involved in exercise to the sleep loss effect. By investigating small-muscle mass exercise, we could examine the components of "arousal" that make mild treadmill walking such an effective antidote to sleepiness after sleep loss. By using respiratory muscles as our small muscle group, we could at the same time approach the question of sleep loss influence on one of the components of respiratory failure.

### Results and Discussion

Twelve young healthy persons volunteered to serve as subjects. All were studied twice in counterbalanced fashion, once after normal sleep and once after a single sleepless night (24 sleepless hours). Both short and long-term maximal ventilation was measured, with all tests maintained isocapnic. Mood and sleepiness was assessed before and during the ventilatory maneuvers. 12-s, 1-min, and 30-min maximal isocapnic voluntary ventilations were measured, as was a 1-min test at 75% of maximal effort.

The results of the study are listed below:

- 1) Sleep loss increased sleepiness (raw score)

<u>Control</u>	<u>Sleep-deprived</u>	<u>p</u>
1.92	4.25	<0.01
<u>±.26</u>	<u>±.39</u>	..

and sleepiness during the 30-min MVV increased above that before the maneuver began.

<u>Before</u>	<u>During</u>	<u>p</u>
4.25	5.00	<0.01
<u>±.39</u>	<u>.43</u>	

While sleepiness was unchanged in the control situation:

<u>Before</u>	<u>During</u>	<u>p</u>
1.92	2.08	NS
<u>±.26</u>	<u>±.31</u>	

Discussion: Prolonged dynamic exercise involving a small muscle mass is ineffective as a means of arousal, even when performed maximally. The failure of this exercise to blunt sleepiness, when treadmill walking did suggest the presence of a muscle-mass dependent effect.

2) Sleep loss worsened mood:

Total mood disturbance (raw score)

	<u>Control</u>	<u>Sleep-deprived</u>	<u>p</u>
Before tests	5.7 ± 8.0	41.6 ± 8.8	<0.07
During 30-min MVV	5.7 ± 6.8	47.8 ± 10.9	<0.07

Discussion: Thus, there was no worsening during the MVV. The mood disturbance, as is typical after sleep loss, was composed largely of increased fatigue and confusion, and decreased vigor. Tension, anxiety, and depression were unaltered.

3) Sleep loss reduced maximal voluntary ventilation ( $l \cdot \text{min}^{-1}$  BTPS):

	<u>Control</u>	<u>Sleep-deprived</u>	<u>p</u>
72-sec MVV	168.8 ± 12.2	156.8 ± 11.5	<0.05
1-min MVV	129.4 ± 7.4	116.9 ± 7.0	<0.01
30-min MVV	100.6 ± 6.2	86.7 ± 6.4	<0.01

Discussion: The change averaged 7-12% and was equally large at short-term as at long-term exercise. This contrasts with previous work suggesting that the ability to perform short-term, high-intensity work is preserved after sleep loss. Interestingly, at every minute of the 30-min MVV, maximal ventilation was significantly lower after sleep deprivation. Also, submaximal ventilation was reduced by sleep loss:

	<u>Control</u>	<u>Sleep loss</u>	<u>p</u>
1 min at 3/4 maximal effort	92 + 6.4	84.9 + 6.0	<0.05

This result suggests that sleep deprivation has skewed the entire perceptive range of voluntary ventilatory effort.

### Conclusions

Small muscle mass exercise is hindered at least as much or more than is work utilizing larger muscle masses when sleep deprivation precedes exercise. This effort, though undetermined is speculated to arise from the reduced stimulus for arousal inherent in small muscle mass work.

### Recommendations

Although sleep loss is relatively ineffective as a direct inhibitor of exercise, this may be exercise intensity and mode dependent. Exercise involving smaller amounts of muscle, even when maximal, may allow psychological sleep loss effects to powerfully undermine performance.

### Literature Cited

1. Martin, B. J., and G. M. Gaddis. Exercise after sleep deprivation. Med. Sci. Sports Exercise. 13: 220-223, 1981.
2. Martin, B. J. Effect of sleep deprivation on tolerance of prolonged exercise. Eur. J. Appl. Physiol. 47: 345-355, 1981.
3. Martin, B. J., and H.-I. Chen. Sleep loss and the sympathoadrenal response to exercise. Med. Sci. Sports Exercise 16: 56-59, 1984.
4. Martin, B. J., P. R. Bender, and H.-I. Chen. Stress hormonal response to exercise after sleep loss. Eur. J. Appl. Physiol. 55: 210-214, 1986.

Publications Supported by Contract - DAMD17-81-C-1023

Martin, B. J., and G. M. Gaddis. Exercise after sleep deprivation. Med. Sci. Sports Exercise 13: 220-223, 1981.

Martin, B. J. Effect of sleep deprivation on tolerance of prolonged exercise. Eur. J. Appl. Physiol. 47: 345-354, 1981.

Martin, B. J., and R. Haney. Self-selected exercise intensity is unchanged by sleep loss. Eur. J. Appl. Physiol. 49: 79-86, 1982.

Martin, B. J. and H.-I. Chen. Sleep loss and the sympathoadrenal response to exercise. Med. Sci. Sports Exercise 16: 56-59, 1984.

Kolka, M. A., R. S. Elizondo, and B. J. Martin. Effect of sleep loss on self-selected exercise intensity during cold exposure. Eur. J. Appl. Physiol. 53: 282-285, 1984.

Martin, B. J., P. R. Bender, and H.-I. Chen. Stress hormonal response to exercise after sleep loss. Eur. J. Appl. Physiol. 55: 210-214, 1986.

DISTRIBUTION LIST

1 copy                   Commander  
U.S. Army Medical Research and Development Command  
ATTN: SGRD-RMI-S  
Fort Detrick, Frederick, Maryland 21701-5012

12 copies                Defense Technical Information Center (DTIC)  
ATTN: DTIC-DDAC  
Cameron Station  
Alexandria, VA 22304-6145

1 copy                    Dean  
School of Medicine  
Uniformed Services University of the Health Sciences  
4301 Jones Bridge Road  
Bethesda, MD 20814-4799

1 copy                    Commandant  
Academy of Health Sciences, U.S. Army  
ATTN: AHS-CDM  
Fort Sam Houston, TX 78234-6100

END

2-87

DTIC