

AD-A176 972

ROBUSTNESS OF T-TEST(U) PITTSBURGH UNIV PA CENTER FOR
MULTIVARIATE ANALYSIS T KARIYA ET AL OCT 86 TR-86-31
AFOSR-TR-87-0031 F49620-85-C-0008

1/1


UNCLASSIFIED

F/G 12/1

NL



END
3-87



Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR- 87-0031	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ROBUSTNESS OF t-TEST		5. TYPE OF REPORT & PERIOD COVERED Technical - October 1986
7. AUTHOR(s) Takeaki Kariya Bimal K. Sinha N. C. Giri		6. PERFORMING ORG. REPORT NUMBER 86-31
8. PERFORMING ORGANIZATION NAME AND ADDRESS Center for Multivariate Analysis Fifth Floor Thackeray Hall University of Pittsburgh, Pittsburgh, PA 15260		9. CONTRACT OR GRANT NUMBER(s) Japan - General C61530012 Air Force - F49620-85-C-0008 NSERC Canada - ECAC Quebec
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research Department of the Air Force Bolling Air Force Base, DC 20332		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 611025, 2304/145
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) <i>same as 11</i>		12. REPORT DATE October 1986
		13. NUMBER OF PAGES 18
		15. SECURITY CLASS. (of this report) Unclassified
		16a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) <i>same as 16</i>		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Elliptically symmetric distribution, Student's t-test, UMP test, UMPI test, UMPS test.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this paper we establish the optimality robustness of Student's t-test mainly without invariance. This generalizes some well-known results of Lehmann and Stein (1949), and Kariya and Eaton (1977).		

DD FORM 1 JAN 73 1473

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

AD-A176 972

MMG FILE COPY

AFOSR-TR- 87-0031

ROBUSTNESS OF t-TEST

Takeaki Kariya *
Hitotsubashi University

Bimal K. Sinha **
University of Maryland Baltimore County
and
University of Pittsburgh

N. C. Giri ***
Universite de Montreal


AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFOSR)
NOTICE OF TRANSMITTAL TO DTIC

This technical report has been reviewed and is
approved for public release IAW AFR 190-12.
Distribution is unlimited.

MATTHEW J. KEEPER
Chief, Technical Information Division

Center for Multivariate Analysis

University of Pittsburgh



Approved for public release;
distribution is unlimited.

87: 2 20 235

ROBUSTNESS OF t-TEST

Takeaki Kariya *
Hitotsubashi University

Bimal K. Sinha **
University of Maryland Baltimore County
and
University of Pittsburgh

N. C. Giri ***
Universite de Montreal



October 1986

Technical Report No. 86-31

Center for Multivariate Analysis
Fifth Floor Thackeray Hall
University of Pittsburgh
Pittsburgh, PA 15260

Accession For	
NTIS GSA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Avail and/or	
Dist	Special

A-1

- * Research supported by a grant from the Ministry of Education of Japan under General C61530012.
- ** Research supported by the Air Force Office of Scientific Research (AFSC) under Contract F49620-85-C-0008. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.
- *** Research supported by a grant from NSERC of Canada and FCAC of Quebec.

ROBUSTNESS OF t-TEST

Takeaki Kariya *
Hitotsubashi University

Bimal K. Sinha **
University of Maryland Baltimore County
and
University of Pittsburgh

N. C. Giri ***
Universite de Montreal

ABSTRACT

In this paper we establish the optimality robustness of Student's t-test mainly without invariance. This generalizes some well-known results of Lehmann and Stein (1949), and Kariya and Eaton (1977).

AMS 1980 Subject Classifications: Primary 62G10; Secondary 62F05.

Key words and phrases: Elliptically symmetric distribution, Student's t-test, UMP test, UMPI test, UMPS test.

* Research supported by a grant from the Ministry of Education of Japan under General C61530012.

** Research supported by the Air Force Office of Scientific Research (AFSC) under Contract F49620-85-C-0008. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

*** Research supported by a grant from NSERC of Canada and FCAC of Quebec.

1. INTRODUCTION

The standard optimum properties like UMP, UMPU, UMPS, UMPI, of the familiar Student's t-test for testing the significance of the mean of a univariate normal population with unknown variance against one-sided or both-sided alternatives are well-known. See, for example, Lehmann (1959). In this paper we show mainly without invariance that the t-test continues to be optimum when normality is replaced by a class of elliptically symmetric distributions, thereby establishing optimality robustness of the t-test. To fix ideas, let $\underline{x} = (x_1, \dots, x_n)'$ $\sim N(\underline{\mu}, \sigma^2 I_n)$, $\sigma^2 > 0$ unknown, and consider the problem of testing $H_0^*: \mu = 0$ versus $H_1^*: \mu > 0$ or $H_2^*: \mu \neq 0$. Define the one-sided t-test and two-sided t-test respectively by

$$\phi_1(\underline{x}) = \begin{cases} 1 & \text{if } v > c \\ 0 & \text{otherwise} \end{cases} \quad (1.1)$$

$$\phi_2(\underline{x}) = \begin{cases} 1 & \text{if } |v| > d \\ 0 & \text{otherwise} \end{cases} \quad (1.2)$$

where $v = \underline{x}'\underline{1}/\|\underline{x}\| \|\underline{1}\|$, $\|\underline{x}\| = (\sum_{i=1}^n x_i^2)^{1/2}$, $\|\underline{1}\| = n^{1/2}$. It is proved in Lehmann and Stein (1948) that

- (a) for $\frac{1}{2} \leq \alpha < 1$, ϕ_1 is UMP for H_0^* versus H_1^* ; however,
- (b) no UMP test exists for H_0^* versus H_1^* for $0 < \alpha < \frac{1}{2}$. On the other hand, it is trivial that
- (c) ϕ_1 is UMPI for H_0^* versus H_1^* while
- (d) ϕ_2 is both UMPS and UMPI for H_0^* versus H_2^* .

In the sequel, we assume that $\underline{x}$ has a density of the form

$$f(\underline{x}|\underline{\mu}, \sigma^2) = \sigma^{-n} q(\|\underline{x} - \underline{\mu}\|^2 / \sigma^2) \quad (1.3)$$

for some $q \in Q$ given by

$$Q = \{q | q: [0, \infty) \rightarrow [0, \infty), \int_{R^n} q(\|x\|^2) dx = 1\}. \quad (1.4)$$

In what follows we mostly assume that both σ^2 and q in (1.3) are unknown, and consequently define the three classes of densities on R^n :

$$F_0(\mu) = \{f | f(x) = q(\|x - \mu\|^2), q \in Q\} \quad (1.5)$$

$$F_1(\mu) = \{f | f \in F_0(\mu), q \text{ nonincreasing}\}$$

$$F_2(\mu) = \{f | f \in F_1(\mu), q \text{ convex}\}.$$

The restrictions of nonincreasingness and convexity imposed on q in (1.5) above are standard in robustness study (see Kariya (1981)). Clearly $F_2(\mu) \subset F_1(\mu) \subset F_0(\mu)$ for all real μ . Denote the density of x by $h(x)$ and consider the testing problems

$$H_0: h \in F_0(0) \text{ versus } K_0: h \in F_0(\mu), \mu > 0 \text{ unknown} \quad (1.6)$$

$$H_1: h \in F_1(0) \text{ versus } K_1: h \in F_1(\mu), \mu > 0 \text{ unknown}$$

$$H_2: h \in F_2(0) \text{ versus } K_2: h \in F_2(\mu), \mu \neq 0 \text{ unknown}.$$

Note the very general nature of both null and alternative hypotheses in (1.6) which are essentially nonparametric. We define by C_i^α the class of level α tests under H_i , $i = 0, 1, 2$, defined as

$$C_i^\alpha = \{\phi \in C^\alpha | E_f \phi \leq \alpha \text{ for all } f \in F_i(0)\}, \quad 0 < \alpha < 1, \quad (1.7)$$

where C^α denotes the class of all level α tests on R^n . It then follows that $C_0^\alpha \supset C_1^\alpha \supset C_2^\alpha$ become $F_2(0) \subset F_1(0) \subset F_0(0)$.

There are some results in the literature for the testing problems similar to (1.6). Lehmann and Stein (1949) proved that $(e)_{\phi_1}$ is UMP for

testing H_0 versus h is $N(\mu 1, \sigma^2 I_n)$, $\mu > 0$ unknown. Kariya and Eaton (1977) proved that $(f)\phi_1$ is UMP for testing H_0 versus K_1 while $(g)\phi_2$ is UMPS for testing H_0 versus K_2 . It is interesting to note the implications of these results. Clearly (f) is stronger than (e) . Moreover, the class of level α tests for which ϕ_1 is UMP in (e) is precisely C_0^α and not $C_N^\alpha = \bigcap_{\sigma^2 > 0} C_N^\alpha(\sigma^2)$ which is relevant in (b) . Here $C_N^\alpha(\sigma^2) = \{\phi \in C^\alpha | E[\phi | N(0, \sigma^2 I_n)] \leq \alpha\}$, and C_N^α is clearly much bigger than C_0^α . The same is true for the optimality properties in (f) and (g) in that they hold only in C_0^α . For example, the UMP property of ϕ_1 in (f) simply means that for any $f(\cdot | \mu) \in F_1(\mu)$ and $\mu > 0$,

$$E[\phi_1 | f(\cdot | \mu)] \geq E[\phi | f(\cdot | \mu)] \quad \text{for all } \phi \in C_0^\alpha \quad (1.8)$$

but it does not mean that for a fixed $h(\cdot | \mu) \in F_1(\mu)$, ϕ_1 is UMP for testing

$$H_1(h): \mu = 0 \text{ versus } K_1(h): \mu > 0 \quad (1.9)$$

in the class of level α tests under the fixed h defined as

$$C^\alpha(h) = \{\phi \in C^\alpha | E[\phi | h(\cdot | 0)] \leq \alpha\}. \quad (1.10)$$

It is, therefore, evident that in the consideration of optimality robustness, it is essential to clearly specify the class in which optimality holds.

It is the object of this paper to strengthen the results in (f) and (g) . We prove in Section 2 that for each fixed $h \in F_1(\mu)$, ϕ_1 is UMP for testing $\mu = 0$ versus $\mu > 0$, i.e. for the problem (1.9), in the class of conditional level α tests given $w = \|x\|^2$, which is a subclass of $C^\alpha(h)$ given in (1.10) above. Also, the class C_0^α is shown to be the intersection of the classes of conditional level α tests over $h \in F_1(0)$. These two

results imply (f). Further, when $\frac{1}{2} \leq \alpha < 1$, ϕ_1 is shown to be UMP in $C_\alpha = \bigcap_{h \in F_1(0)} C^\alpha(h)$. In Section 3, it is shown that, for each fixed $h \in F_2(\mu)$, ϕ_2 is UMP for testing $\mu = 0$ versus $\mu \neq 0$ in the class of conditional similar level α tests satisfying a certain condition. This result is stronger than (g). Finally, in Section 4, some properties of the t-test with invariance are pointed out.

The following result, which is very basic and whose proof is omitted, is useful for our purpose.

PROPOSITION 1. Let $h \in F_0(\mu)$ be the density of $\tilde{x}$, where $h(\tilde{x}) = q(\|\tilde{x} - \mu_1\|^2)$. Then

(1) (v, w) is a sufficient statistic for h .

(2) The joint density of (v, w) is given by

$$g(v, w; \mu) = c_0 q(w - 2\sqrt{w}v\sqrt{n}\mu + n\mu^2) r_0(v) w^{n/2-1}$$

with

$$r_0(v) = 2[B(\frac{1}{2}, \frac{n-3}{2})]^{-1} (1 - v^2)^{(n-3)/2}$$

where $c_0 = \{\Gamma(\frac{1}{2})\}^n / \Gamma(n/2)$ and $B(a, b)$ denotes the Beta function.

(3) When $\mu = 0$, v and w are independent with density $r_0(v)r_1(w)$ where

$$r_1(w) = c_0 w^{n/2-1} q(w).$$

By (1) of the above proposition, without loss of generality we can restrict the class of tests for problems (1.6) to the one based on (v, w) . We denote this class by $\mathcal{D}$ in the rest of the paper.

2. ONE-SIDED TESTING PROBLEMS

As mentioned above, in this section we consider the following one-sided testing problems:

$$\text{For a fixed } h \in F_1(\mu), \quad \mu = 0 \text{ versus } \mu > 0 \quad (2.1)$$

$$H_0: h \in F_0(0) \text{ versus } K_1: h \in F_1(\mu), \quad \mu > 0 \text{ unknown} \quad (2.2)$$

$$H_1: h \in F_1(0) \text{ versus } K_1: h \in F_1(\mu), \quad \mu > 0 \text{ unknown.} \quad (2.3)$$

Note that the classes of level α tests relevant for these problems are respectively $C^\alpha(h)$, C_0^α and C_1^α defined in (1.10) and (1.7). Of course, by Proposition 1, any test function $\phi(x)$ belonging to these classes is essentially of the form $\phi(v, w)$. Let $E_0^V(\cdot)$ denote the expectation of $\cdot$ with respect to the density $r_0(v)$ of v given in (2) of Proposition 1 and let a.a.(w, h) denote "almost all w with respect to the density $r_1(w)$ of w " given in (4) where r_1 depends on h . Also, let

$$\begin{aligned} \mathcal{D}^\alpha(h) &= \{\phi \in \mathcal{D}^\alpha \mid \phi(v, w) \in C^\alpha(h)\}, \quad h \in F_1(0) \\ \mathcal{D}_i^\alpha &= \{\phi \in \mathcal{D}^\alpha \mid \phi(v, w) \in C_i^\alpha\}, \quad i = 0, 1 \end{aligned} \quad (2.4)$$

where $\mathcal{D}^\alpha$ denotes the class of level α tests based on (v, w) .

Our first main result is the following.

THEOREM 2.1. For problem (2.1), the test ϕ_1 defined in (1.1) is UMP in the class of conditional level α tests

$$K^\alpha(h) = \{\phi \in \mathcal{D}^\alpha(h) \mid E_0^V[\phi(v, w)] \leq \alpha \quad \text{a.a.}(w, h)\}. \quad (2.5)$$

Proof. Write $h(\underline{x}) = q(\|\underline{x} - \underline{\mu}_1\|^2)$ in $F_1(\underline{\mu})$ and fix $\underline{\mu} = \underline{\mu}_1 > 0$.

From (2) of Proposition 1, the conditional density of v given w is given by $g(v, w; \underline{\mu}_1) / \int_{-1}^1 g(v, w; \underline{\mu}_1) dv$. Using the nonincreasing property of q and applying the Neyman-Pearson Lemma, it follows that the test with the critical region $v > c(w)$ is MP in the class (2.5). But, when $\underline{\mu} = 0$, v and w are independent from (3) of Proposition 1. Therefore, $c(w)$ is independent of w . Finally, $r_0(v)$ being independent of q , the above test coincides with ϕ_1 and is UMP in (2.5), thus completing the proof.

Remark 2.1. It is important to note that ϕ_1 is UMP only in $K^\alpha(h)$ but not in $\mathcal{D}^\alpha(h)$. In fact, no UMP test exists in $\mathcal{D}^\alpha(h)$ because an MP test for a fixed $\underline{\mu} = \underline{\mu}_1 > 0$ in general depends on both q and $\underline{\mu}_1$. However, the above result implies (f) as demonstrated below.

Let

$$K_i^\alpha = \bigcap_{h \in F_i(0)} K^\alpha(h), \quad i = 0, 1. \quad (2.6)$$

LEMMA 2.1. (1) $\mathcal{D}_0^\alpha = K_0^\alpha$. (2) $\mathcal{D}_0^\alpha = K_1^\alpha$.

Proof. If $\phi \in K_0^\alpha$, then $\phi \in K^\alpha(h)$ for all $h \in F_0(0)$ which implies $E_0^V[\phi(v, w)] \leq \alpha$, a.a.(w.h) for any $h \in F_0(0)$. This immediately gives $E[\phi(v, w)|h] \leq \alpha$ for all $h \in F_0(0)$ so that $\phi \in \mathcal{D}_0^\alpha$. To show the converse, suppose $\phi \in \mathcal{D}_0^\alpha$ and there exists $h_1 \in F_0(0)$ such that h_1 gives positive mass to the set $S = \{w > 0 | E_0^V[\phi(v, w)] > \alpha\}$. Clearly S is independent of any $h \in F_0(0)$ since, by Proposition 1, the density of v under $\underline{\mu} = 0$ is $r_0(v)$, independent of h . Denote by $r_1(w)$ the marginal density of w under h_1 , which is of the form given in (3) of Proposition 1. Since $r_1(w)$ is absolutely continuous with respect to Lebesgue measure, S has a positive Lebesgue

measure, which in turn implies that S contains a bounded nonempty open set A . Now define a density on R^n by $h_2(x) = I_A(\|x\|^2) / \int_{R^n} I_A(\|x\|^2) dx$ where $I_A(\cdot)$ denotes the indicator function of the set A . Obviously, $h_2 \in F_0(0)$ and h_2 gives the whole mass to the set A . Therefore, $E[\phi|h_2] = E[E_0^V[\phi(v,w)]|r_2] > \alpha$ where $r_2(w)$ is the density of w under h_2 . This implies $\phi \notin D_0^\alpha$, a contradiction. This proves (1).

To prove (2), note that $D_0^\alpha \subseteq K_1^\alpha$ by (1) and the fact that $F_1(0) \subseteq F_0(0)$. To show the converse, suppose $\phi \in K_1^\alpha$ which means $E[\phi|h] \leq \alpha$ for all $h \in F_1(0)$. By the completeness of w for the family $\{N(0, c^2 I_n)^{1/2} > 0\}$ whose densities are included in $F_1(0)$, we get $E_0^V[\phi(v,w)] \leq \alpha$, a.e. (Lebesgue), implying $\phi \in D_0^\alpha$. This proves (2).

As a by-product, we get the following interesting facts.

COROLLARY 2.1. For problem (2.1), ϕ_1 is UMP in $K_1^\alpha = D_0^\alpha$. For problem (2.2), ϕ_1 is UMP in D_0^α .

Next we consider problem (2.3). Note that although the alternatives for the two problems (2.2) and (2.3) are the same (K_1), the null hypotheses H_0 and H_1 are different as pointed out below.

LEMMA 2.2. $D_0^\alpha \subseteq D_1^\alpha$.

Proof. Since $F_0(0) \supseteq F_1(0)$, it is clear that $D_0^\alpha \subseteq D_1^\alpha$. To extend this test $\phi \in D_1^\alpha$ but $\phi \notin D_0^\alpha$, let $\phi(w) = 1$ for $1 < w < 2 + a$, ($a > 0$), and $\phi(w) = 0$ otherwise. Then for $f(x) = \exp(-x^2/2)$ in $F_1(0)$, from (2.3) it is easy to see that we get

$$E_\phi[\phi(w)] = \int_1^{2+a} w c_0 w^{n/2-1} dw = c_0(2+a)^{n/2+1}/(1+a) < c_0(1+a)^{n/2+1} = 1$$

since $q(1) \leq 1$ because $q(\int_0^1 \|x\|^2 dx) = 1$ and q is nonincreasing, for sufficiently small $a > 0$, $E_f[:(w)] \leq 1$ for all $f \in F_1$. On the other hand, defining $f_1(x) = I_B(\|x\|^2) / \int_B \|x\|^2 dx$ with $B = \{x \in \mathbb{R}^n : \|x\| \leq a\}$, it is clear that $f_1(x) \in F_0(0)$ but $\notin F_1(0)$, and $E_{f_1}(w) > 1$. This completes the proof.

In view of the above lemma, it follows that a theorem analogous to Theorem 2.1 for problem (2.2), the same need not be true. In fact, we show below that the optimality condition (2.1) holds only for $\frac{1}{2} \leq \alpha \leq 1$. We can further show that (2.1) does not hold for $\alpha < \frac{1}{2}$.

Towards this end, we need the following lemma which is a direct consequence of Lemma 2.1, (2.1), (2.2).

LEMMA 2.2 Let $f_1 \in F_1$ be a function satisfying (2.1) for $\alpha > 0$. Then, for all $f \in F_1$, we have

$$E_f(w) \leq E_{f_1}(w) \quad \text{if and only if} \quad \int_0^1 \|x\|^2 dx \leq \int_0^1 \|x\|^2 dx.$$

$$\left\{ \begin{array}{l} \int_0^1 \|x\|^2 dx \leq \int_0^1 \|x\|^2 dx \\ \int_0^1 \|x\|^2 dx \geq \int_0^1 \|x\|^2 dx \end{array} \right.$$

where $\int_0^1 \|x\|^2 dx = \int_0^1 \|x\|^2 dx$ and $\int_0^1 \|x\|^2 dx = \int_0^1 \|x\|^2 dx$.

Proof. The result is obvious for $\alpha = 1$. For $\alpha < 1$, we have

LEMMA 2.3 Assume $\frac{1}{2} \leq \alpha \leq 1$, and fix $f_1(x) = q_1(\|x\|^2)$ in $F_1(0)$

where $q_1(t) = 1 - \alpha t$ for testing $H_0: f_1(0) \leq 1$ versus $H_1: f_1(0) > 1$ with $\alpha > 0$, f_1 is

$$f_1(x) = \frac{1}{\int_0^1 \|x\|^2 dx}$$

Proof. Fix $\mu = \mu_1 > 0$ and choose $f_1(x) = q_1(\|x - \mu_1 \underline{1}\|^2) = f_1^{\mu_1}(x)$ (say), $f_0(x) = q_1(\|x\|^2 + 2\beta\mu_1\sqrt{n}\|x\| + n\mu_1^2)/J$ in Lemma 2.3, where $0 < \beta < 1$ and $J = \int_{\mathbb{R}^n} q_1(\|x\|^2 + 2\beta\mu_1\sqrt{n}\|x\| + n\mu_1^2)dx > 0$. Note that J can be written as

$$\begin{aligned} J &= \int_{\mathbb{R}^n} q_1((\|x\| + \beta\mu_1\sqrt{n})^2 + \delta)dx \\ &= c_1 \int_0^\infty q_1((r + \beta\mu_1\sqrt{n})^2 + \delta)r^{n-1}dr \\ &= c_1 \int_\tau^\infty q_1(r^2 + \delta)(r - \tau)^{n-1}dr < \infty \end{aligned}$$

since $1 = \int_{\mathbb{R}^n} q_1(\|x\|^2)dx = c_1 \int_0^\infty q_1(r^2)r^{n-1}dr$. Here c_1 is a constant, $\tau = \beta\mu_1\sqrt{n} > 0$ and $\delta = n\mu_1^2(1 - \beta^2) > 0$. Clearly, $f_0(x) \in F_1(0)$ because $q_1(\cdot)$ is nonincreasing in $\|x\|$ for $\tau > 0$. Hence, by Lemma 2.3 with $K = J$ and $v = 0$, the test ψ with critical region

$$q_1(\|x - \mu_1 \underline{1}\|^2) \geq q_1(\|x\|^2 + 2\beta\mu_1\sqrt{n}\|x\| + n\mu_1^2) \quad (2.8)$$

is MP for testing $H_1: h \in F_1(0)$ versus $h = f_1^{\mu_1}$ provided $E_f[\psi] \leq \alpha$ for all $f \in F_1(0)$. Since q_1 is nonincreasing, (2.8) yields the test ϕ_1 with the critical region $v \geq -\beta$. However, by (3) of Proposition 1, the distribution of v when $\mu = 0$ is independent of $h \in F_0(0) \supset F_1(0)$. Hence ϕ_1 is MP for testing H_1 versus $f = f_1^{\mu_1}$ and hence UMP against f_1 because ϕ_1 is independent of the fixed $\mu_1 > 0$. Finally, note that $P\{v > -\beta\} \geq P\{v > 0\} = \frac{1}{2}$ for any $h \in F_1(0)$. Hence the theorem.

Since the UMP test ϕ_1 for a fixed q_1 in $F_1(\mu)$ is independent of q_1 , we immediately have

COROLLARY 2.2. For problem (2.3), ϕ_1 is UMP whenever $\frac{1}{2} \leq \alpha < 1$.

3. TWO-SIDED TESTING PROBLEMS

In this section we deal with the following two testing problems:

$$\text{For a fixed } h \in F_2(\mu), \quad \mu = 0 \text{ versus } \mu \neq 0 \text{ unknown} \quad (3.1)$$

$$H_2: h \in F_2(0) \text{ versus } K_2: h \in F_2(\mu), \quad \mu \neq 0 \text{ unknown.} \quad (3.2)$$

Note that the classes of level α test functions relevant for these two problems are respectively $C^\alpha(h)$ and C_2^α defined in (1.10) and (1.7).

As in Section 2, let

$$\mathcal{D}^\alpha(h) = \{\phi \in \mathcal{D}^\alpha \mid \phi(v, w) \in C^\alpha(h)\}, \quad h \in F_2(0) \quad (3.3)$$

$$\mathcal{D}_2^\alpha = \{\phi \in \mathcal{D}^\alpha \mid \phi(v, w) \in C_2^\alpha\}$$

where $\mathcal{D}^\alpha$ is the class of all level α tests based on (v, w) . Further, let

$$K(v: \mu, w) = g(v, w; \mu) / \int g(v, w; \mu) dv \quad (3.4)$$

be the conditional density of v given w where $g(v, w; \mu)$ is given in (2) of Proposition 1. Note that $K(v: 0, w) = r_0(v)$. For a test function $\phi \in \mathcal{D}^\alpha$, let

$$\pi(\phi, (\mu, w, h)) \equiv E[\phi(v, w) \mid K(\cdot: \mu, w)] = \int_{-1}^1 \phi(v, w) K(v: \mu, w) dv \quad (3.5)$$

be the conditional power of ϕ , given w , which reduces to the conditional size

$$\pi(\phi, (0, w, h)) \equiv E_0^v[\phi(v, w)] = \int_{-1}^1 \phi(v, w) r_0(v) dv \quad (3.6)$$

under $\mu = 0$. Finally, let $KU^\alpha(h)$ and $KS^\alpha(h)$ be respectively the class of conditional unbiased tests of level α in $\mathcal{D}^\alpha$ and the class of conditional similar tests of level α in $\mathcal{D}^\alpha$ for $h \in F_2(\mu)$ defined as

$$\begin{aligned}
KU^\alpha(h) &= \{\phi \in \mathcal{D}^\alpha(h) \mid E_0^V[\phi(v, w)] \leq \alpha \quad \text{a.a.}(w, h), \\
&\quad \pi(\phi, (\mu, w, h)) \geq \alpha \text{ for all } \mu \neq 0 \quad \text{a.a.}(w, h)\} \\
KS^\alpha(h) &= \{\phi \in \mathcal{D}_2^\alpha \mid E_0^V[\phi(v, w)] = \alpha \quad \text{a.a.}(w, h)\}.
\end{aligned}
\tag{3.7}$$

It is easy to verify that for each fixed $h \in F_2(\mu)$, the conditional power $\pi(\phi, (\mu, w, h))$ is continuous a.a.(w, h) at $\mu = 0$, which yields the relation

$$KU^\alpha(h) \subseteq KS^\alpha(h) \text{ for each fixed } h \in F_2(\mu). \tag{3.8}$$

We are now ready to state the first main result of this section.

THEOREM 3.1. For problem (3.1), the two-sided t-test ϕ_2 defined in (1.2) is UMP in the class of tests in $KS^\alpha(h)$ satisfying

$$E_0^V[v\phi(v, w)] = 0 \quad \text{a.a.}(w, h). \tag{3.9}$$

Proof. Write $h(x) = q(\|x - \mu_1\|^2)$ where q is fixed nonincreasing and convex, and consider the problem of testing $\mu = 0$ versus $\mu = \mu_1 \neq 0$ in the conditional density $K(v; \mu, w)$ in (3.4). Obviously $KS^\alpha(h)$ is the class of size α tests for this problem. Now by the generalized Neyman-Pearson lemma, a test ϕ^* which maximizes the conditional power $\pi(\phi, (\mu_1, w, h))$ subject to (3.9) is given by

$$\phi^*(v, w) = \begin{cases} 1 & \text{if } K(v; \mu_1, w) \geq c_1 r_0(v) + c_2 v r_0(v) \\ 0 & \text{otherwise} \end{cases} \tag{3.10}$$

where c_1 and c_2 are constants chosen to satisfy the size α condition $E_0^V[\phi^*] = \alpha$ and (3.9). Using (2) of Proposition 1 and the convexity of q , it follows easily that ϕ^* is equivalent to

$$\phi^*(v, w) = \begin{cases} 1 & \text{if } v > b \text{ or } v < a \\ 0 & \text{otherwise} \end{cases} \quad (3.11)$$

which is clearly independent of μ_1 . From (3) of Proposition 1, due to the symmetry of $r_0(v)$, we get that the constants a and b satisfying $E_0^V(\phi^*) = \alpha$ and (3.9) must obey the relation $-a = b = c$ (say). Therefore $\phi^* = \phi_2$, and hence for any $\phi \in KS^\alpha(h)$ satisfying (3.9) and for any $\mu \neq 0$, we get

$$\pi(\phi_2, (\mu, w, h)) \geq \pi(\phi, (\mu, w, h)) \quad \text{a.a.}(w, h). \quad (3.12)$$

Taking expectations with respect to w in both sides of (3.12) proves the theorem.

In view of the relation in (3.8) and the fact that $\phi_2 \in KU^\alpha(h)$, we immediately conclude

COROLLARY 3.1. For problem (3.1), ϕ_2 is UMP in the class of tests in $KU^\alpha(h)$ satisfying (3.9).

We next consider problem (3.2). Let

$$KU^\alpha = \bigcap_{h \in F_2(0)} KU^\alpha(h), \quad KS^\alpha = \bigcap_{h \in F_2(0)} KS^\alpha(h) \quad (3.13)$$

denote respectively the class of conditional unbiased tests of level α in $\mathcal{D}^\alpha$ and the class of conditional similar tests of level α in $\mathcal{D}^\alpha$ appropriate for this problem. Also let U^α and S^α be respectively the class of unbiased tests of level α in $\mathcal{D}_2^\alpha$ and the class of similar tests of level α in $\mathcal{D}_2^\alpha$ defined as

$$U^\alpha = \{\phi \in \mathcal{D}_2^\alpha \mid \pi(\phi, (0, h)) \leq \alpha \text{ for any } h \in F_2(0)\} \quad (3.14)$$

$$\pi(\phi, (\mu, h)) \geq \alpha \text{ for any } h \in F_2(\mu) \text{ and } \mu \neq 0\}$$

$$S^\alpha = \{\phi \in \mathcal{D}_2^\alpha \mid \pi(\phi, (0, h)) = \alpha \text{ for any } h \in F_2(0)\}.$$

where $\pi(\phi, (\mu, h)) \equiv E[\phi(v, w)]$ is the power function of ϕ under $h \in F_2(\mu)$.

The various relationships amongst these four classes are given below.

LEMMA 3.1. (1) $KU^\alpha \subset U^\alpha$

(2) $U^\alpha \subset S^\alpha$

(3) $KS^\alpha = S^\alpha$.

Proof. (1) follows from the definitions of KU^α and U^α . (2) follows from the easily verifiable fact that for any $\phi \in \mathcal{D}_2^\alpha$ and any $h \in F_2(\mu)$, $\pi(\phi, (\mu, h))$ is continuous in μ . To show (3), note that $KS^\alpha \subset S^\alpha$ follows from the definitions. On the other hand, since $N(\mu) \equiv \{N(\mu, \sigma^2) | \sigma^2 > 0\} \subset F_2(\mu)$ and w is complete for $N(0) \subset F_2(0)$, it follows that $\pi(\phi, (0, h)) = E_0^w[E_0^v(\phi(v, w) | w)] = \alpha$ necessarily implies

$$E_0^v[\phi(v, w)] = \alpha \quad \text{a.a.}(w, h) \quad \text{for all } h \in F_2(0),$$

which shows $S^\alpha \subset KS^\alpha$. This completes the proof of the lemma.

Our main result for problem (3.2) is the following.

THEOREM 3.2. For problem (3.2), the two-sided t-test ϕ_2 is UMP in S^α i.e., UMP similar.

Proof. By Theorem 3.1 and (3) of Lemma 3.1, it is enough to show that any $\phi \in S^\alpha$ satisfies (3.9). Assuming $h \in F_2(\mu)$ is the density of $N(\mu, \sigma^2 I_n)$, it follows that when $\mu = 0$,

$$E_0^v[v\phi(v, w)] = \alpha E_0^v[v] = 0 \quad (3.15)$$

because the derivative of $\pi(\phi, (\mu, h))$ at $\mu = 0$ is zero and because w is complete for $N(0) \equiv \{N(0, \sigma^2 I_n) | \sigma^2 > 0\}$. However, by Proposition 1, (3.15) is true for any $h \in F_2(0)$. This completes the proof.

Since $\phi_2 \in U^\alpha$, by (2) of Lemma 3.1 we immediately get

COROLLARY 3.2. For problem (3.2), ϕ_2 is UMP in U^α , i.e. UMP unbiased.

4. t-TEST WITH INVARIANCE

It is well-known (Kariya (1981)) that when the density of x is given by (1.3), the one-sided t-test ϕ_1 is UMPI for testing $\mu = 0$ versus $\mu > 0$ for any fixed nonincreasing q and the two-sided t-test ϕ_2 is UMPI for testing $\mu = 0$ versus $\mu \neq 0$ for any fixed nonincreasing and convex q . The groups leaving the problems invariant are respectively given by $G_1 \equiv R_+ \times \tilde{O}(n)$ and $G_2 \equiv R_* \times \tilde{O}(n)$, where $R_+ = \{a > 0\}$, $R_* = \{a \in R | a \neq 0\}$, $\tilde{O}(n) = \{\Gamma \in O(n) | \Gamma 1 = 1\}$, and $O(n)$ is the group of $n \times n$ orthogonal matrices. Maximal invariants under G_1 and G_2 with group actions $(\gamma, \Gamma)(x) = \gamma \Gamma x$, $(\gamma, \Gamma) \in G_i$, $i = 1, 2$, are respectively v and $|v|$, and the classes of invariant level α tests are respectively given by

$$J_1^\alpha = \{\phi \in \mathcal{D}^\alpha | \phi \text{ is based on } v \text{ only, } E_0^v[\phi] \leq \alpha\} \quad (4.1)$$

$$J_2^\alpha = \{\phi \in J_1^\alpha | \phi(-v) = \phi(v)\}. \quad (4.2)$$

Since the null distribution of v does not depend on q (vide Kariya (1981)), it follows from (2.4) that

$$J_2^\alpha \subset J_1^\alpha \subset \mathcal{D}_0^\alpha \subset \mathcal{D}_1^\alpha \subset \mathcal{D}^\alpha(h), \quad h \in F_2(0). \quad (4.3)$$

In fact, the following result holds.

LEMMA 4.1. $J_1^\alpha \not\subset \mathcal{D}_0^\alpha$.

Proof. Take any test function $\psi(w) \neq \text{constant}$ and any $\phi(v) \in J_1^\alpha$. Then the test defined by $\psi(w)\phi(v) \in D_0^\alpha$ because $E_0^V[\psi(w)\phi(v)] = \psi(w)E_0^V[\phi(v)] \leq \alpha$. However, $\psi(w)\phi(v) \notin J_1^\alpha$, completing the proof.

Consequently, for the one-sided testing problem, the results in Corollary 2.1 are somewhat stronger than the result that ϕ_1 is UMPI.

An analogous result is obtained below for the two-sided testing problem. Recall the definitions of S^α and J_2^α given in (3.14) and (4.2) respectively.

LEMMA 4.2. $J_2^\alpha \not\subseteq S^\alpha$.

Proof.[#] Note from Proposition 1 that $h \in F_2(0)$, v and w are independent and the densities are respectively given by $r_0(v)$ and $r_1(w)$ where q depends on h . The conditional size condition $E_0^V[\phi(v, w)] = \alpha$ and the conditional similarity condition (3.9) can be expressed as

$$\int \phi(v, w) r_0(v) dv = \alpha \quad \text{a.a. } (w, h \in F_2(0)) \quad (4.4)$$

$$\int v \phi(v, w) r_0(v) dv = 0 \quad \text{a.a. } (w, h \in F_2(0)). \quad (4.5)$$

Now take two tests $\phi_1(v)$ and $\phi_2(v)$ based on v only which are functionally independent and satisfy (4.5) with

$$E_i = \int \phi_i(v) r_0(v) dv, \quad i = 1, 2$$

and $E_1 + E_2 = 0$. Next choose any two tests $\phi_1(w)$ and $\phi_2(w)$ based on w only which satisfy

$$\phi_1(w) + \phi_2(w) = 1$$

Finally, define the test $:\phi(v, w) = \phi_1(v)\phi_1(w) + \phi_2(v)\phi_2(w)$. Clearly $:\phi$ satisfies $E_0^V[:\phi(v, w)] = \alpha$ and $:\phi(\neq \text{constant})$ does depend nontrivially on w . This completes the proof.

[#] Clearly $J_2^\alpha \subset S^\alpha$. We will construct a test $:\phi(v, w) \in S^\alpha$ but $\notin J_2^\alpha$.

REFERENCES

- [1] Kariya, T. (1981). Robustness of multivariate tests. Ann. Statist. 9, 1267-1275.
- [2] Kariya, T. and Eaton, M.L. (1977). Robust tests for spherical symmetry. Ann. Statist. 5, 206-215.
- [3] Lehmann, E.L. (1959). Testing Statistical Hypotheses. John Wiley and Sons, New York.
- [4] Lehmann, E.L. and Stein, C. (1948). Most powerful tests of composite hypotheses. Ann. Math. Stat. 19, 495-516.
- [5] Lehmann, E.L. and Stein, C. (1949). On the theory of some nonparametric hypotheses. Ann. Math. Statist. 20, 28-45.