

AD-A177 893

ON THE VALIDITY OF BEURLING THEOREMS IN POLYDISCS(U)
NORTH CAROLINA UNIV AT CHAPEL HILL CENTER FOR
STOCHASTIC PROCESSES V MANDREKAR SEP 86 TR-154
AFOSR-TR-87-0073 F49620-85-C-0144


1/1

UNCLASSIFIED

F/G 12/1

NL





UNCLASSIFIED

②

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE


1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY NA		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for Public Release; Distribution Unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE NA			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 154		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 87-0073	
6a. NAME OF PERFORMING ORGANIZATION University of North Carolina	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION AFOSR/NM	
6c. ADDRESS (City, State and ZIP Code) Center for Stochastic Processes, Statistics Department, Phillips Hall 039-A, Chapel Hill, NC 27514		7b. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR	8b. OFFICE SYMBOL (If applicable) nm	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620 85 C 0144	
8c. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC		10. SOURCE OF FUNDING NOS.	
		PROGRAM ELEMENT NO. 6.1102F	TASK NO. 2304 A/5
11. TITLE (Include Security Classification) On the validity of Beurling theorems in polydiscs		WORK UNIT NO.	
12. PERSONAL AUTHOR(S) Mandrekar, V.			
13a. TYPE OF REPORT technical preprint.	13b. TIME COVERED FROM 8/85 TO 9/86	14. DATE OF REPORT (Yr., Mo., Day) September 1986	15. PAGE COUNT 4
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	Keywords:	
XXXXXXXXXXXXXXXXXXXX			
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
Let Z be the set of integers. We denote by m, n etc. the elements of Z. Let U be the open unit disc and T the boundary of U in the complex plane $\mathbb{C}$. Let λ, ρ, U_2 and T^2 be the respective cartesian product and σ^2 the normalized Lebesgue measure on T^2. For $p > 0$, we denote by $L^p(T^2, \sigma_2)$ the normalized Lebesgue space of the equivalence class of p-integrable functions.			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☐ DTIC USERS ☐		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL Peggy Ravitch Maj Crowley		22b. TELEPHONE NUMBER (Include Area Code) 913-962-2307 5025	22c. OFFICE SYMBOL AFOSR/NM

DTIC FILE COPY

DTIC
ELECTE
FEB 24 1987

CENTER FOR STOCHASTIC PROCESSES

Department of Statistics
University of North Carolina
Chapel Hill, North Carolina



ON THE VALIDITY OF BEURLING THEOREMS IN POLYDISCS

by

V. Mandrekar

Approved for publication by AFOSR

Technical Report No. 154

September 1986

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFOSR)
NOTICE OF TECHNICAL RESEARCH (AFOSR)
This technical report has been reviewed and is
Approved for public release (AFOSR 190-12).
Distribution is unlimited.
MATTHEW J. KERPER
Chief, Technical Information Division

ON THE VALIDITY OF BEURLING THEOREMS IN POLYDISCS

by

V. Mandrekar*

Department of Statistics & Probability
Michigan State University
East Lansing, MI 48824

and

Center for Stochastic Processes
Department of Statistics
University of North Carolina
Chapel Hill, NC 27514

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By _____	
Distribution/	
Availability Codes	
Dist _____	

A-1



*Supported in part by ONR N00014-85-K-0150 and the Air Force Office of Scientific Research Contract No. F49620 85 C 0144.

Let $\mathbb{Z}$ be the set of integers. We denote by m, n etc. the elements of $\mathbb{Z}$. Let U be the open unit disc and T the boundary of U in the complex plane $\mathbb{C}$. Let $\mathbb{Z}^2, \mathbb{C}^2, U^2$ and T^2 be the respective cartesian product and σ_2 the normalized Lebesgue measure on T^2 . For $p > 0$, we denote by $L^p(T^2, \sigma_2)$ the usual Lebesgue space of the equivalence class of p -integrable functions and $H^p(U^2) = \{f : f : U^2 \rightarrow \mathbb{C} \text{ analytic and } \sup_{0 \leq r \leq 1} \int_T |f_r(\underline{t})|^p d\sigma_2 < \infty\}$. Here $f_r(\underline{t}) = f(z)$ with $z = r\underline{t}$. Let $\underline{z} = (z_1, z_2) = (r_1 e^{i\theta_1}, r_2 e^{i\theta_2})$ and $\underline{t} = (e^{i\theta_1}, e^{i\theta_2})$, then $P(\underline{z}, \underline{t}) = P_{r_1}(\theta_1 - \theta_1) \cdot P_{r_2}(\theta_2 - \theta_2)$ is called Poisson Kernel with

$$P_r(\theta) = \frac{1-r^2}{1-2r\cos\theta+r^2}. \text{ It is known that for } f \in H^p(U^2), \lim_{r \rightarrow 1} f_r(\underline{t}) = f^*(\underline{t})$$

exists and is in $L^p(T^2, \sigma_2)$. For $f \in L^p(T^2, \sigma_2)$, let $f^e(z) = \int_{T^2} P(\underline{z}, \underline{t}) f(\underline{t}) d\sigma_2$, then $f^e \in H^p(U^2)$. In case $p = 2$, $f \in H^2(U^2)$ if $f \in L^2(T^2, \sigma_2)$ and

$f(\underline{t}) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} a_{mn} t_1^m t_2^n$ and $f^e = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} a_{mn} z_1^m z_2^n$. Conversely every $f \in H^2(U^2)$ has this form and $f^*(\underline{t}) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} a_{mn} t_1^m t_2^n$. For further information, see [4].

In [4], Rudin gives an example of a shift-invariant subspace of $H^2(T^2)$ which is not of the form $q \cdot H^2$, where q is an inner function. Our purpose here is to characterize invariant subspaces of the form qH^2 in terms of the action of the shifts on it. We note that subspaces of the form qH^2 can be represented as

$$(1) \quad qH^2 = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \oplus V_1^m V_2^n (M)$$

where M equals the span of q in $H^2(T^2)$ and V_1 is the multiplication by t_1 on $H^2(T^2)$ with $\underline{t} = (t_1, t_2) \in T^2$. It is easy to check that

$M = \{qH^2 \oplus V_1(qH^2)\} \cap \{qH^2 \oplus V_2(qH^2)\}$. As V_1 commutes with V_2 (in short, $V_1 \sim V_2$), we get from (1) and Theorem 4.1 of [2] (see also [5]) that V_1 and V_2 are doubly commuting (i.e. $V_1 \sim V_2, V_1 \sim V_2^*$). In fact, we have

2. Theorem. An invariant subspace $M \neq \{0\}$ of $H^2(T^2)$ is of the form $q \cdot H^2$ with q inner function if and only if V_1 and V_2 are doubly commuting on M .

Proof: Necessity was proved above. To prove the sufficiency we get, in view of Theorem 4.2 ((c) $\Rightarrow$ (b)) [2],

$$(3) \quad M = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \oplus V_1^m V_2^n (R_1^\perp \cap R_2^\perp),$$

where $R_1 = M \ominus V_1 M$, using Theorem 1.2 of [2]. Here we observe that

$\cap V_2^n (R_1^\perp)$, $\cap V_1^m (R_2^\perp)$ and $\cap V_1^m V_2^n (H^2(T^2))$ equal zero. Let $q_1, q_2 \in R_1^\perp \cap R_2^\perp$, then

$\int_T t_1^m t_2^n q_1 \bar{q}_2 d\sigma_2 = 0$ for $m, n > 0$. As $V_2(R_1^\perp) \subseteq (R_1^\perp)$ (Lemma 1.3 and Theorem 4.2 of [2]), we get $\int_T t_2^n q_1 \bar{t}_1^m q_2 d\sigma_2 = 0$ for all $n > 0$ and $m > 0$. Since $\bar{t}_1^m = t_1^{-m}$, by the symmetry of the problem we get for $(m, n) \neq (0, 0)$

$$\int t_1^m t_2^n q_1 \bar{q}_2 d\sigma_2 = 0.$$

Since $q_1 \bar{q}_2 \in L^1(T^2, \sigma_2)$ we get $q_1 \bar{q}_2 = c_1$ a.e. σ_2 . In particular, $|q_1|^2 = c_2$.

Hence $R_1^\perp \cap R_2^\perp$ is one dimensional. Also q generating $R_1^\perp \cap R_2^\perp$ is an inner-function. Assume $|q| = 1$ a.e. choosing q of norm 1. Now (3) gives the result.

Let $f \in H^2(T^2)$ and $M_f = \overline{\text{sp}\{V_1^m V_2^n f : m, n \geq 0\}}$ then M_f is an invariant subspace.

4. Corollary. $M_f = qH^2(T^2)$ if and only if V_1 and V_2 are doubly commuting on M_f .

Following Helson [1], we say that a function g is H -outer if $M_g = H^2(T^2)$.

5. Corollary. A function $f \in H^2(T^2)$ has the property $f = q \cdot g$ with q inner and g H -outer if and only if V_1 and V_2 doubly commute on M_f .

Proof: By Corollary 4, only if part follows as $M_f = qH^2(T^2)$. To prove the converse we note that by Corollary, $M_f = qH^2(T^2)$ giving $f = q \cdot g$, $g \in H^2(T^2)$.

Hence $M_f = q \cdot M_g$ giving g is H -outer.

In ([4], p. 72) a function $f \in H^2(U^2)$ is called outer (we call it R -outer)

if $\log|f(z)| = \int_{T^2} \log|f^*| d\sigma_2$. Given a function $f \in H^2(T^2)$, we denote by $f^e \in H^2(U^2)$ given by $\int_{T^2} P(z,t)f(\underline{t})d\sigma_2$. In this case we note that $(f^e)^* = f$.

It is already known ([4], Theorem 4.4.6) that f is H-outer then f^e is R-outer. From this we get in view of Corollary 4 the following.

6. Corollary. Let g be H-outer then g^e is R-outer and V_1 and V_2 doubly commute on M_g .

We now prove the converse of Corollary 6. Assume now that V_1, V_2 doubly commute on M_f and f^e is R-outer then by Corollaries 5 and 6, the definition of f^e, g^e and the fact that $|q| = 1$ we get $f^e = pg^e$ with $|p| = 1$. Thus we get that the slice function $f_w^e(\lambda) = p_w(\lambda)g_w^e(\lambda)$. Using Lemma 4.4.4(a) of [4] and the uniqueness of outer function ([1], p. 13) we get $p_w(\lambda) = 1$ for all w and λ giving $p = 1$ i.e., $f = g$. Combining this with Corollary 6 gives

7. Corollary. Let $f \in H^2(T^2)$ then $M_f = H^2(T^2)$ if and only if f^e is R-outer and V_1 and V_2 doubly commute.

In view of Theorem 4.2 of [2], we get that Corollary 7 includes Beurling Theorem proved in ([6], Theorem 1.5). Now using essentially classical techniques ([3],[2]) one can derive associated results in prediction theory in [6].

References

1. Helson, H. (1964) Lectures on invariant subspaces, Academic Press, N.Y.
2. Kallianpur, G. and Mandrekar, V. (1983) Non-deterministic random fields and Wold and Halmos decompositions for commuting isometries, Prediction Theory and Harmonic Analysis, (eds. Mandrekar, V. and Salehi, H.) North Holland, Amsterdam.
3. Masani, P. (1962) Shift-invariant spaces and prediction theory 107, 275-290.
4. Rudin, W. (1979) Function Theory in Polydisks, Benjamin, N.Y.
5. Slocinski, M. (1980) On Wold-type decomposition of a pair of commuting isometries, Ann. Pol. Math. 37, 255-262.
6. Soltani, A.R. (1984) Extrapolation and moving average representation for stationary random fields and Beurling's Theorem, Ann. Prob. 12, 120-132.

END

3-87

DTIC