

AD-A177 916

A SPECTROSCOPY OF HELIX MELTING(U) PURDUE UNIV
LAFAYETTE IND DEPT OF PHYSICS E M PRONOFKY 11 MAR 87
N00014-86-K-0252


1/1

UNCLASSIFIED

F/G 7/4

NL





PHOTOCOPY RESOLUTION TEST CHART

AD-A177 916

REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION U			1b RESTRICTIVE MARKINGS NA		
2a SECURITY CLASSIFICATION AUTHORITY NA			3 DISTRIBUTION/AVAILABILITY OF REPORT DISTRIBUTION UNLIMITED		
2b DECLASSIFICATION/DOWNGRADING SCHEDULE NA					
4 PERFORMING ORGANIZATION REPORT NUMBER(S) Purdue Research Foundation			5 MONITORING ORGANIZATION REPORT NUMBER(S) NA		
6a NAME OF PERFORMING ORGANIZATION Purdue Research Foundation		6b OFFICE SYMBOL (If applicable) NA	7a NAME OF MONITORING ORGANIZATION Office of Naval Research		
6c ADDRESS (City, State, and ZIP Code) W. Lafayette, IN 47907			7b ADDRESS (City, State, and ZIP Code) 800 N. Quincy St. Arlington, VA 22217-5000		
8a NAME OF FUNDING/SPONSORING ORGANIZATION Strategic Defense Initiative Organization		8b OFFICE SYMBOL (If applicable)	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-86-K-0252		
8c ADDRESS (City, State, and ZIP Code) Washington, D.C. 20301-7100			10 SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO 63222 c	PROJECT NO	TASK NO
			WORK UNIT ACCESSION NO		
11 TITLE (Include Security Classification) A Spectroscopy of Helix Melting					
12 PERSONAL AUTHOR(S) Prohofskv, Earl William					
13a TYPE OF REPORT Annual		13b TIME COVERED FROM 4/86 TO 3/87		14 DATE OF REPORT (Year Month, Day) 1987-3-11	
15 PAGE COUNT 4					
16 SUPPLEMENTARY NOTATION					
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	DNA HELIX MELTING		
			FREE ELECTRON LASER IRRADIATION		
			homopolymers polymers		
19 ABSTRACT (Continue on reverse if necessary and identify by block number) We have calculated the dependences of the melting of poly(dG)-poly(dC) on vibrational mode excitation level. We find that bands of modes in the 10-100/ cm^{-1} range can alter details of helix melting. Melting is involved in important biological processes. The frequency range of this phenomenon is that of the Santa Barbara FEL. The SBFEL is an ideal device for exploration of this biological phenomenon. We are in collaboration with Prof. S. M. Lindsay's program which is an experimental exploration of the irradiation of DNA at SBFEL. (keyword)					
DTIC FILE COPY DTIC ELECTE MAR 17 1987 S D					
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT ☐ DTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION U		
22a NAME OF RESPONSIBLE INDIVIDUAL M. Marron			22b TELEPHONE (Include Area Code) (202) 696-4038		22c OFFICE SYMBOL ONR

Annual Report

A Spectroscopy of Helix Melting

N00014-86-K-0252

E. W. Prohofsky
Department of Physics
Purdue University
West Lafayette, IN 47907

Our effort is centered on calculations of DNA helix melting and the role of vibrational excitations in such melting. Our early results indicate that vibrational modes in the frequency range from $10 - 100 \text{ cm}^{-1}$ are the dominant element in bringing about helix melting. This frequency range is the same as that covered by the Santa Barbara Free Electron Laser Facility. It is the premier source of fine tuned radiant energy in this regime. We are calculating the expected helix behavior as these modes are excited presumably by FEL irradiation. The theoretical method we have developed indicated that several details of the helix melting were controlled by the excitation level of these vibrational modes.

By melting we mean the usual definition of DNA melting i.e., breaking up of the double helix into single strand DNA. The most fundamental processes such as RNA transcription and DNA replication require this form of melting, as it is essential to the reading of the base sequence message incorporated in the DNA. In these processes a small region of DNA is melted and this melt is propagated along the DNA so that whole genes can be read. The process must be carefully controlled and this control is a likely element of control of gene expression.

In our most sophisticated theoretical model we use a DNA homopolymer, poly(dG)·poly(dC), with a single base pair melted region. We monitor the elongation or propagation of such a melt by calculating the melting of adjacent base pairs. We find that the direction of propagation of the melt is controlled by the level of excitation of specific vibrational modes. Modes at $80 - 100 \text{ cm}^{-1}$ are particularly effective at advancing the melt along the $3' \rightarrow 5'$ direction in the G backbone. Modes below 20 cm^{-1} are particularly effective at advancing the melt in the opposite direction along the double helix. Not only is the direction of melt controlled by the level of vibrational excitation, but also the way in which the melting occurs is controlled by mode excitation. Those modes which cause melting to proceed $3' \rightarrow 5'$ along G also cause melting to originate at the major groove side of the DNA helix. Those modes that elongate the melt in the opposite direction induce melting to originate at the minor groove or opposite side of the helix. Our initial results therefore indicate that details of the melting process are a strong functions of details of the vibrational mode levels of excitation. These mode excitations may be influenced by FEL irradiation. The results quoted are published in J. Biomolec. Struct. and Dyn. 4, 437-442, (1986). A reprint is enclosed with this report.

A second effort during this year was the calculation of the behavior of these $10 - 100 \text{ cm}^{-1}$ modes with temperature down to very low temperature. Experimental data on the low temperature frequency shift became available from FTIR measurements made at the Max Planck Institute in Stuttgart. We have predicted frequency shifts in good agreement with those observed. This work has been

written up in preprint form and has been submitted for publication in Phys. Rev. B.

A preprint is enclosed with this report. The agreement is an indication that our calculations and parameters do correspond well to the DNA double helix.

More progress has been made during this first year but it has been along the line of formulating and coding further calculations for which results are not yet available. We are making progress on calculating the melting behavior of a greater range of DNA base compositions so as to better compare our theoretical results to the observations on FEL irradiation when such data becomes available. We are in close collaboration with Prof. S. M. Lindsay at Arizona State University who is involved in DNA irradiation experiments at the Santa Barbara FEL.

Accession For	
NTIS CRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	



END

4-87

DTIC