

AD-A178 229



Institute Report 230

**ALTERNATIVE METHODS FOR MEASURING VESICANT
AGENT ACTION ON RABBIT SKIN**


Peter Schmid, PhD, June R. Jaeger, BS, George J. Klain,
PhD, John P. Hannon, PhD, Carol A. Bossone, MS,
Michael C. Powanda, PhD

DIVISION OF CUTANEOUS HAZARDS

20030127002

February 1987

DTIC FILE COPY



LETTERMAN ARMY INSTITUTE OF RESEARCH PRESIDIO OF SAN FRANCISCO CALIFORNIA 94129

This document has been approved
for publication and sale, its
distribution is unlimited.

Alternative methods for measuring vesicant agent action on rabbit skin--
Schmid et al.

Reproduction of this document in whole or in part is prohibited except with the permission of the Commander, Letterman Army Institute of Research, Presidio of San Francisco, California 94129. However, the Defense Technical Information Center is authorized to reproduce the document for United States Government purposes.

Destroy this report when it is no longer needed. Do not return it to the originator.

Citation of trade names in this report does not constitute an official endorsement or approval of the use of such items.

In conducting the research described in this report, the investigation adhered to the "Guide for the Care and Use of Laboratory Animals," as promulgated by the Committee on Revision of the Guide for Laboratory Animal Facilities and Care, Institute of Laboratory Animal Resources, National Research Council.

This material has been reviewed by Letterman Army Institute of Research and there is no objection to its presentation and/or publication. The opinions or assertions contained herein are the private views of the author(s) and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense. (AR 360-5)

 9 Feb 77
(Signature and date)

This document has been approved for public release and sale; its distribution is unlimited.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Institute Report No. 230	2. GOVT ACCESSION NO. A178229	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ALTERNATIVE METHODS FOR MEASURING VESICANT AGENT ACTION ON RABBIT SKIN		5. TYPE OF REPORT & PERIOD COVERED Institute Report January to June, 1985
7. AUTHOR(s) Peter Schmid, PhD, June R. Jaeger, BS, George J. Klain, PhD, John P. Hannon, PhD, Carol A. Bossone, MS, Michael C. Powanda, PhD		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Letterman Army Institute of Research Presidio of San Francisco, CA 94129		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS U.S. Medical Research and Development Command Ft. Detrick, MD 21701		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 62734A 3M162734A875 BC WU205
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE February 1987
		13. NUMBER OF PAGES 14
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS: (Continue on reverse side if necessary and identify by block number) mustard gas analog; butyl 2-chloroethyl sulfide, vesicant, rabbit, ¹²⁵ I-labelled albumin, tissue water, edema; erythema; laser Doppler, skin		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) ► Tissue damage to the skin of rabbits exposed to the vapors of butyl 2-chloroethyl sulfide was assessed by conventional means, by measuring the changes in water content of tissue, accumulation of ¹²⁵ I-labelled albumin, and by changes in the signal from a laser Doppler instrument. The content of water and ¹²⁵ I-labelled albumin in tissue significantly increases after exposure. Extravasation of red cells, as measured by a ⁵¹ Cr-labelling technique, appears to be low, indicating that, under the conditions of the experiment, damage to the capillaries appears to be not very important.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ABSTRACT

Tissue damage to the skin of rabbits exposed to the vapors of butyl 2-chloroethyl sulfide was assessed by conventional means, by measuring the changes in water content of tissue, accumulation of ^{125}I -labeled albumin, and by changes in the signal from a laser Doppler instrument. The content of water and ^{125}I -labeled albumin in tissue significantly increases after exposure. Extravasation of red cells, as measured by a ^{51}Cr -labeling technique, appears to be low, indicating that, under the conditions of the experiment, damage to the capillaries appears to be not very important.

Alternative Methods for Measuring Vesicant Agent Action on Rabbit Skin--Schmid

INTRODUCTION

The use of vesicants has long been considered an exceedingly effective means of denying territory and materials to combatants and support personnel who are not prepared to protect themselves against, or to decontaminate themselves after being exposed to, this class of chemical warfare agents. Sulfur mustard (HD) is especially insidious because skin decontamination is unlikely to be effective unless instigated within 10 to 15 minutes after the initial exposure to this vesicant (1).

Recent evidence indicates that changes in cutaneous cellular metabolism occur within a few minutes after exposure to the reputedly less toxic monofunctional sulfur mustard, n-butyl 2-chloroethyl sulfide (BCS), an analog of HD (2), although there may be no visible signs of physiological manifestations on humans for several hours, or days, after exposure to bifunctional mustard (HD). Vesicant action on animals' skin is manifested as edema and erythema, rather than as confluent blisters as seen on human skin.

It is not yet clear how the early biochemical manifestations after exposure to mustard are related to the clinical evidence, i.e., edema, erythema and skin damage, but these early metabolic changes are consistent with rapid fixation of mustard to the skin. Although the early metabolic alterations may prove useful for determining the efficacy of skin-protection formulations, classically, edema and erythema have been used to assess the irritant/injury potential of a variety of compounds, including mustard. Unfortunately, visual observation provides only semi-quantitative data, requires a highly skilled observer, and may result in significant laboratory-to-laboratory variations in the results (3-5). We therefore have undertaken the task to develop methods which will allow objective, user-independent, quantitative analyses of edema and erythema.

Schmid--2

METHODS

Five female New Zealand rabbits, weighing approximately 3-kg, were shaven 24 hrs before the experiment. Each animal was anesthetized with one ml ketamine, injected IM, and maintained with additional injections of 0.5 ml, when needed. Five 19-mm-in-diameter glass cups containing filter paper circles impregnated with 10 μ l (6.55 μ moles) neat BCS (Fig. 1a) were attached by means of double-stick discs to one side of the spine of the shaved dorsum of each anesthetized rabbit for 30 minutes (Fig. 1b). Glass cups containing filter paper, but no BCS, were attached on the opposite side of the spine as controls. At the end of 3 hrs, 125 I-labeled albumin and 51 Cr-labeled erythrocytes were injected intravenously and allowed to circulate and equilibrate for 1 hr. At the end of 4 hrs, a sample of blood was drawn; the rabbits were then euthanized with an overdose of T-61[®] (American Hoechst Corp.), and a large portion of the dorsal skin was removed and mounted on Saran[®]-wrapped cardboard to reduce shrinkage of excised skin (Fig. 1c). The circular sites of application of vesicant and the control site(s) were cut out with a motor-driven boring machine for rubber/cork stoppers equipped with a 22-mm cutting tube (Fig. 1d). Each circular tissue was then placed in a liquid scintillation uncapped vial, weighed immediately, and the vial containing the skin was placed in a vacuum oven, heated to 60°C and evacuated. The samples were removed and weighed periodically on a Mettler balance, Model 440, until a constant weight was attained. Twenty milliliters of 30% aqueous potassium hydroxide were then added to the tissue and heated at 90°C to dissolve the skin. The dpm were counted in a dual-channel gamma isotope counter to assess accumulation of 125 I and 51 Cr.

Two female 3-kg New Zealand rabbits were used to test the relative protection afforded by one layer of battle dress uniform (BDU) or Cortex[®] fabric against BCS. The experimental design was the same as above, except that a layer of fabric was inserted between the BCS-containing filter paper and the rabbit's skin (Fig. 2). On each of the rabbits, we used one layer of fabric per site. Five of the sites on one side were controls, five on the other side contained 10 μ l of neat BCS. At the end of 3 hrs, 125 I-labeled albumin was injected intravenously. No 51 Cr-labeled erythrocytes were used in this test. Harvesting of the skin samples and analyses were performed as described in the preceding paragraph.

A laser Doppler flowmeter (Med Pacific Corp., Seattle, WA) was used to measure alterations in the skin microcirculation of two female 3-kg New Zealand rabbits after their skin was treated with BCS. The experimental procedure used was similar to that described in the first paragraph under METHODS, except no isotopes were injected into either rabbit. An extra 30 minutes, beyond the 30-minute period of exposure to BCS, were allowed to elapse to prevent contamination of the probe of the flowmeter by any BCS residue remaining on the skin surface. The animals were not sacrificed, but were kept anesthetized until the laser Doppler readings had been taken. Triplicate blood flow measurements were made for a duration of 150 seconds each time at each experimental site. The animals were kept alive and unanesthetized overnight with no apparent discomfort. They had free access to food and water. The microcirculation at the skin reaction sites and controls were again measured in triplicate the following morning after the rabbits were anesthetized with 1 ml ketamine IM. The animals were euthanized with T-61[®] immediately after the 24-hr readings. The laser Doppler flowmeter was calibrated and zeroed at the beginning of each experiment. A special holder was attached to the skin with double-sided sticky tape while each site was measured.

RESULTS

The mean water content of treated and untreated skin sites on the rabbits, expressed as a percentage of the total weight of the excised skin, is a measure of the amount of edema. The mean water content varied from 77.9 to 84.4 for vesicant-exposed sites and from 66.0 to 73.1 for control sites (Table 1). Each value in the table represents the mean from five sites. Although there are expected variations from animal to animal, the data clearly show an appreciably higher water content in the BCS sample than in the control sample for the five animals.

Schmid--4

Table 1. Comparison of the Mean Water Content¹ of BCS-treated and Untreated Skin Sites of Each Rabbit

Rabbit No.	Vesicant		Control	
	Mean % ²	S.D. ³	Mean % ²	S.D. ³
1	78.0	3.12	66.0	1.58
2	77.9	1.79	70.9	1.42
3	81.3	0.32	72.9	0.87
4	80.4	1.20	72.5	1.08
5	84.4	1.68	73.1	0.58

¹percentage of the weight of excised skin

²average of five sites on each animal

³standard deviation. N = 5

When the water content data for specific sites (vesicant and control) for the five animals were averaged (Table 2) little variation was found between different sites; however, an appreciably higher water content was found in the BCS-treated sites than in the control sites.

Table 2. Comparison of Site Variation Among Rabbits as Determined by the Mean Water Content¹ of BCS-treated and Untreated Sites

Site No.	Vesicant ²		Control ²	
	Mean % ³	S.D. ⁴	Mean % ³	S.D. ⁴
1	80.9	0.97	72.6	2.61
2	81.0	2.19	71.6	2.61
3	81.4	3.01	70.9	2.05
4	79.5	4.05	70.5	2.60
5	80.5	3.54	69.5	4.23

¹percentage of weight of excised skin

²treated with vesicant on the left side; no treatment on the right-hand side of the spine

³average of five rabbits per each corresponding site

⁴standard deviation. N = 5

Deposition of ^{125}I -labelled albumin and ^{51}Cr -labelled erythrocytes, another measurement of edema and erythema, is shown for vesicant and control sites for albumin only. There were no differences in ^{51}Cr deposition, despite the presence of erythema. The ^{125}I -labelled albumin values are between 3 and 9 times higher in vesicant-treated than in control sites, with an inter-animal variation of the order of 2, from the least to the most responsive animal, when exposed to BCS (Table 3).

Table 3. Comparison of the Deposition of ^{125}I -labelled Albumin¹ at Vesicant and Control Sites of Each Rabbit

Rabbit No.	Vesicant		Control	
	Mean ²	S.D. ³	Mean ²	S.D. ³
1	731	124	96	15
2	428	98	131	10
3	444	53	72	8
4	593	133	107	20
5	990	142	106	16

¹dpm/min/site (22-mm-diameter piece of excised skin)

²average of five sites on each animal

³standard deviation. N = 5

Visual observations of skin reactions (6) on the five rabbits were made 4 hrs post application of butyl mustard and were scored according to the Draize system as described in the footnotes of Table 4 and 5. Control sites showed a value of zero for edema and erythema.

Schmid--6

Table 4. Evaluation of Skin Edema of Rabbits 4 Hours After Butyl Mustard Application Using Draize Scoring System¹

Rabbit No.	Site No.				
	1	2	3	4	5
1	4	4	4	4	4
2	4	3	3	2	2
3	4	4	4	4	4
4	3	3	3	3	3
5	4	4	4	4	4

¹1 = Very slight edema (barely perceptible)

2 = Slight edema (edges of area well defined by definite raising)

3 = Moderate edema (area raised approximately 1 mm)

4 = Severe edema (raised more than 1 mm and extending beyond area of exposure)

Table 5. Evaluation of Skin Erythema of Rabbits 4 Hours After Butyl Mustard Application Using Draize Scoring System¹

Rabbit No.	Site No.				
	1	2	3	4	5
1	2	2	2	1	1
2	3	3	2	1	1
3	1	1	1	1	1
4	3	3	3	3	3
5	2	2	2	2	2

¹1 = Very slight erythema

2 = Well defined erythema

3 = Moderate to severe erythema

4 = Severe erythema (beet redness) to slight eschar formation (injuries in depth)

Despite the limited data, it appears that both BDU fabric and Gortex® provide some protection against BCS vapor, as shown in Table 6.

Table 6. Effect of Fabric in Protecting Skin Against Butyl Mustard (BCS)

Site Treatment	Percent Water Content		¹²⁵ I-labeled Albumin (dpm/site)	
	Mean ¹	S.D. ²	Mean ¹	S.D. ²
Control	72.0	1.4	153	98
Vesicant alone	79.9	3.2	600	205
Vesicant + BDU (0.55 mm thick)	77.4	2.8	249	36
Vesicant + Gortex® (0.45 mm thick)	75.8	2.4	169	54

¹average of four sites (2 on each animal)

²standard deviation. N = 2

Cutaneous microcirculation was also measured with a laser Doppler flowmeter. Fig. 3 illustrates preliminary data from one of two experiments using two different rabbits. The output, the flow value, is given in millivolts: the larger the value, the greater the blood flow (9). The values shown are mean values from one rabbit for 5 control and 5 experimental sites. Within 30 minutes after exposure to BCS, the experimental sites display an almost twofold increase in signal, which persists for at least five hours.

DISCUSSION

The objectives of this pilot study were twofold: (a) to develop a system to quantitate measurements of edema and erythema, and (b) to provide an assay system for assessing the efficacy of formulations that protect the skin against vesicants, particularly, sulfur mustards. The rabbit was chosen as the animal model for our tests because its skin often shows a stronger response to mild

and moderate irritants than does human skin, thereby providing an additional margin of safety (6).

Since edema is the infiltration of predominantly albumin-containing fluid into the site of irritation (7), we measured both the water content of tissue and the accumulation of ^{125}I -labeled albumin. Measurements of the accumulation of ^{125}I -labeled albumin and water content of the tissues provide very sensitive indices of edema. Water content, especially, appears to reflect both intracellular and extravascular fluid accumulation.

Erythema is redness produced by congestion of the capillaries; ^{51}Cr -labeled erythrocytes were used as a measure of capillary stasis and/or damage to capillaries and extravasation of red cells. Under the conditions of our experiments, it appears that very little extravasation occurs, since the amount of ^{51}Cr at skin sites exposed to BCS is not significantly different from that of the control skin. Perhaps by using a higher radioactivity of ^{51}Cr to label the red blood cells, a small amount of activity might be detected at the BCS site; however, Issekutz (8) found that the labeling of erythrocytes with ^{51}Cr does not provide a sensitive measurement of erythema.

After the isotope experiments, erythema was measured by laser Doppler velocimetry. The preliminary data from this instrumentation (Fig. 3) suggest that it may be a useful non-invasive tool to measure changes in microvascular blood flow (9) associated with mustard damage to skin.

Negligible variation in mean water content at various sites on the rabbit skin (Table 2) indicated that, for any given rabbit, there are no large differences in response to butyl mustard, i.e., values close to the shoulder are not different from values close to the hind leg.

Variations in skin response from one rabbit to another are caused by a myriad of factors, i.e., differences in anesthesia, ambient temperature, and differences in stages of the hair growth cycle which affects capillary blood flow. Some of these variations can be minimized through careful control of experimental conditions.

We evaluated the relative protection afforded by a single thickness of battle dress uniform (BDU) fabric and Gortex® material against butyl mustard (BMS). Single layers of swatches of each cloth were placed between the cups containing vesicant and the rabbit's skin for the duration of the exposure (Fig. 2). As a positive control, the skin was exposed to vesicant with no intervening cloth. Despite limited data, it can be seen that both BDU fabric and Gortex® provide some protection against butyl mustard vapor. Conversely, these data also demonstrate that this system can be used to evaluate not only formulations, but also cloth, for their ability to protect against vesicants. These data also emphasize the value of measuring both water content of tissue and accumulation of ^{125}I -labeled albumin for quantitating edema resulting from mustard injury to animals' skin.

REFERENCES

1. Smith HW, Clowes GHA, Marshall EK. On dichloroethylsulfide (mustard gas). IV. The mechanism of absorption by the skin. *J Pharmacol* 1919;13:1-30.
2. Klain GJ, Rowanda MC, Schmid P, Black KE. Metabolic markers of chemically-induced cutaneous irritation/injury. Army Science Conference, West Point, NY, 1984.
3. Draize JH, Woodward G, Calvery HO. Methods for the study of irritation and toxicity of substances applied topically to the skin and mucous membranes. *J Pharmacol Exp Ther* 1944;82:377-419.
4. Weil CS, Scala RA. Study of intra- and interlaboratory variability in results of rabbit eye and skin irritation tests. *Toxicol Appl Pharmacol* 1971;19:276-360.
5. Phillips L, Steinberg M, Milbach HL, Akers WA. A comparison of rabbit and human skin response to certain irritants. *Toxicol Appl Pharmacol* 1972;21:369-382.

Schmid--10

6. Code of Federal Regulations, 1976, Title 16, part 1500.41.
7. Uchida K, Takeuchi Y, Movat HZ. Simple method of quantitation of enhanced vascular permeability. Proc Soc Exp Biol Med 1970;133:1384-1387.
8. Issekutz AC. Quantitation of acute inflammation in the skin: Recent methodological advances and their application to the study of inflammatory reactions. Surv Synth Pathol Res 1983;1:89-110.
9. Nilsson GE, Tenland T, Oberg PA. Evaluation of a laser Doppler flowmeter for measurement of tissue blood flow. IEEE Trans Biomed Eng 1980;27:597-604.




Figure 1. Glass cap containing viscant (top left). Caps attached to dorsal aspect of an anesthetized rabbit (bottom left). Excised skin being attached to saran-wrapped cardboard (top right). Skin sample removed with electric cork borer (bottom right).

Schmid--12





Figure 2. Top: Vesicant-containing cups with BDU (left) and Cortex (right) switches attached. Bottom: Vesicant-containing cups attached to the dorsal aspect of an anesthetized rabbit.

LASER LIGHT PATH SCHEMATIC



LASER DOPPLER FLOWMETRY




Figure 3. Top: Diagram of the laser light path. Bottom: Bar graph representing toxic reaction of rabbit skin with 5 control and 5 vesicant-exposed sites.

Schmid et al--14

OFFICIAL DISTRIBUTION LIST

Commander
US Army Medical Research
and Development Command
ATTN: SGRD-RMS/Mrs. Madigan
Fort Detrick, MD 21701-5012

Defense Technical Information Center
ATTN: DTIC/DDAB (2 copies)
Cameron Station
Alexandria, VA 22304-6145

Office of Under Secretary of Defense
Research and Engineering
ATTN: R&AT (F&LS), Room 3D129
The Pentagon
Washington, DC 20301-3080

The Surgeon General
ATTN: DASG-FLO
Washington, DC 20310

HQ DA (DASG-ZXA)
WASH DC 20310-2300

Commandant
Academy of Health Sciences
US Army
ATTN: HSHA-CDM
Fort Sam Houston, TX 78234-6100

Uniformed Services University
of Health Sciences
Office of Grants Management
4301 Jones Bridge Road
Bethesda, MD 20814-4709

US Army Research Office
ATTN: Chemical and Biological
Sciences Division
PO Box 12211
Research Triangle Park, NC 27709-2211

Director
ATTN: SGRD-UWZ-L
Walter Reed Army Institute
of Research
Washington, DC 20307-5100

Commander
US Army Medical Research Institute
of Infectious Diseases
ATTN: SGRD-ULZ-A
Fort Detrick, MD 21701-5011

Commander
US Army Medical Bioengineering
Research & Development Laboratory
ATTN: SGRD-UBG-M
Fort Detrick, Bldg 568
Frederick, MD 21701-5010

Commander
US Army Medical Bioengineering
Research & Development Laboratory
ATTN: Library
Fort Detrick, Bldg 568
Frederick, MD 21701-5010

Commander
US Army Research Institute
of Environmental Medicine
ATTN: SGRD-UE-RSA
Kansas Street
Natick, MA 01760-5007

Commander
US Army Institute of Surgical Research
Fort Sam Houston, TX 78234-6200

Commander
US Army Research Institute
of Chemical Defense
ATTN: SGRD-UV-AJ
Aberdeen Proving Ground, MD 21010-5425

Commander
US Army Aeromedical Research Laboratory
Fort Rucker, AL 36362-5000

AIR FORCE Office of Scientific
Research (NL)
Building 410, Room A217
Bolling Air Force Base, DC 20332-6448

Commander
USAFSAM/TSZ
Brooks Air Force Base, TX 78235-5000

Head, Biological Sciences Division
OFFICE OF NAVAL RESEARCH
800 North Quince Street
Arlington, VA 22217-5000