

AD-A178 430

EFFECT OF AEROBIC TRAINING ON THE PLASMA ACTH RESPONSE
TO EXERCISE (U) NAVAL HEALTH RESEARCH CENTER SAN DIEGO
CA M J BUONO ET AL. 01 SEP 86 NAUHLTHRSCHC-86-28

1/1


UNCLASSIFIED

I/C 6/14

ML

[REDACTED]

END
DATE
FILMED
4 87



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963-A

AD-A178 430

2

EFFECT OF AEROBIC TRAINING ON THE PLASMA ACTH RESPONSE TO EXERCISE

M. J. BUONO
J. E. YEAGER
A. SUCEC

DTIC
ELECTE
MAR 16 1987
S D

REPORT NO. 86-28



DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

NAVAL HEALTH RESEARCH CENTER

P.O. BOX 85122
SAN DIEGO, CALIFORNIA 92138-9174

NAVAL MEDICAL RESEARCH AND DEVELOPMENT COMMAND
BETHESDA, MARYLAND

87 3 13 116

FILE COPY

EFFECT OF AEROBIC TRAINING ON THE PLASMA ACTH
RESPONSE TO EXERCISE

by

Michael J. Buono*

John E. Yeager**

Anthony A. Sucec*

*San Diego State University Heart Institute
Departments of Biology and Physical Education
San Diego, California
and

**Naval Health Research Center
P.O. Box 85122
San Diego, California



Accession For	
NTIS CRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Report No. 86-28, supported by the Naval Medical Research and Development Command, Department of the Navy, under Research Work Unit N40764.B995.AB-087. The views expressed in this article are those of the authors and do not reflect the official policy or position of the Department of the Navy, Department of Defense, nor the U.S. Government.

Summary

A previous study has shown that plasma adrenocorticotropin (ACTH) secretion during submaximal exercise is increased following training. This finding is in contrast to studies of other hormones such as glucagon, epinephrine and growth hormone, all of which show decreases following training. This study re-examines the effect of training on plasma ACTH levels during exercise. Ten adult volunteers were split into a control and an exercise group. The exercise group underwent a 12 week training program consisting of running 3.5 kilometers per day, three days per week. Before and after the training program all subjects were tested for plasma ACTH response to a 150 Watt workload for 15 minutes. The experimental group showed an 11% increase in Max Vo_2 following training while the control group's Max Vo_2 was unchanged. The experimental group's ACTH response to the submaximal workload was decreased 63% ($p < 0.05$), while the control group's response was unchanged. These data suggest that the ACTH response to exercise is blunted following training. This response is similar to that of other hormones such as glucagon, epinephrine and growth hormone. Such a finding is in agreement with previous studies that have shown that ACTH follows an intensity dependent pattern during exercise.

Introduction

A review of the literature (3,6) reveals that only one study (1) has previously measured plasma adrenocorticotropin hormone (ACTH) during exercise in humans before and after a physical training program. The results of that study (1) suggest that training augments the secretion of ACTH in response to a sub-maximal exercise bout.

Such a finding is interesting in light of two facts. First, most studies report that exercise training usually results in reduced plasma levels of glucagon, epinephrine, norepinephrine, and growth hormone (3,6) in response to the same absolute workload. Thus, the study by Carr et al (1) is not in agreement with these previous findings. Second, most studies (6) report that at the same sub-maximal workload trained individuals have a smaller rise in plasma cortisol than do untrained individuals. Since cortisol secretion is under the control of ACTH, it is difficult to explain how training decreases the cortisol response to exercise while at the same time increasing plasma ACTH levels. Therefore, in light of the above discussion, it was the purpose of this study to re-examine the effect of training on plasma ACTH levels during exercise.

Methods

The subjects for this study were 10 adult volunteers (4 male, 6 female). Written informed consent was obtained from each subject prior to testing.

The 10 subjects were split into a control group and an experimental group, each of which contained 2 males and 3 females. Each subject reported to the laboratory on 4 separate occasions. On the first visit each subject performed a graded treadmill run to exhaustion. Oxygen uptake was measured each min of the run using open-circuit spirometry (Alpha-Technologies, System 4400, Laguna Hills, CA) with the highest value being recorded as maximal O_2 uptake (VO_2 max).

On the second visit, the subjects rested in a seated position for 10 min, following which a 10 ml venous blood sample was obtained from an antecubital vein. Each subject then exercised on a Godart cycle ergometer at 150W for 15 min. A second venous blood sample was obtained immediately upon completion of the exercise bout.

Following the second lab visit the subjects in the experimental group started a 12-week exercise training program. The subjects ran 3.5 km a day, three days a week for the 12 week period. The subjects' exercise heart rate during the training runs was held between 70-85% of their previously determined maximal heart rate. The control group did not participate in the organized exercise sessions. Furthermore, they were instructed not to change their exercise habits over the course of the study.

Following the 12 week training program all of the subjects reported to the laboratory for their third lab visit and again performed a graded treadmill run to exhaustion. On their fourth visit

they repeated the sub-maximal exercise protocol that they performed on their second visit. The second and fourth lab visits were held at approximately the same time of day for each subject.

All of the collected blood samples were measured, in duplicate, for plasma ACTH using the RIA procedures described by Orth (11). Intra-assay coefficients of variation were 3 and 5% respectively, at 80 and 20% B/B₀. The working standard ACTH reference preparation for the assay was synthetic human ACTH (1-39). The sensitivity of the assay was 5 pg/ml.

Independent t-tests were used to test for differences between the two groups. Paired t-tests were used to examine the effect of training on VO₂ max and ACTH. Significance was set at the p < 0.05 level.

Results

The mean ($\pm$ S.E.) descriptive and fitness variables for the two groups are shown in Table 1. As can be seen, there was no significant difference between the two groups in age, ht, wt, or pre-training VO₂ max. The training program, however, produced a significant increase in VO₂ max in the experimental group, while the control group did not change significantly over the 12 week period. The mean 11% increase in VO₂ max in the experimental group is similar in magnitude to previously reported training studies (12).

Table 1

Descriptive Variables for the Experimental and Control Groups Before and After Training					
	Age (Yrs)	HT(cm)	Wt(kg)	VO2Max pre(Lmin ⁻¹)	VO2Max post(Lmin ⁻¹)
Experimental (n=5)	22.1 $\pm$ 0.4	165.6 $\pm$ 1.3	67.6 $\pm$ 1.5	2.98 $\pm$.28	3.20 $\pm$.36*
Control (n=5)	26.6 $\pm$ 0.8	168.0 $\pm$ 1.1	61.7 $\pm$ 2.0	2.70 $\pm$.24	2.78 $\pm$.23

Values are mean $\pm$ S.E.

*pre- vs post-training values are significantly (p<0.05) different

The exercise induced plasma ACTH response for both groups is shown in Fig. 1. As can be seen, the experimental group demonstrated a significant reduction in the delta ACTH response to the 150W exercise bout following the training program. Delta ACTH is simply the post-exercise ACTH value minus the pre-exercise value, and represents the increase in plasma ACTH that resulted from the exercise bout. The control group did not show a significant difference in the delta ACTH response to the exercise bout over the course of the study.




Fig. 1. Mean ($\pm$ S.E.) delta ACTH responses for the two groups before and after the training program.

Discussion

The most important finding of the current study is that the ACTH response to sub-maximal exercise is blunted following a training program. This finding suggests that plasma ACTH responds in a similar fashion to training when compared to many other hormones (3,6). The results of this study, however, are in conflict with the previously published work of Carr et al (1). They found that ACTH levels rose 63% during exercise prior to training and by 116% after two months of aerobic conditioning. A closer examination of their paper, however, reveals that the workloads used to elicit the ACTH response increased from a mean of 70W to 100W during the course of the study. From this data it can be estimated that their subjects would have had to have increased their $\dot{V}O_2$ max over the course of the study 32% to have made the 70W and 100W workloads of equal relative intensity. As suggested by Galbo (6), it seems more likely that the workloads over the course of the study made up an increasing percentage of the subjects $\dot{V}O_2$ max. If the above assumption is correct then the data of Carr et al (1) is consistent with the findings of Farrell et al (4) who have reported that ACTH increases during exercise in an intensity dependent pattern.

In the current study the 150W workload represented a mean ($\pm$ S.E.) of $73\pm 6\%$ of the experimental groups pre-training VO_2 max but only $66\pm 7\%$ of their post-training VO_2 max. These results also support the previously determined intensity dependent pattern of ACTH during exercise (4).

The mechanism for the blunted ACTH response to exercise following training can only be speculated on at this time. However, at least 3 potential explanations seem worthy of discussion. First, previous studies (9) have shown that arginine vasopressin (AVP) can potentiate the effect of hypothalamic corticotropin releasing factor (CRF) on ACTH secretion. Interestingly, it has been reported (2) that training reduces the AVP response to sub-maximal exercise. Thus, the decreased ACTH response to exercise following training may be secondary to a decrease in AVP.

Second, previous investigators (4,5) have suggested that disturbances in intracellular homeostasis during exercise may be sensed by peripheral chemoreceptors which stimulate ACTH secretion. Such a hypothesis is consistent with the results of the current study, since it is known that following training the intracellular homeostasis is less disturbed, as evidenced by smaller changes in muscle lactate and phosphagen levels (10) at the same absolute workload.

Lastly, several animal studies at least suggest that cardiovascular dynamics during exercise may act as a stimulus to ACTH release (8). In the current study the mean ($\pm$ S.E.) exercise heart rate in the experimental group during the 150W workload did decrease significantly from 154 ± 13 bpm to 146 ± 13 bpm. However, there was not a significant correlation ($r = -.034$, NS) in the experimental group between the delta ACTH and delta exercise heart rate responses pre- to post-training. Such a finding is in agreement with others (4,7) who also did not demonstrate a significant relationship between heart rate and ACTH.

In conclusion, the current study shows that the ACTH response to a sub-maximal workload is blunted following training. Such a finding is in agreement with previous work that has shown that ACTH follows an intensity dependent pattern during exercises.

References

1. Carr, D.B., B.A. Bullen, G.S. Skirnar, M.A. Arnold, M. Rosenblatt, I.Z. Beitins, J.B. Martin, and J.W. McArthur. Physical conditioning facilitates the exercise-induced secretion of beta-endorphin and beta-lipotropin in women. New Engl. J. Med. 305:560-563, 1981.
2. Convertino, V.A., L.C. Keil, and J.E. Greenleaf. Plasma volume, renin, and vasopressin responses to graded exercise after training. J. Appl. Physiol. 54:508-514, 1983.
3. Farrell, P.A. Adrenocorticotrophic hormone and exercise. In: Exercise Endocrinology, edited by K. Fotherby and S. Pal. New York: de Gruyter, 1985, pp. 139-156.
4. Farrell, P.A., T.L. Garthwaite, and A.B. Gustafson. Plasma adrenocorticotropin and cortisol responses to sub-maximal and exhaustive exercise. J. Appl. Physiol. 55:1441-1444, 1983.
5. Few, J.D., C.C. Cashmore, and G. Terton. Adrenocortical response to one-leg and two-leg exercise on a bicycle ergometer. Eur. J. Appl. Physiol. 44:167-174, 1980.
6. Galbo, H. Hormonal and metabolic adaptation to exercise. New York: Thieme-Straton, 1983, pp. 46-51.
7. Gambert, S.R., T.L. Garthwaite, C.H. Pontser, E.E. Cook, F. Tristani, E.B. Duthie, D.R. Martinson, T.C. Hange, and D.J. McCarty. Running elevates plasma beta-endorphin immunoreactivity and ACTH in untrained human subjects. Proc. Soc. Exp. Biol. Med. 168:1-4, 1981.
8. Cann, D.S., D.G. Ward, and D.E. Carlson. Neural pathways controlling release of corticotropin (ACTH). In: Interaction within the Brain-Pituitary-Adrenocortical System, edited by M.T. Jones, New York: Academic Press, 1979, pp. 74-86.
9. Jones, M.T. Control of adrenocortical hormone secretion. In: The Adrenal Gland, edited by V. James, New York: Raven Press, 1979, pp. 93-130.
10. Karlsson, J., L. Nurdesjo, L. Jorfeldt, and B. Saltin. Muscle lactate, ATP, and CP levels during exercise after physical training in man. J. Appl. Physiol. 33:199-203, 1972.
11. Orth, D.N. Adrenocorticotrophic hormone (ACTH). In: Methods of Hormone Radioimmunoassay, edited by B.M. Jaffe and H.R. Behrman. New York: Academic Press, 1979, pp. 245-284.
12. Pollock, M. The quantification of endurance training programs. In: Exercise and Sport Sciences Reviews, Vol. 1, edited by J.H. Wilmore, New York: Academic Press, 1973, pp. 155-188.

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE

A178 430

REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION Unclassified			1b RESTRICTIVE MARKINGS None		
2a SECURITY CLASSIFICATION AUTHORITY N/A			3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.		
2b DECLASSIFICATION/DOWNGRADING SCHEDULE N/A					
4 PERFORMING ORGANIZATION REPORT NUMBER(S) NHRC Report No. 86-28			5 MONITORING ORGANIZATION REPORT NUMBER(S)		
6a NAME OF PERFORMING ORGANIZATION Naval Health Research Center		6b OFFICE SYMBOL (If applicable) Code 60		7a NAME OF MONITORING ORGANIZATION Commander, Naval Medical Command	
6c ADDRESS (City, State, and ZIP Code) P. O. Box 85122 San Diego, CA 92138-9174			7b ADDRESS (City, State, and ZIP Code) Dept of the Navy Washington, DC 20372-5120		
8a NAME OF FUNDING/SPONSORING ORGANIZATION Naval Medical Research & Development Command		8b OFFICE SYMBOL (If applicable)		9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c ADDRESS (City, State, and ZIP Code) NMCNCR Bethesda, MD 20814-5044			10 SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO 63764A	PROJECT NO 3M463764B	TASK NO 995.AB-087
			WORK UNIT ACCESSION NO. DA307899		
11 TITLE (Include Security Classification) (U) EFFECT OF AEROBIC TRAINING ON THE PLASMA ACTH RESPONSE TO EXERCISE					
12 PERSONAL AUTHOR(S) Buono, Michael J. **; Yeager, John E., Sucec, Anthony					
13a TYPE OF REPORT Interim		13b TIME COVERED FROM _____ TO _____		14 DATE OF REPORT (Year, Month, Day) 1986 September 1	
15 PAGE COUNT					
16 SUPPLEMENTARY NOTATION **San Diego State University Heart Institute, Depts of Biology & Physical Education, San Diego, California					
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Exertion;		
			Cycle ergometry;		
			Adrenocorticotropin.		
19 ABSTRACT (Continue on reverse if necessary and identify by block number) The purpose of this study was to re-examine the effect of training on plasma ACTH levels during exercise. Ten adult volunteers were split into a control and an experimental group. The experimental group participated in a 12-week training program which resulted in a significant 11% increase in their mean maximal oxygen uptake. The plasma ACTH response to a 150W workload was measured in both groups before and after the training program. The experimental group demonstrated a significant reduction in the delta ACTH response to the workload, while the control group demonstrated an unchanged response over the course of the study. These data suggest that the ACTH response to an absolute, submaximal workload is blunted following training. Such a finding is in agreement with previous studies that have shown that ACTH follows an intensity dependent pattern during exercise. R4 200812					
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT ☐ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT ☐ DTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a NAME OF RESPONSIBLE INDIVIDUAL John Yeager			22b TELEPHONE (Include Area Code) (619) 225-6671		22c OFFICE SYMBOL Code 60

ATE
LMED
-8