

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY N/A since UNCLASSIFIED			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE N/A since UNCLASSIFIED					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) JSR-85-405			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION MITRE Corp. JASON Program Office		6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION Defense Advanced Research Projects Agency	
6c. ADDRESS (City, State, and ZIP Code) 7525 Colshire Dr. McLean, VA 22102			7b. ADDRESS (City, State, and ZIP Code) 1400 Wilson Blvd. Arlington, VA 22209		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Director Defense Nuclear Agency		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F19628-86-C-0001	
8c. ADDRESS (City, State, and ZIP Code) Washington, DC 20305-1000			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
			WORK UNIT ACCESSION NO.		
11. TITLE (Include Security Classification) Early-Time HANE Simulation and Experiment					
12. PERSONAL AUTHOR(S) Cornwall, M.; Hammer, D; LeLevier, R.; Vesecky, J.					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM TO		14. DATE OF REPORT (Year, Month, Day) 870415	
15. PAGE COUNT					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
18	3		HANE High Altitude Bursts JASON Report Nuclear Explosives Nuclear Weapons Communications		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
This report, the sixth in a series on this subject, focuses on the NRL laser facility, new experimental results from it, and computer models for the experiments. The NRL facility continues to be upgraded in experimental and diagnostic capability in a sensible and commendable manner. The experimental plan is good and the experiments will continue to be powerful tools for understanding debris-air coupling processes for early-time. This tool is very appropriate to the collisional regime. We anticipate it to be a substantial contribution to the understanding of collisionless coupling also, both with and without magnetic fields. It remains very important to stay focused on the critical question of early-time debris-air coupling related to late-time spatial structures. We recommend cataloging collisionless phenomena for consideration while continuing to perform collisional experiments. Early-time shadowgraphy with radiating ambient gasses should be considered. While aneurysms remain an interesting phenomena, we caution that they					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT ☐ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL SANDRA E. YOUNG			22b. TELEPHONE (Include Area Code) (202) 325-1079		22c. OFFICE SYMBOL DNA/CSTI

19. ABSTRACT (Continued)

may not be related to the primary HANEs explorations. We recommend consideration of blunt-ended probes as a supplement to spherical obstacles in plasma flow experiments. We recommend the use of a fast multichannel device after the spectrometer and the consideration of two-color interferometry as diagnostic tools. We recommend the purposeful exploration of target variations through the deliberate use of spherical or hemispherical and other non-flat targets of planned inhomogeneity. Finally, we recommend increased correspondence between experimental simulations and the HANE data base. We look forward to two-dimensional modeling efforts and suggest the consideration of special purpose machines or parallel machines before investing in more conventional computers for such modeling.

18. SUBJECT TERMS (Continued)


Ionosphere

Electromagnetic Pulse

JSR-85-405

LIBRARY
RESEARCH REPORTS DIVISION
NAVAL POSTGRADUATE SCHOOL
MONTEREY, CALIFORNIA 93940

Early-Time HANE Simulation and Experiment



MITRE

//

Corporation.

Early-Time HANE Simulation and Experiment

M. Cornwall
D. Hammer
R. LeLevier
J. Vesecky

May 1987

JSR-85-405

Approved for public release, distribution unlimited

JASON
The MITRE Corporation
7525 Colshire Drive
McLean, Virginia 22102-3481

ABSTRACT

This report, the sixth in a series on this subject, focuses on the NRL laser facility, new experimental results from it, and computer models for the experiments. The NRL facility continues to be upgraded in experimental and diagnostic capability in a sensible and commendable manner. The experimental plan is good and the experiments will continue to be powerful tools for understanding debris-air coupling processes for early-time. This tool is very appropriate to the collisional regime. We anticipate it to be a substantial contribution to the understanding of collisionless coupling also, both with and without magnetic fields. It remains very important to stay focussed on the critical question of early-time debris-air coupling related to late-time spatial structures.

We recommend cataloging collisionless phenomena for consideration while continuing to perform collisional experiments. Early-time shadowgraphy with radiating ambient gasses should be considered. While aneurysms remain an interesting phenomena, we caution that they may not be related to the primary HANES explorations. We recommend consideration of blunt-ended probes as a supplement to spherical obstacles in plasma flow experiments. We recommend the use of a fast multichannel device after the spectrometer and the consideration of two-color interferometry as diagnostic tools. We recommend the purposeful exploration of target variations through the deliberate use of spherical or hemispherical and other non-flat targets of planned inhomogeneity. Finally, we recommend increased correspondence between experimental simulations and the HANE data base. We look forward to two-dimensional modeling efforts and suggest the consideration of special purpose machines or parallel machines before investing in more conventional computers for such modeling.

TABLE OF CONTENTS

	<u>Page</u>
1.0 INTRODUCTION AND SUMMARY	1-1
2.0 THE NRL EXPERIMENT	2-1
2.1 Coupling	2-2
2.2 Aneurysms	2-3
2.3 Obstacles	2-4
2.4 Diagnostic Development	2-4
2.5 Target Variations	2-5
2.6 HANE Simulation	2-7
3.0 SUPPORTING THEORY AND COMPUTER SIMULATIONS	3-1
DISTRIBUTION LIST	D-1

LIST OF ILLUSTRATIONS

<u>Table</u>		<u>Page</u>
1-1	EARLY-TIME HANE SIMULATION ISSUES EXPERIMENTAL	1-4
1-2	EARLY-TIME HANE SIMULATION ISSUES THEORY AND COMPUTATION	1-5

1.0 INTRODUCTION AND SUMMARY

In 1985 JASON conducted its sixth summer study on the effects of high altitude nuclear explosives (HANEs) on the ionosphere and ionospheric communications. This year we concentrated on the upgraded NRL laser facility, new experimental results coming from it, and computer simulations designed to model various aspects of the experiment. We also heard briefings on the connection of certain phenomena, observed in STARFISH and CHECKMATE, to the early-time experimental results.

During the five years of JASON involvement in the DNA HANE program, many of the important physics issues have been identified and discussed at length in our earlier reports, so we forego detailed elaboration of the basic physics problems. This year's brief report will focus on current and projected work - both experimental and theoretical, including simulations - designed to address these physics issues.

Our general impression is that the NRL laser experiments are and will continue to be very powerful tools for understanding debris-air coupling processes for a certain part of the time history of the early-time regime. The accompanying theory and simulations, though presently incapable of dealing with all of the relevant physics and

chemistry, are well-matched to the present level of experimental understanding of the collisional regime ($p > 1.5$ Torr). There is much hope that straightforward continuation of the present efforts (including careful consideration of finite UV radiation lifetimes in the coupling process) will lead to a broad understanding of collisional coupling, except possibly for very early times where details of the target-laser interaction are important and at times too large for the debris to be contained in the plasma chamber. However, this understanding will require the use of new diagnostics, some of them planned for the future but not yet in routine use. These include multi-spectral shear interferometry, possibly coupled with Faraday rotation measurements, and the coherent Thomson-scattering laser probe, possibly operated with a streak camera. Obstacles in the way of the blast wave may also be useful in the collisional regime, although these will be more interesting for collisionless shocks.

Both the level of experimental effort and the level of theoretical understanding are lower in the collisionless regime. It is evident that coupling diminishes as the ambient pressure goes down, in general accord with what simulations predict. But the influence of, and mechanisms for generating, magnetic fields are not well understood, nor is it clear why aneurysms do not form when

collisions are reduced. Answering these questions will require more sophisticated two-dimensional hybrid codes.

Since very early-time laser-target interactions could very well lead to persistent structure effects at late times, it is important to study the widest possible array of target geometries, both experimentally and with simulations. Of special interest are back-illuminated targets, and targets designed to mimic CHECKMATE, which may well have produced many debris fragments that, though vaporized, maintained their identity for long times. This possibility, long espoused by Walt Chesnut, was dealt with in some detail in last year's JASON report on HANEs.

These and other issues are summarized in Tables 1-1 and 1-2 and the rest of the report supplies supporting detail. Our overall impression of the early-time HANE program continues to be good. We can envisage substantial understanding of both collisional and collisionless coupling, both with and without magnetic fields, emerging from this program. But the workers in the program must always keep one question foremost in their minds: How is early-time debris-air coupling ultimately related to late-time spatial structures? Answering this question is a formidable task.

TABLE 1-1

EARLY-TIME HANE SIMULATION ISSUES
EXPERIMENTAL

<u>COLLISIONAL</u> ($p > 1.5$ Torr) DEBRIS-AIR COUPLING: AFTER 1 EQUAL-MASS RADIUS, FORMS NORMAL (RAYLEIGH-TAYLOR STABLE) TAYLOR- SEDOV BLAST WAVE. IS IT R-T UNSTABLE EARLIER? WHAT'S INSIDE? ANEURYSMS: BULGES ON BLAST WAVE. FORMED HOW? WHEN? DECAY? IMPACT ON LATE-TIME STRUCTURE? RELATED TO LASER BEAM IN AMBIENT GAS? TO LASER-TARGET INTERACTION?	<u>COLLISIONLESS</u> ($p < 1.5$ Torr) DEBRIS COUPLES POORLY TO AIR: RUNS AHEAD OF BLAST WAVE; DEBRIS ELECTRONS FORM MAGNETIC PULSE. SOME AIR IONS REFLECTED ($V = 2V_D$) NO ANEURYSMS: WHY NOT?
DEBRIS-TARGET INTERACTION: NEED TO STUDY A VARIETY OF TARGETS (BACK-ILLUMINATED; DOUBLE FOIL; CAVITY; FRAGMENT-RICH) UV PROBLEM: RADIATION LIFE TIMES NOT SHORT COMPARED TO DEBRIS TIME SCALES (UNLIKE REAL HANE). USE Xe OR MRC GAS AS AMBIENT. DIAGNOSTICS: MULTI-COLOR SHEARING INTERFEROMETRY: COHERENT THOMSON SCATTER; FARADAY ROTATION; OBSTACLES (FORM COLLISIONLESS SHOCK?)	

TABLE 1-2

EARLY-TIME HANE SIMULATION ISSUES
THEORY AND COMPUTATION

PRESENT EARLY-TIME CODES: 1D HYBRID (FLUID ELECTRONS, USUALLY MASSLESS). DO NOT ADDRESS STRUCTURE ISSUES. WELL-MATCHED TO CURRENT EXPERIMENTS EXCEPT FOR UV PROBLEM, ANEURYSMS.

NEW CODES NEEDED: 2D HYBRID. MAY REQUIRE SPECIAL-PURPOSE ARRAY PROCESSORS. WILL ADDRESS: SPONTANEOUS MAGNETIC FIELDS, SOME ASPECTS OF STRUCTURE FORMATION (ANEURYSM?).

ALSO, CODES WITH FINITE UV LIFETIMES, RADIATION TRANSPORT.

NEED MORE LASER-TARGET SIMULATIONS WITH A VARIETY OF TARGETS.

RELATION OF DEBRIS-AIR COUPLING TO STRUCTURE FORMATION: IS THERE A CLASSICAL R-T INSTABILITY AT ~ 1 EQUAL-MASS RADIUS? DOES IT HEAL LATER (CONDITIONS CHANGE TO STABLE)?

FOR COLLISIONLESS CASE, DOES ANOMALOUS RESISTIVITY EXIST? DOES IT AFFECT STRUCTURE FORMATION?

LESSONS FROM SPACE PHYSICS: ANOMALOUS RESISTIVITY NOT WORKING WELL FOR AURORAS; BOW SHOCK; TAIL NEUTRAL LINE. INERTIAL EFFECTS AND SPATIAL GRADIENTS MORE IMPORTANT. NEED TO RE-THINK NRL-3076 VIEWPOINT.

2.0 THE NRL EXPERIMENT

In this section, we shall briefly present a few thoughts on the HANE simulation experiments underway at NRL. We shall not attempt to summarize them here since there is now plenty of written literature by the NRL group that does that. In general, we feel that the experiments have gone well so far. We especially like the way the NRL group accepts suggestions for experiments from outsiders, and the way they continually expand their diagnostic capability and improve existing diagnostics. With regard to the future, we believe that it is important for the experimentalists to take advantage of the full range of capability of their new apparatus as soon as possible. By this, we mean that as soon as the new diagnostics and the magnetic field coil are checked out, both collisionless and collisional interaction experiments should be run under a variety of target, laser, and magnetic field conditions. We believe that it is as important to begin cataloging collisionless phenomena for consideration by the High Altitude Working Group as it is to complete the studies of the collisional regime. Finally, before going on to specific comments and suggestions on the experiment, we wish to observe that the active role being played by Cliff Prettie in interferometry data analysis and by Bob Stellingwerf in working with the NRL experimentalists on production of narrow ion velocity distributions is a very good

programmatic development. We hope that more of this active involvement by theorists and computationalists with the experimentalists will occur in the future.

2.1 Coupling

In the absence of a magnetic field, the efficiency of coupling of the laser-produced debris plasma to the ambient medium seems to be well explained on the basis of classical collisional processes. However, the physics under which the laser-ablation plasma expanding into the ambient gas and plasma turns into a blast wave could use additional attention along the lines of the work presented in the May 1985, Early Time Working Group meeting at NRL by Lee, Molander and Elton of NRL. Very early time, $t < 40$ ns, shadowgraphy may also provide useful information. Perhaps early time shadowgraphy with radiating ambient gases such as the MRC mixture or just Xe would reveal an unstable debris/air interface.

At pressures too low for collisional coupling to be effective, ≤ 200 mTorr, there is evidence for weak collisionless coupling only with a transverse magnetic field present in the magnetic probe data as well as in the time-of-flight data, but the field strength and experimental volume to properly address collisionless coupling are only now becoming available. In the collisionless regime, we can

expect to require measurements at larger distances and later times than in collisional experiments, and also along the magnetic field lines. All in all, we can expect a variety of interesting phenomena, some of which will probably not be anticipated, and which will no doubt take all of NRL's diagnostic capability and a major effort by the entire Early Time Working Group to understand. We are looking forward to seeing the first results soon.

2.2 Aneurysms

The aneurysm remains as mysterious as when first seen. Is it actually caused directly by the laser? Is it the last stage of an instability that occurs between the laser-ablation plasma and the ambient medium? If so, when does it form? Do self-generated magnetic fields play a role? Is it the opposite--the beginning phase of the break up of the blast wave into a turbulent front? In this case, some later time shadowgraphs are needed. What is its relationship to the classical Raleigh-Taylor instability, if any? These are all interesting questions, and experiments to obtain detailed plasma conditions in aneurysms, or designed to test hypotheses on the origin and nature of aneurysms are to be encouraged. However, we are concerned that the aneurysm is an artifact of collisional laser-disk target ablation experiments, and has no relationship to HANEs.

Unless the new, uniform profile laser beam impinging upon the inside of a curved target produces a debris plasma which forms an aneurysm, we wonder if it is in DNA's best interest to continue to expend so much effort on this phenomenon.

2.3 Obstacles

The use of spherical obstacles to determine plasma flow properties behind the blast wave, as discussed in a briefing to JASON by Guilian, was very interesting. It suggested to us that the same thing might be done in collisionless experiments to determine at least qualitatively very quickly if there is a dissipation mechanism that allows a stationary bow shock to form. We expect that dark field shadowgraphy using the tunable dye laser will readily show up the existence of such a shock, while it will be a very difficult job to determine exactly what the dissipation mechanism is. Therefore, it makes sense to do the former before expanding the large effort necessary to do the latter. Finally, we note that it may even be possible to use a properly aligned, blunt ended probe (e.g., magnetic probe) as the obstacle.

2.4 Diagnostic Development

NRL is in the process of developing several new diagnostic methods and improving others to be able to better diagnose

collisionless interactions and conditions in and behind the blast wave in collisional interactions. We are looking forward to results from both coherent and incoherent Thomson scattering, from resonant shadowgraphy and interferometry, etc. As discussed during the NRL presentations, we suggest that more detailed information can be obtained from Thomson scattering by the use of a fast multichannel device after the spectrometer (e.g., a microchannel plate streak camera, such as the Hamamatsu system, or an optical multichannel analyzer if no temporal dispersion of the scattered light is desired). In addition, the idea of 2 color interferometry in temporally separated probe laser beams, also discussed at the JASON presentations, in order to obtain interferograms at two times, should be seriously considered for implementation. Presumably the mirror needed to separate the two beams spatially after they pass through the interaction region is easy enough to obtain, although we recognize that delivering both doubled and a tripled probe beams to the interaction volume will not be so simple.

2.5 Target Variations

Firstly, we reiterate our suggestion in last year's report that spherical, or perhaps hemispherical, targets should be irradiated from the inside instead of continuing to emphasize flat targets.

Secondly, we suggest attacking the question of structure with target variation. Irregular structure in laboratory simulations of HANE's can arise from plasma instabilities even when the illuminating laser beam and target are optimized to yield a very uniform initial expansion. Irregularities presumably will also arise from inhomogeneities introduced at the very beginning of the laser-target interaction. We recommend that this second cause of irregular structure be purposely explored. In particular we encourage the use of a wide variety of inhomogeneous targets. Target irregularities could include variations in constituents, density and shape. Target illumination and probe radiation aspect angles could also be varied. The idea behind these irregular targets is to simulate a nuclear device which disassembles asymmetrically, e.g., CHECKMATE. The relevant question is, do these irregular targets produce irregular structure in the expansion after the energy release, e.g., jets, aneurysms, turbulence? If so, does the structure heal or grow? A further thought is the possibility of making a nuclear device which produces enhanced structure and hence enhanced nuclear effects, e.g., on satellite communication links. Target shape irregularities are a fertile field for invention. Targets with holes have been tried. Some further suggestions are targets with highly variable thickness, either random or quasi-periodic with constant or varying scale sizes. The idea here is to produce a highly irregular debris cloud,

even breaking off chunks of the target. Targets with randomly rough or wavy surfaces, bosses, holes nearly completely through the target, etc., are possibilities. Density and constituent inhomogeneities could be accomplished by having blobs of heavier or lighter material or voids inside the target material, like a plum pudding.

2.6 HANE Simulation

Ultimately the NRL experiments are intended to help us understand HANE phenomena that are inadequately explained. The new NRL apparatus should make quite reasonable HANE simulation experiments possible in the near future. Therefore, we suggest that certain conditions should be selected on the basis of their relevance to a particular HANE and studied in detail from early time ($t < 20$ ns) to late time (few 100 ns) in a series of shots with all possible diagnostics brought to bear. The goal would be to put together as complete a picture of that case as possible for in-depth consideration by interested parties, including anyone outside the Early Time Working Group who might be interested. For example, NRL shot number 14823 at 200 mT may have had weak coupling that occurred before $t = 40$ ns because the debris speed (375 km/s) is lower than the vacuum debris speed (450 km/s) under the same laser-target conditions. Was there some anomalous slowing down before

$t = 40$ ns? Furthermore, at 200 mT coupling may be some sort of collisional-collisionless hybrid. STARFISH data exhibited aspects of both collisional and collisionless coupling, and the magnetic precursor and the following debris blast wave in NRL shot 14823 are somewhat reminiscent of STARFISH data. Therefore, the conditions of that shot might be appropriate for intensive study.

3.0 SUPPORTING THEORY AND COMPUTER SIMULATIONS

Most of the phenomena of HANEs, as well as the NRL experiments are beyond analytic understanding so computer simulations are important. However, present-day simulations of the collisionless regime run the risk of begging one or both of two critical questions: 1) How does collisionless coupling take place? 2) How is spatial structure generated?

The first question is begged in one standard procedure using MHD (multi-fluid) theory, with plasma-kinetic phenomena modeled by anomalous transport coefficients, in the spirit of Lampe, Manheimer, and Papadopoulos (NRL-3076). In principle, such an approach would be capable of dealing with formation of spatial structure (such as shocks aneurysms), but one would not be at all sure that the coupling had been described correctly. This sort of modified MHD approach is widely used in modeling magnetospheric phenomena (Earth's bow shock, magnetic tail, etc.) and does indeed show spatial structure. However, there are three important problems of magnetospheric physics where the very concept of anomalous resistivity has been seriously questioned, and where plasma-kinetic effects are better understood as resulting from finite inertia and transit-time effects. These are the bow shock, the neutral line in the tail, and the aurora. (We would not be surprised to see the analysis of AMPTE comet formation

also not yield to a simple anomalous-resistivity approach.) In all these cases specific inertial effects have been found which account for the observed phenomena, while anomalous-resistive effects have been difficult or impossible to justify. In the aurora, for example, there is no lack of plasma turbulence or means for generating the turbulence through instabilities, but nearly all the turbulent modes have electric fields $\underline{E}$ which are very nearly perpendicular to the earth's magnetic field $\underline{B}$, so that they are ineffective in impeding motion along $\underline{B}$ as anomalous resistivity would demand. Instead, inertia provides for a finite transit time $\sim \ell/V$, where ℓ is the length of an auroral field line and V a typical particle velocity. To some extent the inverse time V/ℓ plays the role of a "collision" frequency, although there are no actual collisions. Careful examination of inertial effects show that they do not even lead to constitutive equations of the usual form $\underline{J} = \underline{\sigma} \cdot \underline{E}$. For the aurora, one finds $J_{||} \sim (Ne^2/mV) (\phi_E - \phi_I)$ where $J_{||}$ is the current along $\underline{B}$, $\phi_{E,I}$ is the electrostatic potential at the equator and at the ionosphere on the field line, and the rest of the symbols have their usual meaning. Similar modifications are found for the bow shock and neutral line problems.

When finite inertia effects are involved, macroscopic length scales play an important role in determining the effective collision

tensor $\vec{\sigma}$ (or its modification), and one cannot specify $\vec{\sigma}$ on the basis of microscopic homogeneous plasma physics, as in Lampe et al. It may be possible in the near future to construct a scheme similar in spirit to that of Lampe et al., in which anomalous resistivity is, as appropriate, replaced or modified by finite-inertia and transit-time effects depending on macroscopic length scales, and it would then be useful to run MHD codes incorporating this modified scheme. Spatial structures would emerge as a self-consistent result of balancing MHD and plasma-kinetic responses. An example of this is already found in the auroral problem, where this balance correctly predicts the overall latitudinal width of an auroral arc.

The second question, how is spatial structure formed, tends to be begged in hybrid simulations that use fluid (usually massless) electrons and particle ions. This type of simulation fully exercises the capacity of large modern computers, and it is very good at dealing with collisionless coupling, but not very good at dealing with a spectrum of spatial scales, whether externally-imposed or self-consistently generated. The hybrid simulations presented at this year's HANEs briefings were one-dimensional, but we are informed that two-dimensional codes are in the offing, perhaps a year or so away. These 2D codes will be indispensable for understanding intrinsically two-dimensional phenomena such as spontaneous magnetic field

generation and aneurysms, and we strongly urge that they be developed with an eye toward handling structure formation as well as possible. Some thought should be given to the possibility of building a special-purpose array processor that might have unusual powers in this direction. While these machines are very fast, they often have the disadvantage of requiring a relatively long time to program. One such machine is the massively parallel processor (MPP) at NASA Goddard Space Flight Center. This machine contains a few hundred microprocessors that run concurrently. The MPP has the advantage that it is currently in use for similar problems and a user community exists. It may well be that special purpose computers are not cost effective for the task at hand. However, this is the time to consider them, i.e. before resources are already expended on conventional computers. Use of a special purpose machine for lab HANE modelling would provide DNA with a sample of what such machines are capable. This experience would then provide DNA with some idea of their application to other DNA problems with high computational loads.

We have already said that laser-target interactions may profoundly affect early-time structure formation, and continuation of computer simulations in this area will be vital.

Finally, we comment on the role of simulations in the collisional regime, where collisional MHD plus chemistry, radiation, and charge exchange are the important ingredients. Present codes should be modified to incorporate finite UV radiation lifetimes (unimportant in real HANEs, but important for the NRL experiment which is over in less than 1 μ sec). In a typical present-day code, UV radiation is either instantaneous or it can be turned off altogether, and the latter seems to fit experimental data better. Presumably an even better fit would come from properly incorporating finite radiative lifetimes.

DISTRIBUTION LIST

Mr. Saul Amarel
DARPA
1400 Wilson Blvd.
Arlington, VA 22209

Wahab Ali
Naval Research Laboratory
Code: 4700
4555 Overlook Avenue, S.W.
Washington, DC 20375-5000

Russel Armstrong
Mission Research Corporation
One Tara Blvd., Suite 302
Nashua, NH 03062

Dr. Marv Atkins [3]
Deputy Director, Science & Tech.
Defense Nuclear Agency
Washington, D.C. 20305

LTCOL Joseph Bailey
Office of Science & Technology
Policy
Room 5026
New Executive Office Building
17th & Pennsylvania Avenue
Washington, DC 20506

Bob Bigoni
Mission Research Corporation
Drawer 719
Santa Barbara, CA 93102-0719

Steve Bodner
Naval Research Center
Code: 4730
4555 Overlook Avenue, S.W.
Washington, DC 20375-5000

Joe Borovsji
Los Alamos National Lab.
Los Alamos, NM 87545

Walter Chestnut
SRI International
333 Ravenswood Avenue
Menlo Park, CA 92038

Dr. Ron Clark
DARPA/NTO
1400 Wilson Blvd.
Arlington, VA 22209

Dr. John M. Cornwall
University of California
Los Angeles, CA

Glen Dahlbacha
Plasma Research Corporation
2990 Main Street
Alameda, CA 94501

Mr. John Darrah
Sr. Scientist and Technical
Advisor
HQ Space Cmd/XPN
Peterson AFB, CO 80914

**Defense Technical Information [2]
Center
Cameron Station
Alexandria, VA 22314**

RADM Craig E. Dorman
Director, ASW Programs
SPAWAR/PD/80
The Pentagon, Room 5D616
Washington, D.C. 20363-5100

DISTRIBUTION LIST (Continued)

Dr. Robert C. Duncan
DARPA
1400 Wilson Boulevard
Arlington, VA 22209

Mr. John Entzminger
DARPA
1400 Wilson Blvd.
Arlington, VA 22209

H. Carl Fitz
Physical Research, Inc.
134 Holiday court, Suite 309
Annapolis, MD 21401

Mr. Robert Foord [2]
P.O. Box 1925
Washington, D.C. 20505

Peter Gary
Los Alamos National Lab.
Los Alamos, NM 87545

Dr. Larry Gershwin
NIO for Strategic Programs
P.O. Box 1925
Washington, D.C. 20505

The Honorable Richard P. Godwin
Under Secretary of Defense for
Acquisition
The Pentagon, Room 3E1006
Washington, D.C. 20301

Dr. S. William Gouse, Z605
Senior Vice President
and General Manager
The MITRE Corporation
Metrek Division
7525 Colshire Drive
McLean, VA 22102

J. Giuliani
Naval Research Laboratory
Code: 4780
4555 Overlook Avenue, S.W.
Washington, DC 20375-5000

Alan Gregersen
Mission Research Corporation
Drawer 719
Santa Barbara, CA 93102-0719

Jacob Grun
Naval Research Laboratory
Code: 4732
4555 Overlook Avenue, S.W.
Washington, DC 20375-5000

Dr. David A. Hammer
Cornell University
Ithaca, NY

Mr. Jack Hammond
Strategic Defense Initiative
Organization
1717 H. Street
Washington, D.C. 20301

Dr. William Happer
Princeton University
Princeton, NJ 08540

Dr. Edward Harper [2]
SSBN, Security Director
OP-021T
The Pentagon, Room 4D534
Washington, D.C. 20350

Mark Hausman
Mission Research Corporation
Drawer 719
Santa Barbara, CA 93102-0719

DISTRIBUTION LIST (Continued)

Dr. Robert Henderson, W385
Director, JASON Program Office
The MITRE Corporation
7525 Colshire Drive
McLean, VA 22102

Mr. R. Evan Hineman
Deputy Director for Science
& Technology
P.O. Box 1925
Washington, D.C. 20505

Joe Huba
Naval Research Laboratory
Code: 4780
4555 Overlook Avenue, S.W.
Washington, DC 20375-5000

CDR Earl R. Jones
NAVOPIINTCEN
Attn: 06 Dept
4301 Suitland Road
Suitland, MD 20390-5170

Mike Keskinen
Naval Research Laboratory
Code: 4786
4555 Overlook Avenue, S.W.
Washington, DC 20375-5000

Mr. Edwin L. Key
Senior Vice President
for Research and Engineering
The MITRE Corporation
P.O. Box 208
Bedford, MA 01730

Ralph Kilb
Mission Research Corporation
Drawer 719
Santa Barbara, CA 93102-0719

MAJGEN Donald L. Lamberson
Assistant Deputy Chief of Staff
(RD&A) HQ USAF/RD, Rm. 4E334
Washington, D.C. 20330

Dr. Robert E. LeLevier
EOS Incorporated
Pacific Palisades, CA

The MITRE Corporation [5]
7525 Colshire Drive
McLean, VA 22102
ATTN: JASON Library, W002

Conrad Longmire
Mission Research Laboratory
Drawer 719
Santa Barbara, CA 93102-0719

John Lyon
Naval Research Laboratory
Code: 4780
4555 Overlook Avenue, S.W.
Washington, DC 20375-5000

Dr. Gordon MacDonald, Z605
Vice President
and Chief Scientist
The MITRE Corporation
7525 Colshire Drive
McLean, VA 22102

Director [2]
National Security Agency
Fort Meade, MD 20755
ATTN: Mr. Richard Madden, R6

Mr. Charles Mandelbaum
Mail Stop ER-32/G-226 GTN
U.S. Department of Energy
Washington, D.C. 20545

DISTRIBUTION LIST (Continued)

Charles Manka
Naval Research Laboratory
Code: 4732
4555 Overlook Avenue, S.W.
Washington, DC 20375-5000

Ted Mazurek
Mission Research Corporation
Drawer 719
Santa Barbara, CA 93102-0719

Ed McLean
Naval Research Laboratory
Code: 4732
4555 Overlook Avenue, S.W.
Washington, DC 20375-5000

Dr. Al Mense, Chief Scientist
(Acting)
Strategic Defense Initiative
Organization
The Pentagon, Room 3C444
Washington, D.C. 20301

Mr. Mewson
HQ SAC/NRI
Offutt AFB
Nebraska 68113-5001

Dr. Marvin Moss [2]
Technical Director
Office of Naval Research
800 N. Quincy Street
Arlington, VA 22217

Andrew Mostovych
Naval Research Laboratory
Code: 4732
4555 Overlook Avenue, S.W.
Washington, DC 20375-5000

Dr. Julian Nall [2]
P.O. Box 1925
Washington, D.C. 20505

Superintendent (Code 1424)
Naval Postgraduate School
Monterey, CA 93943
Attn: Documents Librarian

Director
National Security Agency
Fort Meade, MD 20755
ATTN: Mr. Edward P. Neuburg
DDR-FANX III

Dr. Robert Norwood [2]
Office of the Deputy Under
Secretary of the Army
Assistant Under Secretary
of the Army
The Pentagon, Room 2E653
Washington, D.C. 20310-0102

BG Malcolm O'Neil
Strategic Defense Initiative
Organization
1717 H. Street
Washington, D.C. 20301

Sid Ossakow
Naval Research Laboratory
Code: 4700
4555 Overlook Avenue, S.W.
Washington, DC 20375-5000

Dennis Papadopoulos
Science Applications Internat.
Corporation
1710 goodridge Drive
McLean, VA 22102

C. Pawley
 Naval Research Laboratory
 Code: 4732
 4555 Overlook Avenue, S.W.
 Washington, DC 20375-5000

T. Payser
 Naval Research Laboratory
 Code: 4732
 4555 Overlook Avenue, S.W.
 Washington, DC 20375-5000

J. Pierre
 Defense Nuclear Agency
 RAAE
 Washington, DC 20305

Major Donald Ponikvar
 U.S. Army
 Strategic Defense Command
 P. O. Box 15280
 Arlington, VA 22215-0150

B. Prasad
 Defense Nuclear Agency
 RAAE
 Washington, DC 20305

Mr. John Rausch
 NAVOPINTCEN Detachment, Suitland
 4301 Suitland Road
 Washington, D.C. 20390

The MITRE Corporation
 Records Resources
 Mail Stop W971
 McLean, VA 22102

Dr. Richard Reynolds
 DARPA
 1400 Wilson Blvd.
 Arlington, VA 22209

Barrie Ripin
 Naval Research Laboratory
 Code: 4732
 4555 Overlook Avenue, S.W.
 Washington, DC 20375-5000

Mr. Alan J. Roberts, Z605
 Senior Vice President
 and General Manager
 Washington C³I Operations
 The MITRE Corporation
 7525 Colshire Drive
 McLean, VA 22102

Mr. Thomas Rona
 Deputy Director
 Office of Science & Tech. Policy
 Old Executive Office Building
 17th & Pennsylvania Ave., N.W.
 Washington, D.C. 20500

Dale Sappenfield
 Los Alamos National Lab.
 Los Alamos, NM 87545

Dr. Phil Selwyn [2]
 Technical Director
 Office of Naval Technology
 800 N. Quincy Street
 Arlington, VA 22217

Mr. Shen Shey
 DARPA
 1400 Wilson Blvd.
 Arlington, VA 22209

Dr. Gordon Smith
 Strategic Defense Initiative
 Organization
 1717 H. Street
 Washington, D.C. 20301

DISTRIBUTION LIST (Continued)

Dr. Joel A. Snow [2]
Director
Science & Tech. Staff
U.S. DOE/ER-6
Washington, D.C. 20585

Dave Sowle
Mission Research Corporation
Drawer 719
Santa Barbara, CA 93102-0719

Jack Sperling
JAYCOR
P.O. Box 85154
San Diego, CA 92138

John Stamper
Naval Research Laboratory
Code: 4732
4555 Overlook Avenue, S.W.
Washington, D.C. 20375-5000

Bob Stellingwerf
Mission Research Corporation
1720 Randolph Road SE
Albuquerque, NM 87106

COMO William O. Studeman
Director of Naval Intelligence
Office of Naval Intelligence
Navy Department (OP-009)
Washington, D.C. 20310

Mr. Alexander J. Tachmindji, Z605
Senior Vice President
The MITRE Corporation
7525 Colshire Drive
McLean, VA 22102

Dr. Vigdor Teplitz
ACDA
320 21st Street, N.W.
Room 4484
Washington, D.C. 20451

Bob Thompson
Austin Research Associates
1901 Rutland Avenue
Austin, TX 78758

Bill White
Mission Research Laboratory
Drawer 719
Santa Barbara, CA 93102-0719

Dan Winske
Los Alamos National Lab.
Los Alamos, NM 87545

Leon A. Wittwer
Defense Nuclear Agency
RAAE
Washington, DC 20305

Joe Workman
Berkeley Research Assoc.
P.O. Box 241
Berkeley, CA 94701

Mr. Leo Young
OUSDRE (R&AT)
The Pentagon, Room 3D1067
Washington, D.C. 20301-3081

Steve Zalesak
Naval Research Laboratory
Code: 4780
4555 Overlook Avenue, S.W.
Washington, DC 20375-5000

Mr. Charles A. Zraket
President
The MITRE Corporation
P.O. Box 208
Bedford, MA 01730

U229851