

AD-R182 516


TORSIONAL MOTION IN AROMATIC MOLECULES CONFORMATIONAL
ANALYSIS OF METHYL- (U) COLORADO STATE UNIV FORT
COLLINS DEPT OF CHEMISTRY P J BREEN ET AL MAY 87
TR-38 N00014-79-C-8647 F/G 28/5

1/1

UNCLASSIFIED

NL

END
8-87
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

8c. ADDRESS (City, State, and ZIP Code)			10 SOURCE OF FUNDING NUMBERS			
CHEMISTRY BRANCH ARLINGTON, VIRGINIA 22217			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) "Torsional Motion in Aromatic Molecules. Conformational Analysis of Methyl-, Ethyl, and n-Propylbenzenes"						
12. PERSONAL AUTHOR(S) P.J. Breen, J.A. Warren, E.R. Bernstein and J.I. Seeman						
13a. TYPE OF REPORT Technical Report		13b. TIME COVERED FROM TO		14. DATE OF REPORT (Year, Month, Day)		15. PAGE COUNT 10
16. SUPPLEMENTARY NOTATION The view, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position policy, or decision, unless so designated by other documentation.						
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)			
FIELD	GROUP	SUB-GROUP				
19. ABSTRACT (Continue on reverse if necessary and identify by block number)						
See attached Abstract Page DTIC ELECTE JUL 10 1987 S D E						
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED			
22a. NAME OF RESPONSIBLE INDIVIDUAL Elliot R. Bernstein			22b. TELEPHONE (Include Area Code) 303-491-6347		22c. OFFICE SYMBOL	

OFFICE OF NAVAL RESEARCH

CONTRACT N00014-79-C-0647

TECHNICAL REPORT #30

"TORSIONAL MOTION IN AROMATIC MOLECULES -
XYLENES AND N-PROPYL TOLUENES"

by

P.J. Breen, J.A. Warren, E.R. Bernstein and J.I. Seeman

Prepared for Publication
in the
Journal of the American Chemical Society

Department of Chemistry
Colorado State University
Fort Collins, Colorado 80523

May 1987

Reproduction in whole or in part is permitted for
any purpose of the United State Government.

This document has been approved for public release
and sale; its distribution is unlimited.

ABSTRACT

Supersonic molecular jet spectroscopy is used to obtain information on the torsional motion in methyl, ethyl and n-propyl substituted benzenes. Dispersed emission and time of flight mass spectral results are evaluated for toluene, the three xylenes, 1,3-diethylbenzene and 3-n-propyltoluene. The aromatic ring methyl groups in m- and p-xylene can be considered as independent free rotors in S_0 and S_1 while in o-xylene they are considerably more hindered in these two states and for S_1 cross kinetic and potential terms must be introduced. The two conformations of 1,3-diethylbenzene (syn and anti, with the ethyl groups perpendicular to the plane of the benzene ring) and the three conformations of the propyl group in 3-n-propyltoluene (anti, syn-gauche and anti-gauche) are observed.

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	




To an ever increasing extent, information regarding torsional vibrations of organic molecules is being provided by theoretical treatments rather than by experimental observation.¹ When experimental data are available, all too frequently they pertain to mole fraction-weighted averages of the individual contributing conformations rather than the individual conformations themselves.² We now demonstrate that supersonic molecular jet spectroscopy can provide a novel means (a) to determine experimental values for torsional potential barriers in S_0 and S_1 , (b) to "freeze out" molecular conformations which have very low (ca. 1-5 kcal mol⁻¹) free energy barriers between themselves, (c) to "count" the number of stable ground state conformations, and by doing so, establish their molecular geometries, and (d) to observe spectroscopic properties of these individual conformations.

Much current interest exists in the conformational analysis of alkyl-substituted benzene substrates, and herein we focus specific attention on the first three members of this class of compound: methyl, ethyl, and *n*-propyl. Both time of flight mass spectra (TOFMS) and dispersed emission (DE) spectra are determined for the systems studied.³ Figure 1 presents the TOFMS of *m*-xylene for the 0⁰ region of the $S_1 \leftarrow S_0$ transition. The various features of this spectrum are attributed to transitions between internal rotor states of the two methyl groups. A double 1-dimensional rigid rotor model can be employed to fit these data.⁴⁻⁶ Parameters of this quantum mechanical model can be found for both the ground (S_0) and excited (S_1) states and are reported in Table 1 for *m*-xylene and the other compounds discussed herein.

The TOFMS and DE spectra of *o*-xylene are considerably different from those of *m*-xylene. The methyl groups of *o*-xylene act as independent hindered rigid rotors in the S_0 state,⁷ whereas both kinetic and potential

interactions between the two neighboring groups can be identified in the S_2 state. The relevant rigid rotor model parameters for S_0 and S_1 states are presented and compared with those for other systems in Table 1.


For n -alkylbenzenes, τ_1 ($C_{\text{ortho}}-C_{\text{ipso}}-C_{\alpha}-C_{\beta}$) is the first torsional angle which defines the orientation of the alkyl group relative to the aromatic ring (c.f. 3). Ethylbenzene (4) is the prototypical example, with a barrier to rotation² about τ_1 of $<5 \text{ kcal mol}^{-1}$. While some recent theoretical and experimental studies have concluded that the orthogonal conformation is preferred for 4, some controversy remains and a number of reports conflict with this assignment.⁸ The TOFMS and DE spectra of 1,3-diethylbenzene (5) reveal two origins, one each for the syn and anti orthogonal conformations, 5s and 5a respectively. Had planar ($\tau_1=0$) or angular conformations ($0<\tau_1<90^\circ$) been obtained, then more than two origins for 5 would have been observed. Importantly, the TOFMS experiment is sufficiently sensitive to molecular symmetry to distinguish between 5s and 5a.



n -Propylbenzene (6) serves as the prototypical example to define the second torsional angle τ_2 (c.f. 3). In their pioneering report, Hopkins, Powers and Smalley had observed two origins in the TOFMS for 6 but could not distinguish between the possible (anti, gauche, or eclipsed) conformations.⁹ Over the last few years, evidence has accumulated from detailed studies of 6

and related compounds that $\text{CH}^{\cdots}\pi$ attractive interactions can stabilize conformations when appropriate molecular geometries are possible.¹⁰ Most studies have assumed that anti and gauche conformations are predominant,¹¹ and this assumption has been found to be consistent with empirical force field (EFF) calculations;¹⁰ however, EFF calculations are not specifically parameterized for $\text{CH}^{\cdots}\pi$ interactions. Indeed, Hopkins, Powers and Smalley⁹ suggested that the eclipsed form could be the preferred conformation for *n*-propylbenzene, pending unambiguous experimental data.

Figure 2 shows the TOFMS of 3-*n*-propyltoluene (7). Three origin transitions, appearing as doublets, are observed. These correspond to the anti conformation 8 and the two gauche conformations, syn-gauche 9s and anti-gauche 9a. Had the eclipsed conformation 10 and the anti conformation 8 been predominant to the exclusion of the gauche conformations, only two origin transitions would have appeared. The doubling of the origins is due to two non-equilibrating methyl rotor states of 8, namely the 0a and 1e states. The doublets at ca. 25 cm^{-1} from each origin are due to propyl torsions, features also noted in the spectra of *n*-propylbenzene and 4-*n*-propyltoluene. The observation of only three origins also substantiates the conclusion derived from the data on the ethylbenzenes that $\tau = 90^\circ$.



These results show: a) supersonic molecular jet spectroscopy is capable of observing specific molecular conformations which have very low barriers to conformational interchange; b) aromatic ring methyl groups in the p-, and m-xylenes can be considered as independent, nearly free rotors in S_0 and S_1 ; c) the aromatic ring methyl groups in o-xylene are considerably more hindered in S_0 and S_1 , and for S_1 , cross kinetic and potential terms must be introduced; d) the barriers for rotation are greater in S_1 than S_0 ; e) the dynamic nature of the motion of the ring methyl groups is contrasted by a more static, locked-in character of the aromatic ethyl and n-propyl substituents; f) aromatic n-alkyl substituents have their first C-C torsion perpendicular to the plane of the aromatic ring; and g) aromatic n-propyl substituents exist in anti and gauche conformations, as observed in these TOFMS/DE experiments.

TABLE I

Summary of the Hamiltonian parameters used to fit the experimental data for the alkylbenzenes.
Values are given in cm^{-1} ^a

	<u>S₀</u>		<u>S₁</u>			
	<u>v₃</u>	<u>v₆</u>	<u>v₃</u>	<u>v₆</u>	<u>χ₆</u> ^b	<u>v'₃</u> ^c
toluene	0	10	0	25	-	-
p-xylene	0	10	0	25	-	-
m-xylene	0	25	81	-30	-	-
o-xylene	425	18	166	0	.72	-25
3-n-propyltoluene	0	23	75	0	-	-

^a 1 kcal mol⁻¹ = 350 cm⁻¹. ^b Kinetic cross term. ^c Potential cross term.

REFERENCES

- (1) (a) Ōsawa, E.; Musso, H. Top. Stereochem. 1982, 13, 117. (b) Burkert, U.; Allinger, N. L. "Molecular Mechanics," A.C.S. Monograph 177, American Chemical Society: Washington, D.C., 1982. (c) Clark, T. "A Handbook of Computational Chemistry", Wiley: New York, 1985. (d) Hehre, W. J.; Radom, L.; Schleyer, P. v. R.; Pople, J. A. "Ab Initio Molecular Orbital Theory", Wiley: New York, 1986.
- (2) (a) Seeman, J. I. Chem. Rev. 1983, 83, 1983. (b) Ōki, M. "Applications of Dynamic NMR Spectroscopy to Organic Chemistry", VCH Publishers: Deerfield Beach, FL, 1985. (c) Jackman, L. M.; Cotton, F. A. "Dynamic NMR Spectroscopy", Wiley: New York, 1975. (d) Chen, J.-S.; Shirts, R. B.; Lin, W.-C. J. Phys. Chem. 1986, 90, 4970. (e) Berg, U.; Liljefors, R.; Roussel, C.; Sandstrom, J. Acc. Chem. Res. 1985, 18, 80.
- (3) Bernstein, E. R.; Law K.; Schauer, M. J. Chem. Phys. 1984, 80, 207.
- (4) Okuyama, K.; Mikami, N.; Ito, M. J. Phys. Chem. 1985, 89, 5617.
- (5) (a) Bunker, P. "Molecular Symmetry and Spectroscopy", Academic Press: London, 1979. (b) Groner P.; Durig, J. R. J. Chem. Phys. 1977, 66, 1856.
- (6) Complete details will be reported in our full papers.
- (7) Our molecular orbital-molecular mechanics calculations⁶ and recent ab initio calculations (Gough, K. M.; Henry, B. R.; Wildman, T. A. J. Mol. Str. 1985, 124, 71) support these experimental findings.
- (8) (a) True, N. S.; Farag, M. S.; Bohn, R. K.; MacGregor, M. A.; Radhakrishnan, J. J. Phys. Chem. 1983, 87, 4622. (b) Kao, J. J. Am. Chem. Soc., in press.

- (9) Hopkins, J. B.; Powers, D. E.; Smalley, R. E. J. Chem. Phys. 1980, 73, 5039.
- (10) (a) Hirota, M.; Abe, K.; Suezawa, H.; Nishio, M. J. Mol. Str. 1985, 126, 455. (b) Hirota, M.; Sekiya, T.; Abe, K.; Tashiro, H.; Karatsu, M.; Nishio, M.; Ōsawa, E. Tetrahedron 1983, 39, 3091.
- (11) Radcliffe, M. D.; Mislow, K. J. Org. Chem. 1984, 49, 2058.


FIGURE CAPTIONS

FIGURE 1 Electronic absorption spectrum of the O_0^0 region of m-xylene, obtained with time-of-flight mass detection (TOFMS). The origin occurs at 36956.3 cm^{-1} . The two features at 3.7 and 7.0 cm^{-1} to lower energy of the origin are due to methyl torsions, and their positions are indicative of differences in the potential barrier to methyl rotation in S_0 and S_1 .


FIGURE 2 Electronic absorption spectrum (TOFMS) of the O_0^0 region of 3-n-propyltoluene. The spectrum contains three origins at 36982.8 , 37040.4 and 37060.1 cm^{-1} , corresponding to three different propyl group conformations. Accompanying each origin peak is a second peak to lower energy which is assigned as due to a methyl torsion. The doublet structures are indicative of differences in the potential barrier height to methyl rotation in S_1 and S_0 .

Wavenumber scale (cm⁻¹):

Wavenumber (cm ⁻¹)
169.0
164.4
75.2
70.2
32.6
29.0
16.9
0
-3.7



WAVENUMBERS (cm⁻¹)



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1

END

8-87

DTIC